## Oxidative Carbon–Carbon Bond Formation via Silyl Bis-enol Ethers: Controlled Cross-Coupling for the Synthesis of Quaternary Centers

Michael D. Clift, Carla N. Taylor, and Regan J. Thomson\*

Department of Chemistry, Northwestern University, 2145 Sheridan Road, Evanston, Illinois 60208

r-thomson@northwestern.edu

Received September 22, 2007

## $Me_{R}$ Si O Ce(NO<sub>3</sub>)<sub>6</sub>(NH<sub>4</sub>)<sub>2</sub> O Me Me NaHCO<sub>3</sub> A T3 examples

MeCN, 0 °C 10 min

ABSTRACT

Unsymmetrical silvl bis-enol ethers have been developed as effective substrates for synthesizing quaternary centers from tetralone derivatives through oxidative carbon-carbon bond formation. The derived products are shown to be highly versatile intermediates that may be used to generate diverse structures such as cyclopentenones, 2*H*-pyrroles, and spirocyclic pyrrolidines.

The oxidative coupling of enolates, first reported in 1935,<sup>1</sup> provides direct access to highly versatile 1,4-dicarbonyl intermediates.<sup>2</sup> A major limitation of this chemistry lies with the lack of methods that allow for selective and controlled heterocoupling of two different enolates. For many cases a mixture of both homo- and heterocoupled products are obtained (Figure 1, eq 1). Recently, Baran and co-workers demonstrated that a number of selective cross-couplings can be achieved when the oxidation potential between the substrates is significantly different, such as between an amide or ester enolate and a ketone enolate.<sup>3</sup> In the case of silyl enol ethers, cross-coupling can typically only be achieved by using an excess of one enol silane.<sup>4</sup> Narasaka's solution

involved the use of enamines, which are preferentially oxidized over enol silanes.<sup>5</sup> MacMillan's recently disclosed catalytic enantioselective oxidative coupling uses enamine catalysis to gain impressive selectivity.<sup>6</sup> To date, no general method for controlled oxidative ketone–ketone cross-coupling has been reported.

up to 81% yield



**Figure 1.** General approach toward controlled oxidative carbon – carbon bond formation.

LETTERS 2007 Vol. 9, No. 22 4667–4669

ORGANIC

<sup>(1)</sup> Ivanoff, D.; Spasoff, A. Bull. Soc. Chim. Fr. 1935, 2, 76-78.

<sup>(2)</sup> For additional early contributions, see: (a) H. Schäfer, A. A. Angew. Chem., Int. Ed. Engl. 1968, 7, 474-475. (b) Brettle, R.; Parkin, J. G.; Seddon, D. J. Chem. Soc. (C) 1970, 1317-1320. (c) Rathke, M. W.; Lindert, A. J. Am. Chem. Soc. 1971, 93, 4605-4606. (d) Dessau, R. M.; Heiba, E. I. J. Org. Chem. 1974, 39, 3457-3459. (e) Ito, Y.; Konoike, T.; Saegusa, T. J. Am. Chem. Soc. 1975, 97, 2912-2914. (f) Ito, Y.; Konoike, T.; Harada, T.; Saegusa, T. J. Am. Chem. Soc. 1977, 99, 1487-1493. (g) Kobayashi, Y.; Taguchi, T.; Tokuno, E. Tetrahedron Lett. 1977, 3741-3742. (h) Frazier, R. H.; Harlow, R. L. J. Org. Chem. 1980, 45, 5408-5411.

<sup>(3)</sup> Baran, P. S.; DeMartino, M. P. Angew. Chem., Int. Ed. 2006, 45, 7083-7086.

As part of our program directed at synthesizing bioactive polycyclic molecules, we became interested in using silyl bis-enol ethers to generate quaternary centers through intramolecular oxidative bond formation. We were encouraged by a report in 1998 by Schmittel and co-workers who revealed that symmetrical silyl bis-enol ethers will undergo oxidative bond formation to generate 1,4-diketones.<sup>7</sup> Despite the potential utility of this process to generate unsymmetrical 1,4-diketones, it has never been fully developed.<sup>8</sup> We now describe the development of an efficient method for selective ketone—ketone oxidative cross-coupling for the synthesis of quaternary centers, which is based on the use of unsymmetrical silyl bis-enol ethers. The basic concept is outlined in eq 2 of Figure 1.

Our initial studies were directed at preparing unsymmetrical silvl bis-enol ethers derived from 2-methyltetralone (6) and a variety of methyl ketones. Rathke and Weipert have reported a general method for the synthesis of unsymmetrical silvl bis-enol ethers based on generating chlorosilanes (5) in situ from enol silylamines (4) and acetyl chloride.<sup>9</sup> Subsequent addition of NaI, triethylamine and the second ketone afforded the desired product in good yield. These conditions, however, led to mixtures of regioisomeric products when ketones with two sets of enolizable positions are used. Therefore, we developed a modified procedure based on Corey's conditions for selective kinetic generation of enol silanes from methyl ketones.<sup>10</sup> Thus, addition of the methyl ketone 3 to a mixture of both LDA and chloro-N.Ndiethylamino-dimethylsilane at -78 °C affords excellent regioselectivity for the desired enol ethers (i.e., 4). Evaporation of the solvent, trituration with pentanes to remove LiCl, and subsequent distillation provides good to excellent yields for a range of enol silylamines **4** (Scheme 1).<sup>11</sup>



The addition of acetyl chloride to a solution of 4 (R = Me) resulted in rapid formation (<10 min) of the corre-

- (6) (a) Jang, H. Y.; Hong, J. B.; MacMillan, D. W. C. J. Am. Chem. Soc. **2007**, *129*, 7004–7005. (b) Beeson, T. D.; Mastracchio, A.; Hong, J. B.; Ashton, K.; MacMillan, D. W. C. Science **2007**, *316*, 582–585.
- (7) Schmittel, M.; Burghart, A.; Malisch, W.; Reising, J.; Sollner, R. J.
   Org. Chem. 1998, 63, 396–400.
- (8) Schmittel, M.; Haeuseler, A. J. Organomet. Chem. 2002, 661, 169-179.
- (9) Rathke, M. W.; Weipert, P. D. Synth. Commun. 1991, 21, 1337-1351.

sponding chlorosilane **5** (R = Me) and diethylacetamide as indicated by <sup>1</sup>H NMR spectroscopy. Enolization of 2-methyltetralone (**6**) with LDA, followed the addition of the chlorosilane **5** (R = Me) generated from **4** (R = Me), resulted in clean formation of the desired unsymmetrical silyl bisenol ether **7a** (Scheme 2). While purification of the silyl bis-



enol ether by flash chromatography may be achieved, we found this to be unnecessary due to the effectiveness of the synthesis. Only 1.1 equiv of enol silane 4 (R = Me) was used in relation to 2-methyltetralone (6). This stoichiometry will have important consequences for the use of this chemistry in carrying out complex fragment couplings in the context of total synthesis.

Our initial exploration of potential oxidants utilized enol silane **7a**.  $Cu(OTf)_2^{2g}$  has been reported to be a good oxidant for TMS-enol ether coupling but in our hands gave <10% of the desired adduct **8a**. Cerium(IV) ammonium nitrate (CAN)/NaHCO<sub>3</sub><sup>4a</sup> was quickly found to be the oxidant of choice, providing diketone **8a** in an overall yield of 71% from 2-methyltetralone (Scheme 2).

As a control experiment, a 1:1 mixture of the TMS-enol silanes derived from 2-methyltetralone (i.e., 9) and acetone (i.e., 10) was exposed to the reaction conditions. The complex reaction mixture that resulted contained little or none of the desired adduct **8a** (Scheme 3).



The scope of the methyl ketone component was explored next and was general for a variety of substrates (Table 1). Aliphatic substrates are well tolerated, providing moderate to good yields of the 1,4-diketone products (entries 1–7). As can be seen, increasing steric hindrance led to a decrease in overall reaction efficiency. While  $\beta$ -substitution on the methyl ketone fragment was well tolerated, providing **8e** in 61% overall yield (Table 1, entry 5),  $\alpha$ -substitution resulted in lower yields. Isopropyl and tertiary butyl substituents afforded the corresponding products in 50% and 41% yield, respectively (Table 1, entries 6 and 7). Aryl substituents

<sup>(4) (</sup>a) Baciocchi, E.; Casu, A.; Ruzziconi, R. *Tetrahedron Lett.* **1989**, *30*, 3707–3710. (b) Fujii, T.; Hirao, T.; Ohshiro, Y. *Tetrahedron Lett.* **1992**, *33*, 5823–5826.

<sup>(5)</sup> Narasaka, K.; Okauchi, T.; Tanaka, K.; Murakami, M. Chem. Lett. 1992, 2099–2102.

<sup>(10)</sup> Corey, E. J.; Gross, A. W. *Tetrahedron Lett.* **1984**, *25*, 495–498. (11) For R = Me and R = t-Bu, enol silylamines prepared according to ref. 9.

 Table 1.
 Substrate Scope for 2-Methyltetralone Derived

 Compounds<sup>a</sup>
 Provide Scope for 2-Methyltetralone Derived



<sup>*a*</sup> Reagents and conditions: (i) 1.5 mmol **6**, LDA (1.1 equiv), -78 °C to 0 °C; then **4** (1.1 equiv)/MeCOCl (1.0 equiv); (ii) CAN (2.0 equiv), NaHCO<sub>3</sub> (4.0 equiv), MeCN, 0 °C. <sup>*b*</sup> Isolated yield.

provided good yields, with both electron-rich and electronpoor systems behaving similarly (entries 8–11). Crossconjugated enol silanes derived from  $\alpha,\beta$ -unsaturated ketones (i.e., **71** and **7m**) were also suitable compounds (Table 1, entries 12 and 13).

Having established the methodology as useful for 2-methyltetralone substrates, we investigated the use of other cyclic aryl ketones (Figure 2). 2-Ethyltetralone was an effective



Figure 2. Variation of substituted enolate component.

reaction partner with both acetone and acetophenone derived systems (11 and 12, respectively). 2-Methylindanone provided only 30% of dione 13 (R = Me), whereas dione 14 (R = Ph) was obtained in 76% yield. Such dramatic differences in yield were also observed when 2-methylben-zosuberone was used as a substrate. To date, 2-methylcy-clohexanone has not proven to be a synthetically useful reaction partner (<10% yield with acetophenone). We are currently engaged in solving this issue in order to fully expand the scope of this methodology.

We were also interested in utilizing the versatility of the 1,4-diketone products by using straightforward reactions to create a diverse array of structures (Scheme 4).



Cyclopentenone 17<sup>12</sup> was accessed from 8a by NaOHmediated aldol condensation (73% yield). Thus, the threestep procedure from 2-methyltetralone (6) represents an efficient strategy for cyclopentenone annulation.<sup>13</sup> Condensation of 8a with ammonium acetate in acetic acid at 100 °C afforded the corresponding 2H-pyrrole 19. We had initially hoped to isolate the 3H-pyrrole, but under the acidic reaction conditions a ring contraction of protonated 3H-pyrrole 18 ensued to generate the spirocycle 19. This interesting transformation presumably proceeds through a selective [1,5]sigmatropic alkyl shift.<sup>14</sup> 2H-Pyrrole 19 was selectively reduced (15:1 dr) to pyrrolidine 20 upon treatment with Pd/C and H<sub>2</sub>. The X-ray crystal structure of the HCl salt of 20 confirmed that hydrogenation occurred anti to the large aromatic ring.<sup>15</sup> We are currently exploring the generality of this ring contraction/hydrogenation sequence in order to prepare stereochemically rich spirocyclic alkaloids

In summary, we have developed unsymmetrical silyl bisenol ethers as effective substrates for crossed-ketone oxidative coupling reactions of tetralone derivatives. Using these compounds provides a general method for the preparation of quaternary centers in a convergent manner. Current and future research is directed at expanding the scope of silyl bis-enol ether-based transformations and applying this chemistry to the synthesis of bioactive polycyclic natural products.

Acknowledgment. Support is provided by Northwestern University (NU) and the NU Analytical Services Laboratory (NSF Grants DMR0114235 and CHE9871268). We gratefully acknowledge Charlotte Stern (Northwestern University) for X-ray crystallography and Prof. Karl Scheidt (Northwestern University) for helpful discussions.

**Supporting Information Available:** Experimental procedures, spectral data, and crystallographic data (PDF, CIF). This material is available free of charge via the Internet at http://pubs.acs.org.

## OL702306Z

(13) For a related strategy, see: Miyashita, M.; Yanami, T.; Yoshikoshi,A. J. Am. Chem. Soc. 1976, 98, 4679–4681.

<sup>(12)</sup> Snyder, L.; Meyers, A. I. J. Org. Chem. 1993, 58, 7507-7515.

<sup>(14)</sup> For related examples, see: (a) Chiu, P.-K.; Sammes, M. P. *Tetrahedron Lett.* **1987**, *28*, 2775–2778. (b) Lui, K. H.; Sammes, M. P. *J. Chem. Soc., Perkins Trans. 1* **1990**, 457–468. (c) Shimkin, A. A.; Shirinian, V. Z.; Nikalin, D. M.; Krayushkin, M. M.; Pivina, T. S.; Troitsky, N. A.; Vorontsova, L. G.; Starikova, Z. A. *Eur. J. Org. Chem.* **2006**, 2087–2092. (15) See Supporting Information for details.