This article was downloaded by: [University of Newcastle, Australia] On: 08 January 2015, At: 08:45 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Synthesis and Characterization of New Schiff Bases 2-(2-(2-(Aryl)Methyleneamino)Phenylthio) Ethylthio)-N-((aryl)Methylene)Benzeneamine

Ali Kakanejadifard^a, Toraj Hayati^a & Azam Rezayat^a ^a Department of Chemistry, Faculty of Science, Lorestan University, Khorramabad, Iran Published online: 02 Jul 2010.

To cite this article: Ali Kakanejadifard , Toraj Hayati & Azam Rezayat (2010) Synthesis and Characterization of New Schiff Bases 2-(2-(2-(Aryl)Methyleneamino)Phenylthio) Ethylthio)-N-((aryl)Methylene)Benzeneamine, Phosphorus, Sulfur, and Silicon and the Related Elements, 185:7, 1511-1515, DOI: <u>10.1080/10426500903107930</u>

To link to this article: http://dx.doi.org/10.1080/10426500903107930

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Phosphorus, Sulfur, and Silicon, 185:1511–1515, 2010 Copyright © Taylor & Francis Group, LLC ISSN: 1042-6507 print / 1563-5325 online DOI: 10.1080/10426500903107930

SYNTHESIS AND CHARACTERIZATION OF NEW SCHIFF BASES 2-(2-(2-(ARYL)METHYLENEAMINO)PHENYLTHIO) ETHYLTHIO)-N-((ARYL)METHYLENE)BENZENEAMINE

Ali Kakanejadifard, Toraj Hayati, and Azam Rezayat

Department of Chemistry, Faculty of Science, Lorestan University, Khorramabad, Iran

A series of Schiff bases containing four to six coordination sites $N_2S_2 X_2(X = O,N)$ 2-(2-(*aryl*)methyleneamino)phenylthio)ethylthio)-N-((*aryl*)methylene)benzeneamine (2*c*–*f*) were prepared from the reaction of 1,2-di(2-aminophenylthio)ethane (1) with aromatic aldehydes. All compounds were characterized by means IR, mass, ¹H and ¹³C NMR spectroscopy, and elemental analysis, and in the case of 2b with a single crystal X-ray diffraction. The X-ray crystal structure of 2b showed that the resonance occurs between aromatic rings, through the C=N bonds of the molecule.

Supplemental materials are available for this article. Go to the publisher's online edition of Phosphorus, Sulfur, and Silicon and the Related Elements to view the free supplemental file.

Keywords 2-(2-(2-(Aryl)methyleneamino)phenylthio)ethylthio)-N-((aryl)methylene)benzeneamine; 1,2-di(2-aminophenylthio)ethane; Schiff base; X-ray

INTRODUCTION

The coordination chemistry of transition metal ions with the N_2S_2 donor sets has received considerable interest in recent years.^{1–5} Quadridentate ligands with the donor sets N_2S_2 are very important for nuclear medicine, because their technetium and rhenium complexes are sufficiently stable in vivo to function as effective radioimaging and therapeutic agents.^{6,7}

The N, S donor ligand and Pd(II), Pt(II) complexes, used to prepare antiviral, antifungal, antimicrobial, antitumor^{8,9} products, and their mechanism of action most probably involves the inhibition of ribonucleotide reductase, converting ribonucleotides to deoxyribonucleotides.^{10,11} Recently, new N₂S₂ Schiff bases **2a–b** were prepared from the reaction of 1,2-di(2-amino phenylthio)ethane (**1**) with aromatic aldehydes.^{12–14} These Schiff bases are suitable for preparation of coordination polymers and oxidative cleavage of the ligand when it was reacted with transition metal ions, rather than forming a simple complex.^{15–17} Considering these properties, we have prepared some new Schiff bases, N₄S₂, N₂O₄, N₂S₄,

Received 9 March 2009; accepted 6 June 2009.

We thank Lorestan University for financial support of this work.

Address correspondence to Ali Kakanejadifard, Department of Chemistry, Lorestan University, Khorramabad, Iran. E-mail: kakanejadi.a@lu.ac.ir

and $N_2O_2S_2$, from the reaction of 1,2-di(2-aminophenylthio)ethane (1) with aromatic aldehydes. The Schiff bases 2 contain four to six coordination sites; they are therefore suitable for the preparation of catalysts or coordination polymers.

RESULTS AND DISCUSSION

The condensation reaction of 1,2-di(2-aminophenylthio)ethane (1) with various substituted aromatic aldehydes in absolute ethanol gave the acyclic Schiff base products (Scheme 1). All prepared compounds **2c–f** are air-stable solids and soluble in acetone, CDCl₃, and THF. The structure of the prepared compounds **2c–f** was confirmed by means of ¹H NMR, ¹³C NMR, IR, mass spectroscopy, and elemental analysis. The compound **2b** was further characterized by single crystal X-ray crystallography (Figure 1).



Scheme 1

The infrared spectra of all products, recorded in the region of 4000–500 cm⁻¹, similar to the total absence of NH₂ in the IR spectra of **1**, together with the appearance of new absorption (C=N) in the range of 1605–1627 cm⁻¹, clearly indicated that a new Schiff base had formed in each case. A band observed in the spectra of **2c–f** in the region of 2835–2923 cm⁻¹ can be assigned to the CH=N group. The IR spectra of **2c–f** show bands at 740–778 cm⁻¹ assigned to the C–S stretching mode. The ¹H NMR spectrum of **2c–f** shows a single ¹H imine resonance at ca. 8.18–8.71 ppm, demonstrating the equivalence of the two imine environments. The absorption signals of aromatic protons appeared in the region of 6.54–7.81 ppm, and the CH₂ group of all Schiff bases appeared in 3.19–3.27 ppm. The ¹³C NMR spectra include imine and methylene carbons, which appeared at 152.21–154.58 ppm and 31.51–31.92 ppm, respectively. The molecular structure of **2b**, which is shown in Figure 1, possesses a center of symmetry and exhibits a helical structure.¹⁸

The bond lengths, angle, and torsion of **2b** show that a resonance occurs between aromatic rings, through the C=N bonds of the molecule. The thiophene ring is a rather planar relative to the aminothiphenole ring, and the imine groups have the *anti* conformation in agreement with the NMR spectra. However, the C(1)–C(1A) bond has a single bond characteristic (Table I).

EXPERIMENTAL

All chemical reagents were obtained from Merck and Fluka and used without further purification. The diamine 1,2-di(2-aminophenylthio)ethane (1) was prepared by the method in the literature.⁸ Melting points were measured with Electrothermal 9200 apparatus and are uncorrected. Elemental analyses were performed on an Elementar Vario EL III elemental analyzer. The FT-IR spectra (potassium bromide) were recorded with a Shimadzu 8400 S spectrometer. Mass spectra were recorded on a Shimadzu QD 5050. ¹H and ¹³C NMR spectra were measured with a Bruker DRX-500 Avance Spectrometer in CDCl₃ as solvent.



Figure 1 ORTEP drawing of 2-(2-(-2-((thiophen-2-yl)methyleneamino)phenylthio) ethylthio)-N-((thiophen-2-yl) methylene)benzenamine (**2b**).

General Procedure for Preparation of Schiff Base from the Reaction of 2-(2-(2-Amino phenylthio)benzeneamine with Aldehyde

To a stirring solution of 1,2-di(2-aminophenylthio)ethane (1) (2.76 g, 10 mmol) in absolute ethanol (25 mL), a solution of aldehyde (20 mmol) was added dropwise for 10 min at room temperature. The mixture was stirred and heated to reflux for 5 h. Then the solvent was evaporated under reduced pressure, and the product was crystallized from acetonitrile.

2-(2-(Furan-2-yl)methyleneamino)phenylthio)ethylthio)-N-((furan-2-yl)methylene) benzeneamine (2c). Yield: 2.12 g (49%), mp 122–123°C; Color:

Bond lengths	Å	Bond angles	(°)	Torsion angles	(°)	
C(1)-C(1)#1	1.524	C(1)#1-C(1)-S(1)	107.55	C(2)-S(1)-C(1)-C(1)#1	-166.07	
S(1)-C(1)	1.811	C(2)-S(1)-C(1)	102.36	C(1)-S(1)-C(2)-C(3)	-4.13	
S(1)-C(2)	1.763	S(1)-C(1)-H(1A)	110.20	C(1)-S(1)-C(2)-C(7)	174.52	
C(2) - C(7)	1.412	C(7)-C(2)-S(1)	115.42	C(8)-N(1)-C(7)-C(6)	53.92	
N(1)–C(7)	1.411	C(3)-C(2)-S(1)	125.63	S(1)-C(2)-C(7)-N(1)	2.73	
N(1)–C(8)	1.282	N(1)-C(7)-C(2)	117.03	C(8)-N(1)-C(7)-C(2)	-130.73	
C(8)–C(9)	1.439	C(8)–N(1)–C(7)	118.94	C(7)-N(1)-C(8)-C(9)	-177.31	
S(2) - C(9)	1.726	N(1)-C(8)-C(9)	121.68	N(1)-C(8)-C(9)-C(10)	-174.31	
C(9) - C(10)	1.3757	N(1)-C(8)-H(8A)	119.20	N(1)-C(8)-C(9)-S(2)	8.43	
C(8)–H(8A)	0.9500	C(8)-C(9)-S(2)	121.50	C(8)-C(9)-C(10)-C(11)	-177.69	
C(1)–H(1A)	0.9900	C(12)-S(2)-C(9)	91.53	S(2)-C(9)-C(10)-C(11)	-0.20	

Table I Selected bond lengths (Å), bond angles (°), and torsion angles (°) for 2b

yellow; FT-IR: 3124–3095 (Ar–H), 2923, 2880 (CH=N), 1623 (C=N), 1571–1436 (C=C), 1022 (C–O), 783 (C–S), 744 (C–S) cm⁻¹; ¹H NMR: $\delta = 8.18$ (s, 2H, CH=N), 7.62 (d, 2H, Ar–H, J = 3.7 Hz), 7.29 (d, 2H, Ar–H, J = 3.4 Hz), 7.20–7.18 (d, 2H, Ph–H, J = 5.2 Hz), 7.15–7.12 (t, 2H, Ar–H), 7.00–6.95(m, 4H, Ph–H), 6.55–6.54 (d, 2H, Ph–H, J = 4.7 Hz), 3.19 (s, 4H, CH₂); ¹³C NMR: δ 152.21, 150.73, 145.88, 130.59, 128.11, 126.72, 126.38, 118.46, 118.26, 118.14, 116.20, 112.24, 31.51. The EI-MS: *m/z* 432(M)⁺, 429, 355, 271; Anal. Calcd. For C₂₄H₂₀N₂O₂S₂: C, 66.64; H, 4.66; N, 6.48; S, 14.83. Found: C, 66.47; H, 4.53; N, 6.34; S, 14.03.

2-(2-(1H-Pyrrol-2-yl)methylene)-2-(2-(1H-pyrrol-2-yl)methyleneamino) phenylthio) ethylthio)benzeneamine (2d). Yield: 2.2 g (51%), FT-IR: 3329 (N-H), 3132–3095 (Ar–H), 2912, 2877 (CH=N), 1650 (C=N), 1606–1477 (C=C), 748 (C–S) cm⁻¹; ¹H NMR: δ 8.17 (s, 2H, CH=N), 7.44 (d, 2H, Ar–H, J = 4.1 Hz), 7.38 (d, 2H, Ar–H, J = 3.3 Hz), 7.23(d, 2H, Ph–H, J = 5.4 Hz), 7.17–7.14 (t, 2H, Ar–H), 7.03 (d, 2H, Ph–H, J = 4.9 Hz), 6.76–6.71 (m, 4H, Ph–H), 6.30 (bs, 2H, NH), 3.18 (s, 4H, CH₂); ¹³C NMR: δ 150.73, 150.18, 136.17, 130.48, 129.91, 126.83, 126.30, 118.49, 117.02, 116.26, 114.88, 31.50. The EI-MS: *m/z* 430(M)⁺, 429(M-1)⁺, 271; Anal. Calcd. For C₂₄H₂₂N₄S₂: C, 66.94; H, 5.15; N, 13.01; S, 14.89. Found: C, 66.62; H, 4.91; N, 12.89; S, 15.02.

2-(2-(Pyridin-2-yl)methyleneamino)phenylthio)ethylthio)-N-((pyridin-2-yl)methyl ene)benzeneamine (2e). Yield: 2.22 g (51%), mp 127–128°C, Color: red; FT-IR: 3153–3062 (Ar–H), 2975, 2877 (N=CH), 1627 (C=N), 1588–1471 (C=C), 777 (C–S), 740 (C–S) cm⁻¹; ¹H NMR: δ 8.53 (s, 2H, CH=N), 8.70 (d, 2H, Py–H, J = 5.3 Hz), 8.28 (d, 2H, Py–H, J = 4.9 Hz), 7.81–7.77 (t, 2H, Py–H, J = 2.8 Hz), 7.38–7.35 (t, 2H, Py–H, J = 3.05 Hz), 7.31–7.19(m, 8H, Ar–H), 3.23 (s, 4H, CH₂); ¹³C NMR: δ 160.73, 154.58, 149.87, 149.62, 136.69, 130.97, 128.25, 127.06, 126.96, 125.27, 122.00, 118.21, 31.53; The EI-MS: m/z 454(M)⁺, 452(M-2)⁺ 431, 429, 355, 271. Anal. Calcd. For C₂₆H₂₂N₄S₂: C, 68.69; H, 4.88; N, 12.32; S, 14.11. Found: C, 68.84; H, 4.71; N, 12.14; S, 14.38.

2-(2-(Pyridin-4-yl)methyleneamino)phenylthio)ethylthio)-N-((pyridin-4-yl)methyl ene)benzeneamine (2f). Yield: 2.80 g (60%), mp 162–162.5°C, Color: yellow; FT-IR: 3058–3018 (Ar–H), 2993, 2887 (CH=N), 1623 (C=N), 1596–1460 (C=C), 817 (C–S), 746 (C–S) cm⁻¹; ¹H NMR: δ 8.37 (s, 2H, N=CH), 8.76 (d, 2H, Py–H, J = 5.6 Hz), 7.77, 7.75 (d, 2H, Py–H, J = 5.6 Hz), 7.32–7.01 (m, 8H, Ar–H), 3.21 (s, 4H, CH₂). ¹³C NMR: δ 158.07, 150.65, 149.74, 142.48, 130.93, 128.12, 127.36, 126.90, 122.39, 118.06, 31.25. The EI-MS: m/z 454(M)⁺, 452(M-2)⁺ 431, 429, 355, 271. *Anal.* Calcd. For C₂₆H₂₂N₄S₂: C, 68.69; H, 4.88; N, 12.32; S, 14.11. Found: C, 68.41; H, 4.68; N, 12.11; S, 14.29.

X-Ray Structure Analysis of 2-(2-(-2-((Thiophen-2-yl)methyleneamino) phenylthio)ethylthio)-N-((thiophen-2-yl)ethylene)benzenamine (2b)

The clear colorless crystal with dimension $0.30 \times 0.20 \times 0.15$ mm³ crystallized from MeCN/MeOH was used for data collection on a Bruker SMART 1000 CCD diffractommeter with graphite monochromated MoK α ($\lambda = 0.71073$ Å). C₂₄H₂₀N₂S₄, FW = 464.66. Monoclinic crystals in a P 21/c Space group, a = 11.179(5) Å, b = 7.730(4) Å, c = 12.608(6) Å, $\alpha = 90^{\circ}$, $\beta = 91.899(12)^{\circ}$, $\gamma = 90^{\circ}$, V = 1088.9(9) Å⁻³. Z = 2, D_{calc} = 1.417 g/cm³. $\mu = 0.451$ mm⁻¹, F (000) = 484, T = 100 (2) K. Of the 12,880 total reflections, 2899 were unique ([R(int) = 0.0294]). The structure was solved by direct methods (SHELEXTL)¹⁹ and refined by full-matrix least-squares on F² (isotropic refinement of the molecule and location of remaining non-hydrogen atoms from a difference Fourier map and subsequent anisotropy refinement on all atoms; H atoms were found after high-angle refinement in a difference Fourier map and their positions included in the final stages of refinement), factors of = 0.0288, wR2 = 0.0751 for 2569 reflections with I < 2σ (I). No significant features, only ripples from 0.337 to -0.257 e Å⁻³, were observed in the final difference map. The nonhydrogen atoms were refined anisotropically.

Complete bond lengths and angles, coordinates, and displacement parameters have been deposited at Cambridge Crystallography Data Center. Supplementary data are available from the CCDC, 12 Union Road, Cambridge CB2 IEZ, UK on request, quoting the deposition numbers 730627 for 2-(2-(-2-((thiophen-2-yl)methyleneamino)phenylthio) ethylthio)-N-((thiophen-2-yl) methylene)benzenamine (K5).

REFERENCES

- 1. P. Chattopadhyay and C. Sinha, Indian J. Chem. Sect. A, 76, 34 (1995).
- 2. P. Chakraborty, S. Chandra, and A. Chakravorty, Inorg. Chim. Acta, 229, 477 (1995).
- 3. J. R. Dilworth and S. Parrot, J. Chem. Soc. Rev, 27, 43 (1998).
- 4. H. Zhang, M. Dai, C. Qi, B. Li, and X. Guo, Appl. Rad. Isotopes, 60, 643 (2004).
- 5. R. K. Home and J. A. Katzenellenbogen, Nucl. Med. Biol., 24, 485 (1997).
- M. Glaser, M. J. Howard, K. Howland, A. K. Powell, M. T. Rae, S. Wocadlo, R. A. Williamson, and P. J. Blower, *J. Chem. Soc.*, *Dalton Trans.*, 3087 (1998).
- 7. P. K. Dhara, B. Dasa, J. M. Lob, and P. Chattopadhyay, Appl. Rad. Isotopes, 62, 729 (2005).
- 8. A. Garoufis, S. K. Hadjikakou, and N. Hadjiliadis, Coord. Chem. Rev., 253, 1384 (2009).
- 9. C. Lodeiro and F. Pina, Coord. Chem. Rev., 253, 1353 (2009).
- E. Bermejo, R. Carballa, A. Castineiras, R. Dominguez, A. E. Liberta, C. Maichelle-Mössmer, M. M. Salberg, and D. X.West, *Eur. J. Inorg. Chem.*, 965 (1999).
- A. G. Quiroga, J. M. Perez, I. Lopez-Solera, J. R. Masaguer, A. Luque, P. Roman, A.Edwaeds, C. Alonso, and C. Navarro-Ranninger, *J. Med. Chem.*, 41, 1399 (1998).
- 12. S. K. Hadjikakou and N. Hadjiliadis, Coord. Chem. Rev., 253, 235 (2009).
- 13. W. X. Zhang, C. Ma, and S. Si, *Trans. Met. Chem.*, **26**, 38 (2001).
- 14. S. Chandra and R. Kumar, Trans. Metal. Chem., 29, 269 (2004).
- G. Rajsekhar, C. P. Rao, P. K. Saarenketo, E. Kolehmainen, and K. Rissanen, *Inorg. Chem. Commun.*, 5, 649 (2002).
- G. Rajsekhara, C. P. Rao, P. Saarenketo, N. Nattinenb, and K. Rissanen, New J. Chem., 28, 75 (2004).
- 17. M. Morshedi, M. Amirnasr, S. Triki, and A. D. Khalaji, *Inorg. Chim. Acta*, 362, 1637 (2009).
- 18. E. Borisova, M. D. Reshetova, and Y. A. Ustrynyuk, Chem. Rev., 107, 46 (2007).
- G. M. Sheldrick, SHELXTL v. 5.10 "Structure Determination Software Suite," Bruker AXS, Madison, Wisconsin, USA (1998).