

## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

### A Simple One-Pot Synthesis of Methyltribromosilane

Michael W. Pitcher<sup>a</sup> & Patricia A. Bianconi<sup>b</sup>

<sup>a</sup> Department of Chemistry , Middle Eastern Technical University , Ankara, Turkey

<sup>b</sup> Department of Chemistry , The University of Massachusetts at Amherst. , Amherst, MA, USA  
Published online: 15 Jun 2007.

To cite this article: Michael W. Pitcher & Patricia A. Bianconi (2007) A Simple One-Pot Synthesis of Methyltribromosilane, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 182:8, 1819-1822, DOI: [10.1080/10426500701323382](https://doi.org/10.1080/10426500701323382)

To link to this article: <http://dx.doi.org/10.1080/10426500701323382>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## A Simple One-Pot Synthesis of Methyltribromosilane

**Michael W. Pitcher**

Department of Chemistry, Middle Eastern Technical University,  
Ankara, Turkey

**Patricia A. Bianconi**

Department of Chemistry, The University of Massachusetts at Amherst,  
Amherst, MA, USA

*A simple one-pot synthesis of methyltribromosilane has been developed. The straightforward reaction of a slight excess of silicon tetrabromide with one equivalent of powdered methylolithium in refluxing pentane produces methyltribromosilane in high yields.*

**Keywords** Halogenated organosilanes; methyltribromosilane; Rochow synthesis; silicon carbide

## INTRODUCTION


Halogenated organosilanes are today used in an enormous variety of industries and in a diverse array of applications. Methyltribromosilane and other halogenated organosilanes are of interest as precursors for the Chemical Vapor Deposition (CVD) of  $\beta$ -SiC films,<sup>1,2</sup> and in the synthesis of novel polysilyne network polymers.<sup>3–5</sup> Methyltribromosilane and other halogenated organosilanes are usually prepared by the Rochow synthesis,<sup>6</sup> or by a method involving dropwise addition of a Grignard reagent to an ether solution containing silicon tetrabromide.<sup>7</sup> The former method requires high temperatures and the use of hydrogen bromide gas, and the later synthesis is complicated by difficult separation of the products. In this article, we report a simple, alternative route to the production of methyltribromosilane.

Received December 8, 2006; accepted February 6, 2007.

Address correspondence to Michael Pitcher, Department of Chemistry, Middle Eastern Technical University, 06531 Ankara, Turkey. E-mail: pitcher@metu.edu.tr

## RESULTS AND DISCUSSION

The simple addition of powdered methyllithium to a refluxing mixture of silicon tetrabromide and pentane produces white crystalline bromosilane products which are easily isolated. NMR studies of the crystalline material confirm the major product to be methyltribromosilane (**1**—Scheme 1), along with smaller quantities of dimethyldibromosilane (**2**), and trimethylbromosilane (**3**)



**SCHEME 1**

((CH<sub>3</sub>)<sub>2</sub>SiBr<sub>2</sub>) (**2**) and trimethylbromosilane ((CH<sub>3</sub>)<sub>3</sub>SiBr) (**3**).<sup>8</sup> <sup>1</sup>H NMR (200 MHz, d<sub>6</sub>-benzene) shows the methyl resonance of **1** at  $\delta$  = 1.00, that of **2** at  $\delta$  = 0.67, and that of **3** at  $\delta$  = .31, in a ratio of 4.1:1.9:0.5 by integration. <sup>13</sup>C NMR (75.5 MHz) shows a methyl resonance at  $\delta$  = 126.6 (triplet, J<sub>C-H</sub> = 96.4 Hz) corresponding to methyltribromosilane (**1**). The other resonances at  $\delta$  = 124.2 (triplet) and  $\delta$  = 121.4 (triplet), corresponding to the methyl groups of the di- and tri-methyl-substituted products are not seen. Elemental analyses: Found (C 5.03%, H 0.91%); Calculated for (CH<sub>3</sub>SiBr<sub>3</sub>) (C 4.24%, H 1.06%). Based on the <sup>1</sup>H NMR, the reaction product therefore consists of 79% CH<sub>3</sub>SiBr<sub>3</sub>, 18% (CH<sub>3</sub>)<sub>2</sub>SiBr<sub>2</sub> and 3% (CH<sub>3</sub>)<sub>3</sub>SiBr. The purity is acceptable, for the intended use as a reagent in the synthesis of poly(methylsilylene).<sup>3</sup> Fractional distillation of the mixture, would enable isolation of pure methyltribromosilane.<sup>6,7</sup> Although this process has not been used to produce any other halogenated organosilanes, it is presumed that this facile synthesis method could be used as a route for the preparation of these compounds, too.

## EXPERIMENTAL

Silicon tetrabromide (99+%) was purchased from Aldrich and used as received. Methylolithium in diethylether (1.4 M) was purchased from Aldrich; the ether was removed by vacuum and the methylolithium used in powder form. Anhydrous pentane (99+%) was purchased from Aldrich and dried over sodium and benzophenone. All work was performed under an argon atmosphere, using either a glove box or schlenk line techniques.

Using a two-necked 300 ml round bottom flask equipped with a stirrer, reflux condenser and powder addition funnel, addition of powdered methylolithium (3 g, 0.1367 mol) to a 5% excess of silicon tetrabromide (50 g, 0.1438 mol) in 250 ml of pentane, was done at a rate sufficient to maintain reflux (about a one-hour addition time, Scheme 1). The reaction mixture was then refluxed for an additional hour in order to ensure complete reaction.

After reflux, the apparatus and products of the synthesis were allowed to cool to room temperature, and the lithium bromide side product was removed by filtration. The bromosilane reaction products can then be isolated from the remaining pentane by crystallization in a dry-ice/acetone bath. After immersion for approximately two hours, the pentane can be cannulaed off, leaving white crystals (28.1 g, 69.1% yield based on methyltribromosilane) behind.

## REFERENCES

- [1] T. H. Kunstmann, H. Angerer, J. Knecht, S. Veprek, N. W. Mitzel, and H. Schmidbaur, *Chem. Mater.*, **7**, 1675 (1995).
- [2] Th. Kunstmann and S. Veprek, *Appl. Phys. Lett.*, **67**, 3126 (1995).
- [3] M. W. Pitcher, *Advanced Materials: The Fabrication, Characterization and Study of Novel Ceramics and Composites*, Ph.D. thesis, University of Massachusetts, Amherst, MA, 2001.
- [4] P. A. Bianconi and T. W. Weidman, *J. Am. Chem. Soc.*, **110**, 2342 (1988).
- [5] M. W. Pitcher, S. J. Joray, and P. A. Bianconi, *Adv. Mater.*, **16**, 706 (2004).
- [6] E. G. Rochow, *J. Am. Chem. Soc.*, **67**, 963 (1945).
- [7] P. A. McCusker and E. L. Reilly, *J. Am. Chem. Soc.*, **75**, 1583 (1953).
- [8] E. V. Van den Berghe and G. P. Van der Kelen, *J. Organomet. Chem.*, **59**, 175 (2004).