HETEROCYCLES, Vol. 86, No. 2, 2012, pp. 1621 - 1635. © 2012 The Japan Institute of Heterocyclic Chemistry Received, 29th August, 2012, Accepted, 1st October, 2012, Published online, 15th October, 2012 DOI: 10.3987/COM-12-S(N)114

# SYNTHESISOF1,4-PHOSPHASILACYCLOHEXA-2,5-DIENESBEARING HYDROGENORCHLORINE ATOMSONATOMS

### Yoshiyuki Mizuhata, Satoshi Morikawa, and Norihiro Tokitoh\*

Institute for Chemical Research, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan tokitoh@boc.kuicr.kyoto-u.ac.jp

Abstract -2,3,5,6-Unsubstituted 1,4-phosphasilacyclohexa-2,5-dienes bearing a functional group on the silicon atom were synthesized *via* the corresponding stannacycle. The structures of newly synthesized silacyclic compounds were determined by X-ray crystallographic analysis.

# Dedicated to Dr. Professor Ei-ichi Negishi, Purdue University on the occasion of his 77th birthday.

The chemistry of  $[4n+2]\pi$ -electron ring systems containing a heavier group 14 element (Si, Ge, Sn, Pb), which we call, "*heavy aromatic compounds*,"<sup>1-3</sup> has attracted much attention from the viewpoint of comparison with the parent aromatic hydrocarbons, which play very important roles in organic chemistry. Although heavy aromatic compounds have been known to be highly reactive and undergo ready dimerization or oligomerization under ambient conditions, we have succeeded in the synthesis of the first stable neutral sila-, germa-, and stanna-aromatic compounds,<sup>2</sup> *i. e.*, heavier analogues of benzene [Si,<sup>4</sup> Ge<sup>5</sup> (Sn: unstable<sup>6</sup>), naphthalene (Si,<sup>7,8</sup> Ge,<sup>9</sup> Sn<sup>10</sup>), anthracene (Si,<sup>11</sup> Ge<sup>12</sup>), and phenanthrene [Si,<sup>13</sup> Ge<sup>12</sup> (Sn: unstable<sup>14</sup>)], by taking advantage of kinetic stabilization using efficient steric protection groups, 2,4,6-tris[bis(trimethylsilyl)methyl]phenyl (Tbt) and 2,6-bis[bis(trimethylsilyl)methyl]-4-[tris(trimethyl-silyl)methyl]phenyl (Bbt). During the course of our studies on heavy aromatic compounds, we have been interested in the corporation of heavier group 14 elements and heteroatoms of other group, that is, the synthesis of heavier congeners of heteroaromatic compounds. As the first target compounds, we designed the 1,4-phosphasilabenzene derivatives, whose synthetic strategies are shown in Scheme 1. For their syntheses, 1,4-phosphasilacyclohexa-2,5-dienes **1** bearing a functional group on the silicon atom

should be key precursors. As for 1,4-phosphasilacyclohexa-2,5-diene skeletons, some benzene- or thiophene-fused derivatives have been reported,<sup>15</sup> but there is no example bearing a functional group on the silicon atom. In addition, 2,3,5,6-unsubstituted derivative is limited to only one example, 4,4-di-methyl-1-phenyl-1,4-phosphasilacyclohexa-2,5-diene, reported by Märkl *et al.*<sup>16</sup> In this paper, we report the construction of 1,4-phosphasilacyclohexa-2,5-diene skeletons bearing hydrogen or chlorine atoms on the silicon atoms together with their properties.



Scheme 1. Synthetic strategies for 1,4-phosphasilabenzene derivatives

It is well known that the stannacycles are good precursors for the construction of heterocycles by the direct exchange of the tin atom with a heteroatom source<sup>17</sup> or by the lithiation with alkyllithium and the subsequent reaction with a heteroatom source.<sup>4,6,18</sup> Therefore, we synthesized 1,4-silastannacyclohexa-2,5-diene **2** according to the route shown in Scheme 2. In order to achieve selective introduction of two ethynyl substituents, we used dimethoxysilane **3** as a starting material. The reaction of **3** with ethynylmagnesium bromide afforded diethynylsilane **4** in 91% yield. Then, the hydrostannylation reaction of **4** using dibutyltin dihydride in heptane afforded **2** in 70% yield. The structures of **2** and **4** were reasonably supported by the <sup>1</sup>H and <sup>13</sup>C NMR and mass spectra along with elemental analysis, and finally determined by X-ray crystallographic analysis. The structure of **2** is shown in Figure 1. This is the first structural determination of a 2,3,5,6-unsubstituted 1,4-silastannacyclohexa-2,5-diene. The central 1,4-silastannacyclohexa-2,5-diene skeleton of **2** has a boat shape, and the deviations of Si1 and Sn1 atoms from the C1–C2–C3–C4 plane are 0.275 and 0.331 Å, respectively. The selected structural parameters of **2** are summarized in Table 1.



Scheme 2. Synthesis of 1,4-silastannacyclohexa-2,5-diene 2



Figure 1. Thermal ellipsoid plot of 2 (50% probability). Hydrogen atoms except for those on the  $SiC_4Sn$  ring and the minor part of the disordered butyl substituent were omitted for clarity.

In order to introduce a phosphorus moiety instead of the dibutyltin unit of 2, we initially attempted the direct exchange using phosphorus trichloride by following the syntheses of heavier congeners of pyridine.<sup>17</sup> However, desired product was not obtained even after several attempts. Next, we tried the transmetalation reaction between tin and lithium and the following subsequent reactions with aryldichlorophophines. As a result of the optimization of reaction conditions, it was found that the reaction of 2 with 3.2 equivalent of n-BuLi in the presence of 3.2 equivalents of tetramethylethylenediamine afforded the corresponding dilithio species effectively. The subsequent addition of dichlorophosphines afforded the corresponding 1,4-phosphasilahexa-2.5-dienes 5 and 6 in moderate yields (Scheme 3). In the case using dichlorophenylphosphine, ca. 1:1 mixture of cis- and trans-isomers 5 was obtained as judged by  ${}^{31}$ P NMR ( $\delta_P = -25.5, -25.7$ ). In the case using Mes\*-substituted dichlorophosphine, on the other hand, only *trans*-isomer **6** was generated, judging from the sole signal in the <sup>31</sup>P NMR ( $\delta_P = -24.5$ ) and the results of its X-ray crystallographic analysis (Figure 2). The bulkiness of Mes\* group is considered to suppress the formation of the corresponding *cis*-isomer. Since 1.4-phosphasilahexa-2.5-dienes 5 and 6 were found to be somewhat unstable and their isolation in enough amount to analyze their chemical properties were difficult, their oxidation by hydrogen peroxide was performed. The reactions proceeded cleanly to afford the corresponding phosphine oxides 7 and 8. At this stage, *cis*- and *trans*-isomers of 7 were isolable. The structures of 7 and 8 were reasonably supported by the <sup>1</sup>H and <sup>13</sup>C NMR and mass spectra together with elemental analysis, and finally determined by X-ray crystallographic analysis (Figure 2). These are the first examples of structural determination of 2,3,5,6-unsubstituted 1,4-phoshasilacyclohexa-2,5-diene. The oxidation of 6proceeded with an inversion fashion. The <sup>31</sup>P NMR signals were observed at 3.4 (*cis*-7), 5.7 (*trans*-7), and 12.6 (trans-8) ppm.



Scheme 3. Synthesis of 1,4-phosphasilacyclohexa-2,5-dienes



**Figure 2.** Thermal ellipsoid plots of *trans*-**6**, *trans*-**7**, *cis*-**7**, and *trans*-**8** [50% probability except for *cis*-**7** (30%)]. Hydrogen atoms except for those on the SiC<sub>4</sub>P ring, solvents, minor parts of the disorder, and one of the two independent molecules of *trans*-**8** were omitted for clarity.

Selected structural parameters of 2, 6, 7, and 8 and their crystallographic data are summarized in Tables 1 and 2, respectively. The structural parameters of *cis*-7 are not accurate due to the severe disorder covering the whole of Tbt group. The conformations of SiC<sub>4</sub>P rings of Mes\*-substituted derivatives, *trans*-6 and 8, were boat shapes similarly to that of stannacycle 2. On the other hand, phenyl-substituted ones, *trans*- and *cis*-7, have almost planar SiC<sub>4</sub>P rings, and the sums of the internal bond angles are 718.8 (*trans*-7) and 719.6° (*cis*-7). While the bond lengths in 6-8 have little differences, there are large differences in their internal angles. Theoretical calculations by using the model compounds bearing hydrogen atoms on the silicon and phosphorus atoms instead of aryl substituents (phosphine 9 and phosphine oxide 10, respectively) showed that the boat shape of 9 and the planar geometry of 10 were global minima (Table 1). Therefore, the boat-shape conformations of 8 were considered to be most likely due to the steric repulsion between bulky substituents and/or the packing forces. The benzene-and thiophene-fused 1,4-phosphasilacyclohexa-2,5-diene were reported to have planar or butterfly-shape towards the Si–P axis.

Table 1.Se	elected bond	lengths	and angles <sup>a</sup>
------------	--------------	---------	-------------------------

(or H)

Tht

C1 - C2

$H^{W^{W^{W^{W^{W^{W^{W^{W^{W^{W^{W^{W^{W^$	$\delta X$ $\varepsilon$ -C3 . <sup>b</sup> t	wo independ	ent molecules.	<sup>c</sup> calcula	ted at B3LYP/	6-31G(d) lev	el.	
	2	trans-6	trans-7	cis-7	trans	$5-8^b$	<b>9</b> <sup>c</sup>	<b>10</b> <sup>c</sup>
Х	$\operatorname{Sn}(n-\operatorname{Bu})_2$	PMes*	P(O)Ph	P(O)Ph	P(O)M	/les*	PH	P(O)H
Si1–C1 (Å)	1.852(6)	1.8679(18)	1.8737(18)	1.832(6)	1.871(4)	1.878(3)	1.8704	1.8727
C1–C2 (Å)	1.344(8)	1.341(2)	1.333(2)	1.341(8)	1.340(5)	1.343(5)	1.3455	1.3434
C2–X (Å)	2.097(6)	1.8067(18)	1.7938(18)	1.776(6)	1.803(4)	1.803(4)	1.8291	1.8189
X–C3 (Å)	2.149(6)	1.7960(18)	1.7919(18)	1.772(7)	1.796(4)	1.800(3)	1.8291	1.8189
C3–C4 (Å)	1.321(7)	1.344(2)	1.337(2)	1.342(8)	1.336(5)	1.324(4)	1.3455	1.3434
C4–Si1 (Å)	1.877(5)	1.8677(18)	1.8612(18)	1.823(7)	1.870(4)	1.876(3)	1.8704	1.8727
$\alpha$ (deg)	111.9(3)	103.57(8)	105.48(8)	106.3(3)	103.72(16)	103.40(15)	106.57	107.81
$\beta$ (deg)	126.3(5)	124.82(14)	125.71(14)	125.8(5)	122.1(3)	122.1(3)	123.65	124.40
$\gamma$ (deg)	124.6(5)	124.77(14)	126.70(14)	126.6(5)	120.0(3)	120.6(3)	127.12	127.56
$\delta$ (deg)	101.5(2)	104.00(8)	108.32(8)	108.1(3)	105.87(16)	105.99(16)	104.50	108.23
$\varepsilon$ (deg)	122.7(4)	125.10(14)	127.15(14)	127.3(5)	120.7(3)	120.6(3)	127.12	127.56
$\zeta$ (deg)	127.6(5)	124.46(14)	125.42(14)	125.5(5)	121.9(3)	122.9(3)	123.65	124.40
$\Sigma(\alpha - \zeta)$ (deg)	714.6	706.7	718.8	719.6	694.3	695.6	712.61	719.96

	2	trans-6	$[trans-7 \cdot H_2O]$	cis- <b>7</b>	[ <i>trans</i> -8·2.25CHCl <sub>3</sub> ]	
Empirical	C <sub>39</sub> H <sub>82</sub> Si <sub>7</sub> Sn	$C_{49}H_{93}PSi_7$	C37H71O2PSi7	C37H69OPSi7	$C_{51.25}H_{95.25}Cl_{6.75}OPSi_7$	
formula						
Formula weight	866.37	909.83	775.54	757.52	1194.41	
Temperature	103(2)	103(2)	103(2)	103(2)	103(2)	
(K)						
Crystal color	colorless	colorless	colorless	colorless	colorless	
Crystal	0.20 x 0.20 x 0.05	0.20 x 0.10 x 0.05	0.30 x 0.20 x 0.10	0.40 x 0.20 x 0.10	0.20 x 0.10 x 0.10	
dimensions						
(mm)						
Crystal system	triclinic	triclinic	monoclinic	orthorhombic	triclinic	
Space group	P-1 (#2)	P-1 (#2)	$P2_1/a$ (#14)	$Pca2_{1}$ (#29)	<i>P</i> -1 (#2)	
a (Å)	12.0429(3)	9.9781(2)	18.1768(2)	12.5275(6)	13.0770(3)	
<i>b</i> (Å)	13.4396(4)	10.92960(10)	10.9821(2)	18.8233(9)	22.6909(5)	
<i>c</i> (Å)	17.2333(7)	28.3562(5)	23.1305(3)	20.1264(13)	23.9992(7)	
lpha (°)	76.083(3)°	99.5806(11)	90	90	92.6189(18)	
eta (°)	74.8668(16)°	92.8543(7)	90.8411(8)	90	99.9179(14)	
γ(°)	69.5949(13)°	107.4502(8)	90	90	100.9656(8)	
$V(\text{\AA}^3)$	2488.88(14)	2892.58(8)	4616.80(12)	4746.0(4)	6864.1(3)	
Ζ	2	2	4	4	4	
$D_{\text{calc}} \left( \mathbf{g} \cdot \mathbf{cm}^{-3} \right)$	1.156	1.045	1.116	1.060	1.156	
$\mu (\mathrm{mm}^{-1})$	0.706	0.221	0.270	0.260	0.457	
$\theta$ range (°)	1.84 to 25.00	1.46 to 25.00	2.17 to 25.00	1.95 to 25.00	1.79 to 25.00	
Independent	8446	10121	8101	7652	24068	
reflections						
$R_{\rm int}$	0.0599	0.0337	0.0310	0.1042	0.0751	
Completeness	96.3	99.5	99.7	97.1	99.6	
to $\theta(\%)$						
No. of	483	608	451	692	1290	
parameters						
Goodness of fit	1.046	1.037	1.021	1.018	1.005	
Final R indices	0.0577	0.0393	0.0301	0.0580	0.0567	
[ <i>I</i> >2 <i>o</i> ( <i>I</i> )]						
R indices (all	0.1384	0.1034	0.0820	0.1446	0.1408	
data)						
Largest diff.	1.030	0.390	0.647	0.238	0.706	
peak (e·Å <sup>3</sup> )						
Largest diff.	-0.928	-0.206	-0.377	-0.212	-0.558	
hole (e·Å <sup>3</sup> )						

 Table 2.
 X-Ray crystallographic data

In order to obtain **5** in a pure form, the reduction of isolated **7** using trichlorosilane was performed (Scheme 4). Unexpectedly, chlorination of Si–H occurred together with the reduction of the phosphine oxide moiety to afford **11**. Although we have not obtained detailed structural information for **11**, a single isomer was generated from each isomer ( $\delta_{si} = -26.4$ ,  $\delta_p = -25.2$ ). The structure of **11** was supported by the reasonable chemical shifts in <sup>29</sup>Si and <sup>31</sup>P NMR and the fact that the reaction of **11** with lithium aluminum hydride afforded hydrosilane **5** again as a mixture of isomers. On the other hand, the reaction of **8** with trichlorosilane did not proceed under similar conditions.



Scheme 4. Reduction and chlorination of 7

By using hydro- or chlorosilanes 5, 6, and 11 here obtained, we performed several reactions expecting the generation of 1,4-phosphasilabenzenium cations (5 or 6 with  $Ph_3C^+TPFPB^-$ , 11 with  $Ag^+TPFPB^-$ , and so on). However, all attempts were unsuccessful so far and gave a complicated mixture. On the other hand, we have a preliminary result of the elimination of phenyl group on the phosphorus atom in 5 by the reaction with lithium dispersion. We think this result suggests the possibility to access 1,4-phosphasilabenzene. At the same time, the ring-closing methodology using stannacycle 2 is considered to be applicable to synthesize other silicon-containing heterocycles. We are currently continuing this study with a scope of various atom combinations.

# **EXPERIMANTAL**

**General procedures.** All experiments were performed under an argon atmosphere unless otherwise noted. Solvents used for the reactions were purified by The Ultimate Solvent System (GlassContour Company).<sup>19</sup> <sup>1</sup>H NMR (300 MHz), <sup>13</sup>C NMR (76 MHz), <sup>31</sup>P NMR (121 MHz), and <sup>119</sup>Sn NMR (111 MHz) spectra were measured in CDCl<sub>3</sub> or C<sub>6</sub>D<sub>6</sub> with a JEOL JNM-AL300 spectrometer. In <sup>1</sup>H NMR, signals due to CHCl<sub>3</sub> (7.25 ppm) and C<sub>6</sub>D<sub>5</sub>H (7.15 ppm) were used as references, and those due to CDCl<sub>3</sub> (77 ppm) and C<sub>6</sub>D<sub>6</sub> (128 ppm) were used in <sup>13</sup>C NMR. <sup>31</sup>P NMR spectra were measured using 85% H<sub>3</sub>PO<sub>4</sub> in water (0 ppm) as an external standard. <sup>119</sup>Sn NMR was measured with NNE technique using SnMe<sub>4</sub> as an external standard. Multiplicity of signals in <sup>13</sup>C NMR spectra was determined by DEPT technique, which was described as s, d, t, and q for compounds **2-4** and as C, CH, CH<sub>2</sub>, and CH<sub>3</sub> with the

coupling with <sup>31</sup>P for phosphorus-containing compounds **5-8** and **11**. GPLC (gel permeation liquid chromatography) was performed on an LC-908, LC-918, or LC-908-C60 (Japan Analytical Industry Co., Ltd.) equipped with JAIGEL 1H and 2H columns (eluent: chloroform or toluene). All melting points were determined on a Yanaco micro melting point apparatus and were uncorrected. Elemental analyses were carried out at the Microanalytical Laboratory of the Institute for Chemical Research, Kyoto University. All theoretical calculations were carried out using the Gaussian 09<sup>20</sup> programs.

**Synthesis of 3.** To a THF solution (30 mL) of TbtBr (5.00 g, 7.91 mmol) at -78 °C was added *t*-BuLi (1.57 M, 12.0 mL, 19.0 mmol). After stirring for 30 min, trimethoxysilane (3.02 mL, 2.09 g, 23.7 mmol) was added to the solution followed by warming up to room temperature overnight. Then, MeOH was added to the mixture, and the reaction mixture was extracted with hexane/H<sub>2</sub>O. The organic layer was dried over MgSO<sub>4</sub> and the solvent was removed to give NMR pure **3** (5.02 g, 7.79 mmol, 99%) as a colorless solid. **3**: mp 163 °C (dec.); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, 298 K): δ 0.01 (s, 36H), 0.25 (s, 18H), 1.30 (s, 1H), 2.32 (s, 2H), 3.59 (s, 6H), 5.14 (s, 1H), 6.20 (s, 1H), 6.33 (s, 1H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>, 298 K); δ 0.5 (q), 0.7 (q), 27.0 (d), 27.4 (d), 30.5 (d), 51.6 (q), 121.7 (d), 124.2 (s), 126.5 (d), 145.4 (s), 151.6 (s), 151.9 (s); <sup>29</sup>Si NMR (60 MHz, CDCl<sub>3</sub>, 298 K); δ -22.2, 1.7, 1.9; HRMS (FAB) *m/z* calcd for C<sub>29</sub>H<sub>66</sub>O<sub>2</sub>Si<sub>7</sub> 642.344 (M<sup>+</sup>), Found: 642.3443 (M<sup>+</sup>); Anal. Calcd for C<sub>29</sub>H<sub>66</sub>O<sub>2</sub>Si<sub>7</sub>: C, 54.13; H, 10.34. Found: C, 53.90; H, 10.21.

**Synthesis of 4.** To a THF solution (20 mL) of **3** (6.05 g, 9.40 mmol) was added ethynylmagnesium bromide in THF (0.5 M, 56.4 mL, 28.2 mmol) at room temperature. The reaction mixture was stirred for 13 h under reflux conditions. Then, NH<sub>4</sub>Cl aq was added to the mixture at 0 °C, and the reaction mixture was extracted with hexane. The organic layer was dried over MgSO<sub>4</sub> and the solvent was removed. The reaction mixture was separated by silica gel chromatography (hexane) to give **4** (5.38 g, 8.53 mmol, 91%) as a colorless solid. **4**: Rf = 0.37 (hexane); colorless crystals; mp 166.5-167.8 °C; <sup>1</sup>H NMR (300 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$  0.04 (br s, 18H), 0.05 (br s, 18H), 0.07 (br s, 18H), 1.33 (s with satellites, 1H, <sup>2</sup>J<sub>HSi</sub> = 10.0 Hz), 2.40 (s, 2H), 2.580 (s, 1H), 2.584 (s, 1H), 5.22 (s with satellites, 1H, <sup>1</sup>J<sub>HSi</sub> = 225 Hz), 6.29 (br s, 1H), 6.41 (br s, 1H); <sup>13</sup>C NMR (75 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$  0.6 (q), 0.7 (q), 1.0 (q), 29.1 (d), 29.6 (d), 30.8 (d), 85.1 (s), 97.0 (d), 118.5 (s), 121.9 (d), 126.8 (d), 146.3 (s), 152.5 (s), 152.9 (s); <sup>29</sup>Si NMR (59 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$  –76.5, 2.2, 2.8; HRMS (FAB) *m/z* calcd for C<sub>31</sub>H<sub>62</sub>Si<sub>7</sub> 630.3236 (M<sup>+</sup>), Found 630.3237; Anal. Calcd for C<sub>31</sub>H<sub>62</sub>Si<sub>7</sub>: C, 58.97; H, 9.99. Found: C, 59.22; H, 9.93.

Synthesis of 2. The heptane solution (35 mL) of 4 (790 mg, 1.25 mmol) and dibutyltin dihydride (1.11 g, 4.75 mmol) was stirred for 18 h under reflux conditions. Then, the solvent was removed, and the reaction mixture was separated by silica gel chromatography (hexane) and GPC (eluent: toluene) to give 2 (759 mg, 0.876 mmol, 70%) as a colorless solid. 2: Rf = 0.87 (hexane); colorless crystals; mp 114.9-115.4 °C; <sup>1</sup>H NMR (300 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  0.16 (br s, 18H), 0.20 (br s, 18H), 0.21 (br s, 18H),

0.86-1.12 (m, 10H), 1.15-1.79 (m, 8H), 1.46 (s, 1H), 2.60 (br s, 1H), 2.63 (br s, 1H), 5.63 (s with satellites, 1H,  ${}^{1}J_{HSi} = 188$  Hz), 6.53 (br s, 1H), 6.63 (br s, 1H), 7.27 (d,  ${}^{3}J = 18.9$  Hz, 2H), 7.44 (d,  ${}^{3}J = 18.9$  Hz, 2H);  ${}^{13}$ C NMR (75 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  1.0 (q), 1.1 (q), 1.5 (q), 11.1 (t), 11.3 (t), 13.88 (q), 13.90 (q), 27.4 (t), 27.6 (t), 29.4 (d), 29.5 (t×2), 29.6 (d), 30.9 (d), 122.3 (d), 125.2 (s), 127.2 (d), 145.2 (s), 151.3 (d), 152.5 (d), 152.4 (s), 153.1 (s);  ${}^{29}$ Si NMR (59 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  -57.2, 2.0, 2.3;  ${}^{119}$ Sn NMR (111 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  -170.9; HRMS (FAB) *m*/*z* calcd for C<sub>39</sub>H<sub>82</sub>Si<sub>7</sub>Sn 866.3823 (M<sup>+</sup>), Found 866.3828; Anal. Calcd for C<sub>39</sub>H<sub>82</sub>Si<sub>7</sub>Sn: C, 54.07; H, 9.54. Found: C, 54.11; H, 9.55.

Synthesis of 7 via 5. To a THF solution (0.78 mL) of 2 (115 mg, 0.133 mmol) and tetramethylethylenediamine (0.0639 mL, 0.426 mmol) was added *n*-BuLi (1.48 M in hexane, 0.288 mL, 0.426 mmol) at -78 °C. After stirring for 15 h at the same temperature, dichlorophenylphosphine (0.0904 mL, 0.666 mmol) was added to the reaction mixture at -78 °C. The reaction mixture was allowed to warm up to -60 °C, and kept for 6 h at the same temperature. Then, H<sub>2</sub>O<sub>2</sub> aq (30%, 0.0367) mL, 1.20 mmol) was added to the reaction mixture at 0 °C. After stirring for 3 h at room temperature, the reaction mixture was extracted with hexane. The organic layer was dried over MgSO<sub>4</sub> and the solvent was removed. The reaction mixture was separated by silica gel chromatography (Et<sub>2</sub>O) to give *cis*-7 (32.9 mg, 0.0434 mmol, 33%) and *trans*-7 (34.8 mg, 0.0459 mmol, 35%) as colorless solids. *cis*-7: Rf = 0.57 (Et<sub>2</sub>O); colorless crystals; mp 154 °C (dec.); <sup>1</sup>H NMR (300 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$  0.04 (br s, 18H), 0.05 (br s, 18H), 0.07 (br s, 18H), 1.35 (s with satellites, 1H,  ${}^{2}J_{SiH} = 5.0$  Hz), 2.13 (br s, 1H), 2.15 (br s, 1H), 5.21 (s with satellites, 1H,  ${}^{1}J_{HSi} = 199$  Hz), 6.32 (br s, 1H), 6.43 (br s, 1H), 6.99-7.25 (m, 4H), 7.46-7.55 (m, 3H), 7.72-7.80 (m, 2H); <sup>13</sup>C{<sup>1</sup>H} NMR (75 MHz, 298 K, CDCl<sub>3</sub>) δ 0.7 (CH<sub>3</sub>), 1.0 (CH<sub>3</sub>), 29.9 (CH), 30.3 (CH), 30.8 (CH), 119.3 (C), 122.0 (CH), 126.9 (CH), 128.7 (CH, d, J<sub>CP</sub> = 11.7 Hz), 130.8 (CH, d,  $J_{CP} = 10.5$  Hz), 131.8 (CH, d,  $J_{CP} = 3.1$  Hz), 132.8 (C, d,  $J_{CP} = 105$  Hz), 141.7 (CH, d,  ${}^{1}J_{CP} = 99.8$ Hz), 145.9 (CH, d,  ${}^{3}J_{CP} = 6.2$  Hz), 146.5 (C), 152.9 (C), 153.2 (C);  ${}^{31}P$  NMR (121 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$ 3.43 ( ${}^{3}J_{PSi} = 33 \text{ Hz}$ ); <sup>29</sup>Si NMR (59 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$  –61.9 (d,  ${}^{3}J_{SiP} = 33 \text{ Hz}$ ), 1.9, 2.3; Anal. Calcd for  $C_{37}H_{70}OPSi_7$ : C, 58.67; H, 9.18. Found: C, 57.77; H, 8.96. *trans*-7: Rf = 0.30 (Et<sub>2</sub>O); colorless crystals; mp 178 °C (dec.); <sup>1</sup>H NMR (300 MHz, 298 K, CDCl<sub>3</sub>) δ 0.026 (br s, 18H), 0.037 (br s, 18H), 0.042 (br s, 18H), 1.35 (s, 1H), 1.92 (br s, 1H), 1.97 (br s, 1H), 5.31 (d with sateslites, 1H,  ${}^{4}J_{HP} = 5.9$  Hz,  ${}^{1}J_{\text{HSi}} = 202 \text{ Hz}$ , 6.31 (br s, 1H), 6.43 (br s, 1H), 6.93-7.36 (m, 4H), 7.46-7.58 (m, 3H), 7.73-7.78 (m, 2H); <sup>13</sup>C{<sup>1</sup>H} NMR (75 MHz, 298 K, CDCl<sub>3</sub>) δ 0.67 (CH<sub>3</sub>), 0.72 (CH<sub>3</sub>), 1.0 (CH<sub>3</sub>), 30.3 (CH), 30.6 (CH), 30.9 (CH), 120.0 (C), 122.0 (CH), 126.9 (CH), 128.7 (CH, d,  $J_{CP} = 12.3$  Hz), 130.7 (CH, d,  $J_{CP} = 10.5$  Hz), 132.1 (CH, d,  $J_{CP} = 2.5$  Hz), 132.6 (C, d,  $J_{CP} = 105.9$  Hz), 141.4 (CH, d,  $J_{CP} = 99.8$  Hz), 146.5 (C), 149.1 (CH, d,  $J_{CP} = 5.5$  Hz), 152.5 (C), 152.7 (C); <sup>31</sup>P NMR (121 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$  5.65 (<sup>3</sup> $J_{PSi} = 35$  Hz); <sup>29</sup>Si NMR (59 MHz, 298 K, CDCl<sub>3</sub>)  $\delta$  –61.6 (d, <sup>3</sup>J<sub>SiP</sub> = 35 Hz), 1.9, 2.1; HRMS (FAB) *m/z* calcd for C<sub>37</sub>H<sub>70</sub>OPSi<sub>7</sub> [(M+H)<sup>+</sup>]: 757.3549, found 757.3525; Anal. Calcd for C<sub>37</sub>H<sub>72</sub>O<sub>2</sub>PSi<sub>7</sub> (+H<sub>2</sub>O): C, 57.30; H,

## 9.23. Found: C, 57.58; H, 9.14.

In several attempts, before adding  $H_2O_2$  aq, removing the solvents and filtration through Celite<sup>\*</sup> with hexane were performed to give a mixture containing **5** mainly. However, separation of isomers each other from the mixture was difficult. **5** (a mixture of *cis*- and *trans*-isomers): a colorless solid; <sup>1</sup>H NMR (300 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>, only for characteristic peaks)  $\delta$  5.48 (Si–H, d, 1H, <sup>4</sup>J<sub>HP</sub> = 17.8 Hz), 5.82 (Si–H, d with satellites, 1H, <sup>4</sup>J<sub>HP</sub> = 3.3 Hz, <sup>1</sup>J<sub>HSi</sub> = 192 Hz); <sup>31</sup>P NMR (121 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  –25.7, –25.5; <sup>29</sup>Si NMR (59 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  –61.4 (d, <sup>3</sup>J<sub>SiP</sub> = 11.0 Hz), –57.0 (d, <sup>3</sup>J<sub>SiP</sub> = 12.2 Hz), 2.0, 2.1, 2.3.

Synthesis of *trans*-6. To a THF solution (0.78 mL) of **2** (790 mg, 0.912 mmol) and tetramethylethylenediamine (0.438 mL, 2.92 mmol) was added *n*-BuLi (1.48 M in hexane, 2.05 mL, 3.03 mmol) at -78 °C. After stirring for 11 h at the same temperature, Mes\*PCl<sub>2</sub> (1.580 g, 4.56 mmol) was added to the reaction mixture at -78 °C. The reaction mixture was allowed to warm up to -30 °C, and kept for 24 h at the same temperature. Then, the solvents were removed at -30 °C, and the reaction mixture was put into a glovebox filled with argon. To the reaction mixture was added *n*-hexane, and the resulting suspension was filtered with Celite. The solvent was removed in vacuo. Fortunately, the single crystal of *trans*-6 was obtained from the mixture at this stage. However, the isolation of pure *trans*-6 in enough amount to analyze its chemical properties was difficult in spite of a number of times of recrystallization in a glovebox. Therefore, the mixture was separated by GPC (eluent: toluene) to give *trans*-6 (632 mg, 0.694 mmol, 76%, if pure) contaminated with some impurities. *trans*-6: <sup>1</sup>H NMR (300 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  0.18 (s, 18H), 0.24 (br s, 36H), 1.29 (s, 9H), 1.49 (s, 1H), 1.64 (br s, 18H), 2.60 (br s, 2H), 5.97 (br s, 1H), 6.46 (dd, *J* = 16.5, 22.3 Hz, 2H), 6.56 (br s, 1H), 6.66 (br s, 1H), 7.12-7.40 (m, 2H), 7.62 (s, 1H), 7.63 (s, 1H); <sup>31</sup>P NMR (121 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  –24.5.

**Synthesis of** *trans-8.* To a THF solution (0.84 mL) of *trans-6* (130 mg, 0.143 mmol, if pure) was added H<sub>2</sub>O<sub>2</sub> aq (30%, 0.0438 mL, 1.43 mmol) at 0 °C. After stirring for 12 h at room temperature, the solvents were removed in vacuo. The reaction mixture was separated by silica gel chromatography (*n*-hexane:Et<sub>2</sub>O = 2:1) to give *trans-8* (108 mg, 0.117 mmol, 82%) as a colorless solid. *trans-8*: Rf = 0.60 (*n*-hexane:Et<sub>2</sub>O = 2:1); colorless crystals; mp 242 °C (dec.); <sup>1</sup>H NMR (300 MHz, 298 K, CDCl<sub>3</sub>) δ 0.03 (br s, 36H), 0.04 (br s, 18H), 1.29 (s, 9H), 1.34 (s, 1H), 1.48 (s, 18H), 1.83 (br s, 1H), 1.92 (br s, 1H), 5.36 (d, 1H, <sup>4</sup>*J*<sub>HP</sub> = 6.1 Hz, satellite, <sup>1</sup>*J*<sub>HSi</sub> = 211 Hz), 6.32 (br s, 1H), 6.44 (br s, 1H), 7.07 (dd, *J* = 41.7, 16.5 Hz, 2H), 7.34 (s, 1H), 7.35 (s, 1H), 7.49 (dd, *J* = 30.2, 16.5 Hz, 2H); <sup>13</sup>C{<sup>1</sup>H} NMR (75 MHz, 298 K, CDCl<sub>3</sub>) δ 0.8 (CH<sub>3</sub>), 1.1 (CH<sub>3</sub>), 30.1 (CH), 30.3 (CH), 30.8 (CH), 31.0 (CH<sub>3</sub>), 33.8 (CH<sub>3</sub>), 34.5 (CH), 40.1 (C, d, *J*<sub>CP</sub> = 3.1 Hz), 121.1 (C), 122.0 (CH), 123.8 (CH, *J*<sub>CP</sub> = 11.3 Hz), 126.3 (C, d, *J*<sub>CP</sub> = 106.8 Hz), 127.1 (CH), 143.2 (CH, d, *J*<sub>CP</sub> = 4.2 Hz), 145.8 (C), 149.4 (CH, d, *J*<sub>CP</sub> = 98.6 Hz), 151.0 (C, *J*<sub>CP</sub> = 3.2 Hz), 152.2 (C), 152.6 (C), 159.3 (C); <sup>31</sup>P NMR (121 MHz, 298 K, CDCl<sub>3</sub>) δ 12.6 (<sup>3</sup>*J*<sub>PSi</sub> = 35 Hz); <sup>29</sup>Si NMR (59 MHz, 298 K, CDCl<sub>3</sub>) δ -62.0 (d, <sup>3</sup>*J*<sub>SiP</sub> = 35 Hz), 1.9, 2.1; HRMS (APPI-TOF) *m/z* calcd for C<sub>49</sub>H<sub>94</sub>OPSi<sub>7</sub>

[(M+H)<sup>+</sup>]: 925.5422, found 925.5500; Anal. Calcd for C<sub>49</sub>H<sub>93</sub>OPSi<sub>7</sub>: C, 63.57; H, 10.12. Found: C, 63.60; H, 10.28.

**Synthesis of 11.** To a C<sub>6</sub>D<sub>6</sub> solution (1.55 mL) of *cis*-7 (140 mg, 0.186 mmol) in an NMR tube (5¢) filled with argon was added trichlorosilane (0.0934 mL, 0.924 mmol) at room temperature. The NMR tube was sealed with screw-type cap, and then heated at 80 °C. Starting material was consumed after 1.5 h, resulting in the formation of **11** almost quantitatively. Recrystallization form *n*-hexane at –40 °C afforded **11** (77.0 mg, 0.0992 mmol, 53%). When using *trans*-7 as a starting material, the same compound was generated. **11**: <sup>1</sup>H NMR (300 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  0.15 (br s, 18H), 0.24 (br s, 18H), 0.27 (br s, 18H), 1.47 (s, 1H), 2.78 (s, 2H), 6.52 (br s, 1H), 6.63 (br s, 1H), 6.77-7.15 (m, 7H), 7.40-7.45 (m, 2H); <sup>13</sup>C{<sup>1</sup>H} NMR (75 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  1.0 (CH<sub>3</sub>), 1.1 (CH<sub>3</sub>), 1.5 (CH<sub>3</sub>), 29.1 (CH), 29.4 (CH), 31.1 (CH), 122.9 (CH), 124.6 (C, d, *J*<sub>CP</sub> = 6.2 Hz), 127.7 (CH), 128.8 (CH, d, *J*<sub>CP</sub> = 8.6 Hz), 130.1 (CH), 133.4 (C, d, *J*<sub>CP</sub> = 4.9 Hz), 133.9 (CH, d, *J*<sub>CP</sub> = 5.6 Hz), 134.1 (CH, d, *J*<sub>CP</sub> = 21.6 Hz), 146.7 (C), 148.2 (CH, d, *J*<sub>CP</sub> = 16.1 Hz), 152.9 (C), 153.4 (C); <sup>31</sup>P NMR (121 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  -25.2 (satellite, <sup>3</sup>*J*<sub>PSi</sub> = 9 Hz); <sup>29</sup>Si NMR (59 MHz, 298 K, C<sub>6</sub>D<sub>6</sub>)  $\delta$  -26.4 (d, <sup>3</sup>*J*<sub>SiP</sub> = 9 Hz), 2.3, 2.8.

**Reaction of 11 with lithium aluminum hydride.** To a THF solution (0.65 mL) of **11** (25.4 mg, 0.0327 mmol) was added lithium aluminum hydride (1.7 mg, 0.045 mmol) at 0 °C. The reaction mixture was warmed up to room temperature. After stirring for 1 h, the solvent was removed, and the reaction mixture was put into a glovebox filled with argon. To the reaction mixture was added *n*-hexane, and the resulting suspension was filtered with Celite to afford almost pure **5** (a mixture of isomers, 24.5 mg, 0.0330 mmol, if pure).

**Reaction of 5 with Ph<sub>3</sub>C<sup>+</sup>TPFPB<sup>-</sup>**. In a glovebox filled with argon,  $C_6D_6$  (0.60 mL) was added to the mixture of **5** (a mixture of isomers, 17.8 mg, 0.0240 mmol) and Ph<sub>3</sub>C<sup>+</sup>TPFPB<sup>-</sup> (22.1 mg, 0.0240 mmol) in an NMR tube (5 $\phi$ ) at room temperature. The NMR tube was sealed with screw-type cap. Starting material was almost consumed after heating at 80 °C for 51 h. Although the generation of Ph<sub>3</sub>CH was observed, the <sup>29</sup>Si and <sup>31</sup>P NMR spectra were complicated ( $\delta_p$ : -15.2, -15.0, 6.7, 7.0;  $\delta_{Si}$ : -62.0, -61.9, -61.4, -61.3).

**Reaction of 6 with Ph\_3C^+TPFPB^-.** In a glovebox filled with argon,  $C_6H_6$  (1.90 mL) solution of **6** (60.6 mg, 0.0666 mmol) was added to  $Ph_3C^+TPFPB^-$  (61.4 mg, 0.0666 mmol) in a glass tube (10 $\phi$ ) at room temperature. The NMR tube was degassed and sealed. During heating at 70 °C for 15 h, the reaction mixture was separated to two phases. The tube was opened in a glovebox, and the solvent was removed. Although the generation of  $Ph_3CH$  was observed, many signals were observed around -44 and -23 ppm in the <sup>31</sup>P NMR. In addition to these signals, characteristic peak was observed at +221.4 ppm. Unfortunately, we could not determine the structure of the corresponding compound.

**Reaction of 11 with Ag<sup>+</sup>TPFPB<sup>-</sup>.** In a glovebox filled with argon, a  $C_6D_6$  solution (0.77 mL) of 11

(30.0 mg, 0.0387 mmol) was added to Ag<sup>+</sup>TPFPB<sup>-</sup> (39.5 mg, 0.0387 mmol) in an NMR tube (5 $\phi$ ) at room temperature. The NMR tube was sealed with screw-type cap. Starting material was consumed after 1 h, resulting in the complicated mixture ( $\delta_p = 6.5$ , 8.2, and many signals appeared between -19 and -26 ppm).

**Reaction of 5 with lithium dispersion.** In a glovebox filled with argon, lithium dispersion (3.6 mg, 0.52 mmol) was added to a THF solution (0.64 mL) of **5** (a mixture of isomers, 38 mg, 0.051 mmol) in an NMR tube (5 $\phi$ ) at room temperature. The NMR tube was sealed with screw-type cap. Starting material ( $\delta_p = -25.1, -25.3$  in THF) was consumed after 3 h, resulting in the quantitative formation of novel species ( $\delta_p = -17.4$  in THF). The non-decoupling <sup>31</sup>P NMR showed tt pattern (J = 44.3, 11.7 Hz), suggesting the retain of the –C(H)=C(H)–P–C(H)=C(H)– moiety. The <sup>1</sup>H NMR signals in 5-8 ppm were observed at 5.69 (dd, J = 15.9, 11.7 Hz, 2H), 6.49 (s, 1H, Si–H), 6.58 (br s, 1H, *m*-Tbt), 6.67 (br s, 1H, *m*-Tbt), 7.73 (dd, J = 44.3, 15.9 Hz, 2H) ppm. The detachment of the phenyl group and formation of P–Li species are most reasonable. However, this species decomposed by condensation, and several attempts of trapping reaction by MeOH or benzyl chloride afforded a complicated mixture.

General procedure for X-ray crystallographic analysis of compounds 2, *trans*-6, [*trans*-7·H<sub>2</sub>O], *cis*-7, and [*trans*-8·2.25CHCl<sub>3</sub>]. The intensity data were collected on a Rigaku/MSC Mercury CCD diffractometer with graphite monochromated MoK $\alpha$  radiation ( $\lambda = 0.71070$  Å). A single crystal suitable for X-ray analysis was mounted on a glass fiber. The structures were solved by a direct method (SHELXS-97)<sup>21</sup> and refined by full-matrix least-squares procedures on  $F^2$  for all reflections (SHELXL-97).<sup>21</sup> All hydrogen atoms except for Si–H were placed using AFIX instructions, while all other atoms were refined anisotropically. Deposition numbers CCDC-898201 (2), 898202 (*trans*-6), 898205 (*trans*-7·H<sub>2</sub>O), 898204 (*cis*-7), and 898203 (*trans*-8·2.25CHCl<sub>3</sub>). Free copies of the data can be obtained via http://www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK; Fax: +44 1223 336033; e-mail: deposit@ccdc.cam.ac.uk).

## ACKNOWLEDGEMENTS

This work was supported by Grants-in-Aid for Creative Scientific Research (No. 17GS0207), Scientific Research (B) (No. 22350017), Scientific Research on Innovative Areas (No. 23108707, " $\pi$ -Space"), Young Scientist (B) (No. 21750042), and the Global COE Program (B09, "Integrated Materials Science") from the Ministry of Education, Culture, Sports, Science and Technology, of Japan. Y.M. thanks Mitsubishi Chemical Corp., the Society of Synthetic Organic Chemistry, Japan, and Kinki Invention Center, for financial support.

### **REFERENCES AND NOTES**

- M. Haaf, T. A. Schmedake, and R. West, Acc. Chem. Res., 2000, 33, 704; V. Y. Lee, A. Sekiguchi, M. Ichinohe, and N. Fukaya, J. Organomet. Chem., 2000, 611, 228; A. Sekiguchi and V. Y. Lee, Chem. Rev., 2003, 103, 1429; V. Y. Lee and A. Sekiguchi, Angew. Chem. Int. Ed., 2007, 46, 6596; V. Y. Lee and A. Sekiguchi, Organometallic Compounds of Low-Coordinate Si, Ge, Sn and Pb: From Phantom Species to Stable Compounds; Wiley, 2010; Chapter 6.
- 2. N. Tokitoh, Acc. Chem. Res., 2004, 37, 86.
- A. Sekiguchi, M. Tsukamoto, and M. Ichinohe, *Science*, 1997, **275**, 60; S. B. Choi, P. Boudjouk, and P. R. Wei, *J. Am. Chem. Soc.*, 1998, **120**, 5814; T. Nishinaga, Y. Izukawa, and K. Komatsu, *J. Am. Chem. Soc.*, 2000, **122**, 9312; M. Saito, R. Haga, and M. Yoshioka, *Chem. Commun.*, 2002, 1002; M. Saito, R. Haga, and M. Yoshioka, *Chem. Lett.*, 2003, **32**, 912; M. Saito, R. Haga, M. Yoshioka, K. Ishimura, and S. Nagase, *Angew. Chem. Int. Ed.*, 2005, **44**, 6553; M. Ichinohe, M. Igarashi, K. Sanuki, and A. Sekiguchi, *J. Am. Chem. Soc.*, 2005, **127**, 9978; V. Y. Lee, R. Kato, M. Ichinohe, and A. Sekiguchi, *J. Am. Chem. Soc.*, 2005, **127**, 13142; S. Ishida, T. Nishinaga, R. West, and K. Komatsu, *Chem. Commun.*, 2005, 778; M. Driess, S. Yao, M. Brym, and V.-C. Wullen, *Angew. Chem. Int. Ed.*, 2006, **45**, 6730; R. Kinjo, M. Ichinohe, A. Sekiguchi, N. Takagi, M. Sumimoto, and S. Nagase, *J. Am. Chem. Soc.*, 2007, **129**, 7766; M. Igarashi, M. Ichinohe, and A. Sekiguchi, *J. Am. Chem. Soc.*, 2007, **129**, 12660; M. Saito, M. Sakaguchi, T. Tajima, K. Ishimura, S. Nagase, and M. Hada, *Science*, 2010, **328**, 339.
- K. Wakita, N. Tokitoh, R. Okazaki, and S. Nagase, *Angew. Chem. Int. Ed.*, 2000, **39**, 634; K. Wakita, N. Tokitoh, R. Okazaki, N. Takagi, and S. Nagase, *J. Am. Chem. Soc.*, 2000, **122**, 5648; A. Shinohara, N. Takeda, T. Sasamori, T. Matsumoto, and N. Tokitoh, *Organometallics*, 2005, **24**, 6141; N. Tokitoh, K. Wakita, T. Matsumoto, T. Sasamori, R. Okazaki, N. Takagi, M. Kimura, and S. Nagase, *J. Chin. Chem. Soc.*, 2008, **55**, 487.
- N. Nakata, N. Takeda, and N. Tokitoh, J. Am. Chem. Soc., 2002, 124, 6914; N. Nakata, N. Takeda, and N. Tokitoh, J. Organomet. Chem., 2003, 672, 66; N. Nakata, N. Takeda, and N. Tokitoh, Angew. Chem. Int. Ed., 2003, 42, 115.
- 6. Y. Mizuhata, N. Noda, and N. Tokitoh, Organometallics, 2010, 29, 4781.
- N. Tokitoh, K. Wakita, R. Okazaki, S. Nagase, P. v. R. Schleyer, and H. Jiao, J. Am. Chem. Soc., 1997, 119, 6951; K. Wakita, N. Tokitoh, R. Okazaki, S. Nagase, P. v. R. Schleyer, and H. Jiao, J. Am. Chem. Soc., 1999, 121, 11336; K. Wakita, N. Tokitoh, and R. Okazaki, Bull. Chem. Soc. Jpn., 2000, 73, 2157.
- N. Takeda, A. Shinohara, and N. Tokitoh, *Organometallics*, 2002, 21, 4024; A. Shinohara, N. Takeda, T. Sasamori, and N. Tokitoh, *Bull. Chem. Soc. Jpn.*, 2005, 78, 977.

- N. Nakata, N. Takeda, and N. Tokitoh, *Organometallics*, 2001, 20, 5507; N. Nakata, N. Takeda, and N. Tokitoh, *Chem. Lett.*, 2002, 818; N. Nakata, N. Takeda, and N. Tokitoh, *Organometallics*, 2003, 22, 481.
- Y. Mizuhata, T. Sasamori, N. Takeda, and N. Tokitoh, J. Am. Chem. Soc., 2006, 128, 1050; Y. Mizuhata, T. Sasamori, N. Nagahora, Y. Watanabe, Y. Furukawa, and N. Tokitoh, Dalton Trans., 2008, 4409.
- 11. N. Takeda, A. Shinohara, and N. Tokitoh, *Organometallics*, 2002, **21**, 256; A. Shinohara, N. Takeda, and N. Tokitoh, *J. Am. Chem. Soc.*, 2003, **125**, 10804.
- T. Sasamori, K. Inamura, W. Hoshino, N. Nakata, Y. Mizuhata, Y. Watanabe, Y. Furukawa, and N. Tokitoh, *Organometallics*, 2006, 25, 3533.
- 13. N. Tokitoh, A. Shinohara, T. Matsumoto, T. Sasamori, N. Takeda, and Y. Furukawa, *Organometallics*, 2007, **26**, 4048.
- 14. Y. Mizuhata, N. Takeda, T. Sasamori, and N. Tokitoh, Chem. Lett., 2005, 128, 1088.
- a) Y. Ren and T. Baumgartner, *Chem. Asian J.*, 2010, **5**, 1918; b) Y. Ren, T. Linder, and T. Baumgartner, *Can. J. Chem.*, 2009, **87**, 1222; c) H. Tsuji, S. Komatsu, Y. Kanda, T. Umehara, T. Saeki, and K. Tamao, *Chem. Lett.*, 2006, **35**, 758; d) N. K. Skvortsov and S. V. Toldov, *Russ. J. General Chem.*, 1994, **64**, 1583; e) A. Ishii, T. Tsuchiya, J. Nakayama, and M. Hoshino, *Tetrahedron Lett.*, 1993, **34**, 2347.
- 16. G. Märkl, H. Baier, and R. Liebl, Synthesis, 1977, 842.
- a) A. J. Ashe, T. R. Diephouse, and M. Y. El-Sheikh, J. Am. Chem. Soc., 1982, 104, 5693; b) G.
  Märkl and F. Kneidl, Angew. Chem., 1974, 86, 745; c) A. J. Ashe and M. D. Gordon, J. Am. Chem.
  Soc., 1972, 94, 7596; d) A. J. Ashe, J. Am. Chem. Soc., 1971, 93, 6690.
- G. Märkl, H. Baier, R. Liebl, and D. S. Stephenson, *Liebigs Ann. Chem.*, 1981, 870; G. Märkl, H. Baier, and R. Liebl, *Liebigs Ann. Chem.*, 1981, 919.
- A. B. Pangborn, M. A. Giardello, R. H. Grubbs, R. K. Rosen, and F. J. Timmers, *Organometallics*, 1996, 15, 1518.
- Gaussian 09, Revision A.02, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli,

1635

J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2009.

21. G. M. Sheldrick, Acta Cryst., 2008, A64, 112.