



Pergamon

Tetrahedron Letters 41 (2000) 2425–2428

TETRAHEDRON
LETTERS

Indium compound-catalyzed deoxygenative allylation of aromatic ketones by a hydrosilane–allylsilane system

Makoto Yasuda, Yoshiyuki Onishi, Takeshi Ito and Akio Baba*

Department of Applied Chemistry, Faculty of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

Received 5 January 2000; revised 20 January 2000; accepted 21 January 2000


Abstract

A combination of chlorodimethylsilane and allyltrimethylsilane effectively promoted the deoxygenative allylation of aromatic ketones in the presence of a catalytic amount of an indium compound to give the terminal alkenes. Indium trihalide or metallic indium showed high catalytic activity. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: indium and compounds; silicon and compounds; allylation; deoxygenation; reduction.

Synthetic applications of metallic indium have attracted much attention during the last decade¹ due to its low ionic potential² and stability under aqueous conditions.³ The synthetic potential of indium(III) halides is also increasing.⁴ However, they are used in an analogous manner to that of conventional Lewis acids such as aluminum halides and boron halides. There are few synthetic applications peculiar to indium compounds. We have recently reported the InCl_3 -catalyzed reductive Friedel–Crafts alkylation^{5,6} where InCl_3 promotes the hydrosilylation of carbonyl compounds with chlorodimethylsilane and the successive deoxygenation led to the carbocationic intermediate which reacts with aromatics. When an excess amount of chlorodimethylsilane was added, the cation was reduced to the corresponding hydrocarbons as previously reported.⁷ If the resulting carbocation could be trapped by carbon nucleophiles other than aromatics, a novel protocol of deoxygenative functionalization of ketones could be developed.

In this communication, we report a case of trapping the cation by an allyl group, where indium trichloride characteristically catalyzed the deoxygenative allylation of ketones by using allylsilanes and chlorodimethylsilane.



At first, we examined the catalysts in the reaction of acetophenone, chlorodimethylsilane and allyltrimethylsilane (Table 1). Indium trichloride effectively promoted the deoxygenative allylation to furnish

* Corresponding author.

4-phenyl-1-pentene (**1**) in 86% yield for 2 h (entry 1). It is interesting that even metallic indium gave a moderate yield (41%, entry 4) which could be improved up to 62% under a longer reaction time (entry 5). In contrast, neither AlCl_3 , a representative Friedel–Crafts catalyst, nor $\text{Sc}(\text{OTf})_3$, a characteristic promoter for reductive Friedel–Crafts reaction,^{6c–e} gave the desired allylated product (entries 6 and 7). Of all the Lewis acids examined, ZnCl_2 , $\text{BF}_3 \cdot \text{OEt}_2$, and CuCl_2 did not show any catalytic effect at all under similar conditions. These results indicate that indium trichloride is an indispensable catalyst for this type of allylation.




Table 1

Entry	Catalyst	Solvent	Yield/ % ^a	Entry	Catalyst	Solvent	Yield/ % ^a
1	InCl_3	CH_2Cl_2	86	8	InCl_3	$\text{ClCH}_2\text{CH}_2\text{Cl}$	63
2	InBr_3	CH_2Cl_2	60	9	InCl_3	cyclohexane	0
3	InI_3	CH_2Cl_2	85	10	InCl_3	DMF	0
4	In	CH_2Cl_2	41	11	InCl_3	THF	0
5 ^b	In	CH_2Cl_2	62	12	InCl_3	CH_3CN	35
6	AlCl_3	CH_2Cl_2	1	13	InCl_3	CHCl_3	16
7	$\text{Sc}(\text{OTf})_3$	CH_2Cl_2	< 0.2	14	InCl_3	CCl_4	0

^a Yields were determined by GLC. ^b The reaction was carried out for 16 hours.

The choice of solvent was also found to be definitely significant in this deoxygenative allylation as shown in Table 1 (entries 1 and 8–14). The reaction with acetophenone proceeded only when dichloromethane or 1,2-dichloroethane was used as a solvent. In contrast, the use of cyclohexane, DMF or THF completely suppressed the reaction, where the starting ketone was recovered (entries 9–11). Even other haloalkanes such as carbon tetrachloride and chloroform were unsuitable solvents (entries 13 and 14).




Table 2


Entry	Ar	R	Conditions	Product	Yield/ % ^a
1	Ph	Et	rt, 2 h	2	67
2	Ph	Ph	rt, 2 h	3	99
3	Ph	$\text{CH}_2\text{CH}_2\text{Cl}$	rt, 1 h → 60 °C, 3 h	4	70
4	<i>p</i> -Cl-C ₆ H ₄	Me	rt, 2 h	5	84
5	<i>p</i> -EtOCO-C ₆ H ₄	Me	rt, 1 h → 60 °C, 3 h	6	44
6	<i>p</i> -O ₂ N-C ₆ H ₄	Ph	rt, 1 h → 60 °C, 2 h	7	75
7	PhCH_2CH_2	Me	rt, 2 h	complicated mixture	
8	Ph	H	rt, 2 h	complicated mixture	

^a Yields were determined by GLC.

We then explored the generality of the InCl_3 -catalyzed deoxygenative allylation of aromatic ketones and the results are summarized in Table 2. The allylation of various types of aromatic ketones proceeded in moderate to excellent yields. Benzophenone, particularly, gave a quantitative yield at ambient temperature. The functionalities such as nitro, ester or chlorine were tolerated under these reducing conditions even at high temperature to give the corresponding olefins selectively (entries 3–6). The reaction with alkylketone gave a complicated mixture including a trace amount of a deoxygenative allylation product (entry 7). A harshly exothermic reaction took place with the use of benzaldehyde to give a complicated mixture (entry 8), in which side reactions such as direct allylation of the aldehyde, hydrosilylation and reductive Friedel–Crafts alkylation would simultaneously occur.⁵

A general procedure is as follows: acetophenone (2.0 mmol) was added to a mixture of indium trichloride (0.1 mmol), chlorodimethylsilane (2.2 mmol) and allyltrimethylsilane (2.2 mmol) in dichloromethane (10 mL) at ambient temperature. The reaction mixture was stirred for 2 h, quenched by water (20 mL), and extracted with ether (20 mL \times 3). After the organic layer was dried over MgSO_4 and concentrated in vacuo, the residue was purified by chromatography (eluted by hexane) on silica gel to give the product **1**.

We propose a plausible reaction path as illustrated in Scheme 1. At first, Lewis acid-promoted hydrosilylation of ketone takes place to form a silyl ether **8**. Then desiloxylation is accelerated by InCl_3 (via **8a** or **8b**) to generate the cation **9** (or a cationic species),⁸ which is readily trapped by allyltrimethylsilane. It is pointed out that the reaction (Table 1, entry 1) was accompanied with only a small amount (8%) of ethylbenzene formed by the trapping of the cation **9** with chlorodimethylsilane instead of allyltrimethylsilane. The stability of the cation intermediate should be an important factor, because *p*-nitrobenzophenone, which is facily susceptible to hydrosilylation showed more sluggish reaction than benzophenone. This hypothesis is also supported by the fact that aromatic ketones gave higher yields than alkylketones or aldehydes. The diphenylmethyl cation derived from benzophenone, in particular, is very stable to accomplish this reaction course.



Scheme 1. Plausible reaction paths of the deoxygenative allylation

Another possible path in which the allylation is followed by deoxygenative reduction (*path 2*) can be considered. However, no direct allylation of ketones with allyltrimethylsilane was observed in the presence of an InCl_3 catalyst. In addition, no formation of double allylated compounds, which should be facily formed by the second allylation,⁹ was observed. These results rule out *path 2*.

In conclusion, a novel deoxygenative allylation of aromatic ketones, namely, a selective transfunctionalization from carbonyl oxygen into hydrogen and an allyl group, was achieved by using allyltrimethylsilane and chlorodimethylsilane in the presence of a catalytic amount of an indium catalyst. The

disclosure of the detailed role of indium trichloride and further extension of this method are now in progress.

Acknowledgements

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture, Japan.

References

1. (a) Chao, L.-C.; Rieke, R. D. *J. Org. Chem.* **1975**, *40*, 2253–2255. (b) Araki, S.; Butsugan, Y. *J. Org. Chem.* **1988**, *53*, 1831–1833. (c) Cintas, P. *Synlett* **1995**, 1087–1096 and references cited therein.
2. Metal indium mediated S.E.T.-type pinacol coupling reaction and reduction: (a) Kalyanam, N.; Rao, G. V. *Tetrahedron Lett.* **1993**, *34*, 1647. (b) Lim, H. J.; Keum, G.; Kang, S. B.; Chung, B. Y.; Kim, Y. *Tetrahedron Lett.* **1998**, *39*, 4367–4368. (c) Moody, C. J.; Pitts, M. R. *Synlett* **1998**, 1028. (d) Moody, C. J.; Pitts, M. R. *Synlett* **1998**, 1029.
3. Li, C.-J.; Chan, T.-H. *Tetrahedron* **1999**, *55*, 11149–11176 and references cited therein.
4. For the recent reports of InX_3 -catalyzed reaction: (a) Loh, T.-P.; Xu, K.-C.; Ho, D. S.-C.; Sim, K.-Y. *Synlett* **1998**, 369–370. (b) Kobayashi, S.; Busujima, T.; Nagayama, S. *Tetrahedron Lett.* **1998**, *39*, 1579–1582. (c) Babu, G.; Perumal, P. T. *Tetrahedron* **1998**, *54*, 1627–1638. (d) Ranu, B. C.; Hajra, A.; Jana, U. *Org. Lett.* **1999**, *1*, 1141–1143.
5. (a) Miyai, T.; Onishi, Y.; Baba, A. *Tetrahedron Lett.* **1998**, *39*, 6291–6294. (b) Miyai, T.; Onishi, Y.; Baba, A. *Tetrahedron* **1999**, *55*, 1017–1026.
6. For reductive Friedel–Crafts alkylation of aromatics: (a) Hashimoto, Y.; Hirata, K.; Kihara, N.; Hasegawa, M.; Saigo, K. *Tetrahedron Lett.* **1992**, *33*, 6351–6354. (b) Hashimoto, Y.; Hirata, K.; Kagoshima, H.; Kihara, N.; Saigo, K. *Tetrahedron* **1993**, *49*, 5969–5978. (c) Tsuchimoto, T.; Hiyama, T.; Fukuzawa, S. *Chem. Commun.* **1996**, 2345–2346. (d) Fukuzawa, S.; Tsuchimoto, T.; Hiyama, T. *J. Org. Chem.* **1997**, *62*, 151–156. (e) Tsuchimoto, T.; Tobita, K.; Hiyama, T.; Fukuzawa, S. *J. Org. Chem.* **1997**, *62*, 6997–7005.
7. Miyai, T.; Ueba, M.; Baba, A. *Synlett* **1999**, 182–184.
8. Mukaiyama et al. reported the InCl_3 - Me_3SiCl -catalyzed desiloxylation from *O*-trimethylsilyl monothioacetal, in which indium trichloride acts as a unique promoter: (a) Mukaiyama, T.; Ohno, T.; Nishimura, T.; Han, J. S.; Kobayashi, S. *Chem. Lett.* **1990**, 2239–2242. (b) Mukaiyama, T.; Ohno, T.; Nishimura, T.; Han, J. S.; Kobayashi, S. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 2524–2527.
9. (a) Ishii, A.; Kotera, O.; Saeki, T.; Mikami, K. *Synlett* **1997**, 1145–1146. (b) Aggarwal, V. K.; Vennall, G. P. *Synthesis* **1998**, 1822–1826.