FULL PAPER

Dipalladium complexes with bridging diorganosilyl ligands. Synthesis, structure, and properties of $[LPd(\mu-SiH(R)Ph)]_2$ (R = Ph or Me; L = PMe₃, PEt₃ or PMePh₂)

Yong-Joo Kim,*^a Sang-Chul Lee,^a Jong-Il Park,^a Kohtaro Osakada,*^b Jun-Chul Choi^b and Takakazu Yamamoto^b

- ^a Department of Chemistry, Kangnung National University, Kangnung 210-702, Korea. E-mail: yjkim@knusun.kangnung.ac.kr
- ^b Research Laboratory of Resources Utilization, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan

Received 27th September 1999, Accepted 30th November 1999

The complexes *trans*-PdEt₂L₂ (L = PMe₃, PEt₃ or PMePh₂) reacted with equimolar or twofold molar amounts of H₂SiPh₂ to produce various bimetallic palladium(II) complexes with bridging diphenylsilyl ligands, [(Me₃P)-Pd(μ -SiHPh₂)]₂ 1, (Me₃P)Pd(μ -SiHPh₂)Pd(PMe₃)₂ 2 and [LPd(μ -SiHPh₂)]₂ (L = PEt₃ 3 or PMePh₂ 4), in moderate to good yields. Similar reactions using H₂SiMePh afforded the complexes [LPd(μ -SiHMePh)]₂ (L = PMe₃ 5 or PEt₃ 6). The isolated complexes were characterized by NMR spectroscopy and elemental analyses. Molecular structures of 3 and 5 have been determined by X-ray crystallography and suggest an agostic interaction between the μ -SiHPh₂ or μ -SiHMePh group and the Pd. The NMR study of the complexes revealed the bridging co-ordination of the silyl ligand with the Si–H–Pd three-center, two-electron bond in solution similar to that in the solid state. Complex 2 undergoes exchange of the PMe₃ ligands on the NMR timescale. The reactions of 2 with CO and HSPh afforded 1 and Pd(SPh)₂(PMe₃)₂, respectively.

Bridging co-ordination of SiHR₂ and SiH₂R groups to two or more transition metal centers is common to many transition metals and, in most cases, exhibits an agostic interaction between the SiH group and the metal center, as schematically depicted in (A).¹ The reaction of diorganosilanes with low valent transition metal complexes often gives (A) or diorganosilylene co-ordinated dinuclear metal complexes (B) that are formed *via* the metal promoted Si–H bond activation of (A).



The chemistry of silylpalladium complexes² has received particular attention in relation to the mechanism of palladiumcatalysed hydrosilylation and bis-silylation of alkenes and alkynes as well as related reactions of organosilanes^{3,4} since they probably involve intermediate palladium complexes with organosilyl ligands. Reports on palladium complexes with organosilyl ligands are fewer in number than those on silylplatinum complexes. In particular, there have been few reports on the dinuclear palladium complexes with bridging diorganosilyl (SiHR₂) ligands, whereas analogous dinuclear platinum complexes have been reported using many phosphine ligands.^{5,6} Studies on the structure and chemical properties of palladium complexes with bridging diorganosilyl ligands are intriguing because the complexes appear to be less stable and more readily reactive than the reported platinum analogues.

In this paper we report the preparation of several dinuclear palladium complexes with bridging diorganosilyl ligands from the reaction of diorganosilane with diethylpalladium complexes as well as results of NMR and X-ray studies of the complexes. Part of this work has been reported preliminarily.⁷

Results and discussion

Preparation of diorganosilyl bridged dinuclear palladium complexes

The reaction of *trans*-PdEt₂(PMe₃)₂ with equimolar H₂SiPh₂ in THF or hexane at 55 °C proceeds smoothly to give a mixture of dinuclear palladium complexes $[(Me_3P)Pd(\mu-SiHPh_2)]_2$ 1 and $(Me_3P)Pd(\mu-SiHPh_2)_2Pd(PMe_3)_2$ 2, eqn. (1). The two products



are isolated by fractional crystallization. The reaction in a 1:2 molar ratio also led to the formation of complexes. The complexes *trans*-PdEt₂L₂ (L = PEt₃ or PMePh₂) react with H₂SiPh₂ and with H₂SiMePh in a similar way to give the dinuclear complexes [LPd(μ -SiHPh₂)]₂ (L = PEt₃ **3** 60%; PMePh₂ **4** 52%), eqn. (2), and [LPd(μ -SiHMePh)]₂ (L = PMe₃ **5** 39%; PEt₃ **6** 30%), eqn. (3), respectively. Products of similar reactions using H₃SiPh and H₂SiEt₂ did not contain these dinuclear palladium complexes due to the decomposition of initially formed palladium complexes during the reaction.

Complexes 1-4 with bridging diphenylsilyl ligands are thermally more stable than the methyl(phenyl)silyl bridged com-



plexes 5 and 6 which undergo decomposition gradually at room temperature in solution. Reactions (2) and (3) gave the symmetric dinuclear complexes 3-6 as the sole isolated products, whereas the product of the reaction of PdEt₂(PMe₃)₂ with H_2SiPh_2 contains 2 with three PMe₃ ligands as the major product. The yield of the symmetric product 1 with a structure similar to 3-6 is much lower. The compact PMe₃ ligand tends to form unsymmetric dinuclear palladium(I) complexes by coordination of the two phosphine ligands to one Pd, as already noted in the structural study of (Me₃P)Pd(µ-CH₂CHCH₂)-(µ-SPh)Pd(PMe₃)₂.⁸ The more bulky PEt₃ and PMePh₂ ligands do not form the unsymmetric product with a sterically congested palladium center, as shown in reaction (2). The methyl-(phenyl)silyl ligand which is more electron donating than the diphenylsilyl ligand does not allow co-ordination of two PMe₃ ligands to Pd and gives the symmetric product 5 exclusively.

The above reactions involve activation of Pd–C and Si–H bonds, possibly accompanied by the evolution of ethylene and ethane. Since *trans*-PdEt₂(PR₃)₂ undergoes β -hydrogen elimination and ensuing reductive elimination of ethane to form Pd(CH₂=CH₂)(PR₃)₂,⁹ one of the plausible paths for the formation of complexes **1–6** involves initial formation of the ethylene-co-ordinated palladium(0) complexes and their further reaction with diorganosilane to give Pd(SiHR'₂)₂(PR₃)₂ or PdH(SiHR'₂)(PR₃)₂, which then undergoes intermolecular bridging of the metal center through diorganosilyl ligands. To evaluate this possibility, we have examined the reaction of equimolar and twofold molar amounts of H₂SiPh₂ or H₂SiMePh with Pd(CH₂=CHPh)L₂ (L = PMe₃, PEt₃ or PMePh₂) which has a similar structure to that of Pd(CH₂=CH₂)(PMe₃)₂, eqn. (4). The reactions at 55 °C in THF produce

$$\begin{array}{c} Ph, Ph \\ H^{-}Si \\ H^{-}Pd \\ L \\ H^{-}Ph \end{array} + H_{2}SiPh_{2} \longrightarrow \begin{array}{c} Ph, Ph \\ H^{-}Si \\ L \\ H^{-}Pd \\ H^{-}Ph \end{array} + H_{2}SiPh_{2} \\ L \\ H^{-}Pd \\ Ph \end{array}$$

complexes 1–6 in moderate to good yields. In the presence of a twofold molar amount or more of H_2SiPh_2 the reaction proceeds to completion to give the products in high yields. Previously, Schubert and Müller¹⁰ reported the reaction of

Previously, Schubert and Muller¹⁰ reported the reaction of H_2SiPh_2 with $PdMe_2(PMePh_2)_2$ in a 2:1 molar ratio at room temperature to give *cis*-Pd(SiHPh_2)_2(PMePh_2)_2 which is thermally unstable but has been characterized spectroscopically. Similar bis(organosilyl)palladium complexes have been isolated using bulky or chelating diphosphine ligands.¹¹ These results suggest that mononuclear bis(diorganosilyl)palladium complexes can be formed *via* the reaction of diorganosilane with dialkylpalladium complexes. Scheme 1 depicts possible reaction paths for the formation of dinuclear palladium complexes with bridging diorganosilyl ligands assumed on the basis of the above mentioned results. Coupling of bis(diorganosilyl)palladium(II) complexes with palladium(0) complexes leads to



Fig. 1 An ORTEP¹² drawing of $[(Et_3P)Pd(\mu-SiHPh_2)]_2$ 3 showing the atom-labeling scheme and 30% probability thermal ellipsoids. Atoms with asterisks are crystallographically equivalent to those with the same number without asterisks. Selected bond lengths (Å) and angles (°): Pd1–Pd1* 2.699(2), Pd1–Si1 2.332(2), Pd1–Si1* 2.396(2), Pd1–P1 2.301(2), Pd1–H26 2.04 and Si1–H26 1.60; Pd1–Pd1*–P1 159.51(5), Pd1–Si1–Pd1* 69.60(5), Si1–Pd1–Si1 10.40(5), Pd1–Pd1*–Si1 56.32(6), Pd1=Si1 54.08(4), Pd1–Si1–H26 57, Pd1*–Si1–H26 126, Pd1–Si1–C7 120.6(2), Pd1*–Si1–C7 116.1(2), Pd1–Si1–C1 115.7(2), Pd1*–Si1–C1 121.5(2) and C1–Si1–C7 108.5(2).



the formation of dinuclear product (i), while an alternative path involving dimerization of hydrido(silyl)palladium(II) complexes also accounts for the product. At present we cannot compare the plausibilities of the reaction paths based on the experimental results.

X-Ray crystallography and NMR spectra

Figs. 1 and 2 show the molecular structures of complexes 3 and 5 determined by X-ray crystallography. Molecules of both contain a crystallographic center of symmetry at the midpoint of the Pd-Pd bond. The structural parameters are compared with those of the preliminarily reported 1^7 and with the platinum analogue $[(R'_3P)Pt(\mu-HSiR_2)]_2^{.5}$ The Pd–Pd distances are 2.699(2) and 2.707(1) Å, suggesting the presence of a metalmetal bond. The lengths of the two Pd-Si bonds in 3 and 5 differ significantly (2.396(2) and 2.332(2) Å, and 2.387(1) and 2.333(2) Å, respectively) as they do in the structure of 1 (2.328(2) and 2.386(2) Å). The SiH hydrogens are located in the final electron density map and are included in the plane consisting of two Pd and two Si. These structural features suggest the contribution of an Si-H-Pd three-center two-electron bond to the co-ordination of diphenylsilyl or methyl(phenyl)silyl ligands to two Pd, which causes elongation of the Pd-Si bond compared with the Pd–Si single bond. The Pd–H bond of 3 (2.04 Å) and 5 (1.85 Å) suggests an interaction of the Si-H hydrogen



Fig. 2 An ORTEP drawing of $[(Me_3P)Pd(\mu-SiHMePh)]_2$ 5 showing the atom-labeling scheme and 30% probability thermal ellipsoids. Atoms with asterisks are crystallographically equivalent to those with the same number without asterisks. Selected bond lengths (Å) and angles (°): Pd1–Pd1* 2.707(1), Pd1–Si1 2.387(1), Pd1–Si1* 2.333(2), Pd1–P1 2.297(2), Pd1–H18 1.85 and Si1–H18 1.60; Pd1–Pd1*P=1 155.18(4), Pd1–Si1–Pd1* 69.98(4), Si1–Pd1–Si1* 110.02(4), Pd1–Pd1*– Si1 54.07(4), Pd1*–Pd1–Si1 55.95(4), Pd1–Si1–H18 120.7, Pd1*–Si1– H18 50.7, Pd1–Si1–C4 126.2(2), Pd1*–Si1–C4 118.3(2), Pd1–Si1–C5 112.1(2), Pd1*–Si1–C5 118.5(2) and C4–Si1–C5 107.0(3).

with the Pd, although these bonds are somewhat larger than those in **1** due to insufficient precision in the determination of the position of hydrogen in the crystallography. A platinum analog, $[(R'_3P)Pt(\mu-HSiR_2)]_2$, has a similar structure to that of **1** but with a larger difference between the Pt–Si bond lengths (2.420(2) and 2.324(2) Å) and a short H–Pt distance (1.78 Å).⁵

Complexes 1 and 3–6 exhibit NMR signals from the Si–H hydrogen and phosphorus nuclei at similar positions. The ³¹P-{¹H} NMR spectrum of 3 at room temperature contains a single resonance at δ 15.3. The presence of ²⁹Si satellite signals with two different coupling constants ($J(^{31}P^{29}Si) = 69$ and 27 Hz) is consistent with the unsymmetrical co-ordination of two Si atoms to Pd. The ¹H NMR spectrum at 25 °C exhibits a signal due to the SiH hydrogen at δ 1.76 flanked with satellite signals ($J(^{1}H^{29}Si) = 78$ Hz). This coupling constant is smaller than that of $J(^{1}H^{29}Si)$ in organic molecules and suggests partial activation of the Si–H bond by Pd to form an Si–H–Pd three-center two-electron bond. Complex 3 does not show any significant changes in NMR spectra in the temperature range from 25 to -90 °C and has a structure in solution similar to that found in the crystals.

The NMR spectra of unsymmetric complex 2 indicate fluxional behavior arising from exchange of the PMe₃ ligands. The ³¹P-{¹H} NMR spectrum at -90 °C contains a doublet $(\delta - 33.0, J(^{31}P^{31}P) = 23 \text{ Hz}, J(^{31}P^{29}\text{Si}) = 42 \text{ Hz})$ and a triplet $(\delta - 25.9, J(^{31}P^{31}P) = 23 \text{ Hz}, J(^{31}P^{29}\text{Si}) = 129 \text{ Hz})$ in a 2:1 peak area ratio. The ¹H NMR spectrum at -90 °C exhibits a doublet $(\delta 1.24, ^{3}J(^{31}P^{1}H) = 8, J(^{29}\text{Si}^{1}H) = 84 \text{ Hz})$ and a triplet $(\delta - 2.34, ^{3}J(^{31}P^{1}H) = 11, J(^{29}\text{Si}^{1}H) = 86 \text{ Hz})$ of equal intensity, indicating that the two SiH hydrogens are magnetically inequivalent at that temperature. These ¹H and ³¹P NMR spectra show coalescence of these signals when the temperature is raised and exhibit a single resonance for the phosporous nuclei of the PMe₃ ligands and for the SiH hydrogens at 25 °C. The changes in the NMR spectra are brought about by intra- or intermolecular exchange of the co-ordinated PMe₃.

Chemical properties of the complexes

In connection with gaining insight into the chemical properties of the complexes and their relationship to several synthetic organic reactions catalysed by palladium complexes, we have examined reactions of phenylacetylene and diphenylacetylene with **2**. The reactions of **2** with an equimolar or twofold molar amount of these acetylenes did not occur and resulted in stoichiometric recovery of the starting complex. The reaction of **2** with CO (1 atm) liberates PMe₃ ligand and affords **1** in 79% isolated yield, eqn. (5), but does not give any other products



from further carbonylation of the complex. Many organopalladium complexes with tertiary phosphine ligands were reported to react with CO to give multinuclear carbonyl palladium compounds, $Pd(CO)_m(PR_3)_n$, via elimination of the organic ligands. The inertness of the palladium complex toward carbonylation suggests the stable co-ordination of the bridging diphenylsilyl ligands to two Pd. Treatment of **2** with a twofold molar amount of HSPh in diethyl ether at room temperature results in *trans*-Pd(SPh)₂(PMe₃)₂ 7, eqn. (6), which was identi-

Ph Ph
H—Si PMe₃
Me₃P-Pd
$$\xrightarrow{}$$
 Pd $\xrightarrow{}$ Pd $\xrightarrow{}$ Pd $\xrightarrow{}$ Pd $\xrightarrow{}$ Pd $\xrightarrow{}$ Pd $\xrightarrow{}$ Pd (SPh)₂(PMe₃)₂ (6)
Si-H PMe₃
Ph Ph
2 7

fied by comparison of the NMR data with those of authentic complexes. Poor yields of the complex are partly due to insufficient amount of PMe₃ ligands to complete the above reaction.

In summary, this study has provided dinuclear palladium(I) complexes with bridging diorganosilyl ligands which have no precedent in organopalladium chemistry. The complexes are stabilized by two symmetrically bridging diorganosilyl ligands which form a σ bond with one Pd and are bonded to another Pd in a three-center two-electron manner. This dinuclear framework is stable once it is formed and does not react with carbon monoxide. The inert co-ordination of the diorganosilyl ligands contrasts with the bridging co-ordination of the diorganosilyl ligands in other late transition metal complexes which undergo facile and reversible Si–H bond activation by the metal center and exhibit fluxional NMR spectra at around room temperature.

Experimental

General, materials, and measurements

All manipulations of the air-sensitive compounds were performed under a N₂ or argon atmosphere with use of a standard Schlenk technique. Solvents were distilled from sodium– benzophenone and stored under a nitrogen atmosphere. The complexes *trans*-PdEt₂L₂ and *trans*-PdMe₂L₂ (L = PMe₃, PEt₃ or PMePh₂) were prepared according to the literature methods.⁹ H₂SiPh₂ and H₂SiMePh were obtained from Aldrich Co. and used without further purification.

Elemental analyses were carried out by analytical laboratories at the Basic Science Institute and at SangJi university in Korea. IR spectra were recorded on a Hitachi 270-30 spectrophotometer, NMR (¹H, ¹³C-{¹H} and ³¹P-{¹H}) spectra on Bruker 400 MHz and JEOL Lambda 300 MHz spectrometers. Chemical shifts were referenced to internal Me₄Si or to external 85% H₃PO₄.

Preparations

Complexes 1 and 2. To a Schlenk flask containing *trans*-PdEt₂(PMe₃)₂ (1.470 g, 4.64 mmol) at 0 °C were added H₂SiPh₂ (0.856 g, 4.64 mmol) and THF (7 cm³) in that order. The mixture was heated at 55 °C for 30 min to give a deep red solution. After the solution was stirred at room temperature for 1 h, the solvent was evaporated under reduced pressure to give an orange residue. The addition of hexane and ensuing storage of the residue for several hours at -30 °C produced an orange solid which was collected by filtration, washed with hexane, and recrystallized with THF-hexane to give complex 2 as red crystals (1.18 g, 63%). The filtrate was evaporated under reduced pressure to give a yellow solid from which 1 was obtained as pale yellow crystals after recrystallization from ether (0.053 g, 3%). Data for 1: mp 105 °C (decomp.) (Found: C, 49.08; H, 5.77. $C_{15}H_{20}PPdSi$ requires C, 49.25; H, 5.51%); δ_H (400 MHz, CD₂Cl₂ at 25 °C) 7.63 (8 H, m, C₆H₅), 7.31 (12 H, m, C₆H₅), 1.77 (2 H, s, SiH, $J(^{29}Si^{1}H) = 77$ Hz) and 1.19 (18 H, br, CH₃); (at 0 °C) 7.61 (8 H, m, C₆H₅), 7.33 (4 H, m, C₆H₅), 7.31 (8 H, m, C₆H₅), 1.76 (2 H, d, SiH, $J({}^{31}P^{1}H) = 7$, $J({}^{29}Si^{1}H) = 80$ Hz) and 1.17 (18 H, br, CH₃); (at -30 °C) 7.60 (8 H, m, C₆H₅), 7.31 (12 H, m, C₆H₅), 1.72 (2 H, d, SiH, $J({}^{31}P^{1}H) = 7$, $J({}^{29}Si^{1}H) = 77$ Hz) and 1.14 (18 H, br, CH₃); (at -70 °C) 7.56 (8 H, m, C₆H₅), 7.29 (12 H, m, C₆H₅), 1.66 (2 H, br, SiH, $J(^{29}Si^{1}H) = 82$ Hz) and 1.10 (18 H, br, CH₃); (at -90 °C) 7.53 (8 H, m, C₆H₅), 7.30 $(12 \text{ H}, \text{ m}, \text{C}_6\text{H}_5), 1.62 (2 \text{ H}, \text{ br}, \text{SiH}, J(^{29}\text{Si}^1\text{H}) = 80 \text{ Hz}) \text{ and } 1.08$ (18 H, br, CH₃); δ_P (160 MHz in CD₂Cl₂) (at 25 °C) –34.4 (br, $v_{1/2} = 43$ Hz); (at 0 °C) -26.5 (s, $J(^{29}Si^{31}P) = 74$ Hz); (at -30 °C) -25.8 (s, $J(^{29}Si^{31}P) = 74$ and 28 Hz); (at -70 °C) -25.1 (s, $J(^{29}Si^{31}P) = 74$ Hz); (at -90 °C) -24.7 (s, $J(^{29}Si^{31}P) = 74$ Hz). Data for 2: mp 109 °C (decomp.) (Found: C, 48.75, H, 6.27. $C_{33}H_{49}P_3Pd_2Si_2$ requires C, 49.08; H, 6.11%); δ_H (400 MHz, CD₂Cl₂) (at 25 °C) 7.63 (8 H, m, C₆H₅), 7.29 (12 H, m, C₆H₅), 1.00 (27 H, d, CH₃, J(PH) = 7) and 0.59 (2 H, br, SiH, $J({}^{29}Si^{1}H)$ = 79 Hz); (at 0 °C) 7.63 (8 H, m, C_6H_5), 7.28 (12 H, m, C_6H_5), 0.93 (27 H, s, CH₃) and 0.15 (2 H, br, SiH, $J(^{29}Si^{1}H) = 76$ Hz); (at -30 °C) 7.59 (8 H, m, C₆H₅), 7.29 (12 H, m, C₆H₅), 0.87 (27 H, s, CH₃) and -0.17 (2 H, br, SiH, $v_{1/2} = 53$ Hz); (at -70 °C) 7.55 (8 H, br, C₆H₅), 0.99 (10 H, br, CH₃ and SiH), 0.68 (18 H, br, CH₃) and -2.33 (1 H, br, SiH, $v_{1/2} = 84$ Hz); (at -90 °C) 7.53 (8 H, m, C₆H₅), 7.27 (12 H, m, C₆H₅), 1.24 (1 H, d, SiH, $J(^{31}P^{1}H) = 8$ Hz, $J(^{29}Si^{1}H) = 84$ Hz), 1.03 (9 H, s, CH₃), 0.63 $(18 \text{ H}, \text{ s}, \text{CH}_3), -2.34 (1 \text{ H}, \text{ t}, \text{SiH}, J(^{31}\text{P}^1\text{H}) = 11, J(^{29}\text{Si}^1\text{H}) = 86$ Hz); (160 MHz in CD₂Cl₂) (at 25 °C) -34.4 (br, $v_{1/2} = 43$ Hz); (at 0 °C) -33.3 (br, $v_{1/2} = 43$ Hz); (at -30 °C) -32.2 (br, $v_{1/2} =$ 102 Hz); (at -70 °C) -26.4 (br, $v_{1/2} = 98$) and -33.6 (br, $v_{1/2} = 81$ Hz); (at -90 °C) -25.9 (t, $J({}^{31}P{}^{31}P) = 23$, $J({}^{29}Si{}^{31}P) = 129$) and -33.0 (t, $J({}^{31}P{}^{31}P) = 23$, $J({}^{29}Si{}^{31}P) = 42$ Hz).

Complexes 3-6. To a Schlenk flask containing trans- $PdEt_2(PEt_3)_2 \ (0.171 \ g, \ 0.43 \ mmol)$ at 0 °C were added H_2SiPh_2 (0.079 g, 0.43 mmol) and THF (3 cm^3) in that order. The mixture was heated at 55 °C for 30 min to give a deep orange solution which was stirred for an additional 1 h at room temperature. The solvent was evaporated under reduced pressure to give an orange residue. The addition of hexane and ensuing storage of the product for several hours at -30 °C produced an orange solid which was collected by filtration, washed with hexane, and recrystallized from ether-hexane to give complex 3 as pale yellow crystals (0.105 g, 60%) (Found: C, 52.70; H, 6.74. C₁₈H₂₆PPdSi requires C, 53.01; H, 6.43%); δ_H (300 MHz, in CD₂Cl₂) 0.85 (18 H, br quintet due to virtual coupling, PCH₂CH₃), 1.50 (12 H, m, PCH₂CH₃), 1.76 (2 H, br d, Si-H, $J({}^{1}\mathrm{H}^{29}\mathrm{Si}) = 78$, $J(\mathrm{HP}) = 4.8$ Hz), 7.32, 7.67 (20 H, m, Ph); $\delta_{P}({}^{31}P-{}^{1}H)$ from external 85% H₃PO₄ at 120 MHz in CD₂Cl₂) 15.3 (s, $J({}^{31}P{}^{29}Si) = 69, 27$ Hz).

Complexes **4–6** were prepared analogously: **4** (52%) (Found: C, 61.61; H, 5.02. $C_{25}H_{24}PPdSi$ requires C, 61.29; H, 4.94); $\delta_{\rm H}$ (300 MHz, in CD₂Cl₂) 1.67 (6 H, s, PCH₃Ph₂), 2.15 (2 H, br s, SiH) and 7.17–7.49 (40 H, m, Ph); $\delta_{\rm P}(^{31}P\{-^{1}H\})$ from external 85% H₃PO₄ at 120 MHz in CD₂Cl₂) 7.89 (s); ($J(^{31}P^{29}Si)$ not obtained due to broading of signal; **5** (39%) (Found: C, 39.27; H, 5.66. $C_{10}H_{18}PPdSi$ requires C, 39.54; H, 5.97%); $\delta_{\rm H}$ (300

MHz, in CD₂Cl₂) 0.92 (6 H, br s, SiHCH₃Ph), 1.07 (2 H, br t, SiH, *J*(HP) = 2.0 Hz), 7.20, 7.44 (10 H, m, Ph); δ_{P} (external 85% H₃PO₄ standard at 120 MHz in CD₂Cl₂) -24.3 (s); **6** (30%) (Found: C, 45.37; H, 6.79. C₁₃H₂₄PPdSi requires C, 45.15; H, 7.00%); δ_{H} (300 MHz, in CD₂Cl₂) 0.80 (18 H, br quintet due to virtual coupling, PCH₂CH₃), 0.93 (6 H, br s, SiHCH₃Ph), 0.99 (2 H, br s, SiH), 1.42 (12 H, m, PCH₂CH₃), 7.13, 7.40 (10 H, m, Ph); $\delta_{P}(^{31}P-\{^{1}H\}$ from external 85% H₃PO₄ at 120 MHz in CD₂Cl₂) 16.7 (s, *J*(³¹P²⁹Si) = 72, 27 Hz).

Reaction of Pd(CH₂=CHPh)L₂ with H₂SiPh₂ or H₂SiMePh

To a Schlenk flask containing *trans*-PdEt₂(PMe₃)₂ (0.244 g, 0.77 mmol) at 0 °C were added styrene (0.322 g, 3.09 mmol) and THF (3 cm³) in that order. The mixture was heated at 55 °C for 3 h to give a pale yellow solution. The solution was cooled to room temperature and H₂SiPh₂ (0.142 g, 0.77 mmol) added. The initial red solution instantly turned dark red. After stirring for 3 h the solvent was removed under reduced pressure to give a dark red residue. The addition of hexane and ensuing storage of the product for several hours at -30 °C produced an orange solid which was collected by filtration and washed with hexane to give complex **2** (0.257 g, 82%).

Reactions of $Pd(CH_2=CHPh)L_2$ (L = PMe₃, PEt₃ or PMePh₂) with equimolar or twofold molar amounts of H₂SiPh₂ or H₂SiMePh were analogously carried out to give the corresponding complexes.

Reactions of complex 2

With alkyne. To an ether (5 cm^3) solution of complex 2 (0.106 g, 0.13 mmol) was added diphenylacetylene (0.024 g, 0.13 mmol) at room temperature. The initial red solution slowly turned dark red. After stirring for 4 h the solvent was removed under reduced pressure to give a dark red residue. The addition of hexane and ensuing storage of the product for several hours at -30 °C enabled the recovery of 2 (0.93 g, 87%). Reaction of 2 with a twofold molar amount of phenylacetylene proceeded analogously to give the starting material (92%).

With CO. Complex 2 (0.145 g, 0.18 mmol) was dissolved in toluene (3 cm³) at room temperature. After evacuation of the system, CO (1 atm) was introduced and the initial red solution turned dark red. After stirring for 1 h the solvent was evaporated under reduced pressure to give an oily residue. The addition of hexane (2 cm³) and ensuing storage of the residue for several hours at -30 °C produced an orange solid which was collected by filtration and washed with hexane (1 cm³ × 3) to give 1 (0.104 g, 79%).

With HSPh. To an ether (6 cm³) solution of complex 2 (0.125 g, 0.15 mmol) was added benzenethiol (0.034 g, 0.30 mmol) at room temperature. The initial red solution slowly turned orange accompanied by gradual formation of the solid product. After stirring for 1 h the reaction mixture was stored at -30 °C for several hours to cause complete precipitation of a yellow solid which was collected by filtration, washed with hexane (2 cm³ × 2) and dried *in vacuo* to give 7 (0.032 g, 43%). $\delta_{\rm H}$ (300 MHz, in CD₂Cl₂) 1.31 (18 H, s, P(CH₃)₃, J(PH) = 3.4 Hz), 6.80 (1 H, t, Ph), 7.19 (2 H, t, Ph) and 7.68 (2 H, d, Ph). $\delta_{\rm P}$ (³¹P-{¹H} from external 85% H₃PO₄ at 120 MHz in CD₂Cl₂) -13.1 (s). Complex 7 was also obtained from the reaction of *trans*-PdMe₂(PMe₃)₂ with an excess of benzenethiol and identified using the spectroscopic data.

Crystallography

Crystals of complexes **3** and **5** suitable for X-ray diffraction study were obtained by recrystallization from ether–hexane and mounted in glass capillary tubes under argon. Intensities were collected for Lorentz-polarization effects on a Rigaku AFC-5R automated four-circle diffractometer by using Mo-K α radiation ($\lambda = 0.71069$ Å) and the $\omega - 2\theta$ scan method, and an empirical absorption correction (ψ scan) was applied. Calculations were carried out by using the program package TEXSAN¹³ on a DEC Micro VAX-II computer. Atomic scattering factors were obtained from ref. 14. A full matrix least-squares refinement was used for non-hydrogen atoms with anisotropic thermal parameters. Positions of the SiH hydrogens were determined by the Fourier-difference technique, while the other hydrogens were located by assuming the ideal geometry. The hydrogens were included in the structure calculation without further refinement of the parameters.

Crystal data. $C_{36}H_{52}P_2Pd_2Si_2$ 3, M = 815.77, triclinic, space group $P\bar{1}$ (no. 2), a = 10.242(9), b = 11.447(6), c = 9.725(8) Å, $a = 112.13(5)^\circ$, $\beta = 115.65(6)^\circ$, $\gamma = 78.56(6)^\circ$, U = 951(1) Å³, T = 298 K, Z = 1, μ (Mo-K α) = 1.116 mm⁻¹, 3544 reflections measured, 3340 unique ($R_{int} = 0.012$), from which 2835 with $I > 3.00\sigma(I)$ were used in refinement. Final *R*1 and *wR*2 0.038 and 0.045.

 $C_{20}H_{36}P_2Pd_2Si_2$ **5**, M = 607.46, monoclinic, space group $P2_1/n$ (no. 14), a = 8.419(4), b = 16.222(4), c = 10.240(4) Å, $\beta = 108.58(3)^\circ$, U = 1325.7(9) Å³, T = 298 K, Z = 2, μ (Mo-K α) = 1.572 mm⁻¹, 3319 reflections measured, 3152 unique ($R_{int} = 0.042$), from which 2114 with $I > 3.00\sigma(I)$ were used in refinement. Final R1 and wR2 0.033 and 0.036.

CCDC reference number 186/1754.

See http://www.rsc.org/suppdata/dt/a9/a907745f/ for crystallographic files in cif. format.

Acknowledgements

This work was supported by the Basic Science Research Institute Program (No. BSRI-98-3440), Korean Ministry of Education.

References

- Recent reviews on transition metal silyl complexes: T. D. Tilley, in *The Silicon–Heteroatom Bond*, eds. S. Patai and Z. Rappoport, J. Wiley and Sons, New York, 1991; U. Schubert, *Transition Met. Chem.*, 1991, 16, 136; H. Tobita and H. Ogino, *Adv. Organomet. Chem.*, 1998, 42, 223; J. Y. Corey and J. Braddock-Wiking, *Chem. Rev.*, 1999, 99, 175.
- M. D. Curtis and J. Greene, J. Am. Chem. Soc., 1978, 100, 6362;
 T. R. Bierschenk, M. A. Guerra, T. J. Juhlke, S. B. Larson and R. J. Lagow, J. Am. Chem. Soc., 1987, 109, 4855; M. A. Guerra and R. J. Lagow, J. Chem. Soc., Chem. Commun., 1990, 65; Y. Pan, J. T. Mague and M. J. Fink, J. Am. Chem. Soc., 1993, 115, 3842;
 F. Ozawa, M. Sugawara and T. Hayashi, Organometallics, 1994, 13, 3237; S. Shimada, M. Tanaka and M. Shiro, Angew. Chem., Int. Ed. Engl., 1996, 35, 1856; Y. Tanaka, H. Yamashita, S. Shimada and M. Tanaka, Organometallics, 1997, 16, 3246; M. Murakami,

- Reviews: I. Ojima, in *The Chemistry of Organic Silicon Compounds*, eds. S. Patai and Z. Rappoport, John Wiley, New York, 1989, ch. 25; J. L. Speier, *Adv. Organomet. Chem.*, 1979, **17**, 407; P. Braunstein and M. Knorr, *J. Organomet. Chem.*, 1995, **500**, 21.
 H. Okinoshima, K. Yamamoto and M. Kumada, *J. Organomet.*
- H. Okinoshima, K. Yamamoto and M. Kumada, J. Organomet. Chem., 1975, 86, C27; K. Tamao, T. Hayashi and M. Kumada, J. Organomet. Chem., 1976, 114, C19; K. Tamao, S. Okazaki and M. Kumada, J. Organomet. Chem., 1978, 146, 87; H. Sakurai, Y. Kamiyama and Y. Nakadaira, J. Am. Chem. Soc., 1975, 97, 931; Chem. Lett., 1975, 887; H. Sakurai and T. Imai, Chem. Lett., 1975, 891; H. Sakurai, Y. Eriyama, Y. Kamiyama and Y. Nakadaira, J. Organomet. Chem., 1984, 264, 229; H. Watanabe, M. Kobayashi, K. Higuchi and Y. Nagai, J. Organomet. Chem., 1980, 186, 51; D. Seyferth, E. W. Goldman and J. Escudié, J. Organomet. Chem., 1984, 271, 337; T. Kobayashi, T. Hayashi, H. Yamashita and M. Tanaka, Chem. Lett., 1989, 467; Y. Ito, M. Suginome and M. Murakami, J. Org. Chem., 1991, 56, 1948.
- 5 L. Sanow, M. Chai, K. Galat, P. Rinaldi, W. Youngs and C. A. Tessier, 30th Organosilicon Symposium, London, Ontario, May 1977, Abstract P69; L. M. Sanow, M. Chai, D. B. McConnville, K. J. Galat, R. S. Simons, P. L. Rinaldi, W. J. Youngs and C. A. Tessier, *Organometallics*, in the press; M. Auburn, M. Ciriano, J. A. K. Howard, M. Murray, N. J. Pugh, J. L. Spencer, F. G. A. Stone and P. Woodward, *J. Chem. Soc., Dalton Trans.*, 1980, 659; Y. Levchinsky, N. P. Rath and J. Braddock-Wilking, *Organometallics*, 1999, **18**, 2583.
- 6 Other dinuclear platinum complexes with bridging silylene or silyl ligands: E. A. Zarate, C. A. Tessier-Youngs and W. J. Youngs, J. Am. Chem. Soc., 1988, 110, 4068; J. Chem. Soc., Chem. Commun., 1989, 577; A. B. Anderson, P. Shiller, E. A. Zarate, C. A. Tessier-Youngs and W. J. Youngs, Organometallics, 1989, 8, 2320; M. J. Michalczyk, C. A. Recatto, J. C. Calabrese and M. J. Fink, J. Am. Chem. Soc., 1992, 114, 7955; R. H. Heyn and T. D. Tilley, J. Am. Chem. Soc., 1992, 114, 1917.
- 7 Y.-J. Kim, S.-C. Lee, J.-I. Park, K. Osakada, J.-C. Choi and T. Yamamoto, *Organometallics*, 1998, **17**, 4929.
- 8 K. Osakada, Y. Özawa and A. Yamamoto, J. Organomet. Chem., 1990, 399, 341.
- 9 F. Ozawa, T. Ito, Y. Nakamura and A. Yamamoto, J. Organomet. Chem., 1979, 168, 375; F. Ozawa, T. Ito and A. Yamamoto, J. Am. Chem. Soc., 1980, 102, 6457; F. Ozawa, T.-I. Son, S. Ebina, K. Osakada and A. Yamamoto, Organometallics, 1992, 11, 171.
- 10 U. Schubert and C. Müller, J. Organomet. Chem., 1989, 373, 165.
- 11 Y. Pan, J. Mague and M. J. Fink, Organometallics, 1992, 11, 3495.
- 12 C. K. Johnson, ORTEP II, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.
- 13 TEXSAN, Structure Analysis Package, Molecular Structure Corporation, Houston, TX, 1985.
- 14 International Tables for X-Ray Crystallography, Kynoch Press, Birmingham, 1974.

Paper a907745f