


2-Phospha-4-silabicyclo[1.1.0]butane as a Reactive Intermediate**

J. Chris Slootweg, Frans J. J. de Kanter, Marius Schakel, Andreas W. Ehlers, Barbara Gehrhus, Martin Lutz, Allison M. Mills, Anthony L. Spek, and Koop Lammertsma*


Bicyclo[1.1.0]butane remains a fascinating compound. Stretching of its central C–C σ -bond culminates in a planar singlet diradical transition structure for inversion,^[1] whereas stretching of the peripheral bonds leads to valence isomerization.^[2] Bridging heteroatoms affect both processes. They enable the coexistence of the puckered bicyclic and planar diyl structures, and have thereby sparked intensive studies dominated by recent discoveries of diphosphorus analogues. Niecke et al. synthesized a crystalline 2,4-diphosphacyclobutane-1,3-diyl **A**^[3] that isomerizes photolytically to 2,4-diphos-




phabicyclo[1.1.0]butane **B**^[4] and thermally to 1,2-dihydro-1,2-diphosphete **C**.^[5] A still more congested diyl **A** that does not isomerize was reported by Yoshifuji and co-workers.^[6] The group of Bertrand obtained both stable planar diyl and puckered forms for the isoelectronic 1,3-dibora-2,4-diphosphoniocyclobutanes.^[7] 2,4-Diphosphabicyclo[1.1.0]butane is puckered when the phosphorus atoms are *endo,endo* substituted as in dihydro diphosphabenzvalene **1**, which Mathey and co-workers obtained by an *intramolecular* phosphinidene addition to the C=C bond of a 1*H*-phosphirene.^[8] Without geometrical constraints, such additions invariably lead to 1,2-

diphosphetes of type **C** by isomerization or direct C–P insertion.^[9]

Data on 2,4-disilabicyclo[1.1.0]butanes^[10] is limited to two disilabenzvalenes^[11] and is absent for the corresponding planar diyls. Only a single 2-silabenzvalene is known, generated by photolysis of a silabenzene.^[12] For the synthesis of the 2,4-disila derivatives Ando et al. applied valence isomerization of bis(silirene) that involves an intramolecular silylene addition to the C=C bond of a silirene.^[11] The use of stable silylenes might simplify this approach to a broader array of hetero derivatives, which we decided to explore for the unknown 2-phospha-4-silabicyclo[1.1.0]butanes **4**.



Reaction of silylene Si{1,2-(NCH₂tBu)₂C₆H₄} (Si(NN); **3**)^[13] with 1*H*-phosphirene **2a**^[14] in *n*-hexane at room temperature results in red crystalline 2,3-dihydro-1,3-phosphasilete **5a** instead of the desired **4** (Scheme 1).^[15] Evidently, two



Scheme 1. Synthesis of 2,3-dihydro-1,3-phosphasiletes **5**.

silylenes are involved, one in forming the novel ring structure and the other as a substituent after transfer of a phenyl group. No intermediate was detected by ³¹P NMR spectroscopy, either at low temperatures or with different stoichiometries of the reactants. The W(CO)₅-complexed phosphalkene part of **5a** is characterized by the downfield absorptions at $\delta(^{31}\text{P}) = 278.3$ ppm ($J(\text{P,W}) = 254.0$ Hz) and $\delta(^{13}\text{C}) = 197.3$ ppm ($J(\text{C,P}) = 3.5$ Hz). A single-crystal X-ray analysis confirmed the structure of **5a** (Figure 1).^[16] Its four-membered ring deviates from planarity owing to steric congestion, which is reflected in the torsion angle of 11.6(2)° for C1–P1–C2–Si1. The P1–C1 and P1–C2 bond lengths (1.689(4) and 1.835(4) Å, respectively) are normal, but the short Si1–C1 bond (1.841(5) Å) suggests some delocalization of charge.

To lower the migratory aptitude of the phosphorus substituent, we replaced the phenyl for a methyl group and used 1*H*-phosphirene **2b**^[14] instead of **2a** for the addition of silylene **3**. This had the desired effect, giving the crystalline zwitterion **6b** (68%), which only rearranged at elevated temperatures (75 °C, 2 h) to 2,3-dihydro-1,3-phosphasilete **5b** (Scheme 1). The molecular structure of **6b** (Figure 2,^[16] the crystal contains two independent molecules with similar geometries) shows a unique fused tricyclic ring structure which is retained in solution ($\delta(^{31}\text{P}) = -6.8$ ppm ($J(\text{P,W}) =$

[*] J. C. Slootweg, Dr. F. J. J. de Kanter, Dr. M. Schakel, Dr. A. W. Ehlers, Prof. Dr. K. Lammertsma
Department of Organic and Inorganic Chemistry
Faculty of Sciences, Vrije Universiteit
De Boelelaan 1083, 1081 HV, Amsterdam (The Netherlands)
Fax: (+31) 20-444-7488
E-mail: lammert@chem.vu.nl

Dr. B. Gehrhus
Department of Chemistry
School of Life Science, University of Sussex
Brighton, BN1 9QJ (UK)

Dr. M. Lutz, Dr. A. M. Mills, Prof. Dr. A. L. Spek
Bijvoet Center for Biomolecular Research
Crystal and Structural Chemistry, Utrecht University
Padualaan 8, 3584 CH, Utrecht (The Netherlands).

[**] The Netherlands Organization for Scientific Research (NWO/CW) is acknowledged for partial support.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.




Figure 1. Structure of **5a** in the crystal (displacement ellipsoids drawn at the 50% probability level; hydrogen atoms and the hexane solvent molecule are omitted for clarity). Selected bond lengths [Å], angles [°], and torsion angles [°]: W1–P1 2.4492(11), P1–C1 1.689(4), P1–C2 1.835(4), Si1–C1 1.841(5), Si1–C2 1.908(4); C1–P1–C2 92.9(2), P1–C2–Si1 86.13(17), C1–Si1–C2 85.96(18), P1–C1–Si1 92.7(2); C1–P1–C2–Si1 11.6(2).





Figure 2. Structure of **6b** in the crystal, one of the two crystallographically independent molecules is shown (displacement ellipsoids drawn at the 50% probability level; hydrogen atoms omitted for clarity). Selected bond lengths [Å], angles [°], and torsion angles [°]: W1–P1 2.5363(12), P1–C1 1.793(4), P1–C2 1.863(4), Si1–N1 1.725(4), Si1–N2 1.923(4), Si1–C1 1.750(4), Si1–C2 1.906(4), Si2–N2 1.933(4), Si2–N3 1.710(4), Si2–N4 1.705(4), Si2–C2 1.828(4), N1–C10 1.409(6), N1–C16 1.471(6), N2–C11 1.471(5), N2–C21 1.510(5), C1–C4 1.455(6); C1–P1–C2 90.79(19), P1–C2–Si1 85.70(18), C1–Si1–C2 90.70(19), P1–C1–Si1 92.7(2); C2–P1–C1–Si1 –2.4(2), P1–C1–C4–C5 8.2(8).

240.0 Hz), $\delta(^{29}\text{Si}) = -11.2$ ($^2J(\text{Si},\text{P}) = 17.2$ Hz, SiNN) and -33.4 ppm ($^2J(\text{Si},\text{P}) = 18.4$ Hz, SiNN^+). The negatively charged C1 is in a planar environment and is surrounded by shortened C1–P1 (1.793(4) Å), C1–C4 (1.455(6) Å), and C1–Si1 (1.750(4) Å) bonds that can be attributed to negative hyperconjugation, which is supported by the shielding of Si1

($\delta(^{29}\text{Si}) = -33.4$ ppm).^[17] Furthermore, the cationic N2 lengthens the N2–Si and N2–C bonds.^[18]

We assume that the reaction starts by addition of the silylene to the C=C bond of the phosphirene to give 2-phospha-4-silabicyclo[1.1.0]butane **4b** as a transient intermediate, followed by attack of a second silylene to the bridgehead carbon atom in a manner analogous to the reaction of nucleophiles with bicyclo[1.1.0]butane (Scheme 2).^[19] The resulting **7b** carries a silylium cation,



Scheme 2. Mechanism for the formation of **6b**.

which requires stabilization^[20] by coordination with the lone pair of electrons of a nearby nitrogen atom.^[21] This hypothesis was tested with SCS-MP2/6-311 + G** energy calculations on B3LYP/6-31G* optimized model structures (labeled **B**),^[22] which are devoid of $\text{W}(\text{CO})_5$, carry H instead of Np substituents, and have the nitrogen atoms linked by ethylene instead of phenyl groups. This simplification is validated by the similarity in optimized structures between **4B** and that containing the Np and $\text{W}(\text{CO})_5$ groups. Formation of **4B** from the reactants $\text{C}_2\text{H}_4\text{N}_2\text{Si}$ and $\text{C}_3\text{H}_5\text{P}$ is exothermic by 17.5 kcal mol^{−1} with an additional 8.5 kcal mol^{−1} for the addition of a second silylene to give **7B** (Figure 3). Lewis base stabilization





Figure 3. Relative SCS-MP2/6-311 + G**//B3LYP/6-31G* energies (in kcal mol^{−1}) for the rearrangement of **6B** to the thermodynamically more stable **5B** with selected bond lengths [Å].

of the silylium cation gives the kinetically favored zwitterion **6B** ($\Delta E = 21.7$, $\Delta E^\ddagger = 2.1$ kcal mol^{−1}), whereas a 1,3-methyl shift from phosphorus to silicon results in the thermodynamically favored phosphasilete **5B** ($\Delta E = 49.6$, $\Delta E^\ddagger = 6.3$ kcal mol^{−1}). Such 1,3-shifts to an electron-deficient silicon are very rare for both the methyl and phenyl groups.^[23,24]

To reduce the attractiveness of the transient 2-phospha-4-silabicyclo[1.1.0]butane for attack by a second silylene, a methyl group was introduced at the bridgehead position by treating 1*H*-phosphirene **2c**^[14] with **3**. This had the desired effect, but mainly valence isomerization occurred to the isomeric 1,2-dihydro-1,2-phosphasiletes **8a** and **8b**,^[25] together with the formation of some **5c** (5:2:2), because of the elevated temperatures (80 °C, 13 h) required for the reaction to take place (Scheme 3). After fractional crystallization, **8a** could be obtained as a yellow solid for which only single resonances were observed at $\delta(^{31}\text{P}) = 15.2$ ppm



Scheme 3. Isomerization of **4c**.

($^1J(\text{P},\text{W}) = 205.0 \text{ Hz}$) and $\delta(^{29}\text{Si}) = -17.1 \text{ ppm}$ ($^1J(\text{Si},\text{P}) = 45.6 \text{ Hz}$) confirming the stoichiometry of the reaction. The structure of **8a** was established unequivocally by a single-crystal X-ray structural analysis (Figure 4),^[16] which shows




Figure 4. Structure of **8a** in the crystal (displacement ellipsoids drawn at the 50% probability level; hydrogen atoms are omitted for clarity). Selected bond lengths [Å], angles [°], and torsion angles [°]: W1–P1 2.5269(6), P1–Si1 2.3074(8), P1–C1 1.854(2), P1–C10 1.841(2), Si1–N1 1.726(2), Si1–N2 1.7249(19), Si1–C2 1.863(2), C1–C2 1.360(3), C1–C4 1.489(3), C2–C3 1.507(3); C1–P1–Si1 73.30(7), P1–Si1–C2 76.89(7); P1–C1–C2–Si1 7.46(17).

that the four-membered ring is nearly planar with a torsion angle of $7.46(17)^\circ$ for P1–C1–C2–Si1 and that the P1–Si1 bond (2.3074(8) Å) is slightly elongated owing to steric congestion.

SCS-MP2/6-311+G** calculations on B3LYP/6-31G* model structures containing carbon substituents^[22] show that intermediate 2-phospha-4-silabicyclo[1.1.0]butane **4C** can isomerize to the thermodynamically preferred 1,2-dihydro-1,2-phosphasiletes **8A** and **8B** with barriers of only 21.3 and 21.6 kcal mol⁻¹, respectively. This similarity in $\Delta E^\ddagger$ values is reflected in the two experimentally observed isomers. The valence isomerization of **4C** to **8** is symmetry-allowed and can be described as a $[\sigma 2s + \sigma 2a]$ process.^[26] Such a pathway is unprecedented for the analogous isomerization of bicyclo[1.1.0]butane, for which *s-trans*-1,3-butadiene is the favored product.

In conclusion, thermally stable silylene **3** reacts with 1*H*-phosphirenes **2** to give the first isolated 2,3-dihydro-1,3-

phosphasiletes **5**, zwitterion **6b**, and 1,2-dihydro-1,2-phosphasiletes **8** with the novel 2-phospha-4-silabicyclo[1.1.0]butanes **4** being a reactive intermediate in the reaction, the course of which can be tuned by changing the substituents.

Experimental Section

5a: Complex **2a**^[14] (38.6 mg, 72.3 μmol) in *n*-hexane (5 mL) was added slowly to a solution of **3**^[13] (39.7 mg, 144.6 μmol) in *n*-hexane (2 mL) at -50°C , warmed to ambient temperature, and stirred for 16 h. Filtration, concentration, and cooling at -20°C affording red crystals of **5a** (56 mg, 66%): m.p. 245°C (decomp.); ^{29}Si NMR (79.5 MHz, C_6D_6): $\delta = -5.1$ (d, $^2J(\text{Si},\text{P}) = 6.9 \text{ Hz}$; Ph–SiNN), -13.5 ppm (d, $^2J(\text{Si},\text{P}) = 31.8 \text{ Hz}$; SiNN); $^{31}\text{P}\{^1\text{H}\}$ NMR (101.3 MHz, CDCl_3): $\delta = 278.3 \text{ ppm}$ ($^1J(\text{P},\text{W}) = 254.0 \text{ Hz}$); HRMS (EI, 70 eV): m/z (%): 1082 (2) [M^+]; calcd for $\text{C}_{51}\text{H}_{63}\text{O}_5\text{N}_4\text{Si}_2\text{PW}$: 1082.35840; found: 1082.35376.

5b: Complex **2b**^[14] (12.3 mg, 26.0 μmol) and **3** (14.3 mg, 52.1 μmol) were heated at 75°C for 2 h in C_6D_6 (0.5 mL). Evaporation to dryness, extraction into *n*-hexane, and subsequent filtration and cooling at -20°C gave red crystals of **5b** (15 mg, 57%): m.p. 150°C (decomp.); ^{29}Si NMR (79.5 MHz, C_6D_6): $\delta = 4.8$ (d, $^2J(\text{Si},\text{P}) = 4.2 \text{ Hz}$; Me–SiNN), -13.8 ppm (d, $^2J(\text{Si},\text{P}) = 33.2 \text{ Hz}$; SiNN); $^{31}\text{P}\{^1\text{H}\}$ NMR (101.3 MHz, C_6D_6): $\delta = 276.6 \text{ ppm}$ ($^1J(\text{P},\text{W}) = 246.1 \text{ Hz}$); HRMS (EI, 70 eV): m/z (%): 1020 (5) [M^+]; calcd for $\text{C}_{46}\text{H}_{61}\text{O}_5\text{N}_4\text{Si}_2\text{PW}$: 1020.34277; found: 1020.3391.

6b: Complex **2b** (20.4 mg, 43.2 μmol) in *n*-hexane (5 mL) was added to a solution of **3** (23.7 mg, 86.3 μmol) in *n*-hexane (2 mL) at ambient temperature and stirred for 2 h. Concentration and cooling at 0°C affording yellow crystals of **6b** (30 mg, 68%): m.p. 149°C (decomp.); ^{29}Si NMR (79.5 MHz, C_6D_6): $\delta = -33.4$ (d, $^2J(\text{Si},\text{P}) = 18.4 \text{ Hz}$; SiNN⁺), -11.2 ppm (d, $^2J(\text{Si},\text{P}) = 17.2 \text{ Hz}$; SiNN); $^{31}\text{P}\{^1\text{H}\}$ NMR (101.3 MHz, C_6D_6): $\delta = -6.8 \text{ ppm}$ ($^1J(\text{P},\text{W}) = 240.0 \text{ Hz}$).

8: Complex **2c**^[14] (44.9 mg, 81.9 μmol) and **3** (33.7 mg, 122.8 μmol) were heated at 80°C for 13 h in C_6D_6 (0.5 mL). $^{31}\text{P}\{^1\text{H}\}$ NMR (101.3 MHz, C_6D_6): $\delta = 12.0$ ($^1J(\text{P},\text{W}) = 198.5 \text{ Hz}$, 19% yield of **8b** by NMR spectroscopy), 15.2 ($^1J(\text{P},\text{W}) = 205.6 \text{ Hz}$, 50% yield of **8a** by NMR spectroscopy), 308.1 ppm (br s, 19.5% yield of **5c** by NMR spectroscopy). Evaporation to dryness, extraction into *n*-hexane, and subsequent cooling at -80°C gave yellow crystals of **8a** (30 mg, 45%): m.p. 157°C (decomp.); ^{29}Si NMR (79.5 MHz, C_6D_6): $\delta = -17.1 \text{ ppm}$ (d, $^1J(\text{Si},\text{P}) = 45.6 \text{ Hz}$; SiNN); $^{31}\text{P}\{^1\text{H}\}$ NMR (101.3 MHz, C_6D_6): $\delta = 15.2 \text{ ppm}$ ($^1J(\text{P},\text{W}) = 205.0 \text{ Hz}$); HRMS (EI, 70 eV): m/z (%): 822 (80) [M^+]; calcd for $\text{C}_{36}\text{H}_{39}\text{O}_5\text{N}_2\text{SiPW}$: 822.1876; found: 822.1880. **5c:** MS (EI, 70 eV): m/z (%): 1096 (30) [M^+].

Received: March 4, 2004 [Z54183]

Keywords: phosphorus heterocycles · reaction mechanisms · reactive intermediates · substituent effects · valence isomerization

- [1] K. A. Nguyen, M. S. Gordon, J. A. Boat, *J. Am. Chem. Soc.* **1994**, *116*, 9241–9249.
- [2] K. A. Nguyen, M. S. Gordon, *J. Am. Chem. Soc.* **1995**, *117*, 3835–3847.
- [3] E. Niecke, A. Fuchs, F. Baumeister, M. Nieger, W. W. Schoeller, *Angew. Chem.* **1995**, *107*, 640–642; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 555–557.
- [4] E. Niecke, A. Fuchs, M. Nieger, *Angew. Chem.* **1999**, *111*, 3213–3216; *Angew. Chem. Int. Ed.* **1999**, *38*, 3028–3031.
- [5] O. Schmidt, A. Fuchs, D. Gudat, M. Nieger, W. Hoffbauer, E. Niecke, W. W. Schoeller, *Angew. Chem.* **1998**, *110*, 995–998; *Angew. Chem. Int. Ed.* **1998**, *37*, 949–952.

- [6] H. Sugiyama, S. Ito, M. Yoshifuji, *Angew. Chem.* **2003**, *115*, 3932–3934; *Angew. Chem. Int. Ed.* **2003**, *42*, 3802–3804.
- [7] a) D. Scheschkewitz, H. Amii, H. Gornitzka, W. W. Schoeller, D. Bourissou, G. Bertrand, *Science* **2002**, *295*, 1880–1881; b) D. Scheschkewitz, H. Amii, H. Gornitzka, W. W. Schoeller, D. Bourissou, G. Bertrand, *Angew. Chem.* **2004**, *116*, 595–597; *Angew. Chem. Int. Ed.* **2004**, *43*, 585–587.
- [8] M. White, L. Ricard, F. Mathey, *Organometallics* **2003**, *22*, 4825–4828.
- [9] a) N. H. Tran Huy, L. Ricard, F. Mathey, *Organometallics* **1997**, *16*, 4501–4504; b) B. Wang, K. A. Ngyen, G. N. Srinivas, C. L. Watkins, S. Menzer, A. L. Spek, K. Lammertsma, *Organometallics* **1999**, *18*, 796–799.
- [10] Only one Si₂C₂-ring is known: G. Fritz, S. Wartenessian, E. Matern, W. Hömle, H. G. Von Schnering, *Z. Anorg. Allg. Chem.* **1981**, *475*, 87–108.
- [11] a) W. Ando, T. Shiba, T. Hidaka, K. Morihashi, O. Kikuchi, *J. Am. Chem. Soc.* **1997**, *119*, 3629–3630; b) Y. Kabe, K. Ohkubo, H. Ishikawa, W. Ando, *J. Am. Chem. Soc.* **2000**, *122*, 3775–3776.
- [12] K. Wakita, N. Tokitoh, R. Okazaki, N. Takagi, S. Nagase, *J. Am. Chem. Soc.* **2000**, *122*, 5648–5649.
- [13] B. Gehrhus, M. F. Lappert, J. Heinicke, R. Boese, D. Bläser, *J. Chem. Soc. Chem. Commun.* **1995**, 1931–1932.
- [14] A. Marinetti, F. Mathey, *Organometallics* **1984**, *3*, 456–461. Synthesis of **2b** and **2c** was performed in analogy to **2a**; see Supporting Information.
- [15] So far, only one 1λ³-1,3-phosphasilete was mentioned: M. Driess, H. Pritzkow, *Phosphorus Sulfur Silicon Relat. Elem.* **1993**, *76*, 57–60.
- [16] Crystal structure determinations: X-ray intensities were measured on a Nonius KappaCCD diffractometer with rotating anode ($\lambda = 0.71073 \text{ \AA}$) at a temperature of 150(2) K. The structures were refined with SHELXL97^[27] against F² of all reflections. Illustrations, structure calculations and checking was performed with the PLATON package.^[28] **5a** (C₅₁H₆₃N₄O₅P-Si₂W-0.6C₆H₁₄): $M_w = 1134.76$, red block, $0.06 \times 0.18 \times 0.21 \text{ mm}^3$, monoclinic, $P2_1$ (No. 4), $a = 10.5532(1)$, $b = 16.0049(2)$, $c = 17.2900(2) \text{ \AA}$, $\beta = 100.5233(5)$, $V = 2871.21(6) \text{ \AA}^3$, $Z = 2$, $\rho = 1.313 \text{ g cm}^{-3}$; 51129 measured reflections, 13052 unique reflections ($R_{\text{int}} = 0.050$); structure solution with automated Patterson methods;^[29] 643 refined parameters, 46 restraints; $R1/wR2$ [$I > 2\sigma(I)$]: 0.0338/0.0755, $R1/wR2$ (all reflections) = 0.0472/0.0794; GoF = 1.072; Flack $\times$ parameter: $-0.024(4)$. **6b** (C₄₆H₆₁N₄O₅P-Si₂W): $M_w = 1020.99$, yellow block, $0.18 \times 0.24 \times 0.30 \text{ mm}^3$, triclinic, $P\bar{1}$ (No. 2), $a = 10.6476(3)$, $b = 21.2390(8)$, $c = 22.6862(5) \text{ \AA}$, $\alpha = 74.495(2)$, $\beta = 89.5698(19)$, $\gamma = 82.814(3)^\circ$, $V = 4903.0(3) \text{ \AA}^3$, $Z = 4$, $\rho = 1.383 \text{ g cm}^{-3}$. The crystal appeared to be non-merohedrally twinned with a 180° rotation about $uvw = [100]$ as twin operation. The intensity data were evaluated with EvalCCD,^[30] taking this twin relation into account; 55937 measured reflections, 22336 unique reflections ($R_{\text{int}} = 0.058$); structure solution with direct methods;^[31] 1064 refined parameters, 388 restraints; bond lengths and angles of the two independent molecules were restrained to be the same; $R1/wR2$ [$I > 2\sigma(I)$]: 0.0499/0.1070, $R1/wR2$ (all reflections) = 0.0767/0.1216; GoF = 1.128; the refinement^[24] of the twin fraction resulted in 0.794(1):0.206. **8a** (C₃₆H₃₉N₂O₅PSiW): $M_w = 822.60$, yellow plate, $0.06 \times 0.36 \times 0.42 \text{ mm}^3$, monoclinic, $P2_1/c$ (No. 14), $a = 19.1665(1)$, $b = 10.6482(1)$, $c = 20.5181(1) \text{ \AA}$, $\beta = 119.8343(4)^\circ$, $V = 3632.53(5) \text{ \AA}^3$, $Z = 4$, $\rho = 1.504 \text{ g cm}^{-3}$; 58693 measured reflections, 8329 unique reflections ($R_{\text{int}} = 0.046$); structure solution with automated Patterson methods;^[29] 422 refined parameters, no restraints; $R1/wR2$ [$I > 2\sigma(I)$]: 0.0204/0.0455, $R1/wR2$ (all reflections) = 0.0289/0.0488; GoF = 1.038. CCDC-216769 (**5a**), -216770 (**6b**), and -232111 (**8a**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB21EZ, UK; fax: (+44) 1223-336-033; or deposit@ccdc.cam.ac.uk).
- [17] M. Sebastian, M. Nieger, D. Szieberth, L. Nyulászi, E. Niecke, *Angew. Chem.* **2004**, *116*, 647–651; *Angew. Chem. Int. Ed.* **2004**, *43*, 637–641.
- [18] R. Bettenhausen, W. Milius, W. Schnick, *Chem. Eur. J.* **1997**, *3*, 1337–1341.
- [19] S. Hoz in *The Chemistry of the Cyclopropyl Group, Part 2* (Ed.: Z. Rappoport), Wiley, New York, **1987**, Chapter 19.
- [20] K. -C. Kim, C. A. Reed, D. W. Elliott, L. J. Mueller, F. Tham, L. Lin, J. B. Lambert, *Science* **2002**, *297*, 825–827.
- [21] R. Corriu, G. Lanneau, C. Priou, *Angew. Chem.* **1991**, *103*, 1153–1155; *Angew. Chem. Int. Ed. Engl.* **1991**, *30*, 1130–1132.
- [22] a) Gaussian 98 (Revision A.7), M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, C. Gonzalez, M. Challacombe, P. M. W. Gill, B. G. Johnson, W. Chen, M. W. Wong, J. L. Andres, M. Head-Gordon, E. S. Replogle, J. A. Pople, Gaussian, Inc., Pittsburgh, PA, **1998**; b) A. D. Becke, *Phys. Rev. A* **1988**, *38*, 3098–3100; c) C. Lee, W. Yang, R. G. Parr, *Phys. Rev. B* **1988**, *37*, 785–789; d) S. Grimme, *J. Chem. Phys.* **2003**, *118*, 9095–9102; see Supporting Information.
- [23] a) K. Ueno, K. Nakano, H. Ogino, *Chem. Lett.* **1996**, 459–460, and references therein; b) K. Ueno, M. Sakai, H. Ogino, *Organometallics* **1998**, *17*, 2138–2140.
- [24] C. Eaborn, A. Kowalewska, W. Stanczyk, *J. Organomet. Chem.* **1998**, *560*, 41–46, and references therein.
- [25] Only one other synthesis of 1,2-dihydro-1,2-phosphasiletes was reported: a) S. Haber, R. Boese, M. Regitz, *Angew. Chem.* **1990**, *102*, 1523–1525; *Angew. Chem. Int. Ed. Engl.* **1990**, *29*, 1436–1438; b) S. Haber, M. Schmitz, U. Bergsträßer, J. Hoffmann, M. Regitz, *Chem. Eur. J.* **1999**, *5*, 1581–1589.
- [26] J. C. Slootweg, A. W. Ehlers, K. Lammertsma, *Phosphorus Sulfur Silicon Relat. Elem.* **2004**, *179*, 803–807.
- [27] G. M. Sheldrick, SHELXL-97, Program for crystal structure refinement, University of Göttingen, Germany, **1997**.
- [28] A. L. Spek, *J. Appl. Crystallogr.* **2003**, *36*, 7–13.
- [29] P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, S. Garcia-Granda, R. O. Gould, J. M. M. Smits, C. Smykalla, The DIRDIF99 program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands, **1999**.
- [30] A. J. M. Duisenberg, L. M. J. Kroon-Batenburg, A. M. M. Schreurs, *J. Appl. Crystallogr.* **2003**, *36*, 220–229.
- [31] G. M. Sheldrick, SHELXS-97, Program for crystal structure solution, University of Göttingen, Germany, **1997**.