

### Sulfur Dioxide Mediated One-Pot, Three- and Four-Component Syntheses of Polyfunctional Sulfonamides and Sulfonic Esters: Study of the Stereoselectivity of the Ene Reaction of Sulfur Dioxide

Laure C. Bouchez, Srinivas Reddy Dubbaka, Māris Turks, and Pierre Vogel\* Institute of Chemical Sciences and Engineering, Swiss Federal Institute of Technology (EPFL), BCH, CH-1015 Lausanne-Dorigny, Switzerland

pierre.vogel@epfl.ch

Received June 7, 2004

The ene reaction of sulfur dioxide with enoxysilanes or with allylsilanes generates silyl sulfinates that can be brominated (Br<sub>2</sub> or NBS) or chlorinated (NCS or Cl<sub>2</sub>) to produce the corresponding sulfonyl halides. They react with primary and secondary amines or alcohols to give the corresponding sulfonamides and sulfonic esters, respectively. The hetero-Diels-Alder addition of sulfur dioxide to 1-oxy- or 1,3-dioxy-1,3-dienes generates zwitterions that add to enoxysilanes or allylsilanes giving silyl sulfinates that can be converted in situ into polyfunctional sulfonamides or sulfonic esters. This realizes quick access to libraries of complicated sulfonamides and sulfonic esters applying one-pot, three- and four-component methods.

#### Introduction

The sulfonamides constitute an important class of drugs (the sulfa drugs), with several types of pharmacological agents possessing antibacterial, anticarbonic anhydrase, diuretic hypoglycemic antithyroid, antihypertensive, antiinflammatory, and antiviral properties.<sup>1,2</sup> Recently, structurally novel sulfonamide derivatives have shown substantial antitumor activities,<sup>3,4</sup> or are caspase-1<sup>5</sup> inhibitors.<sup>6</sup> The majority of sulfonamides are prepared from the reaction of sulfonyl chlorides with ammonia and primary or secondary amines.<sup>7</sup> Arenesulfonyl chlorides are prepared from arenes by electrophilic substitution with ClSO<sub>3</sub>H or from arenesulfonic acids by reaction with PCl<sub>5</sub>,<sup>8</sup> POCl<sub>3</sub>,<sup>9</sup> or COCl<sub>2</sub>.<sup>10</sup> Other methods imply the reactions of arenediazonium salts with SO<sub>2</sub>/CuCl<sub>2</sub>,<sup>11</sup> the oxidation of thioesters<sup>12</sup> or sulfenyl chlorides,<sup>13</sup> or the reaction of organolithium<sup>14</sup> or organomagnesium<sup>15</sup> reagents with  $SO_2Cl_2$  or  $SO_2 + Cl_2$ . Alkanesulfonyl chlorides can be obtained by reaction of the corresponding

(5) For examples of caspase-1 inhibitors see: Harter, W. G.; Albre-cht, H.; Brady, K.; Bradley, C.; Dunbar, J.; Gilmore, J.; Hays, S.; Kostlan, C. R.; Lunney, B.; Walker, N. *Bioorg. Med. Chem. Lett.* **2004**, 14, 809.

(6) Drew, J. Science 2000, 287, 1960.

(7) (a) Cremlyn, R. Organosulfur Chemistry: An Introduction; J. Wiley and Sons.: New York, 1996; pp 224–225. (b) Anderson, K. K. In Sulfonic Acids and Their Derivatives in Comprehensive Organic Chemistry, Barton, D. H. R., Ollis, W. D., Jones, D. N., Eds.; Pergamon (8) Heumann, K.; Köchlin, P. Chem. Ber. 1882, 15, 1114.

(9) (a) Barco, A.; Benetti, S.; Pollini, G. R.; Taddia, R. Synthesis 1974, 877. (b) Fujita, S. Synthesis 1982, 423.
(10) Blank, H. H.; Pfister, T. German Offen. 1979; Chem. Abstr. 1979, 91, 107804.

<sup>(1) (</sup>a) Navia, M. A. Science 2000, 288, 2132. (b) Mohamed, S. K. Afinidad **2000**, *57*, 451. (c) Supuran, C. T.; Scozzafava, A.; Casini, A. *Med. Res. Rev.* **2003**, *23*, 146. (d) Supuran, C. T.; Casini, A.; Scozzafava, A. Med. Res. Rev. 2003, 23, 535 and references cited therein. (e) Patel, O. G.; Mberu, E. K.; Nzila, A. M.; Macreadie, I. G. Trends Parasitol. 2004, 20, 1. (f) Kumar, A.; Katiyar, S. B.; Agarwal, A.; Chauhan, P. M. S. Drugs Future 2003, 28, 242.

<sup>(2)</sup> For selected examples of biological activities see: (a) Thais-rivongs, S.; Janakiraman, M. N.; Chong, K.-C.; Tomish, P. K.; Dolak, L. A.; Turner, S. R.; Stronbach, J. W.; Lynn, J. C.; Horng, M.-M.; Hinshaw, R. R.; Watenpaugh, K. D. *J. Med. Chem.* **1996**, *39*, 2400. (b) Chazalette, C.; Rivière-Baudet, A.; Scozzafava, F.; Abbate, Z. B.; Maarouf, C. T.; Supurau, J. *J. Enzyme Inhib.* **2001**, *16*, 125. (c) Scozava, A.; Banciu, M. D.; Popescu, A.; Claudin, C. T.; *J. Enzyme Inhib.* 2000, 15, 533. (d) Song, X.; He, H. T.; Siahaan, T. J. Org. Lett. 2002, 4, 549. (e) Dyatkin, A. B.; Hoekstra, W. J.; Kinney, W. A.; Kontoyianni, M.; Santulli, R. J.; Kimball, E. S.; Carolyn-Fisher, M.; Prouty, S. M.; Abraham, W. M.; Andrade-Gordon, P.; Hlasta, D. J.; He, W.; Hornby, P. J.; Damiano, B. P.; Maryanoff, B. E. Biorg. Med. Chem. Lett. 2004, 14, 591. (f) Johansson, A.; Poliakov, A.; Akerblom, E.; Wiklund, K.; Lindeberg, G.; Winiwarter, S.; Danielson, U. H.; Samuelsson, B.; Halberg, A. *Bioorg. Med. Chem.* **2003**, *11*, 2551. (g) Tellew, J. E.; Baska, R. A. F.; Beyer, S. M.; Carlson, K. E.; Cornelius, L. A.; Fadnis, L.; Gu, Z.; Kunst, B. L.; Kowala, M. C.; Monshizadegan, H.; Murugesan, N.; Z.; Kunst, B. L.; Kowala, M. C.; Monshizadegan, H.; Murugesan, N.; Ryan, C. S.; Valentine, M. T.; Yang, Y.; Macor, J. E. *Bioorg. Med. Chem. Lett.* 2003, 13, 1093. (h) Boss, C.; Bolli, M. W.; Weller, T.; Fiscli, W.; Clozel, M. *Bioorg. Med. Chem. Lett.* 2003, 13, 951. (i) Su, D.-S.; Markowitz, M. K.; DiPardo, R. M.; Murphy, K. L.; Harrell, C. M.; O'Malley, S. S.; Ransom, R. W.; Chang, R. S. L.; Ha, S.; Hess, F. J.; Pettibone, D. J.; Mason, G. S.; Boyce, S.; Freidinger, R. M.; Bock, M. G. J. Am. Chem. Soc. 2003, 125, 7516. (j) Singh, S. K.; Reddy, P. G.; Rao, K. S.; Lohray, B. B.; Misra, P.; Rajjak, S. A.; Rao, Y. K.; Venkateswarlu, A. *Bioorg. Med. Chem. Lett.* 2004, 14, 499. (k) Pastorekova, S.; Casini, A.; Scozzafava, A.; Vullo, D.; Pastorek, J.; Sumuran, C. T. *Bioorg. Med. Chem. Lett.* 2004, 10 Watson P. Bastorekova, S.; Casini, A.; Scozzalava, A.; Vinio, D.; Pastorek, J.;
 Supuran, C. T. *Bioorg. Med. Chem. Lett.* 2004, *14*, 869. (*J*) Watson, R.;
 J.; Batty, D.; Baxter, A. D.; Hannah, D. R.; Owen, D. A.; Montana, J. G. *Tetrahedron Lett.* 2002, *43*, 683. (m) Zhong, Z.; Bibbs, J. A.; Yuan, W.; Wong, C. H. *J. Am. Chem. Soc.* 1991, *113*, 2259.

<sup>(3) (</sup>a) Owa, T.; Yoshino, H.; Okauchi, T.; Yoshimatsu, K.; Ozawa, Y.; Sugi, N. H.; Nagasu, T.; Koyanagi, N.; Kitoh, K. *J. Med. Chem.* **1991**, *42*, 3789. (b) Funahashi, Y.; Sugi, N. H.; Semba, T.; Yamamoto, Y.; Hamaoka, S.; Tsukakara-Tamai, N.; Ozawa, Y.; Tsuruoka, A.; Nara, K.; Takahashi, K.; Okabe, T.; Kamata, J.; Owa, T.; Ueda, N.; Haneda, T.; Yonaga, M.; Yoshimatsu, K.; Wakabayashi, T. Cancer Res. 2002, 62, 6116.

<sup>(4) (</sup>a) Abbate, F.; Casini, A.; Owa, T.; Scozzafava, A.; Supuran, C. **1.** Bioorg. Med. Chem. Lett. **2004**, *14*, 217. (b) Winum, J.-Y.; Vullo, D.; Casini, A.; Montero, J.-L.; Scozzafava, A.; Supuran, C. T. J. Med. Chem. **2003**, *26*, 5471. (c) Takagi, M.; Honmura, T.; Watanabe, S.; Yamaguchi, R.; Nogawa, M.; Nishimura, I.; Katoh, F.; Matsuda, M.; Aidaka, H. Invest. New Drugs 2003, 21, 387. (d) Scozzafava, A.; Owa, T.; Mas-trolorenzo, A.; Supuran, C. T. Curr. Med. Chem. 2003, 10, 925 and references cited therein. (e) Terret, C.; Zanetta, S.; Roche, W.; Schellens, J. H. M.; Faber, M. N.; Wanders, J.; Ravic, M.; Droz, J. P. Eur. J. Cancer 2003, 39, 1097.

alkane with SO<sub>2</sub> and Cl<sub>2</sub> under radical conditions.<sup>16</sup> All these methods<sup>17</sup> are relatively harsh (acidic, basic) and cannot be applied to polyfunctional substrates. Recently the direct synthesis of sulfonamides and sulfonic esters from sulfonic acids<sup>18</sup> and a one-pot synthesis of sulfonamides from Grignard reagents and SO<sub>2</sub><sup>18,19</sup> has been reported. Both procedures are amenable to aromatic and heteroaromatic sulfonamides.

We disclose here unprecedented one-pot three- and four-component synthesis of sulfonamides and sulfonic esters that allow the easy preparation of polyfunctional derivatives that can contain keto  $\beta$ -alkoxyketone, allylic ether, alkenyl ester moieties, or/and  $\beta$ , $\gamma$ -unsaturation and up to three stereogenic centers. The methods rely on recently discovered ene reactions of SO<sub>2</sub> with allylsilanes and enoxysilanes, on one hand,<sup>20</sup> and on our reaction cascade combining electron-rich 1,3-dienes and alkenes with sulfur dioxide, on the other hand.<sup>21-23</sup> Sulfonamides with the functional and stereochemical complexities of those described in this work would require multistep procedures applying available methods.<sup>24</sup>

### **Results and Disscusion**

One-Pot, Three-Component Synthesis. In 1975, Grieco et al.<sup>25</sup> reported the ene reaction of sulfur dioxide with enoxysilanes derived from carboxylic esters. We have found that silvl enol ethers derived from acylic and cyclic ketones also undergo the ene reaction with SO<sub>2</sub> under the promotion of Lewis acids with the generation of the corresponding silyl  $\beta$ -oxoalkanesulfinates **3** (Scheme 1). As for trialkyltin sulfinates that can be oxidized into the corresponding sulfonyl halides by halogenosis by (Cl<sub>2</sub>,  $Br_2$ , or  $I_2$ )<sup>26,27</sup> we found that silvl sulfinates **3** are readily

(12) Klamman, D.; Drahowzal, F. *Monatsh. Chem.* **1952**, *83*, 463. (13) (a) Sohechter, M. S.; Haller, H. L. *J. Am. Chem. Soc.* **1941**, *63*, 1764. (b) Banks, R. E.; Haszeldine, R. N.; Reppin, A. J. Chem. Soc.

1966, 1171. (c) Dickman, E.; Brasha, P. Isr. J. Chem. 1969, 7, 589. (14) (a) Quast, M.; Kies, F. Synthesis 1974, 490. (b) Hamda, T.; Yonemitsu, O. Synthesis 1986, 852. (c) Bhattacharya, S. N.; Eaborn,

C.; Walton, D. P. M. J. Chem. Soc. 1968, 1265.

(15) (a) Field, M.; Rieck, H.-P. Chem. Ztg. 1976, 100, 391. (b) Scott,
 R. B.; Lutz, R. E. J. Org. Chem. 1954, 19, 830. (c) Cherbuliez, E.;
 Schrauder, O. Helv. Chim. Acta 1923, 6, 249.
 (16) (a) Deichtinger, H. Chem. Ber. 1963, 86, 3068. (b) Berthold, H.;

Huenecke, H.; Bughardt, D.; Hampel, M.; Helwig, D.; Kopinke, F. D.; Krebes, S.; Niegel, H.; Pritzkow, W. J. Prakt. Chem. 1979, 32, 279.
 (17) For other methods, see: (a) Sheehan, J. C.; Zoller, U.; Ben Ishai,

 (1) For other methods, see: (a) Sheenah, J. C.; Zoher, O.; Bent Shah,
 D. J. Org. Chem. 1974, 39, 1817. (b) Poshkus, A. C.; Herweh, J. E.;
 Magnotta, F. A. J. Org. Chem. 1963, 28, 2766. (c) Glass, R. S.; Swedo,
 R. J. Synthesis 1977, 798. (d) Buchholt, H. C.; Senning, A. Acta Chem.
 Scand. 1970, 24, 2255. (e) Zhang, J.; Shi, Y. Tetrahedron Lett. 2000,
 (d) Scandard S. K. Scantheris 1977, 20. (c) Dicklerg, K. Köh, 41, 8075. (f) Gupta, S. K. Synthesis 1977, 39. (g) Dicklore, K.; Küle, E.; Anders, B. Angew. Chem. 1965, 77, 429.

(18) Caddick, S.; Wilden, J. D.; Judd, D. B. J. Am. Chem. Soc. 2004, 126. 1024-1025.

(19) Pandya, R.; Murashima, T.; Tedeschi, L.; Barrett. A. G. M. J. Org. Chem. 2003, 68, 8274-8276.

(20) (a) Bouchez, L.; Vogel, P. *Synthesis* **2002**, 225. (b) Bouchez, L. C.; Dubbaka, S. R.; Turks, M. European Patent Appl. No. 03003611.5, 2003. (c) Bouchez, L. C.; Dubbaka, S. R.; Turks, M.; Vogel, P. Abstracts of Papers; 226th National Meeting of the American Chemical Society, Sept 7-11, 2003, New York; American Chemical Society: Washington, DC, 2003; ORGN-301.

(21) (a) Deguin, B.; Roulet, J. M.; Vogel, P. Tetrahedron Lett. **1997**, 38, 6197. (b) Roulet, J.-M.; Puhr, G.; Vogel, P. Tetrahedron Lett. **1997**, 38, 6201. (c) Narkevitch, V.; Schenk. K.; Vogel, P. Angew. Chem., Int. Ed. 2000, 39, 1806.

(22) (a) Narkevitch, V.; Megevand, S.; Schenk, K.; Vogel, P. J. Org. *Chem.* **2001**, *66*, 5080. (b) Huang, X.; Vogel, P. *Synthesis* **2002**, 232. (23) Turks, M.; Fonquerne, F.; Vogel, P. *Org. Lett* **2004**, *6*, 1053.





converted into the corresponding sulfonyl chlorides 4 either by chlorination  $(Cl_2)$  or by treatment with Nchlorosuccinimide (NCS). The corresponding sulfonyl bromides can also be prepared by treatment of silyl sulfinates 3 by bromine (Br<sub>2</sub>) or N-bromosuccinimide (NBS). Without purification, the crude sulfonyl halides react with primary and secondary amines in the presence of Et<sub>3</sub>N or pyridine to give the corresponding sulfonamides 5-18 (entries 1-14, Table 1).

To illustrate the method we reacted the trimethylsilyl enol ethers of methyl propanoate,28 pentan-3-one,28 acetophenone, cyclohexanone<sup>29</sup> and camphor<sup>30</sup> with an excess of SO<sub>2</sub> containing 5 mol % of (t-Bu)Me<sub>2</sub>SiOSO<sub>2</sub>CF<sub>3</sub>, as acid promoter, in  $CH_2Cl_2$  at -78 °C. Under these conditions only acyclic compounds 2a and 2b were converted into **3a** and **3b** in a few hours at -78 °C. With cyclic enol ethers **2d** and **2e**, ene reactions with SO<sub>2</sub> were only possible at higher temperatures such as 40 °C but with low yield due to decomposition of starting materials. We found that changing CH<sub>2</sub>Cl<sub>2</sub> for CH<sub>3</sub>CN dramatically enhanced the rates of the reaction. In the case of enoxysilanes 2d and 2e reactions were converted after a few hours at -78 °C. A mixture of SO<sub>2</sub>/CH<sub>3</sub>CN/t-Bu)Me<sub>2</sub>-

(25) Grieco, P. A.; Bothsho, G. V. D. Org. Chem. 2000, 63, 6060.
 (25) Grieco, P. A.; Boxler, D. Synth. Commun. 1975, 315. See also:
 (a) Sergeev, V. N.; Shipov, A. G.; Zaitseva, G.-S.; Baukov, Y. Zh.
 Obshch. Khim. 1979, 2753. (b) Sergeev, V. N.; Shipov, A. G.; Zaitseva,
 G.-S.; Baukov, Y. Chem. Abstr. 1980, 92, 146839.
 (26) King, M. D.; Sue, R. E.; White, R. H.; Young, D. J. Tetrahedron

Lett. 1997. 38. 4493.

(27) For the oxidation of sulfinic acid derivatives, see: Hoyle, J. In The Chemistry of Sulphinic Acids, Esters and their Derivatives; Patai, S., Ed.; J. Wiley & Sons: New York, 1999; Chapter 14, pp 453-474.

(28) Rajanbabu, T. V. J. Org. Chem. 1984, 49, 5080.
 (29) Basso, E. A.; Kaiser, C.; Rittner, R.; Lambert, J. B. J. Org.

Chem. 1993, 58, 1066. (30) Vedejs, E. V.; Pnbish, J. R. J. Org. Chem. 1988, 53, 1593.

<sup>(11) (</sup>a) Meerwein, H.; Dittmar, G.; Göllner, R.; Hafner, K.; Mensch, Steifort, O. Chem. Ber. 1957, 90, 841. (b) Chapman, N. B.; Clark, K.; Sawhney, S. N. J. Chem. Soc. 1968, 518.

<sup>(24)</sup> For examples of sulfonamide multistep syntheses see: (a) Gallant, M.; Carriere, M. C.; Chateauneuf, A.; Denis, D.; Gareau, G. Y. C.; Greig, G.; Juteau, H.; Lachance, N.; Lacombe, P.; Lamontagne, S.; Metters, K. M.; Rochette, C.; Ruel, R.; Slipetz, D.; Sawyer, N.;
 Tremblay, N.; Labelle, M. *Biorg. Med. Chem. Lett.* 2002, *12*, 2583. (b)
 Song, X.; He, T. J.; Siahaan, H. T. *Org. Lett.* 2002, *4*, 549. (c) Katritzky,
 A. R.; Yao, Y.; Denisko, O. V. *J. Org. Chem.* 2000, *65*, 8063.

TABLE 1.One-Pot, Three-Component Synthesis ofSulfonamides 5–18 and Sulfonic Esters 19–22

entry	enoxysilanes	products	product No	yield
1	2a	Meo S NEt2	5	65
2	2a	MeO NH(Bn)	6	70
3	2a	MeO S N(Me)(Bn)	7	65
4	2a	Meo C S N O O O	8	61
5	2b	NEt <sub>2</sub>	9	88
6	2c	Ph NEt <sub>2</sub>	10	67
7	2c	Ph S NH(Bn)	11	65
8	2c	Ph S N(Me)(Bn)	12	72
9	2c	Ph C S P C C	13	57
10	2d	O O O Š×NEt <sub>2</sub>	14	traces (67) <sup>a</sup>
11	2e	H SO <sub>2</sub> NEt <sub>2</sub>	15	69
12	2e		16	60
13	2e	H SO <sub>2</sub> N(Me)(Bn)	17	75
14	2e		18	58
15	2c	Ph S O	19	68 <sup>b</sup>
16	2c	Ph S OEt	20	65
17	2c	Ph S o	21	30
18	2c	Ph S OPh	22	35
<sup>a</sup> Pyridine as base. <sup>b</sup> Without base.				

## SCHEME 2. One-Pot, Three-Component Syntheses of Allyl Sulfonamides and Allyl Sulfonic Esters



SiOSO<sub>2</sub>CF<sub>3</sub> does not freeze at -78 °C. The excess of SO<sub>2</sub> and the solvent were evaporated from -78 °C to room temperature in vacuo before the oxidative workup at -20 °C with Cl<sub>2</sub> (or NCS) or Br<sub>2</sub> (or NBS) and subsequently treated with primary and secondary amines such as Et<sub>2</sub>NH, BnNH<sub>2</sub>, Bn(Me)NH, and *L*-*t*-BuOOC-CH(Me)NH<sub>2</sub> to give the corresponding sulfonamides **14**–**18** (entries 10–14, Table 1),<sup>31</sup> or with alcohols such as MeOH, EtOH, *i*-PrOH, or phenol to provide sulfonic esters **19–22** (entries 15–18, Table 1).

Product from the ene reaction of SO<sub>2</sub> with (2-methylpropen-1yl)trimethylsilane<sup>20</sup> **23** also underwent halogenolysis with NCS and subsequent reaction with  $Et_2NH$ , BnNH<sub>2</sub> to give the corresponding sulfonamides **24** and **25** (Scheme 2), or with alcohols such as MeOH to give the methyl sulfonate **26** (Scheme 2).

The structures of compounds 5-26 in Table 1 and Scheme 2 were established by their spectral data (see Supporting Information).

In the case of the camphor-derived sulfonamide **15**, its <sup>1</sup>H NMR spectra confirmed the endo configuration of the sulfonamide moiety as it showed relatively large vicinal coupling constants <sup>3</sup>J(H-3,H-4) = 4.0 Hz<sup>32</sup> as well as <sup>4</sup>J(H-3,H-5*exo*) = 1.4 Hz.<sup>32</sup> In the presence of MeONa in CD<sub>3</sub>-OD, sulfonamide **15** was rapidly equilibrated with the deuterated **D**-**15** (Scheme 3). Subsequent addition of H<sub>2</sub>O to the above CD<sub>3</sub>OD/CD<sub>3</sub>ONa solution of **D**-**15** led to the recovery of **15** showing that the exchange occurs without epimerization and that the endo configuration of the sulfonamide corresponds to the product of thermodynamic control. Thus, steric repulsions between the Me-C(7) and the *exo*-SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup> group make the exo sulfonamides less stable than their endo stereomers.

Interestingly, when running the ene reaction  $SO_2 + 2e$  in a NMR tube, the <sup>1</sup>H NMR spectrum of the intermediate silyl sulfinate **3e** showed <sup>3</sup>*J*(H-3,H-4) = 1.1 Hz consistent with an exo configurated<sup>32</sup> sulfinic ester (Scheme 3). Halogenation of **3e** with either Br<sub>2</sub> or NBS gave **27**, the <sup>1</sup>H NMR spectrum of which showed <sup>3</sup>*J* (H-

<sup>(31)</sup> For other approaches to the synthesis of N-alkyl-2-oxo-alkane-sulfonamides see: (a) Leclerc, M.; Brienne, M.-J. Tetrahedron Lett. **1990**, 31, 3875. (b) Vega, J. A.; Molina, R.; Alajarin, R.; Vaguero, J. J.; Garcia-Navio, J. L.; Alvarez-Builla, J. Tetrahedron Lett. **1992**, 33, 3677. (c) Vega, J. A.; Majarin, R.; Vaguero, J. J.; Garcia-Navio, J. L.; Alvarez-Builla, J. Tetrahedron Lett. **1993**, 54, 3589. (d) Hendrickson, J. B.; Bergeron, R. Tetrahedron Lett. **1970**, 5, 345. (e) Truce, W. E.; Veiesen, C. W. J. Am. Chem. Soc. **1953**, 75, 2525. (f) Bender, A.; Günther, D.; Willims, L.; Winger, R. Synthesis **1985**, 66. (g) Bender, A.; Günther, D.; Winger, R. Liebigs Ann. Chem. **1985**, 579. (h) Thompson, M. E. Synth. Commun. **1988**, 733.

<sup>(32)</sup> Joela, H. Org. Magn. Reson. 1977, 9, 338 and references cited therein.

# SCHEME 3. Kinetic versus Thermodynamic Control



4, H-3) = 4.0 Hz. This demonstrates that the SO<sub>2</sub>Br moiety in **27** occupies the endo position. The acidity of the reaction medium is thus capable of inducing exo  $\rightarrow$  endo isomerization of the sulfonyl moiety. To confirm the exo configuration of silyl sulfinate **3e**, we treated it with Bu<sub>4</sub>NF. This generated the corresponding tetrabutylammonium sulfinate, which was then reacted with 1-bromobutan-2-one giving the exo sulfone **28**. This compound was isolated in 92% yield and was fully characterized by its spectral data, including 2D NOESY <sup>1</sup>H NMR.

The formation of the exo silyl sulfinate **3e** demonstrates that the exo face of enoxysilane **2e** is preferred for the ene-reaction although the exo silyl sulfinate **3e** must be destabilized by steric factors as for the corresponding sulfonyl bromide and sulfonamides. We attribute the kinetic exo selectivity of the ene reaction of **2e** to the nonplanarity of the bicyclo[2.2.1]hept-2-ene alkene moiety that polarizes its  $\pi$ -electrons toward the exo face.<sup>33</sup>

**Diastereoselectivity of the Ene Reaction of Sulfur Dioxide.** In the case of enoxysilane **2d**, the formation of the cyclohexanone-derived sulfinate **3d** was monitored by <sup>1</sup>H and <sup>13</sup>C NMR. When running the reaction in CD<sub>3</sub>CN in the presence of a catalytical amount of the acid promoter (TBSOTf, 0.05 equiv) and an excess of SO<sub>2</sub> (15 equiv), the corresponding sulfinate **3d** was formed after 1 h at -60 °C and, to our surprise, only one single diastereoisomer was observed.

We decided to perform a second experiment in which the Lewis acid was suppressed and  $SO_2$  was used as cosolvent (3:1,  $SO_2/CD_3CN$ ). In this case, after 2 h at -40 °C, 75% of the starting material had been converted into a 6:4 mixture of two diastereoisomers. After 36 h at room temperature, the reaction was complete and the ratio of

### SCHEME 4. Diastereoselectivity of the Ene Reaction of SO<sub>2</sub> with Enoxysilanes



SCHEME 5. C-C Bond Formation through Umpolung with Sulfur Dioxide: Four-Component Synthesis of Sulfones



the two diastereoisomeric sulfinates was ca. 5:4. This experiment demonstrates that in acetonitrile, even if it takes a longer time, the ene reaction of  $SO_2$  may occur without promotion by a strong (and destructive) Lewis acid. To exemplify this feature, we performed the ene reaction of  $SO_2$  with enoxysilanes **2f** and **2g** without Lewis acid.

In addition, the same observations were made with the camphor-derived enoxysilane 2e. When running the reaction in CD<sub>3</sub>CN, in the presence of a catalytical amount of Lewis acid (TBSOTf, 0.05 equiv) and an excess of SO<sub>2</sub> (15 equiv), the formation of the corresponding sulfinate 3e was complete after 20 min at -78 °C and a 95:5 mixture of two diastereoisomeric sulfinates ((RSs))-3e and (RRs)-3e) was obtained. The 95:5 product ratio did not change on longer standing at -78 °C or on increasing the temperature (decomposition above 20 °C). A second experiment without TBSOTf was performed with SO<sub>2</sub> + 2e. After 2 h at -40 °C, 2e was fully converted into a 95:5 mixture of two diastereoisomeric sulfinates. The product ratio changed from 95:5 to 75:25 after 6 h at -40 °C and finally reached 1:1 after 12 h at this same temperature and stayed unchanged after staying longer at -40 °C. Obviously, in both cases, 2d and 2e, the

<sup>(33)</sup> Jones, G. R.; Caldarelli, S.; Vogel, P. *Helv. Chim. Acta* **1997**, *80*, 59 and references therein.



#### **SCHEME 6.** Example of One-Pot, Four-Component Syntheses of Sulfonamides and Sulfonic Esters



diastereoisomers of **3d** and **3e** arise from the chirality of the sulfur atom in the silvl sulfinates (Scheme 4).

Our experiments demonstrate that the ene reaction of SO<sub>2</sub> with **2e** occurs under kinetic control giving a major silyl sulfinate. It then equilibrates with the thermodynamic diastereoisomeric sulfinate ((*RSs*)-**3e**  $\leftrightarrow$  (*RRs*)-**3e**). Both diastereoisomeric (*RSs*)-**3e** and (*RRs*)-**3e** have *exo*-SO<sub>2</sub>SiMe<sub>3</sub> groups. This isomerization probably implies the migration of the silyl group from one oxygen to the other of the SO<sub>2</sub>SiMe<sub>3</sub> moiety. The same hypotheses can be retained for the reaction of SO<sub>2</sub> with monocyclic enol ether **2d**. Distinction between diastereomeric sulfinate pairs (*Ss*)-**3d** and (*Rs*)-**3d**, and (*RSs*)-**3e** and (*RRs*)-**3e** cannot be possible (Scheme 4). The relative structures of sulfinates **3d** and **3e** were deduced from their spectral data (see Supporting Information).

**One-Pot, Four-Component Synthesis of Sulfonamides and Sulfonic Esters.** Recently we reported<sup>21,22</sup> a new C–C bond-forming reaction  $29 + 30 \rightarrow 31$  (Scheme 5), involving a cascade of reactions starting with the hetero-Diels–Alder addition of SO<sub>2</sub> to 1,3-dienyl ether **29**, giving the corresponding sultine **32**.<sup>34</sup> The latter is ionized into zwitterionic intermediate **33**, which then adds to enoxysilanes (oxyallylation) to give silyl sulfinate **34** that can be alkylated or allylated in situ providing the corresponding sulfones.<sup>21,22</sup> We have extended this reaction cascade to allylsilanes and to 1,3-dioxydienes.<sup>23</sup> We show now that the silyl sulfinates **34** and analogues obtained as intermediates in those reactions can be converted, as the simpler silyl sulfinates **3** and **23**, into polyfunctional sulfonamides or sulfonic esters.

As an illustration of the method we have combined (*E*)-1-methoxybutadiene (**35**) with enoxysilane **2c**, (*E*,*E*)-1benzyloxy-2-methylpenta-1,3-diene (**36**) with enoxysilane

<sup>(34) (</sup>a) Roversi, E.; Scoppelliti; R.; Solari, E.; Estoppey, R.; Vogel,
P.; Braña, P.; Menendez, B.; Sordo, J. A. *Chem. Eur. J.* 2002, *8*, 1336.
(b) Markovic, D.; Roversi, E.; Scoppelliti, R.; Vogel, P.; Meana, R.;
Sordo, J. A. *Chem. Eur. J.* 2003, *9*, 4911.

**2b**, (E,E)-1-benzyloxy-2-methylpenta-1,3-dien-3-yl acetate (**37**) with the trimethylsilyl enol ether of acetone (**38**), and racemic 1,3-dioxydienes **39a** and **39b** with allyltrimethylsilane (**40**) <sup>23</sup> in the presence of Lewis acid (LA). All the intermediate trimethylsilyl sulfinates (**41**-**44**) obtained in this way were converted into the corresponding sulfonyl chlorides by treatment with NCS at -20 °C in CH<sub>2</sub>Cl<sub>2</sub> and, without isolation, they were reacted by representative amines in pyridine at -20 °C giving sulfonamides (**45a**, **46a**, **b**, **47a**, **b**, **48a**, **b**). Alternatively, quenching with alcohol/pyridine mixtures provided the corresponding sulfonic esters, as exemplified with **45b** (Scheme 6).

### Conclusion

This work discloses efficient one-pot, three-component and four-component syntheses of sulfonamides and sulfonic esters. Polyfunctional systems are obtained under smooth reaction conditions and a high molecular diversity can be reached. Some of the keto-sulfonamides obtained can be seen as peptidomimetics. We have demonstrated also that the ene reaction of sulfur dioxide with enoxysilanes can be stereoselective as the hetero-Diels-Alder addition under conditions of kinetic control. Moreover, this kinetic selectivity is not controlled by the relative stability (Dimroth principle) of the two isomeric silyl sulfinate products of the pericyclic reaction. Investigations aiming at a further expansion of this new chemistry are actively pursued in our laboratory and will be reported in due course.

**Acknowledgment.** This project was supported by F. Hoffmann-La Roche Ltd (Roche Research Foundation Grant), by the Swiss National Science Foundation (Grant No. 200020-100002), and the Office Federal de l'Education et de la Sciences (OFES) (Grant No. COI. 0071, COST D13/010/01). We also thank Mr. Martial Rey and Franscisco Sepúlveda for technical help.

**Supporting Information Available:** Experimental procedures, unknown compounds characterization, and references to known compounds as well as all spectral data. This material is available free of charge via the Internet at http://pubs.acs.org.

JO049047J