

Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

An Expeditious Synthesis of Benzoxazine-2-Thione c-Nucleosides Via Cu(OTf)₂-Mediated Dehydrazinative β -Glycosylation

Lal Dhar S. Yadav^a & Vijai K. Rai^a

^a Green Synthesis Lab, Department of Chemistry, University of Allahabad, Allahabad, India

Published online: 26 Nov 2008.

To cite this article: Lal Dhar S. Yadav & Vijai K. Rai (2008) An Expeditious Synthesis of Benzoxazine-2-Thione c-Nucleosides Via Cu(OTf)₂-Mediated Dehydrazinative β -Glycosylation, Nucleosides, Nucleotides and Nucleic Acids, 27:12, 1227-1237, DOI: [10.1080/15257770802458212](https://doi.org/10.1080/15257770802458212)

To link to this article: <http://dx.doi.org/10.1080/15257770802458212>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

AN EXPEDITIOUS SYNTHESIS OF BENZOXAZINE-2-THIONE C-NUCLEOSIDES VIA $\text{Cu}(\text{OTf})_2$ -MEDIATED DEHYDRAZINATIVE β -GLYCOSYLATION

Lal Dhar S. Yadav and Vijai K. Rai

Green Synthesis Lab, Department of Chemistry, University of Allahabad, Allahabad, India

□ A novel expeditious synthetic protocol for 1,3-benzoxazine-2-thione C-nucleosides via $\text{Cu}(\text{OTf})_2$ -mediated dehydrazinative β -glycosylation of 4-hydrazino-2H-benz[e]-1,3-oxazine-2-thiones with unprotected D-ribose is reported.

Keywords 1,3-Benzoxazine-2-thione C-nucleosides; unprotected D-ribose; non-nucleoside reverse transcriptase inhibitor (NNRTI)

INTRODUCTION

Efavirenz (Sustiva), a benzoxazinone derivative, is a non-nucleoside reverse transcriptase inhibitor (NNRTI) that has been approved by the U.S. Food and Drug administration (FDA) and is presently in clinical use for the treatment of AIDS. The strategy for the fight against HIV by developing more efficacious drugs than Efavirenz has been the prime driving force for ingenious benzoxazinone derivatizations.^[1–6] Moreover, our interest in the family of C-nucleosides also resides with the naturally occurring antibiotics. Several of these have potent antiviral, anticancer and antitumour activity and, as a result of C-C glycosidic linkage, they are resistant to enzymatic hydrolysis which is the major factor in the degradation of N-nucleosides.^[7]

The formation of C-C bond at the anomeric carbon, especially C-glycosylation of bioactive heterocycles, is a field of increasing interest in synthetic organic chemistry. The most common methods for the C-C bond formation at the anomeric centre involve nucleophilic attack on the

Received 8 July 2008; accepted 20 August 2008.

We sincerely thank the Sophisticated Analytical Instrument Facility (SAIF), Punjab University, Chandigarh, for providing microanalyses and spectra.

Address correspondence to Lal Dhar S. Yadav, Green Synthesis Lab, Department of Chemistry, University of Allahabad, Allahabad-211 002, U. P., India. E-mail: ldsyadav@hotmail.com

intrinsically electrophilic anomeric carbons.^[8–11] In most cases, these procedures require specific, awkward reaction conditions and generally suffer from low yields. Although microwave (MW)-assisted heterocyclisations have gained significant interest for over a decade,^[12,13] the first example of MW-assisted solvent-free synthesis of nucleoside analogues was reported very recently.^[14]

Considering the above points and in pursuing our work on the development of organic syntheses in solvent-free MW-assisted conditions,^[15–19] we present herein the first report on Cu(OTf)₂-mediated dehydrazinative β -glycosylation yielding benzoxazine-2-thione *C*-nucleosides. The key element in our approach is the novel utilization of salicylaldehyde as a bifunctional building block whose application to the construction of various benzo-fused oxygen heterocycles of chemical and biological interest is well documented.^[20–23]

RESULTS AND DISCUSSION

1,3-Oxazine-2-thiones **4** were synthesized in 78–89% yields by MW irradiation of an intimate solvent-free mixture of salicylaldehydes **1**, thiosemicarbazide **2**, and Cu(OTf)₂ (0.30 equivalent) in an open vessel under air. The reaction proceeds via domino cycloisomerization of in situ formed *N*-phenylthiosemicarbazones **3** to 4-hydrazino-1,3-oxazine-2-thiones **3'** and dehydrazination of compounds **3'** to target oxazines **4** (Scheme 1). Here, Cu(OTf)₂ plays a dual role of Lewis acid and oxidant. Only 0.30 equivalent of Cu(OTf)₂ is sufficient to complete the reaction because Cu(I) formed is oxidized to Cu(II) by air under the reaction conditions and a redox cycle is established. A plausible mechanism of dehydrazination of **3'** into **4** using Cu(II) is depicted in Scheme 2. Although the present dehydrazinative cyclization to afford 1,3-oxazine-2-thione **4** is our new finding, similar dehydrazination reactions using Cu(II) have already been reported in the literature.^[24–27]

In our initial experiments for the synthesis of compounds **4**, we used CeCl₃·7H₂O/NaI as Lewis acid catalyst but compounds **3'** rather than compounds **4** were isolated in 73–87% yields. Compounds **3'** on further treatment with copper(II) sulfate on alumina support afforded the target oxazines **4** in 55–67% yields. In order to improve the yields and synthesize target 1,3-oxazine-2-thiones **4** expeditiously from salicylaldehydes **1** in a one-pot procedure, we relied upon the significant advantages of Cu(OTf)₂, which acts not only as an efficient Lewis acid catalyst for cycloisomerization of compounds **3** to **3'** but also as copper(II) source for the desired dehydrazination of 4-hydrazino-1,3-oxazine-2-thiones **3'** to afford **4**. Other mineral catalysts, viz. CeCl₃·7H₂O, K-10 clay, and acidic, neutral, or basic alumina were far less effective resulting in either no reaction (in the case of basic alumina) or relatively moderate (36–50%, in case of CeCl₃·7H₂O, K-10




Downloaded by [University of Memphis Libraries] at 03:18 15 April 2015


Downloaded by [University of Memphis Libraries] at 03:18 15 April 2015

Downloaded by [University of Memphis Libraries] at 03:18 15 April 2015

Downloaded by [University of Memphis Libraries] at 03:18 15 April 2015



Downloaded by [University of Memphis Libraries] at 03:18 15 April 2015



SCHEME 3 Tentative mechanism for β -glycosylation of 1,3-benzoxazine-2-thione 4

with appropriate reference data.^[30–32] In fact, selective irradiation of the H-1' signal of **9** significantly increased the intensity of the H-4' signal (2.0–2.8%), while no intensity enhancement of the H-3' signal was observed; this is consistent with the β -glycosidic linkage. Furthermore, the comparison of $J_{1',2'}$ values of **9** ($J_{1',2'} = 5.4$ – 5.8 Hz) with those of similar systems in the literature ($J_{1',2'} = 5.4$ – 5.9 Hz) also supported a β -glycosidic linkage.^[30–32] These analyses permit assignment of compounds **9** as the β -D-ribofuranoside anomer.

In summary, we have developed a novel expeditious synthetic protocol for benzoxazine-2-thione *C*-nucleosides via dehydrazinative β -glycosylation. Furthermore, the Knoevenagel-type condensation presented in this paper represents a very convenient method for the preparation of pure β -*C*-glycosides in one step directly from the unprotected sugar and a compound containing an activated methylene group.

EXPERIMENTAL

Melting points were determined by open glass capillary method and are uncorrected. IR spectra in KBr were recorded on a Perkin-Elmer 993 IR spectrophotometer. ¹H NMR spectra were recorded on a Bruker WM-40 C (400 MHz) FT spectrometer in DMSO-*d*₆ using TMS as internal reference. ¹³C NMR spectra were recorded on the same instrument at 100 MHz in DMSO-*d*₆ and TMS was used as internal reference. Mass (EI) spectra

were recorded on a JEOL D-300 mass spectrometer. Elemental analyses were carried out in a Coleman automatic carbon, hydrogen and nitrogen analyzer. A Chemical Laboratory Microwave Oven (Model; BP-310/50, 230 volt, 50 Hz power input) was used for all experiments. All chemicals used were reagent grade and were used as received without further purification. Silica gel-G was used for TLC.

3-Aryl-2H-benz[e]-1,3-oxazine-2-thiones **4**; General Procedure

Thoroughly mixed salicylaldehyde **1** (when $R^1 = R^2 = H$; 0.12 g, 1.00 mmol), *N*-arylthiosemicarbazide **2** (when Ar = 4-ClC₆H₄; 0.20 g, 1.00 mmol) and Cu(OTf)₂ (0.11 g, 0.30 mmol) were taken in a 20 mL open vial and subjected to MW irradiation for 7–12 minutes at 90°C under air. After completion of the reaction as indicated by TLC (hexane-AcOEt, 8:2, v/v), the product was extracted with dichloromethane (3 × 25 mL), the extract was filtered and the filtrate was evaporated under reduced pressure to leave the crude product, which was recrystallized from ethanol to obtain an analytically pure sample of **4** as yellowish needles.

4a: yellowish needles; m.p. 170–172°C. IR (KBr): ν 3011, 1595, 1585, 1450 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 6.56 (d, 1H, $J = 13$ Hz, axial H of CH₂), 6.63 (d, 1H, $J = 13$ Hz, equatorial H of CH₂), 6.71–7.09 (m, 6H_{arom}), 7.28–7.35 (m, 2H_{arom}). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 64.5, 113.1, 114.5, 118.2, 120.5, 122.7, 128.5, 129.6, 130.5, 150.2, 166.2, 192.3. EIMS $m/z = 275, 277$ (M⁺, M⁺ + 2). Anal. calcd for C₁₄H₁₀ClNOS: C, 60.98; H, 3.66; N, 5.08. Found: C, 60.67; H, 3.91; N, 5.26.

4b: yellowish needles; m.p. 159–161°C. IR (KBr): ν 3010, 1600, 1585, 1455 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 6.53 (d, 1H, $J = 13$ Hz, axial H of CH₂), 6.65 (d, 1H, $J = 13$ Hz, equatorial H of CH₂), 6.69–7.13 (m, 3H_{arom}), 7.19–7.41 (m, 4H_{arom}). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 64.2, 113.5, 114.7, 118.8, 120.6, 122.5, 128.1, 129.7, 130.2, 150.1, 166.4, 192.5. EIMS $m/z = 355$ (M⁺). Anal. calcd for C₁₄H₉BrClNOS: C, 47.41; H, 2.56; N, 3.59. Found: C, 47.68; H, 2.71; N, 3.45.

4c: yellowish needles; m.p. 143–145°C. IR (KBr): ν 3012, 1601, 1581, 1456 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 6.55 (d, 1H, $J = 13$ Hz, axial H of CH₂), 6.64 (d, 1H, $J = 13$ Hz, equatorial H of CH₂), 6.68–7.05 (m, 2H_{arom}), 7.21–7.48 (m, 4H_{arom}). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 64.2, 113.2, 114.1, 118.5, 120.2, 121.9, 128.2, 129.5, 130.3, 150.3, 166.2, 192.1. EIMS $m/z = 435$ (M⁺). Anal. calcd for C₁₄H₈Br₂ClNOS: C, 38.78; H, 1.86; N, 3.23. Found: C, 38.44; H, 2.07; N, 2.91.

4d: yellowish needles; m.p. 149–151°C. IR (KBr): ν 3011, 1598, 1580, 1455 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 6.54 (d, 1H, $J = 13$ Hz, axial H of CH₂), 6.66 (d, 1H, $J = 13$ Hz, equatorial H of CH₂), 6.75–7.11 (m, 3H_{arom}), 7.19–7.39 (m, 4H_{arom}). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 64.5, 113.0, 114.2, 118.4, 120.5, 122.2, 128.5, 129.2, 130.1, 150.2, 166.0, 192.2.

EIMS $m/z = 309, 311$ (M^+, M^++2). Anal. calcd for $C_{14}H_9Cl_2NOS$: C, 54.21; H, 2.92; N, 4.52. Found: C, 54.49; H, 2.78; N, 4.35.

4e. yellowish needles; m.p. 183–185°C. IR (KBr): ν 3010, 1605, 1581, 1455 cm^{-1} . 1H NMR (400 MHz, $DMSO-d_6$): δ 6.59 (d, 1H, $J = 13$ Hz, axial H of CH_2), 6.67 (d, 1H, $J = 13$ Hz, equatorial H of CH_2), 6.75–7.12 (m, $2H_{arom}$), 7.31–7.39 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, $DMSO-d_6$): δ 64.3, 113.0, 114.4, 118.8, 120.2, 122.9, 129.0, 130.3, 135.5, 151.1, 166.5, 192.1. EIMS $m/z = 343, 345$ (M^+, M^++2). Anal. calcd for $C_{14}H_8Cl_3NOS$: C, 48.79; H, 2.34; N, 4.06. Found: C, 49.03; H, 2.49; N, 4.31.

4f: yellowish needles; m.p. 170–180°C. IR (KBr): ν 3010, 1603, 1587, 1450 cm^{-1} . 1H NMR (400 MHz, $DMSO-d_6$): δ 3.73 (s, 3H, OMe), 6.52 (d, 1H, $J = 13$ Hz, axial H of CH_2), 6.68 (d, 1H, $J = 13$ Hz, equatorial H of CH_2), 6.70–7.11 (m, $6H_{arom}$), 7.18–7.35 (m, $2H_{arom}$). ^{13}C NMR (100 MHz, $DMSO-d_6$): δ 53.5, 64.0, 113.2, 114.7, 118.5, 120.2, 122.7, 128.3, 129.5, 130.3, 150.5, 166.3, 192.3. EIMS $m/z = 273$ (M^+). Anal. calcd for $C_{15}H_{13}NO_2S$: C, 66.40; H, 4.83; N, 5.16. Found: C, 66.69; H, 4.58; N, 5.27.

4g: yellowish needles; m.p. 155–157°C. IR (KBr): ν 3009, 1595, 1583, 1452 cm^{-1} . 1H NMR (400 MHz, $DMSO-d_6$): δ 3.76 (s, 3H, OMe), 6.53 (d, 1H, $J = 13$ Hz, axial H of CH_2), 6.67 (d, 1H, $J = 13$ Hz, equatorial H of CH_2), 6.75–7.11 (m, $3H_{arom}$), 7.28–7.42 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, $DMSO-d_6$): δ 53.1, 64.3, 113.1, 114.0, 118.3, 120.1, 122.3, 128.2, 129.2, 130.5, 150.2, 166.4, 192.5. EIMS $m/z = 351$ (M^+). Anal. calcd for $C_{15}H_{12}BrNO_2S$: C, 51.44; H, 3.45; N, 4.00. Found: C, 51.19; H, 3.61; N, 3.85.

4h: yellowish needles; m.p. 138–140°C. IR (KBr): ν 3012, 1599, 1588, 1455 cm^{-1} . 1H NMR (400 MHz, $DMSO-d_6$): δ 3.75 (s, 3H, OMe), 6.51 (d, 1H, $J = 13$ Hz, axial H of CH_2), 6.65 (d, 1H, $J = 13$ Hz, equatorial H of CH_2), 6.78–7.15 (m, $2H_{arom}$), 7.21–7.33 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, $DMSO-d_6$): δ 53.3, 64.2, 113.3, 114.3, 118.2, 120.4, 122.2, 128.5, 129.3, 130.2, 150.5, 166.1, 192.2. EIMS $m/z = 331$ (M^+). Anal. calcd for $C_{15}H_{11}Br_2NO_2S$: C, 41.98; H, 2.58; N, 3.26. Found: C, 42.32; H, 2.89; N, 3.39.

4i: yellowish needles; m.p. 163–165°C. IR (KBr): ν 3010, 1605, 1579, 1449 cm^{-1} . 1H NMR (400 MHz, $DMSO-d_6$): δ 3.76 (s, 3H, OMe), 6.52 (d, 1H, $J = 13$ Hz, axial H of CH_2), 6.68 (d, 1H, $J = 13$ Hz, equatorial H of CH_2), 6.71–7.09 (m, $3H_{arom}$), 7.15–7.34 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, $DMSO-d_6$): δ 53.8, 64.4, 113.5, 114.6, 118.7, 120.2, 122.5, 128.1, 129.2, 130.1, 150.8, 166.7, 192.3. EIMS $m/z = 307$ (M^+). Anal. calcd for $C_{15}H_{12}ClNO_2S$: C, 58.92; H, 3.96; N, 4.58. Found: C, 58.59; H, 3.67; N, 4.87.

4j: yellowish needles; m.p. 134–136°C. IR (KBr): ν 3010, 1600, 1582, 1452 cm^{-1} . 1H NMR (400 MHz, $DMSO-d_6$): δ 3.72 (s, 3H, OMe), 6.51 (d, 1H, $J = 13$ Hz, axial H of CH_2), 6.66 (d, 1H, $J = 13$ Hz, equatorial H of CH_2), 6.79–7.11 (m, $2H_{arom}$), 7.28–7.39 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, $DMSO-d_6$): δ 53.5, 64.3, 113.2, 114.5, 118.6, 120.4, 122.4, 128.2, 129.5, 130.7, 150.6, 166.3, 192.7. EIMS $m/z = 341$ (M^+). Anal. calcd for $C_{15}H_{11}Cl_2NO_2S$: C, 52.95; H, 3.26; N, 4.12. Found: C, 52.71; H, 3.49; N, 3.95.

3-Aryl-4- β -D-ribofuranosyl-3,4-dihydro-2H-benz[e]-1,3-oxazine-2-thiones **9; General Procedure**

To a suspension of 1,3-benzoxazine-2-thione **4** (when Ar = 4-ClC₆H₄, R¹ = R² = H; 0.28 g, 1.00 mmol) in THF (10 mL) was added slowly a suspension of sodium hydride (0.12 g, 5.00 mmol) in THF (10 mL) with stirring at room temperature. After the addition was completed and the evolution of hydrogen gas (effervescence) had ceased, the reaction mixture was stirred at 60°C for 45 minutes and then cooled to room temperature. Next, D-ribose (0.30 g, 2.00 mmol) was added to it and further stirred for 4–5 hours at 60°C. After completion of the reaction as indicated by TLC (hexane/AcOEt, 9:1, v/v), glacial acetic acid (2.0 mL) was added followed by water (10 mL) and the product was extracted with dichloromethane (3 \times 25 mL). The extract was filtered and the filtrate was evaporated under reduced pressure to leave the crude product, which was recrystallized from ethanol to obtain an analytically pure sample of **9** as yellowish powder.

9a. Yellowish powder; m.p. 229–231°C. $[\alpha]_D^{[27]} + 46.4^\circ$ (c 1.5, MeOH). IR (KBr): ν 3391, 3013, 1605, 1585, 1456 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.53 (ddd, 1H, *J* = 3.4, 5.3, 11.6 Hz, H_a-5'), 3.61 (ddd, 1H, *J* = 3.4, 5.7, 11.6 Hz, H_b-5'), 4.05 (ddd, 1H, *J* = 4.2, 3.5, 3.4 Hz, H-4'), 4.37 (dd, 1H, *J* = 5.6, 6.4 Hz, H-1'), 4.49 (ddd, 1H, *J* = 4.6, 5.6, 6.8 Hz, H-2'), 4.67 (ddd, 1H, *J* = 4.6, 5.2, 6.8 Hz, H-3'), 5.15 (dd, 1H, *J* = 5.3, 5.7 Hz, OH-5'), 5.31 (d, 1H, *J* = 5.2 Hz, OH-3'), 5.58 (d, 1H, *J* = 5.6 Hz, OH-2'), 6.55 (d, 1H, *J* = 6.4 Hz, H-4), 6.75–7.05 (m, 6H_{arom}), 7.21–7.38 (m, 2H_{arom}). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 37.5, 62.2, 72.8, 73.3, 76.6, 78.3, 83.5, 113.2, 114.5, 119.0, 121.5, 122.9, 130.3, 132.0, 151.8, 166.5, 192.2. MS *m/z* (FAB, NBA) 408 [MH⁺]. Anal. calcd for C₁₉H₁₈ClNO₅S: C, 55.95; H, 4.45; N, 3.43. Found: C, 56.21; H, 4.69; N, 3.27.

9b. Yellowish powder; m.p. 205–207°C. $[\alpha]_D^{[27]} + 46.8^\circ$ (c 1.5, MeOH). IR (KBr): ν 3390, 3009, 1602, 1581, 1450 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.55 (ddd, 1H, *J* = 3.4, 5.3, 11.6 Hz, H_a-5'), 3.60 (ddd, 1H, *J* = 3.4, 5.7, 11.6 Hz, H_b-5'), 4.08 (ddd, 1H, *J* = 4.2, 3.5, 3.4 Hz, H-4'), 4.34 (dd, 1H, *J* = 5.6, 6.4 Hz, H-1'), 4.47 (ddd, 1H, *J* = 4.6, 5.6, 6.8 Hz, H-2'), 4.68 (ddd, 1H, *J* = 4.6, 5.2, 6.8 Hz, H-3'), 5.18 (dd, 1H, *J* = 5.3, 5.7 Hz, OH-5'), 5.30 (d, 1H, *J* = 5.2 Hz, OH-3'), 5.55 (d, 1H, *J* = 5.6 Hz, OH-2'), 6.56 (d, 1H, *J* = 6.4 Hz, H-4), 6.69–7.08 (m, 3H_{arom}), 7.18–7.35 (m, 4H_{arom}). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 38.2, 62.5, 72.8, 73.5, 76.6, 78.7, 83.8, 113.2, 114.9, 119.9, 121.2, 122.5, 130.1, 131.5, 152.0, 166.5, 192.3. MS *m/z* (FAB, NBA) 486 [MH⁺]. Anal. calcd for C₁₉H₁₇BrClNO₅S: C, 46.88; H, 3.52; N, 2.88. Found: C, 46.52; H, 3.77; N, 3.02.

9c. Yellowish powder; m.p. 195–197°C. $[\alpha]_D^{[27]} + 43.9^\circ$ (c 1.5, MeOH). IR (KBr): ν 3390, 3011, 1603, 1586, 1449 cm⁻¹. ¹H NMR (400 MHz,

DMSO- d_6): δ 3.52 (ddd, 1H, $J = 3.4, 5.3, 11.6$ Hz, H_a-5'), 3.64 (ddd, 1H, $J = 3.4, 5.7, 11.6$ Hz, H_b-5'), 4.09 (ddd, 1H, $J = 4.2, 3.5, 3.4$ Hz, $H-4'$), 4.33 (dd, 1H, $J = 5.6, 6.5$ Hz, $H-1'$), 4.48 (ddd, 1H, $J = 4.6, 5.6, 6.8$ Hz, $H-2'$), 4.65 (ddd, 1H, $J = 4.6, 5.2, 6.8$ Hz, $H-3'$), 5.21 (dd, 1H, $J = 5.3, 5.7$ Hz, $OH-5'$), 5.29 (d, 1H, $J = 5.2$ Hz, $OH-3'$), 5.56 (d, 1H, $J = 5.6$ Hz, $OH-2'$), 6.53 (d, 1H, $J = 6.5$ Hz, $H-4$), 6.71–7.05 (m, $2H_{arom}$), 7.21–7.35 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, DMSO- d_6): δ 38.0, 62.1, 72.5, 73.4, 76.3, 78.5, 83.5, 113.1, 114.5, 119.7, 121.1, 122.2, 130.7, 131.2, 152.9, 166.2, 192.1. MS m/z (FAB, NBA) 566 [MH^+]. Anal. calcd for $C_{19}H_{16}Br_2ClNO_5S$: C, 40.34; H, 2.85; N, 2.48. Found: C, 40.59; H, 2.99; N, 2.31.

9d. Yellowish powder; m.p. 211–213°C. $[\alpha]_D^{[27]} + 44.4^\circ$ (c 1.5, MeOH). IR (KBr): ν 3389, 3010, 1601, 1583, 1452 cm^{-1} . 1H NMR (400 MHz, DMSO- d_6): δ 3.56 (ddd, 1H, $J = 3.4, 5.3, 11.6$ Hz, H_a-5'), 3.62 (ddd, 1H, $J = 3.4, 5.7, 11.6$ Hz, H_b-5'), 4.07 (ddd, 1H, $J = 4.2, 3.5, 3.4$ Hz, $H-4'$), 4.35 (dd, 1H, $J = 5.5, 6.4$ Hz, $H-1'$), 4.51 (ddd, 1H, $J = 4.6, 5.5, 6.8$ Hz, $H-2'$), 4.67 (ddd, 1H, $J = 4.6, 5.2, 6.8$ Hz, $H-3'$), 5.19 (dd, 1H, $J = 5.3, 5.7$ Hz, $OH-5'$), 5.33 (d, 1H, $J = 5.2$ Hz, $OH-3'$), 5.59 (d, 1H, $J = 5.6$ Hz, $OH-2'$), 6.55 (d, 1H, $J = 6.4$ Hz, $H-4$), 6.70–7.12 (m, $3H_{arom}$), 7.19–7.35 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, DMSO- d_6): δ 38.5, 62.5, 72.2, 73.1, 76.2, 78.2, 83.7, 113.5, 114.3, 120.2, 121.5, 122.7, 129.9, 131.3, 152.5, 166.1, 192.5. MS m/z (FAB, NBA) 444 [MH^+]. Anal. calcd for $C_{19}H_{17}Cl_2NO_5S$: C, 51.59; H, 3.87; N, 3.17. Found: C, 51.32; H, 3.65; N, 3.39.

9e. Yellowish powder; m.p. 256–258°C. $[\alpha]_D^{[27]} + 46.9^\circ$ (c 1.5, MeOH). IR (KBr): ν 3393, 3010, 1588, 1455 cm^{-1} . δ 3.48 (ddd, 1H, $J = 3.7, 5.4, 11.6$ Hz, H_a-5'), 3.65 (ddd, 1H, $J = 3.7, 5.8, 11.6$ Hz, H_b-5'), 4.08 (ddd, 1H, $J = 4.1, 3.7, 3.7$ Hz, $H-4'$), 4.49 (dd, 1H, $J = 5.5, 6.4$ Hz, $H-1'$), 4.56 (ddd, 1H, $J = 4.5, 5.5, 6.9$ Hz, $H-2'$), 4.74 (ddd, 1H, $J = 4.1, 5.1, 6.9$ Hz, $H-3'$), 5.16 (dd, 1H, $J = 5.4, 5.8$ Hz, $OH-5'$), 5.32 (d, 1H, $J = 5.1$ Hz, $OH-3'$), 5.61 (d, 1H, $J = 5.5$ Hz, $OH-2'$), 6.58 (d, 1H, $J = 6.4$ Hz, $H-4$), 6.82–7.01 (m, $2H_{arom}$), 7.29–7.43 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, DMSO- d_6): δ 38.1, 62.5, 72.5, 73.8, 76.5, 78.4, 83.7, 113.5, 114.8, 119.8, 120.6, 122.7, 130.1, 131.6, 152.5, 166.7, 192.5. MS m/z (FAB, NBA) 475 [MH^+]. Anal. calcd for $C_{19}H_{16}Cl_3NO_5S$: C, 47.87; H, 3.38; N, 2.94. Found: C, 47.58; H, 3.53; N, 3.09.

9f. Yellowish powder; m.p. 221–223°C. $[\alpha]_D^{[27]} + 47.8^\circ$ (c 1.5, MeOH). IR (KBr): ν 3391, 3012, 1607, 1580, 1457 cm^{-1} . 1H NMR (400 MHz, DMSO- d_6): δ 3.53 (ddd, 1H, $J = 3.4, 5.3, 11.6$ Hz, H_a-5'), 3.61 (ddd, 1H, $J = 3.4, 5.7, 11.6$ Hz, H_b-5'), 3.75 (s, 3H, OMe), 4.06 (ddd, 1H, $J = 4.2, 3.5, 3.4$ Hz, $H-4'$), 4.34 (dd, 1H, $J = 5.8, 6.4$ Hz, $H-1'$), 4.53 (ddd, 1H, $J = 4.6, 5.8, 6.8$ Hz, $H-2'$), 4.63 (ddd, 1H, $J = 4.6, 5.2, 6.8$ Hz, $H-3'$), 5.24 (dd, 1H, $J = 5.3, 5.7$ Hz, $OH-5'$), 5.32 (d, 1H, $J = 5.2$ Hz, $OH-3'$), 5.54 (d, 1H, $J = 5.6$ Hz, $OH-2'$), 6.57 (d, 1H, $J = 6.4$ Hz, $H-4$), 6.68–7.13 (m, $6H_{arom}$), 7.22–7.41 (m, $2H_{arom}$). ^{13}C NMR (100 MHz, DMSO- d_6): δ 38.1, 53.5, 62.2, 72.6, 73.6, 76.5, 78.3, 83.5, 113.2, 114.5, 120.1, 121.3, 122.5, 130.3, 131.0, 152.3, 166.7, 192.2.

MS m/z (FAB, NBA) 404 $[MH^+]$. Anal. calcd for $C_{20}H_{21}NO_6S$: C, 59.54; H, 5.25; N, 3.47. Found: C, 59.79; H, 5.58; N, 3.31.

9g. Yellowish powder; m.p. 238–240°C. $[\alpha]_D^{[27]} + 42.3^\circ$ (c 1.5, MeOH). IR (KBr): ν 3390, 3011, 1601, 1583, 1449 cm^{-1} . 1H NMR (400 MHz, DMSO- d_6): δ 3.57 (ddd, 1H, $J = 3.4, 5.3, 11.6$ Hz, H_a-5'), 3.63 (ddd, 1H, $J = 3.4, 5.7, 11.6$ Hz, H_b-5'), 3.74 (s, 3H, OMe), 4.05 (ddd, 1H, $J = 4.2, 3.5, 3.4$ Hz, $H-4'$), 4.31 (dd, 1H, $J = 5.4, 6.3$ Hz, $H-1'$), 4.50 (ddd, 1H, $J = 4.6, 5.4, 6.8$ Hz, $H-2'$), 4.62 (ddd, 1H, $J = 4.6, 5.2, 6.8$ Hz, $H-3'$), 5.17 (dd, 1H, $J = 5.3, 5.7$ Hz, OH-5'), 5.31 (d, 1H, $J = 5.2$ Hz, OH-3'), 5.52 (d, 1H, $J = 5.6$ Hz, OH-2'), 6.51 (d, 1H, $J = 6.3$ Hz, H-4), 6.62–7.11 (m, $3H_{arom}$), 7.21–7.41 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, DMSO- d_6): δ 38.3, 53.9, 62.4, 72.5, 73.3, 76.1, 78.6, 83.3, 113.6, 114.2, 120.3, 121.2, 123.2, 130.2, 131.5, 152.9, 166.5, 192.4. MS m/z (FAB, NBA) 482 $[MH^+]$. Anal. calcd for $C_{20}H_{20}BrNO_6S$: C, 49.80; H, 4.18; N, 2.90. Found: C, 49.61; H, 4.05; N, 3.17.

9h. Yellowish powder; m.p. 248–250°C. $[\alpha]_D^{[27]} + 45.1^\circ$ (c 1.5, MeOH). IR (KBr): ν 3392, 3013, 1599, 1581, 1450 cm^{-1} . 1H NMR (400 MHz, DMSO- d_6): δ 3.54 (ddd, 1H, $J = 3.4, 5.3, 11.6$ Hz, H_a-5'), 3.61 (ddd, 1H, $J = 3.4, 5.7, 11.6$ Hz, H_b-5'), 3.79 (s, 3H, OMe), 4.01 (ddd, 1H, $J = 4.2, 3.5, 3.4$ Hz, $H-4'$), 4.32 (dd, 1H, $J = 5.6, 6.4$ Hz, $H-1'$), 4.49 (ddd, 1H, $J = 4.6, 5.6, 6.8$ Hz, $H-2'$), 4.60 (ddd, 1H, $J = 4.6, 5.2, 6.8$ Hz, $H-3'$), 5.21 (dd, 1H, $J = 5.3, 5.7$ Hz, OH-5'), 5.34 (d, 1H, $J = 5.2$ Hz, OH-3'), 5.51 (d, 1H, $J = 5.6$ Hz, OH-2'), 6.50 (d, 1H, $J = 6.4$ Hz, H-4), 6.67–7.15 (m, $2H_{arom}$), 7.18–7.39 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, DMSO- d_6): δ 38.2, 53.1, 62.3, 72.4, 73.5, 76.7, 78.1, 83.6, 113.4, 114.6, 119.5, 121.6, 123.3, 130.1, 131.8, 151.8, 166.3, 192.2. MS m/z (FAB, NBA) 562 $[MH^+]$. Anal. calcd for $C_{20}H_{19}Br_2NO_6S$: C, 42.80; H, 3.41; N, 2.50. Found: C, 42.55; H, 3.63; N, 2.72.

9i. Yellowish powder; m.p. 224–226°C. $[\alpha]_D^{[27]} + 46.8^\circ$ (c 1.5, MeOH). IR (KBr): ν 3389, 3013, 1602, 1585, 1455 cm^{-1} . 1H NMR (400 MHz, DMSO- d_6): δ 3.57 (ddd, 1H, $J = 3.4, 5.3, 11.6$ Hz, H_a-5'), 3.63 (ddd, 1H, $J = 3.4, 5.7, 11.6$ Hz, H_b-5'), 3.72 (s, 3H, OMe), 4.03 (ddd, 1H, $J = 4.2, 3.5, 3.4$ Hz, $H-4'$), 4.29 (dd, 1H, $J = 5.4, 6.4$ Hz, $H-1'$), 4.47 (ddd, 1H, $J = 4.6, 5.4, 6.8$ Hz, $H-2'$), 4.61 (ddd, 1H, $J = 4.6, 5.2, 6.8$ Hz, $H-3'$), 5.18 (dd, 1H, $J = 5.3, 5.7$ Hz, OH-5'), 5.29 (d, 1H, $J = 5.2$ Hz, OH-3'), 5.50 (d, 1H, $J = 5.6$ Hz, OH-2'), 6.53 (d, 1H, $J = 6.4$ Hz, H-4), 6.63–7.09 (m, $3H_{arom}$), 7.21–7.38 (m, $4H_{arom}$). ^{13}C NMR (100 MHz, DMSO- d_6): δ 38.4, 53.3, 62.0, 72.3, 73.1, 76.8, 78.5, 83.8, 113.2, 114.0, 119.9, 121.4, 122.9, 130.3, 131.3, 152.5, 166.8, 192.5. MS m/z (FAB, NBA) 438 $[MH^+]$. Anal. calcd for $C_{20}H_{20}ClNO_6S$: C, 54.86; H, 4.60; N, 3.20. Found: C, 55.02; H, 4.39; N, 2.93.

9j. Yellowish powder; m.p. 260–262°C. $[\alpha]_D^{[27]} + 48.5^\circ$ (c 1.5, MeOH). IR (KBr): ν 3391, 3012, 1605, 1580, 1453 cm^{-1} . 1H NMR (400 MHz, DMSO- d_6): δ 3.56 (ddd, 1H, $J = 3.4, 5.3, 11.6$ Hz, H_a-5'), 3.65 (ddd, 1H, $J = 3.4, 5.7, 11.6$ Hz, H_b-5'), 3.77 (s, 3H, OMe), 4.02 (ddd, 1H, $J = 4.2, 3.5, 3.4$ Hz, $H-4'$), 4.34 (dd, 1H, $J = 5.8, 6.4$ Hz, $H-1'$), 4.50 (ddd, 1H, J

TABLE 1 Synthesis of products 4 and 9

Product	Time (minutes) ^a	Yield(%) ^{b,c}	product	Time (hours) ^d	Yield(%) ^{b,c}
4a	7	78	9a	3	75
4b	8	87	9b	5	87
4c	8	89	9c	3	85
4d	10	79	9d	4	76
4e	12	88	9e	4	86
4f	10	85	9f	5	85
4g	9	80	9g	5	81
4h	7	86	9h	3	87
4i	11	85	9i	4	87
4j	10	79	9j	5	77

^aMicrowave irradiation time at 90°C.^bYield of isolated and purified products.^cAll compound gave C, H, and N analyses within $\pm 0.34\%$ and satisfactory spectral (IR, ¹H NMR, ¹³C NMR, and EIMS) data.^dTime for oil-bath heating at 60°C.

= 4.6, 5.8, 6.8 Hz, H-2'), 4.63 (ddd, 1H, J = 4.6, 5.2, 6.8 Hz, H-3'), 5.22 (dd, 1H, J = 5.3, 5.7 Hz, OH-5'), 5.32 (d, 1H, J = 5.2 Hz, OH-3'), 5.51 (d, 1H, J = 5.6 Hz, OH-2'), 6.56 (d, 1H, J = 6.4 Hz, H-4), 6.68–7.12 (m, 2H_{arom}), 7.19–7.42 (m, 4H_{arom}). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 38.0, 53.4, 62.4, 72.8, 73.7, 76.3, 78.8, 83.5, 113.1, 114.3, 120.2, 121.0, 122.8, 129.8, 131.6, 152.4, 166.5, 192.3. MS m/z (FAB, NBA) 472 [MH⁺]. Anal. calcd for C₂₀H₁₉Cl₂NO₆S: C, 50.86; H, 4.05; N, 2.97. Found: C, 50.63; H, 4.26; N, 3.17 (Table 1).

REFERENCES

1. Klásek, A.; Kořistek, K.; Polis, J.; Košmrlj, J. Synthesis of novel 3-acyloxy-1,3-dihydro-2H-indol-2-ones and Isomeric 4-acyl-1,4-dihydro-3,1-benzoxazin-2-ones: Double rearrangement of 3-hydroxyquinoline-2,4(1H,3H)-diones. *Tetrahedron* **2000**, 56, 1551–1560.
2. Mindl, J.; Hrabík, O.; Sterba, V.; Kavalek, J. Cyclization of substituted phenyl *N*-(2-hydroxybenzyl)carbamates in aprotic solvents. Synthesis of 4H-1,3-benzoxazin-2-ones. *Coll. Czech. Chem. Commun.* **2000**, 65, 1262–1272.
3. Waxman, L.; Darke, P.L. The herpesvirus proteases as targets for antiviral chemotherapy. *Antivir. Chem. Chemother.* **2000**, 11, 1–22.
4. Girgis, A.S. New 1,3-Benzoxazin-2-ones or thiones of molluscicidal activity. *Pharmazie* **2000**, 426–428.
5. Patel, M.; Ko, S.S.; Mc Hugh, R.J., Jr.; Markwalder, J.A.; Srivastava, A.S.; Cordova, B.C.; Klabe, R.M.; Erickson-Viitanen, S.; Trainor, G.L.; Seitz, S.P. Synthesis and evaluation of analogs of Efavirenz (SUSTIVA) as HIV-1 reverse transcriptase inhibitors. *Bioorg. Med. Chem. Lett.* **1999**, 9, 2805–2810.
6. Patel, M.; Mc Hugh, R.J., Jr.; Cordova, B.C.; Klabe, R.M.; Erickson-Viitanen, S.; Trainor, G.L.; Koo, S.S. Synthesis and evaluation of benzoxazinones as HIV-1 reverse transcriptase inhibitors. Analogs of efavirenz (SUSTIVA). *Bioorg. Med. Chem. Lett.* **1999**, 9, 3221–3226.
7. Fox, J.J.; Watanabe, K.A.; Klein, R.S.; Chu, C.K.; Tam, S.Y.-K.; Reichman, U.; Hirota, K.; Hwang, J.-S.; de Las Heras, F.G.; Wempen, I. *Chemistry and Biology of Nucleosides and Nucleotides* (Harmon, R.E.; Robins, R.K.; Townsend, L.B., Eds.), Academic Press, 1978, p. 415.
8. Beau, J.M.; Gallagher, T. Glycosyl nucleophiles. *Top Curr. Chem.* **1997**, 187, 1–54.

9. Postema, M.H.D. Recent developments in the synthesis of *c*-glycosides. *Tetrahedron* **1992**, 48, 8545–8599.
10. Levy, D.E.; Tang, C. *The Chemistry of C-Glycosides*, Pergamon, Oxford, 1995.
11. Postema, M.H.D. *C-Glycoside Synthesis*, CRS Press, Boca Raton, FL, 1995.
12. Katritzky, A.R.; Singh, S.K. Microwave-assisted heterocyclic synthesis. *Arkivoc* **2003**, xiii, 68–86.
13. Varma, R.S. Solvent-free synthesis of heterocyclic compounds using microwaves. *J. Heterocycl. Chem.* **1999**, 36, 1565–1571.
14. Guezguez, R.; Bougrin, K.; Li Akrai, K.; Benhida, R. A highly efficient microwave-assisted solvent-free synthesis of α - and β -2'-deoxy-1,2,3-triazolyl-nucleosides. *Tetrahedron Lett.* **2006**, 47, 4807–4811.
15. Yadav, L.D.S.; Rai, A.; Rai, V.K.; Awasthi, C. Carbohydrates to functionalized pyridines: A new synthetic approach via enol-driven ring transformations. *Synlett* **2008**, 0529–0534.
16. Yadav, L.D.S.; Awasthi, C.; Rai, V.K.; Rai, A. biorenewable and mercaptoacetylating building blocks in the biginelli reaction: Synthesis of thiosugar-annulated dihydropyrimidines. *Tetrahedron Lett.* **2007**, 48, 4899–4902.
17. Yadav, L.D.S.; Rai, V.K. Thiourea to bicyclic scaffolds: highly regio- and stereoselective routes to dithiazolopyrimidines. *Tetrahedron* **2007**, 63, 6924–6931.
18. Yadav, L.D.S.; Rai, V.K. Three-component coupling strategy for the expeditious synthesis of novel 4-aminobenzoxazinone N-nucleosides. *Tetrahedron Lett.* **2006**, 47, 395–397.
19. Yadav, L.D.S.; Yadav, S.; Rai, V.K. Green protocol for annulation of the *s*-triazine ring on thiazoles using a three-component coupling strategy. *Green Chem.* **2006**, 8, 455–458.
20. O' Callaghan, C.N.; Mc Murry, T.B.M. 2-Hydroxy-2-methyl-2H-1-benzopyran-3-carboxamide derivatives produced by knoevenagel condensation. *J. Chem. Res. (S)* **1997**, 78–79.
21. Bagnell, L.; Cablewski, T.; Strauss, C.R.; Trainor, R.W. Reactions of allyl phenyl ether in high-temperature water with conventional and microwave heating. *J. Org. Chem.* **1996**, 61, 7355–7359.
22. Varma, R.S.; Kumar, D.; Liesen, P.J. Solid state synthesis of 2-arylbenzo[*b*]furans, 1,3-thiazoles and 3-aryl-5,6-dihydroimidazo[2,1-*b*][1,3]thiazoles from α -tosyloxyketones using microwave irradiation. *J. Chem. Soc., Perkin Trans. 1* **1998**, 4093–4096.
23. Bandgar, B.P.; Uppalla, L.S.; Kurule, D.S. Solvent-free one-pot rapid synthesis of 3-carboxycoumarins using focused microwaves. *Green Chem.* **1999**, 1, 243–245.
24. Yadav, L.D.S.; Kapoor, R. Novel salicylaldehyde-based mineral supported expeditious synthesis of benzoxazin-2-ones. *J. Org. Chem.* **2004**, 69, 8118–8120.
25. Yadav, L.D.S.; Yadav, B.S.; Rai, V.K. A novel salicylaldehyde based mineral supported expedient synthesis of benzoxazinone nucleosides. *Tetrahedron Lett.* **2004**, 45, 5351–5353.
26. Mannen, S.; Itano, H.A. Stoichiometry of the oxidation of arylhydrazines with ferricyanide quantitative measurements of absorption spectra of aryl diazenes. *Tetrahedron* **1973**, 29, 3497–3502.
27. Chattaway, F. D. CXXII.—The oxidation of hydrazines by free oxygen. *J. Chem. Soc., Trans.* **1907**, 1323–1330.
28. Ohru, H.; Fox, J.J. Nucleosides LXXXI. An approach to from 2,3-O-isopropylidene-5-O-trityl- β -ribofuranosyl chloride. *Tetrahedron Lett.* **1973**, 14, 1951–1954.
29. Rosemeyer, H.; Toth, G.; Seela, F. Assignment of anomeric configuration of D-ribo-, arabino-, 2'-deoxyribo-, and 2',3'-dideoxyribonucleosides by NOE difference spectroscopy. *Nucleosides Nucleotides* **1989**, 8, 587–597.
30. Franchetti, P.; Cappellacci, L.; Grifantini, M.; Barzi, A.; Nocentini, G.; Yang, H.; O'Connor, A.; Jayaram, H.N.; Carrell, C.; Goldstein, B.M. Furanfuran and thiophenfuran: Two novel tiazofuran analogs. Synthesis, Structure, antitumor activity, and interactions with inosine monophosphate dehydrogenase. *J. Med. Chem.* **1995**, 38, 3829–3837.
31. Dudfield, P.J.; Le, V.-D.; Lindell, S.D.; Rees, C.W. Synthesis of C-ribosyl 1,2,4-triazolo[3,4-*f*][1,2,4]triazines as inhibitors of adenosine and AMP deaminases. *J. Chem. Soc. Perkin Trans 1* **1999**, 2937–2942.
32. Šilhár, P.; Pohl, R.; Votruba, I.; Hocek, M. Synthesis and cytostatic activity of novel 6-(difluoromethyl)purine bases and nucleosides. *Synthesis* **2006**, 1848–1852.