

the steric hindrance of cyclohexyl groups on the phosphorus atoms prevented the formation of the dinuclear structure. The structure of **7b** determined by X-ray structure analysis shows that intramolecular dehydrocoupling reaction has taken place to form a Si–Si bond (Figure 2).¹⁰ Although the mechanism of the formation of **7b** is not clear at the moment, an assumption of the intermediacy of tris(silyl)(hydrido)palladium(IV) **6b** can explain the formation of **7b**.¹¹ Scheme 1 shows two plausible pathways of the formation of **7b** from **6b**. Reductive elimination on **6b** to form a Si–Si bond provides **9** and successive oxidative addition of Si–H bond and elimination of H₂ from **10** give **7b**. Another possibility is the concerted mechanism; simultaneous elimination of H₂ and Si migration from Pd to another Si via transition state **11** gives **7b**.





Figure 2. Molecular structure of complex **7b** (50% probability level). Hydrogen atoms attached on carbon atoms are omitted for clarity. Selected bond distances (Å) and angles (°): Pd1–Si1, 2.348(3); Pd1–Si2, 2.356(3); Pd1–P1, 2.333(2); Pd1–P2, 2.315(2); Si2–Si3, 2.333(4); Si1–Pd1–Si2, 75.43(10); Si1–Pd1–P2, 171.56(10); Si2–Pd1–P1, 173.09(10); P1–Pd1–P2, 87.44(8); Pd1–Si2–Si3, 103.3(1).



Scheme 1. A plausible mechanism for the formation of **7b**.

We are grateful to the Japan Science and Technology Corporation (JST) for financial support through the CREST (Core Research for Evolutional Science and Technology) program and for a postdoctoral fellowship to W.C.

Dedicated to Professor Hideki Sakurai on the occasion of his 70th birthday.

References and Notes

- For recent reviews, see: M. S. Eisen, in "The Chemistry of Organic Silicon Compounds," ed. by Z. Rappoport and Y. Apeloig, Wiley, New York (1998), Vol. 2, Part 3, Chap. 35, p 2037; H. Ogino and H. Tobita, *Adv. Organomet. Chem.*, **42**, 223 (1998); J. Y. Corey and J. Braddock-Wilking, *Chem. Rev.*, **99**, 175 (1999).
- S. Shimada, M. Tanaka, and K. Honda, *J. Am. Chem. Soc.*, **117**, 8289 (1995); S. Shimada, M. Tanaka, and M. Shiro, *Angew. Chem., Int. Ed. Engl.*, **35**, 1856 (1996); S. Shimada, M. L. N. Rao, and M. Tanaka, *Organometallics*, **18**, 291 (1999); S. Shimada, M. L. N. Rao, T. Hayashi, and M. Tanaka, *Angew. Chem. Int. Ed.*, **40**, 213 (2001).
- G.-R. Sun, Ph. D. Thesis, Kyoto University, Kyoto, Japan, 1995; private communications from K. Tamao.
- K. Tamao, H. Yao, Y. Tsutsumi, H. Abe, T. Hayashi, and Y. Ito, *Tetrahedron Lett.*, **31**, 2925 (1990); K. Tamao, T. Hayashi, Y. Ito, and M. Shiro, *Organometallics*, **11**, 2099 (1992).
- Preparation of **2** via **4b**: A pentane solution of ^tBuLi (1.54 M, 137 mL, 0.21 mol) was added to PhSi(NMe₂)₂-(NMeCH₂CH₂NMe₂)₃ (**12**) (47.7 g, 0.162 mol) in 200 mL of hexane at 0 °C, and the mixture was stirred overnight at room temperature. To the resulting solution cooled to –40 °C was added SiHCl₃ (13.0 g, 0.096 mol), and the solution was allowed to warm to room temperature. Successive addition of dry MeOH (100 mL) and SiCl₄ (30 mL, as a source of HCl) at 0 °C, filtration of the resulting white precipitates, and removal of volatiles under vacuum gave crude **4b** as a pale-yellow waxy solid (²⁹Si NMR (C₆D₆, δ) –13.52 (d, ¹J(H,Si) = 219, Si–H)). LiAlH₄ (21 g, 0.55 mol) was added to an ether solution (500 mL) of crude **4b** at 0 °C, and the mixture was refluxed overnight. Filtration of insoluble materials, evaporation of the solvent, and bulb-to-bulb distillation afforded **2** as a colorless liquid (4.3 g, 22.0% based on **12**). ¹H NMR (C₆D₆, δ) 4.32 (s, 6H, SiH₃), 5.31 (s, 2H, SiH₂), 7.04 (m, 4H), 7.47 (m, 4H); ²⁹Si (C₆D₆, δ) –61.1 (qd, ¹J(H,Si) = 201, ²J(H,Si) = 5.9, SiH₃), –38.4 (tt, ¹J(H,Si) = 200, ²J(H,Si) = 5.0, SiH₂); Anal. Calcd for C₁₂H₁₆Si₃: C, 58.95; H, 6.60%. Found: C, 58.79; H, 6.57%.
- To a toluene solution of Pd(PEt₃)₄ and dmpe, depe, or dcpe was added **2** at 0 °C. Then, the mixture was stirred at room temperature for 20 min. The products were isolated by recrystallization from toluene (for **5a** and **7b**) or Al₂O₃ column chromatography (for **5b**).
- 5a**: ²⁹Si{¹H} NMR: (C₄D₈O, δ) –23.8 (tt, J(Si,P) = 11, 79), 79.8 (tt, J(Si,P) = 11, 128). **5b**: ²⁹Si{¹H} NMR: (C₇D₈, δ) –22.8 (tt, J(Si,P) = 10, 76), 82.4 (tt, J(Si,P) = 11, 125). **7b**: ²⁹Si NMR (C₆D₆, δ) 23.8 (dd, ²J(P–Si) = 13, 143 Hz, CSiHC), –31.4 (dd, ²J(P–Si) = 13, 142 Hz, PdSiHSiH₂), –61.8 (s, SiH₂SiH).
- Examples of μ -silylene-bridged dinuclear palladium complexes, see: a) M. Sugimoto, Y. Kato, N. Takeda, H. Oike, and Y. Ito, *Organometallics*, **17**, 495 (1998). b) Y.-J. Kim, S.-C. Lee, J.-I. Park, K. Osakada, J.-C. Choi, and T. Yamamoto, *Organometallics*, **17**, 4329 (1998). c) Y.-J. Kim, S.-C. Lee, J.-I. Park, K. Osakada, J.-C. Choi, and T. Yamamoto, *J. Chem. Soc., Dalton Trans.*, **2000**, 417.
- Crystal Data for **5a**-toluene: C₃₁H₅₂P₄Pd₂Si₃, fw = 845.70, *a* = 18.759(1), *b* = 15.071(2), *c* = 14.609(1) Å, β = 113.008(5)°, *V* = 3801.7(5) Å³, monoclinic, space group C2/c, *Z* = 4, *D*_{calc} = 1.477 g·cm^{–3}, *R*₁ = 0.040 (for 3894 reflections with *I* > 2σ(*I*)), *wR*₂ = 0.140 (for all data (4366 reflections)).
- Crystal Data for **7b**: C₃₀H₆₀P₂PdSi₃, fw = 769.50, *a* = 20.080(2), *b* = 9.696(3), *c* = 20.275(2) Å, *V* = 3947(1) Å³, orthorhombic, space group *Pna*2₁, *Z* = 4, *D*_{calc} = 1.295 g·cm^{–3}, *R*₁ = 0.047 (for 3164 reflections with *I* > 2σ(*I*)), *wR*₂ = 0.135 (for all data (4650 reflections)).
- Tilley has reported that a tris(silyl)(hydrido)platinum(IV) complex can be an intermediate for the dehydrocoupling reaction of phenylsilane: R. H. Heyn and T. D. Tilley, *J. Am. Chem. Soc.*, **114**, 1917 (1992).