

Syncarpamide, a New Antiplasmodial (+)-Norepinephrine Derivative from *Zanthoxylum syncarpum*

Samir A. Ross,^{*,†,‡} Gazi N. N. Sultana,[†] Charles L. Burandt,[†] Mahmoud A. ElSohly,^{†,§} Jannie P. J. Marais,[†] and Daneel Ferreira[†]

National Center for Natural Products Research, Research Institute of Pharmaceutical Sciences, Department of Pharmacognosy, and Department of Pharmaceutics, School of Pharmacy, University of Mississippi, University, Mississippi 38677-1848

Received September 16, 2003


A new (+)-norepinephrine derivative, syncarpamide (**1**), along with a known coumarin, (+)-*S*-marmesin (**2**), and one known alkaloid, decarine (**3**), have been isolated from the stem of *Zanthoxylum syncarpum*. The structure of compound **1** was elucidated on the basis of 1D and 2D NMR, MS, IR, optical rotation, and CD analyses. Its absolute stereochemistry was elucidated by synthesis of its enantiomer and subsequent comparison of CD data. Characterizations of compounds **2** and **3** were based on spectral analysis and comparison with reported data. Compounds **1** and **3** showed antiplasmodial activity, with IC₅₀ values of 2.04 and 1.44 μ M against *Plasmodium falciparum* D₆ clone and 3.06 and 0.88 μ M against *P. falciparum* W₂ clone, respectively. Compound **3** showed cytotoxicity at 56.42 μ M, whereas compound **1** was not cytotoxic at 10.42 μ M. Compound **1** was tested for hypotensive activity, but no activity was observed. Compound **2** showed no antiplasmodial or antimicrobial activities.

The genus *Zanthoxylum* (Rutaceae) comprises some 200 species distributed worldwide. Plants of this genus exhibit a variety of biologically active secondary metabolites including alkaloids, lignans, and coumarins with febrifuge, sudorific, and diuretic properties.¹ The bark of *Zanthoxylum integrifolium* (Merr.) Merr. (Rutaceae) has been utilized in traditional medicine by Ya-Mei aborigines in Lanyu Island in Taiwan as a remedy for snakebite² and dyspepsia and as an aromatic tonic in fever.³ In Sri Lanka, the genus *Zanthoxylum* is represented by three species, viz., *Z. caudatum*, *Z. rhetsa*, and *Z. tetraspermum*.⁴ Of these, *Z. tetraspermum* (*Sinhala*-katukeena) is employed in folklore medicine for the treatment of rheumatism and some forms of diarrhea.⁵ Pure compounds with antimicrobial activity have been obtained from the root bark of *Z. gillettii*.⁶ The bark is pungent, and sticks prepared from it are used for preventing toothache.⁷ The stem bark of *Z. liebmannaum* (Engelm.) P. Wilson is used in Mexican traditional medicine for the treatment of stomach aches, amebiasis, and intestinal parasites and as a local anesthetic agent.⁸

Z. syncarpum Tul. is an evergreen tree occurring in both North and South America.⁹ Although there are several reports on the chemical constituents of different species of *Zanthoxylum*,^{3–8} only two papers dealt with the constituents of *Z. syncarpum*.^{10,11} These phytochemical studies revealed the presence of two alkaloids, skimmianine and *cis*-*N*-methylcanadine, two coumarins, xanthotoxin and isopimpinellin, a diterpene, centipedic acid, the flavonoid ternatin, the triterpene lupeol, β -sitosterol glucoside,¹⁰ and an essential oil.¹¹

This paper deals with the isolation and characterization of a new alkaloid (**1**) and also the first report of the presence of **2** and **3** in *Z. syncarpum*. The isolation of these types of alkaloids and coumarins is of chemotaxonomic importance, as structurally related alkaloids have been reported earlier from *Fagara* and *Zanthoxylum* species.^{12,13}

Stems of *Z. syncarpum* were air-dried, ground, and sieved through a 1.0 mm mesh. An ethanolic extract of this material was carefully chromatographed over a silica gel column. The collected fractions were further purified by repeated chromatography over open silica gel columns, reversed-phase HPLC, and crystallization to afford compounds **1**, **2**, and **3**.



Syncarpamide (**1**) was obtained as light brown solid, $[\alpha]_D^{25} +12.5^\circ$, which gave a positive test with Dragendorff's reagent. The molecular formula was established as C₂₈H₂₇NO₅ by HRESIMS with 16 degrees of unsaturation. Analysis of ¹H NMR data, supported by ¹H–¹H COSY spectra,


* To whom correspondence should be addressed. Tel: (662) 915-5928. Fax: (662) 915-5587. E-mail: sross@olemiss.edu.

[†] National Center for Natural Products Research.

[‡] Department of Pharmacognosy.

[§] Department of Pharmaceutics.

revealed the presence of two pairs of *trans* olefinic systems at δ 6.38 (1H, d, J = 15.5 Hz), 7.63 (1H, d, J = 15.5 Hz), and δ 6.50 (1H, d, J = 16.0 Hz), 7.72 (1H, d, J = 16.0 Hz). The IR absorption band at 1713 cm^{-1} indicated the presence of a carbonyl group, and corresponding carbon peaks at δ 166.0 and 166.5 in the ^{13}C NMR spectra were reminiscent of an amide and an ester bond in **1**. In the HMBC spectra, the first pair of *trans* olefinic protons showed cross-peaks with the carbonyl carbon at δ 166.0 and the second pair with the carbon at δ 166.5. Analysis of ^1H and ^{13}C NMR data also revealed that **1** contained a 1,3,4-trisubstituted and two monosubstituted benzene rings. HMBC connectivity linked each pair of olefinic protons to a 1-substituted benzene ring. Two *O*-methyl singlets at δ 3.87 and 3.90 of the 1, 3, 4-trisubstituted benzene rings were observed in the ^1H NMR spectrum. Cross-peaks in the HMBC spectrum indicated that these methoxy groups are attached to the aromatic carbons at δ 149.1 and 149.2, respectively. One more spin system at δ_{H} 6.00 (1H, t, J = 6.4 Hz), δ_{C} 74.7, and δ_{H} 3.88 (2H, overlapped with methoxy signals), δ_{C} 44.7 was observed. The singlet aromatic resonance of the 1,3,4-trisubstituted benzene ring at δ_{H} 6.96 (δ_{C} 109.8) showed a cross-peak in the HMBC spectra to the oxymethine signal at δ_{H} 6.00 of the aliphatic spin system. The characteristic deshielded secondary carbon at δ_{C} 44.7 of the latter spin system indicated that it was connected to the nitrogen atom. A deshielded oxymethine proton resonance at δ_{H} 6.00 also indicated *O*-cinnamoylation at this position. Thus the structure of syncarpamide was elucidated as shown in **1**. It bears some resemblance to the naturally occurring aegline **4**, which was obtained in racemic form.^{13,14}



The positive specific rotation of syncarpamide (**1**) indicated that it was probably a derivative of (+)-*S*-norepinephrine. To confirm the absolute configuration of (**1**), we opted to derivatize commercially available (–)-*R*-norepinephrine (**5**) with a view to comparing its chiroptical properties with those of syncarpamide (**1**). Thus, methylation of the sparingly soluble (–)-*R*-norepinephrine (**5**) with ethereal diazomethane in a large excess of MeOH afforded (–)-*R*-3,4-di-*O*-methylnorepinephrine (**6**). Treatment with cinnamoyl chloride (2.1 equiv) in pyridine gave the *N,O*-dicinnamoyl derivative (**7**). Its CD spectrum exhibited sequential high-amplitude positive and negative Cotton effects in the 290–300 and 260–270 nm regions for the $n \rightarrow \pi^*$ and $\pi \rightarrow \pi^*$ transition of the α,β -unsaturated carbonyl chromophores, respectively. The CD spectrum of syncarpamide (**1**) showed a mirror image type relationship to that of **7** (Figure 1), hence unequivocally confirming the *S*-norepinephrine configuration of the new naturally occurring (+)-norepinephrine derivative (**1**).

Compound **2** was identified as (+)-*S*-marmesin by comparison of IR, HREIMS, and 1D and 2D NMR spectral data with published values.^{15,16}

Compound **3** was identified as decarine by comparison of IR, HREIMS, and 1D and 2D NMR spectral data with published values.^{17,18} Decarine was also isolated from *Z. decaryi*¹⁸ and *Z. viride*.¹⁹

Syncarpamide (**1**) and decarine (**3**) showed strong in vitro antiparasmodial activity²¹ against *Plasmodium falciparum*,




Figure 1. CD spectra of syncarpamide **1** and its (–)-*R*-norepinephrine enantiomer **7**.

having IC_{50} values of 2.04 and 1.44 μM against the D₆ strain and 3.06 and 0.88 μM against the W₂ strain, respectively.^{22,23} These compounds showed no cytotoxicity, except decarine at high concentration (56.42 μM). Decarine isolated from *Z. tetraspermum* and *Z. coudatum* showed significant antibacterial activity.²⁰ Marmesin (**2**) was not active in either the antimicrobial or antiparasmodial screen.

Experimental Section

General Experimental Procedures. Optical rotations were measured on a RUDOLPH Research Autopol IV automatic polarimeter. UV spectra were recorded using a Hewlett-Packard 8453 instrument. IR spectra were obtained using an AATI Mattson Genesis Series FTIR. ^1H and ^{13}C NMR spectra as well as 2D spectra (COSY, HMQC, and HMBC) were recorded on a Bruker DRX 400 spectrometer operating at 400 MHz for ^1H and 100 MHz for ^{13}C using the solvent peaks as internal references. EIMS was acquired on a GC/MS: Hewlett-Packard 5890A gas chromatography (GC) interface with 5970A selection detector (MSD). The GC/MS system was fitted with a DB-1 column (15 m \times 0.25 mm, and 0.25 μm film thickness, J & W Scientific, Folsom, CA), and HRESIMS and ESIMS data were acquired on a Bruker BioAPEX 30es mass spectrometer. TLC was performed on Merck TLC plates precoated with Si₆₀ F₂₅₄ or Si₆₀ RP18 F₂₅₄ with visualization by spraying with 5% H₂SO₄ in EtOH and heating or by Dragendorff's reagent. HPLC was carried out on a Waters Millennium system with a 996 photodiode array detector. CD spectra were recorded in CHCl₃ (ca. 0.1 mg/mL CHCl₃) on a JASCO J-715 spectrometer; scan parameters: bandwidth (2.0 nm), sensitivity (10 mdeg), response (4 s), scan speed (50 nm/min), step resolution (0.1 nm).

Plant Material. Specimens of *Z. syncarpum* were identified and collected in Lara State, Venezuela, by Dr. Charles L. Burandt in February 2002. Dried and ground stems were used for the isolation. A voucher specimen, BUR230202 (**2**), is deposited at the private herbarium of Dr. Burandt.

Extraction and Bioassay. The powdered, air-dried stems (1.2 kg) were exhaustively extracted with MeOH (3 \times 2 L). The combined MeOH extracts were evaporated, and the residue (63 g, 5.3% yield of dry plant weight) was chromatographed on a silica gel column (380 \times 78 mm, 540 g of silica 32–63 μm , 100:0:0–0:0:100 hexanes–EtOAc–MeOH step gradient) to give eight fractions, C₁–F₁, C₁–F₂, C₁–F₃, C₁–F₄, C₁–F₅, C₁–F₆, C₁–F₇, and C₁–F₈ (1 L each). Combined fractions C₁–F₂ to C₁–F₄ (6.23 g) were further chromatographed on silica gel (280 \times 35 mm column, 160 g of silica 32–63 μm , eluted with 100:0–0:100 CHCl₃–MeOH step gradient) to give 10 fractions. Fraction C₂–F₅ (1.83 g) was further chromatographed on silica gel (150 \times 35 mm column, 60 g of silica 32–63 μm , eluted with 98:2:0–0:0:100 hexane–EtOAc–MeOH step gradient) to give fractions F₅–1, F₅–2, F₅–3, F₅–4, F₅–5, and F₅–6. Fraction F₅–3 was crystallized from an EtOAc–hexane mixture

to afford compound **3** (44.0 mg). Fraction F₅-4 was crystallized from a MeOH–EtOAc mixture to give compound **2** (8.5 mg), and the mother liquid was subjected to reversed-phase HPLC (Prodigy ODS-3, 5 μ m, 21.2 \times 250 mm column, 70:30 (v/v) MeOH–H₂O, 14 mL min⁻¹, photodiode-array detection monitored at 220 nm) separation to afford compound **1** (5.9 mg).

L-Norepinephrine and cinnamoyl chloride were purchased from Acros Organics, NJ. Diazald, 99% (*N*-methyl-*N*-nitroso-*p*-toluenesulfonamide) was purchased from Aldrich Chemical Co. Inc.

Cytotoxicity Assay. In Vitro Cytotoxic Activity. The level of toxicity of each sample was determined by measuring the effect on a fibroblast cell line from African green monkey kidney (VERO, nontransformed). For initial (primary) evaluation, extracts and fractions were screened at a single concentration (100 μ g/mL). Follow-up secondary assays were conducted at three concentrations (4.76, 1.58, and 0.52 μ g/mL), using a culture-treated 96-well microplate 16. For secondary assay, IC₅₀ values were determined from logarithmic graphs of growth inhibition values. The cytotoxic agents doxorubicin and 5-fluorouracil were used as positive controls, while DMSO was used as the negative (vehicle) control.

Antimicrobial Assay. Preliminary antimicrobial activities of the crude extracts/fractions and the IC₅₀/MIC values of compounds were determined using a modified 96-well microplate assay protocol.²⁴ The test organisms used were ATCC strain of *Candida albicans* B311 (#90028), *Cryptococcus neoformans* (#90113), *Staphylococcus aureus* (#6535), and *S. aureus* (# 33591). Amphotericin B and rifampin were used as positive controls, with DMSO as a negative control.

Syncarpamide (1): light brown solid, $[\alpha]_D^{25} +12.50^\circ$ (*c* 0.08, CHCl₃); IR (film) ν_{\max} 3472, 2961, 2923, 1713, 1604, 1449, 1268, 757 cm⁻¹; UV λ_{\max} (MeOH) 242, 250, 278 nm; CD (CHCl₃) $[\theta]_{295.4} +10.5 \times 10^3$, $[\theta]_{263.9} -12.9 \times 10^3$; ¹H NMR (CDCl₃, 400 MHz) δ 3.87 (3H, s), 3.88 (2H, overlapped, H-8), 3.90 (3H, s), 6.00 (1H, t, *J* = 6.4 Hz, H-7), 6.08 (1H, br s), 6.38 (1H, d, *J* = 15.5 Hz, H-8''), 6.50 (1H, d, *J* = 16.0 Hz, H-8'), 6.86 (1H, d, *J* = 8.0 Hz, H-5), 6.96 (1H, s, H-2), 7.00 (1H, d, *J* = 8.0 Hz, H-6), 7.35 (1H, m, H-3'), 7.35 (1H, m, H-4'), 7.35 (1H, m, H-5'), 7.35 (1H, m, H-3''), 7.35 (1H, m, H-4''), 7.35 (1H, m, H-5''), 7.49 (1H, m, H-2'), 7.49 (1H, m, H-6'), 7.49 (1H, m, H-2), 7.49 (1H, m, H-6''), 7.63 (1H, d, *J* = 15.5 Hz, H-7''), 7.72 (1H, d, *J* = 16.0 Hz, H-7'); ¹³C NMR (CDCl₃, 100 MHz) δ 44.7 (t, C-8), 55.9 (q, OCH₃-4), 56.0 (q, OCH₃-3), 74.7 (d, C-7), 109.8 (d, C-2), 111.2 (d, C-5), 117.6 (d, C-8), 119.1 (d, C-6), 120.3 (d, C-8''), 127.9 (d, C-2''), 127.9 (d, C-6''), 128.2 (d, C-2'), 128.2 (d, C-6'), 128.8 (d, C-3'), 128.8 (d, C-5'), 128.9 (d, C-3''), 128.9 (d, C-5''), 129.8 (s, C-1), 130.3 (d, C-4'), 130.6 (d, C-4''), 134.2 (s, C-1'), 134.7 (s, C-1''), 141.6 (d, C-7'), 145.8 (d, C-7''), 149.2 (s, C-4), 149.2 (s, C-3), 166.0 (s, C-9'), 166.5 (s, C-9''); HRESIMS *m/z* 480.1821 (calcd for C₂₈H₂₇NO₅Na, [M + Na]⁺, 480.1787).

Synthesis of (–)-*R*-Norepinephrine Derivative (7). Diazomethane was added to a solution of l-norepinephrine **5** (100 mg) in MeOH (200 mL), and the mixture was cooled to –15 °C. The mixture was kept 72 h at –15 °C. A second batch of diazomethane was added, and the mixture was kept for a further 72 h at –15 °C. The solvent was evaporated with a strong stream of N₂.

The crude methylated product (**6**) was dried and dissolved in pyridine (2 mL). Cinnamoyl chloride (207 mg, 2.1 equiv) was added, and the reaction mixture was gently heated (40 to 45 °C) for 16 h. The excess pyridine was removed by a strong stream of N₂. The mixture was separated by preparative TLC using CHCl₃–MeOH, 99:1, to give the (–)-*R*-norepinephrine

derivative (**7**). Its ¹H NMR spectrum was identical to that of the natural product (**1**), CD (CHCl₃) $[\theta]_{298.6} -12.5 \times 10^3$, $[\theta]_{264.3} +23.5 \times 10^3$.

Marmesin (2): white crystals, $[\alpha]_D^{25} +9.63^\circ$ (*c* 0.166, CHCl₃); IR, UV, HREIMS, and ¹H and ¹³C NMR data identical to literature value.¹⁶

Decarine (3): yellow crystals; UV, HREIMS, ¹H, ¹³C NMR data identical to reference values.^{17–19}

Acknowledgment. The authors acknowledge the United States Department of Agriculture, Agriculture Research Service Specific Cooperative Agreement No. 58-6408-2-0009, for partial support of this work. The authors are also grateful to Dr. D. G. Nagle and Dr. C. F. Hossain (Department of Pharmacognosy, School of Pharmacy, University of Mississippi) for allowing the use of the HPLC facility and Dr. D. C. Dunbar (NCNPR, University of Mississippi) for the HRESIMS analysis.

References and Notes

- Talapatra, S. K.; Dutta, S.; Talapatra, B. B. *Phytochemistry* **1973**, *12*, 729–730.
- Ko, F. N.; Hsiao, G.; Chen, I. S.; Wu, S. J.; Teng, C. M. *Biochem. Pharmacol.* **1993**, *46*, 1165–1173.
- The Wealth of India: Raw Materials*; PID, Council of Scientific and Industrial Research (CSIR): New Delhi, 1976; Vol. II, pp 18–19.
- Stone, B. C. In Dassanayake, M. D., Fosberg, R., Eds. *A Revised Handbook to the Flora of Ceylon*; Oxford and IBH Publishing: New Delhi, 1985; Vol. 5, p 406.
- Jayaweera, D. M. A. In *Medicinal Plants Used in Sri Lanka*. National Science Council: Sri Lanka, 1982; Part 5, p 39.
- Weenen, H.; Nkunya, M. H. H.; Bray, D. H.; Mwasumbi, L. B.; Kinabo, L. S.; Kilimali, V. A.; Wijnberg, J. B. *Planta Med.* **1990**, *56*, 371–373.
- Kalia, N. K.; Sing, B.; Sood, R. P. *J. Nat. Prod.* **1999**, *62*, 311–312.
- Reyes, B.; Navarrete, A.; Sixtos, C.; Aguirre, E.; Jimenez, S.; Estrada, E. *Rev. Mex. Cienc. Farm.* **1991**, *21*, 30–34.
- De Morais, S. M.; Facundo, V. A.; Braz Filho, R. *J. Essent. Oil Res.* **2002**, *14*, 274–275.
- Facundo, V. A.; De Morais, S. M.; Braz Filho, R.; Matos, I. J. de A.; Souza, R. T. *Rev. Bras. Farm.* **1997**, *78*, 57–59.
- Facundo, V. A.; De Morais, S. M.; Machado, M. I. L.; Matos, F. J. de A.; Da Frota, L. C. M. *J. Essent. Oil Res.* **1999**, *11*, 426–428.
- Dadson, B. A.; Minta, A. *J. Chem. Soc., Perkin Trans. 1* **1976**, *2*, 146–147.
- Swinehart, J. A.; Stermitz, F. R. *Phytochemistry* **1980**, *19*, 1219–1223.
- Chatterjee, A.; Bose, S.; Srimany, S. K. *J. Org. Chem.* **1959**, *24*, 687–690.
- Warren, S. *Can. J. Chem.* **1971**, *49*, 2297–2301.
- Elgamal, M. H. A.; Elewa, N. H.; Elkhaisy, E. A. M.; Duddeck, H. *Phytochemistry* **1979**, *18*, 139–143.
- Krane, B. D.; Fagbule, M. O.; Shamma, M. *J. Nat. Prod.* **1984**, *47*, 1–43.
- Vaquette, J.; Pousset, J.-L.; Paris, R.-R.; Cave, A. *Phytochemistry* **1974**, *13*, 1257–1259.
- Waterman, P. G. *Phytochemistry* **1975**, *14*, 843–844.
- Nissanka, A. P. K.; Karunaratne, V.; Bandara, B. M. R.; Kumar, V.; Nakanishi, T.; Nishi, M.; Inada, A.; Tillekeratne, L. M. V.; Wijesundara, D. S. A.; Gunatilaka, A. A. L. *Phytochemistry* **2001**, *56*, 857–861.
- Dou, J.; McChesney, J. D.; Sindelar, R. D.; Goins, D. K.; Walker, L. A. *J. Nat. Prod.* **1996**, *59*, 73–76.
- Makler, M. T.; Ries, J. M.; Williams, J. A.; Bancroft, J. E.; Piper, R. C.; Gibbins, B. L.; Hinriches, D. J. *Am. J. Trop. Med. Hyg.* **1993**, *48*, 739–741.
- Makler, M. T.; Hinriches, D. J. *Am. J. Trop. Med. Hyg.* **1993**, *48*, 205–210.
- National Committee for Clinical Laboratory Standards. *Methods for Dilution. Antimicrobial Susceptibility Test for Bacteria that Grow Aerobically. Approved Standard M7-A*, 4th ed.; National Committee for Clinical Laboratory Standards: Wayne, PA, 1997.

NP030417T