


Synthesis, Cocaine Receptor Affinity, and Dopamine Uptake Inhibition of Several New 2 β -Substituted 3 β -Phenyltropanes

Shreekrishna V. Kelkar,[†] Sari Izenwasser,[§]
Jonathan L. Katz,[§] Cheryl L. Klein,[‡]
Naijue Zhu,[‡] and Mark L. Trudell^{*†}

Department of Chemistry, University of New Orleans,
New Orleans, Louisiana 70148,
Psychobiology Laboratory, NIDA Intramural Research
Program, P.O. Box 5180, Baltimore, Maryland 21224,
and Department of Chemistry, Xavier University of
Louisiana, New Orleans, Louisiana 70125


Received June 27, 1994

Cocaine (1) is believed to mediate its pharmacologic effects through the occupation of high-affinity saturable stereoselective binding sites associated with biogenic amine uptake mechanisms.^{1–5} The reinforcing effects of cocaine have been shown to be associated with the inhibition of dopamine uptake; however, the mechanistic details of dopamine uptake inhibition are unclear at this time.^{6–8} It is conceivable that occupation of the cocaine receptor results in the physical blockade of dopamine at the dopamine reuptake site or that allosteric modulation of the dopamine transporter occurs such that dopamine uptake is not possible.^{3,9,10} In order to obtain a greater understanding of the mechanism by which cocaine affects dopamine uptake inhibition and elicits its reinforcing effects, a study of the structure–activity relationships (SAR) of cocaine and cocaine-related derivatives has been initiated in these laboratories.



The 2 β -carbomethoxy-3 β -phenyltropanes 2 (WIN 35,065-2),^{11,12} 3 (WIN 35,428),^{11,12} and 4 (RTI-55)^{12,13} are among the most potent tropane derivatives for cocaine receptors. From the SAR of these compounds and 2 β -substituted cocaine derivatives it is clear that the substitution at C2 of the tropane nucleus has a profound effect on the affinity of the cocaine receptor ligands.⁸ Recent reports by Carroll *et al.*,^{14,15} Davies *et al.*,^{16,17} and Kozikowski *et al.*¹⁸ have demonstrated that a variety of 2 β -carbonyl- or 2 β -unsaturated 3 β -phenyltropane derivatives exhibit high affinity for cocaine receptors. However, the nature of the interaction between the 2 β -substituents and the binding site has not been clearly identified. To explore the nature of the interaction between the cocaine receptor and high-affinity ligands as well as attempt to identify the proximal effect of electron-rich/unsaturated 2 β -substituents, a series of 2 β -substituted 3 β -phenyltropanes were synthesized and tested *in vitro* for cocaine receptor affinity and for dopamine uptake inhibition.

Scheme 1^a



^a Reagents: (i) LiAlH₄, Et₂O, 0 °C; (ii) (COCl)₂, DMSO, Et₃N, CH₂Cl₂, –78 °C; (iii) (CH₃O)₂POCH₂CO₂CH₃, LiCl, *i*Pr₂NEt, CH₃CN, 25 °C.

The 2 β -substituted 3 β -phenyltropane derivatives 5–10 were prepared from 2. N-Methyl-2 β -carbomethoxy-3 β -phenyltropane 2 was prepared from cocaine according to the modified procedure of Carroll *et al.*¹³ As illustrated in Scheme 1, LiAlH₄ reduction of the 2 β -ester group of 2 afforded the 2 β -(hydroxymethyl)-3 β -phenyltropane 5.¹¹ Swern oxidation of 5 to the corresponding aldehyde followed by subsequent olefination, furnished the α,β -unsaturated ester 6 in 65% overall yield.¹⁹ The mild conditions of the Masamune–Rousch olefination procedure afforded the ester 6 as a single crystalline isomer resulting from no epimerization of the labile intermediate 2 β -aldehyde.²⁰ The structure of 6 was determined by NMR and the 2 β -stereochemistry was unequivocally confirmed by X-ray crystallographic analysis (Figure 1).

With stereochemistry at the 2 β -position no longer susceptible to epimerization, the 2 β -derivatives 7–10 were prepared in a straightforward fashion from the unsaturated ester 6 (Scheme 2). Hydrogenation of 6 afforded saturated ester 7 in quantitative yield, while reduction of 6 gave the allyl alcohol 8 (95% yield). Conversion of ester 7 into 9 was achieved by reduction with LiAlH₄. Finally, Swern oxidation/Masamune–Rousch olefination of 9 furnished the unsaturated ester 10 (89% yield).

The five 2 β -substituted 3 β -phenyltropane derivatives 6–10 were tested for their ability to displace bound [³H]-3 from rat caudate–putamen tissue.²¹ In addition, the compounds 6–10 were tested for their ability to inhibit high affinity uptake of [³H]dopamine into rat caudate–putamen tissue.²¹ The *K*_i values reported in Table 1, are the dissociation constants derived for the unlabeled ligands. The linear portion of the [³H]dopamine uptake inhibition curves were analyzed using standard analysis of variance and linear regression techniques.

Both cocaine and 2 modeled better for two binding sites than for one, and therefore two *K*_i values are given in the table.^{22,23} In contrast, compounds 6–10 did not model for two sites better than for one. The affinities of compounds 6–10 are generally similar to the high affinity values obtained for 1 and 2. The saturated alcohol 9 exhibited a 2-fold increase in potency in both *in vitro* paradigms compared to the esters 6, 7, and 10 and the unsaturated alcohol 8. This result was somewhat surprising since most potent 2 β -substituted 3 β -aryl tropane derivatives derived from either cocaine or 3 β -phenyltropanes reported to date possess an unsaturated moiety such as an ester,^{11–14} a ketone,^{16,17} an oxadiazole,¹⁵ or a vinyl group¹⁸ directly bound to the tropane ring system. Moreover, the 2 β -hydroxymethyl analog 11, homoesters, and C2-unsubstituted 3 β -aryl tropanes have been reported to have diminished activity at cocaine receptors.^{11,14}

[†] University of New Orleans.

[‡] Xavier University of Louisiana.

[§] NIDA Intramural Research Program.





Figure 1. X-ray crystal structure of compound **6** (free base).

Scheme 2^a



^a Reagents: (i) 10% Pd/C, MeOH; (ii) LiAlH₄, Et₂O, 0 °C; (iii) (COCl)₂, DMSO, Et₃N, CH₂Cl₂, -78 °C; (iv) (CH₃O)₂POCH₂CO₂CH₃, LiCl, *i*Pr₂NEt, CH₃CN, 25 °C.

The high affinity of the saturated alcohol **9** suggests that the flexible nature of the saturated hydroxypropyl unit allows the ligand to better conform to the structural constraints imposed by this region of the cocaine binding site. The other analogs **6–8** and **10** possess some degree of unsaturation which increases the rigidity of the side chain and perhaps decreases the ability of the ligand to adopt a higher affinity conformation at the cocaine receptor. Despite recent studies by Kozikowski *et al.*¹⁸ which have shown that 2β-vinyl-3β-aryltropane analogs **12–15** possess high affinity for cocaine receptors, clearly unsaturation does not facilitate binding in this system. Moreover, since the unsaturated alcohol **8** demonstrates

Table 1. *K_i* Values for Displacement of Receptor Bound [³H]-**3** and IC₅₀ Values for Inhibition of [³H]Dopamine Uptake^a


analog	<i>K_i</i> (nM)	IC ₅₀ (nM)
1 ^b	32 ± 5	405 ± 91
	388 ± 221	
2 ^b	33 ± 17	373 ± 100
	314 ± 222	
6	22 ± 2	123 ± 65
7	23 ± 2	166 ± 68
8	26 ± 3	159 ± 43
9	11 ± 1	64 ± 32
10	20 ± 2	203 ± 77

^a All values are the mean ± SEM of three experiments performed in triplicate. ^b The binding data for these drugs are reproduced from ref 23 and were collected under conditions identical to the present ones.

no greater affinity than esters **6**, **7**, and **10**, the hydroxyl group of **9** probably has little effect or at most a nonspecific effect on the affinity of the ligand.

The compounds **6**, **7**, **8**, and **10** were synthesized in order to evaluate the effect of the proximity of an unsaturated group and/or an electron-rich group relative to the tropane ring on cocaine receptor affinity. Interpretation of the *in vitro* affinity data indicates that the proximity of an unsaturated group or an electron-rich group to the tropane nucleus has little effect on the ligand affinity. Despite the fact that the functional groups were displaced by as many as four carbon atoms from the tropane nucleus, the *K_i* values of the analogs **6**, **7**, **8**, and **10** were all equivalent to that of **2**. It is equally noteworthy that the cocaine binding site was able to accommodate the large ester side chain of **10** with no effect on the affinity of the ligand. This result is consistent with data obtained by Carroll *et al.* in which large 2β-substituents of 3β-phenyltropane and cocaine derivatives demonstrated high affinity for cocaine binding sites.^{14,15}

From the results of this study and from known SAR of 2β-substituted 3β-phenyltropane and cocaine derivatives,^{14–18} it can be concluded that the region of the cocaine binding site which is occupied by the 2β-substituent must either be a large cleft in the dopamine transporter protein or is exterior to the binding site where the 2β-substituent lies above the surface of the dopamine transporter protein. Either case would explain the ability of the binding site to accommodate the large side chains of potent ligands. Secondly, this region appears to be relatively insensitive to electrostatic and lipophilic interactions. This is consistent with only the slight difference in affinity observed for compounds **2**, **6–10**. The proposed nature of this region of the cocaine binding is in agreement with the models recently proposed by Kozikowski *et al.*¹⁸ and Srivastava and Crippen.²⁵



In summary, it is the stereochemical orientation of substituents at C2 which is the primary requirement for high-affinity binding at cocaine receptors, while the steric bulk and the lipophilic character of the 2β-

substituents exhibit minimum effects on the affinity of the ligand. The results of this study are in complete agreement with these observations. Moreover, the SAR reported herein supports the hypothesis that electrostatic interactions (including hydrogen bonds) between 2 β -substituents and the cocaine binding site are of minimal importance while hydrophobic interactions may contribute significantly to the free energy of binding and lead to the enhanced potency of ligands.¹⁵ Further studies aimed at the elucidation of the substituent effects of high-affinity 2 β -substituted 3 β -phenyltropane derivatives in both *in vitro* and *in vivo* systems is warranted and are the subject of current investigations.

Acknowledgment. Mr. Brett Heller provided technical assistance in the conduct of radioligand displacement and dopamine uptake assays. We are grateful to the National Institute on Drug Abuse (NIDA First Award) for the financial support of this research. The Louisiana Board of Reagents are acknowledged for allocating funds for the purchase of the NMR spectrometer (ENH-53, 1990–1991) and the FT-IR spectrometer [LEQSF(1993–1994)-ENH-TR-41].

Supplementary Material Available: Experimental procedures, physical data for compounds 6–10, and X-ray crystallographic data for compound 6 (12 pages). Ordering information is given on any current masthead page.

References

- Reith, M. E. A.; Serksen, H.; Lajtha, A. Saturable [³H]Cocaine Binding in Central Nervous System of Mouse. *Life Sci.* **1980**, *27*, 1055–1062.
- Calligaro, D. O.; Eldefrawi, M. E. Central and Peripheral Cocaine Receptors. *J. Pharmacol. Exp. Ther.* **1987**, *243*, 61–67.
- Madras, B.; Fahey, M. A.; Bergman, J.; Canfield, D. R.; Speakman, R. D. Effects of Cocaine and Related Drugs in Nonhuman Primates. I. [³H]Cocaine Binding Sites in Caudate-Putamen. *J. Pharmacol. Exp. Ther.* **1989**, *252*, 131–141.
- Calligaro, D. O.; Eldefrawi, M. E. High Affinity Stereospecific Binding of [³H]Cocaine in Striatum and Its Relationship to the Dopamine Transporter. *Membrane Biochem.* **1988**, *7*, 87–106.
- Ritz, M. C.; Cone, E. J.; Kuhar, M. J. Cocaine Inhibition of Ligand Binding at Dopamine, Norepinephrine and Serotonin Transporters: A Structure-Activity Study. *Life Sci.* **1990**, *46*, 635–645.
- Ritz, M. C.; Lamb, R. J.; Goldberg, S. R.; Kuhar, M. J. Cocaine Receptors on Dopamine Transporters are Related to Self-Administration of Cocaine. *Science* **1987**, *237*, 1219–1223.
- Kuhar, M. J.; Ritz, M. C.; Boja, J. W. The Dopamine Hypothesis of the Reinforcing Properties of Cocaine. *Trends Neurosci.* **1991**, *14*, 299–302.
- For a recent review see: Carroll, F. I.; Lewin, A. H.; Boja, J. W.; Kuhar, M. J. Cocaine Receptor: Biochemical Characterization and Structure-Activity Relationships of Cocaine Analogues at the Dopamine Transporter. *J. Med. Chem.* **1992**, *35*, 969–981.
- Zimanyi, I.; Lajtha, A.; Reith, M. E. A. Comparison of Characteristics of Dopamine Uptake and Mazindol Binding in Mouse Striatum. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1989**, *340*, 626–632.
- Kreuger, B. K. Kinetics and Block of Dopamine Uptake in Synaptosomes from Rat Caudate Nucleus. *J. Neurochem.* **1990**, *55*, 260–267.
- Clarke, R. L.; Daum, S. J.; Gambino, A. J.; Aceto, M. D.; Pearl, J.; Levitt, M.; Cumiskey, W. R.; Bogado, E. F. Compounds Affecting the Central Nervous System. 4. 3 β -Phenyltropane-2-carboxylic Esters and Analogues. *J. Med. Chem.* **1973**, *16*, 1260–1267.
- Carroll, F. I.; Gao, Y.; Rahman, M. A.; Abraham, P.; Parham, K.; Lewin, A. H.; Boja, J. W.; Kuhar, M. J. Synthesis, Ligand Binding, QSAR and CoMFA Study of 3 β -(*p*-Substituted phenyl)-tropane-2 β -carboxylic Acid Methyl Esters. *J. Med. Chem.* **1991**, *34*, 2719–2725.
- Boja, J. W.; Patel, A.; Carroll, F. I.; Rahman, M. A.; Philip, A.; Lewin, A. H.; Kopajtic, T. R.; Kuhar, M. J. [²⁵I]RTI-55: A Potent Ligand for Dopamine Receptors. *Eur. J. Pharmacol.* **1991**, *194*, 133–4.
- Lewin, A. H.; Gao, Y.; Abraham, P.; Boja, J. W.; Kuhar, M. J.; Carroll, F. I. Synthesis of 2 β -Substituted Analogues of Cocaine. Synthesis and Inhibition of Binding to the Cocaine Receptor. *J. Med. Chem.* **1992**, *35*, 135–140.
- Carroll, F. I.; Gray, J. L.; Abraham, P.; Kuzemko, M. A.; Lewin, A. H.; Boja, J. W.; Kuhar, M. J. 3-Aryl-2-(3'-substituted-1'2'4'-oxadiazol-5'-yl)tropane Analogues of Cocaine: Affinities at the Cocaine Binding Site at the Dopamine, Serotonin and Norepinephrine Transporters. *J. Med. Chem.* **1993**, *36*, 2886–2890.
- Davies, H. M. L.; Saikali, E.; Sexton, T.; Childers, S. R. Novel 2-Substituted Cocaine Analogs: Binding Properties at Dopamine Transport Site in Rat Striatum. *Eur. J. Pharmacol. Mol. Pharmacol. Sect.* **1993**, *244*, 93–97.
- Davies, H. M. L.; Saikali, E.; Huby, N. J.; Gilliat, V. J.; Matasi, J. J.; Sexton, T.; Childers, S. R. Synthesis of 3 β -Acyl-3 β -aryl-8-azabicyclo[3.2.1]octanes and Their Binding Affinities at Dopamine and Serotonin Transport Sites in Rat Striatum and Frontal Cortex. *J. Med. Chem.* **1994**, *37*, 1662–1688.
- Kozikowski, A. P.; Roberti, M.; Xiang, L.; Bergmann, J. S.; Callahan, P. M.; Cunningham, K. A.; Johnson, K. M. Structure-Activity Relationship Studies of Cocaine: Replacement of the C-2 Ester Group by Vinyl Argues Against H-Bonding and Provides an Esterase-Resistant High-Affinity Cocaine Analogue. *J. Med. Chem.* **1992**, *35*, 4764–4766.
- Mancuso, A. J.; Swern, D. Activated Dimethyl Sulfoxide: Useful Reagents for Synthesis. *Synthesis* **1981**, 165–185.
- Blanchette, M. A.; Choy, W.; Davis, J. T.; Essensfeld, A. P.; Masamune, S.; Roush, W. R.; Sakai, T. Horner-Wadsworth-Emmons Reaction: Use of Lithium Chloride and an Amine for Base-Sensitive Compounds. *Tetrahedron Lett.* **1984**, *25*, 2183–2186.
- For details see supplementary material and Izenwasser, S.; Rosenberger, J. G.; Cox, B. M. The Cocaine Analog WIN 35,428 Binds to Two Sites in Fresh Rat Caudate-Putamen Under the Correct Assay Conditions. *Life Sci./Pharmacol. Lett.* **1993**, *52*, PL141–145. Izenwasser, S.; Werling, L. L.; Cox, B. M. Comparison of the Effects of Cocaine and Other Inhibitors of Dopamine Uptake in Rat Striatum, Nucleus Accumbens, Olfactory Tubercle and Medial Prefrontal Cortex. *Brain Res.* **1990**, *530*, 303–309. Munson, P. J.; Rodbard, D. Ligand: A Versatile Approach for the Characterization of Ligand-Binding Systems. *Anal. Biochem.* **1980**, *107*, 220–239.
- Izenwasser, S.; Newman, A. H.; Katz, J. L. Cocaine and Several Sigma Receptor Ligands Inhibit Dopamine Uptake in Rat Caudate-Putamen. *Eur. J. Pharmacol.* **1993**, *243*, 201–205.
- Izenwasser, S.; Terry, P.; Heller, B.; Witkin, J. M.; Katz, J. L. Differential Relationships Among Dopamine Transporter Affinities and Stimulant Potencies of Various Uptake Inhibitors. *Eur. J. Pharmacol.*, in press.
- Baker, B. R. *Design of Active-Site Directed Irreversible Enzyme Inhibitors*; John Wiley and Sons: New York, **1967**; p 24.
- Srivastava, S.; Crippen, G. M. Analysis of Cocaine Receptor Site Ligand Binding by Three-Dimensional Voronoi Site Modeling Approach. *J. Med. Chem.* **1993**, *36*, 3572–3579.