



A Concise Synthesis of 2-Amino-1,2,3,4-tetrahydronaphthalene-6,7-diol ('6,7-ADTN') from Naphthalene-2,3-diol

by Süleyman Göksu, Cavit Kazaz, Yaşar Sütbeyaz*, and Hasan Seçen*

Department of Chemistry, Faculty of Arts and Sciences, Atatürk University, TR-25240 Erzurum

2-Amino-1,2,3,4-tetrahydronaphthalene-6,7-diol (**2**; 6,7-ADTN) was synthesized starting from naphthalene-2,3-diol in seven steps and with an overall yield of 44%. Methylation of naphthalene-2,3-diol with dimethyl sulfate, followed by *Friedel–Crafts* acylation with AcCl, gave 2-acetyl-6,7-dimethoxynaphthalene. 2-Acetyl-6,7-dimethoxynaphthalene was converted to 6,7-dimethoxynaphthalene-2-carboxylic acid by a haloform reaction. *Birch* reduction of the carboxylic acid with 4 mol-equiv. of Na in liquid ammonia afforded 1,2,3,4-tetrahydro-6,7-dimethoxynaphthalene-2-carboxylic acid, from which **2** was obtained by a *Curtius* reaction, followed by hydrogenolysis and demethylation.

Introduction. – Dopamine (**1**) [1], a hormone-like substance, is an important neurotransmitter. When present in normal quantities, it facilitates critical brain functions. 2-Amino-1,2,3,4-tetrahydronaphthalene-6,7-diol (= '2-amino-6,7-dihydroxy-1,2,3,4-tetrahydronaphthalene', 6,7-ADTN; **2**) is a compound suggested to act as a potential agonist at the dopamine receptors [2] and to stimulate helix dopamine receptors [3]. It also has a strong central stimulant action [4] and is as active as dopamine in producing adenylate cyclase activity [5][6]. It has been suggested that 6,7-ADTN (**2**) interacts with the dopamine receptor with slightly greater affinity than dopamine itself [7].




Thrift [8a], starting from veratrol, performed the first synthesis of 6,7-ADTN (**2**)¹⁾. *Cannon et al.* [9] described the second method for the preparation of **2** starting from ethyl 3-(3,4-dimethoxyphenyl)propanoate. *Horn et al.* [10] developed a procedure for the synthesis of **2** starting from (3,4-dimethoxyphenyl)acetic acid. *Narula* and *Schuster* [11] synthesized **2** from 2-(3,4-dimethoxyphenyl)ethanol via functionalized aryllithium reagents. *Nordlander et al.* [12] developed the first asymmetric synthesis of (*R*)-6,7-ADTN by using veratrol and D-aspartic acid. (*R*)-1,2,3,4-tetrahydro-6,7-dimethoxynaphthalene-2-amine, *O*-methylated 6,7-ADTN, was synthesized by *Charlton et al.* [13] starting from 2-amino-4,5-dimethoxybenzoic acid. These syntheses of **2** were performed in 5–9 steps with overall yields ranging from 12 to 52%. In this paper, we report a convenient synthesis of 6,7-ADTN (**2**) with a moderate overall yield (44%) starting

¹⁾ For a modified synthesis, see [8b].

from naphthalene-2,3-diol (**3**), a commercially available naphthalene derivative, which was used in the synthesis of **2** for the first time (*Scheme*).

Scheme



i) Me_2SO_4 , K_2CO_3 , acetone, reflux; 94%. ii) AcCl , AlCl_3 , 1,2-dichloroethane, 0° ; 88%. iii) Br_2 , NaOH , then 6M HCl , 90%. iv) Na , liq. NH_3 , then 37% HCl ; 90%. v) $(\text{PhO})_2\text{P}(\text{O})\text{N}_3$, Et_3N , C_6H_6 , reflux; then PhCH_2OH , reflux; 86%. vi) H_2 , Pd-C , EtOH , CHCl_3 ; 96%. vii) 48% HBr , reflux; 80%.

Results and Discussion. – The synthesis started with naphthalene-2,3-diol (**3**). Methylation of **3** with Me_2SO_4 in the presence of K_2CO_3 gave 2,3-dimethoxynaphthalene (**4**). Our literature search showed that the *Friedel–Craft* acetylation of **4** has been performed by reacting with AcCl in PhNO_2 to give 1-(6,7-dimethoxynaphthalen-2-yl)ethanone (**5**) [14] [15]. Despite moderate yields, these procedures involve some difficulties related to the removal of solvent and a series of workup procedures. Another procedure [16], again performed in PhNO_2 with Ac_2O , gave **5** in low yield. Our method for the preparation of **5** was to react **4** with AcCl in 1,2-dichloroethane. Thus, this simple method gave **5** in high yield. Acetyl compound **5** was converted to carboxylic acid **6** by a haloform reaction (Br_2/NaOH). The most critical step of our synthesis was the *Birch* reduction of 6,7-dimethoxynaphthalene-2-carboxylic acid (**6**). We assume that the *Birch* reaction of **6** proceeded by the reduction of the electron-deficient ring of **6**. Indeed, the reaction of **6** with 4 mol.-equiv. of Na in liquid NH_3 afforded tetrahydronaphthalene derivative **7** in high yield (90%). The conversion of the carboxylic acid **7** to carbamate **8** was performed by a known procedure *via Curtius* rearrangement, followed by treatment with PhCH_2OH . Hydrogenolysis of **8** in MeOH in the presence of CHCl_3 gave a pure dimethoxy derivative of 6,7-ADTN hydrochloride **9** as the sole product. The demethylation of **9** to hydrobromide of 6,7-ADTN, **10**, was performed as described in the literature.

In summary, with relatively little synthetic effort, we have achieved a concise synthesis of 6,7-ADTN (**2**) in seven steps starting from commercially available naphthalene-2,3-diol (**3**) (overall yield of 44%).

Experimental Part

General. Column chromatography (CC): silica gel (60 mesh, *Merck*). Prep. thick-layer chromatography (PLC): 1-mm of silica gel 60 PF (*Merck*) on glass plates. M.p.: *Thomas-Hoover* cap. melting-point apparatus; uncorrected. IR Spectra: from solns., 0.1-mm cells, with a *Perkin-Elmer* spectrophotometer. ^1H - and ^{13}C -NMR spectra: 200 (50)-MHz *Varian* spectrometer; δ in ppm.

2,3-Dimethoxynaphthalene (4). A 500-ml, three-necked flask fitted with a condenser and a 50-ml dropping funnel was charged with anh. K_2CO_3 (38 g, 275 mmol), 300 ml of acetone, and naphthalene-2,3-diol (**3**; 20 g, 125 mmol). Me_2SO_4 (34.6 g, 26.6 ml, 275 mmol) was added under stirring from the dropping funnel to the mixture over a period of 2 min. The mixture warmed appreciably and began to reflux after an additional 5 min. When the spontaneous boiling subsided (15–20 min after the addition of the Me_2SO_4), the stirred mixture was heated gently under reflux for 15 h. The precipitate was filtered off, and the acetone was evaporated. The residue was dissolved in CH_2Cl_2 , and the soln. was washed with H_2O (2×40 ml) and dried (Na_2SO_4). Removing the solvent under reduced pressure gave **4** (22.2 g, 94%). White solid. M.p. 110–112° (CH_2Cl_2 /hexane; [15]: 113–116°). ^1H -NMR (200 MHz, CDCl_3): 7.70 (AA' of AA'BB', 2 H); 7.34 (BB' of AA'BB', 2 H); 7.13 (s, 2 H); 4.00 (s, 6 H). ^{13}C -NMR (50 MHz, CDCl_3): 151.5; 131.2; 128.3; 126.2; 108.3; 57.8.

1-(6,7-Dimethoxynaphthalene-2-yl)ethanone (5). To a stirred soln. of **4** (22 g, 117.0 mmol) in 350 ml of dry 1,2-dichloroethane was added freshly distilled AcCl (11.0 g, 140.1 mmol) in one portion and AlCl_3 (42.0 g, 315 mmol) in small portions over 1 h at 0°. The mixture was stirred for 12 h at 0°, and then 300 g of ice and 100 ml of conc. HCl were added. The org. phase was separated, and the aq. phase was extracted with CH_2Cl_2 (2×150 ml). The org. phases were combined and dried (Na_2SO_4). Evaporation of the solvents and chromatography of the residue through a short silica-gel column (10 g; CH_2Cl_2) gave **5** (23.7 g, 88%). White crystals. M.p. 104–106° (EtOH; [15]: 104–106°). ^1H -NMR (200 MHz, CDCl_3): 8.20 (br. s, 1 H); 7.76 (A of AB, dd, $J = 8.5, 1.3$, 1 H); 7.59 (B of AB, br. d, $J = 8.5$, 1 H); 7.08 (s, 1 H); 7.00 (s, 1 H); 3.91 (s, 6 H); 2.59 (s, 3 H). ^1H -NMR: in agreement with the data given in [16]. ^{13}C -NMR (50 MHz, CDCl_3): 199.8; 153.5; 152.0; 135.0; 133.9; 130.3; 128.5; 124.6; 109.5; 108.0; 57.8 (2 C); 28.2.

6,7-Dimethoxynaphthalene-2-carboxylic Acid (6). To a stirred soln. of NaOH (34.89 g, 870 mmol) in 120 ml of H_2O was added Br_2 (42.6 g, 266.2 mmol) over 30 min at 0°. To this mixture was added a soln. of **5** (20.00 g, 87.0 mmol) in 60 ml of THF slowly over 30 min at 0°. The mixture was warmed to r.t. and stirred for 8 h. The org. phase was separated in a separatory funnel and dispatched. To the aq. soln. were added ice (200 g) and 20% NaHSO_3 soln. (100 ml). After acidification of the aq. soln. with 37% HCl ($\text{pH} \leq 3$), the solidified acid **6** was filtered with suction and dried at 60° (18.2 g, 90%). White crystals, solidified. M.p. 238–240° ([15]: 240–243°). ^1H -NMR (200 MHz, $(\text{D}_6)\text{DMSO}$): 12.65 (br. s, 1 H); 8.42 (br. s, 1 H); 7.79 (m, 2 H); 7.48 (s, 1 H); 7.36 (s, 1 H); 3.90 (s, 3 H); 3.89 (s, 3 H). ^{13}C -NMR (50 MHz, $(\text{D}_6)\text{DMSO}$): 169.5; 152.9; 151.6; 133.2; 130.6; 129.8; 128.1; 127.8; 125.3; 109.5; 108.1; 57.4 (2 C).

1,2,3,4-Tetrahydro-6,7-dimethoxynaphthalene-2-carboxylic Acid (7). To a stirred soln. of **6** (10 g, 43.1 mmol) in 300 ml of liq. NH_3 at -70° were added small thinly cut pieces of Na (4.0 g, 174.0 mmol) over 1 h under N_2 atmosphere. The mixture was additionally stirred for 1 h at the same temp. Then, 20 ml of H_2O in 30 ml of EtOH was added dropwise over 1 h under N_2 . After the NH_3 and EtOH were evaporated, ice (100 g) was added. After acidification of the aq. soln. with 37% HCl ($\text{pH} \leq 3$), the solidified acid **7** was filtered with suction and dried at 40° (9.2 g, 90%). White crystals. M.p. 135–137° (CH_2Cl_2 /hexane; [17]: 136.5–137.5°). ^1H -NMR (200 MHz, CDCl_3): 11.76 (br. s, 1 H); 6.58 (s, 1 H); 6.57 (s, 1 H); 3.83 (s, 6 H); 3.01–2.93 (m, 2 H); 2.89–2.68 (m, 3 H); 2.26–2.18 (m, 1 H); 1.95–1.75 (m, 1 H). ^1H -NMR: in agreement with the data given in [13]. ^{13}C -NMR (50 MHz, CDCl_3): 183.8; 149.4; 149.3; 129.4; 128.4; 114.0; 113.8; 57.9 (2C); 41.9, 32.9; 30.0; 27.7.

Benzyl N-(1,2,3,4-Tetrahydro-6,7-dimethoxynaphthalene-2-yl)carbamate (8). Carbamate **8** was prepared from **7** according to the procedure described by *Charlton et al.* [13] in a yield of 86%. M.p. 123–125° ([13]: 130–131.5°). ^1H -NMR (200 MHz, CDCl_3): 7.34 (m, 5 H); 6.56 (s, 1 H); 6.51 (s, 1 H); 5.09 (br. s, 2 H); 5.02 (d, $J = 8$, 1 H); 4.00 (m, 1 H); 3.82 (s, 6 H); 3.01 (dd, $J = 16.2, 4.5$, 1 H); 2.81–2.76 (m, 2 H); 2.56 (dd, $J = 16.2, 8.0$, 1 H); 2.05–1.99 (m, 1 H); 1.79–1.69 (m, 1 H). ^1H -NMR: in agreement with the data given in [13]. ^{13}C -NMR (50 MHz, CDCl_3): 157.8; 149.6; 149.4; 138.6; 130.5; 130.1, 129.3; 128.9; 127.8; 114.3; 113.7; 68.6; 57.9 (2 C); 48.9; 37.5; 30.9; 28.7.

2-Amino-1,2,3,4-tetrahydro-6,7-dimethoxynaphthalene Hydrochloride (9). By a little modification, the procedure described in [18] for the reduction of 4-azido-3-hydroxybutanenitrile was applied to **8** to give **9**. Into a 100-ml flask were placed Pd/C (50 mg) and **8** (0.50 g, 1.47 mmol) in MeOH (30 ml) and CHCl_3 (1 ml). A balloon filled with H_2 gas (3 l) was fitted to the flask. The mixture was deoxygenated by flushing with H_2 and then hydrogenated for 24 h at r.t. The catalyst was removed by filtration. Concentration gave **9** (0.34 g, 95%). Beige

solid. M.p. 217–219° (MeOH/Et₂O; [8]: 220–221°). ¹H-NMR (200 MHz, D₂O): 6.62 (s, 2 H); 3.67 (s, 3 H); 3.66 (s, 3 H); 3.64–3.40 (m, 1 H); 2.96 (dd, *J* = 16.1, 4.8, 1 H); 2.73–2.60 (m, 3 H); 2.12–2.02 (m, 1 H); 1.81–1.66 (m, 1 H). ¹H-NMR: in agreement with the data given in [13]. ¹³C-NMR (50 MHz, D₂O): 151.4; 151.1; 132.0; 129.0; 116.8; 116.4; 60.3 (2 C); 52.1; 36.8; 31.1; 30.3.

2-Amino-1,2,3,4-tetrahydronaphthalene-6,7-diol Hydrobromide (10). This compound was prepared from **9** by the procedure described by Cannon *et al.* [9] in a yield of 80%. M.p. 267–269° (MeOH/Et₂O; [8]: 270–271°). ¹H-NMR (200 MHz, D₂O): 6.48 (s, 2 H); 3.45–3.31 (m, 1 H); 2.85 (dd, *J* = 16.1, 4.8, 1 H); 2.62–2.49 (m, 3 H); 2.06–1.97 (m, 1 H); 1.74–1.58 (m, 1 H). ¹³C-NMR (50 MHz, D₂O): 146.8; 146.5; 131.3; 128.4; 120.3; 119.9; 51.9; 36.4; 31.0; 30.0.

REFERENCES

- [1] J. G. Cannon, *Prog. Drug Res.* **1985**, 29, 303.
- [2] G. N. Woodruff, *Comp. Gen. Pharmacol.* **1971**, 2, 439.
- [3] R. M. Pinder, D. A. Buxton, G. N. Woodruff, *J. Pharm. Pharmacol.* **1972**, 24, 903.
- [4] G. N. Woodruff, A. O. Elkhawad, R. M. Pinder, *Eur. J. Pharmacol.* **1974**, 25, 80.
- [5] J. G. Cannon, B. Costall, P. M. Laduron, J. E. Leysen, R. J. Naylor, *Biochem. Pharmacol.* **1978**, 27, 1417.
- [6] J. W. Kebabian, D. B. Calme, *Nature* **1979**, 277, 93.
- [7] M. S. Abel, Y. Clement-Cormier, *Pharmacologist* **1978**, 20, 241.
- [8] a) R. I. Thrift, *J. Chem. Soc. C*, **1967**, 288; b) E. Schroetter, B. Buente, H. Schick, H. Niedrich, *Pharmazie* **1981**, 36, 88.
- [9] J. G. Cannon, T. Lee, H. D. Goldman, B. Costall, R. J. Naylor, *J. Med. Chem.* **1977**, 20, 1111.
- [10] A. S. Horn, C. J. Grol, D. Dijkstra, *J. Med. Chem.* **1978**, 21, 825.
- [11] A. P. S. Narula, D. I. Schuster, *Tetrahedron Lett.*, **1981**, 22, 3707.
- [12] J. E. Nordlander, M. J. Payne, F. G. Njoroge, V. M. Vishvanath, G. R. Han, G. D. Laikos, M. A. Balk, *J. Org. Chem.* **1985**, 50, 3619.
- [13] J. L. Charlton, K. Koh, G. L. Plourde, *Can. J. Chem.* **1990**, 68, 2028.
- [14] N. P. Buu-Hoi, D. Lavit, *J. Org. Chem.* **1956**, 21, 21.
- [15] K. Y. Zee-Cheng, W. H. Nyberg, C. C. Cheng, *J. Heterocyclic Chem.*, **1972**, 9, 805.
- [16] Y. Naito, M. Sugiura, Y. Yamaura, C. Fukaya, K. Yokoyama, Y. Nakagawa, T. Ikeda, M. Senda, T. Fujita, *Chem. Pharm. Bull.* **1991**, 39, 1736.
- [17] H. L. Holmes, K. M. Mann, *J. Am. Chem. Soc.* **1947**, 69, 2000.
- [18] M. E. Jung, T. J. Shaw, *J. Am. Chem. Soc.* **1980**, 102, 6304.

Received May 31, 2003