

Ligands for Brain Cholinergic Channel Receptors: Synthesis and *in Vitro* Characterization of Novel Isoxazoles and Isothiazoles as Bioisosteric Replacements for the Pyridine Ring in Nicotine

David S. Garvey,^{*,†} James T. Wasicak, Richard L. Elliott, Suzanne A. Lebold, Ann-Marie Hettinger, George M. Carrera, Nan-Horng Lin, Yun He, Mark W. Holladay, David J. Anderson, Evelyn D. Cadman, Joanna L. Raszkievicz, James P. Sullivan, and Stephen P. Arneric

Neuroscience Research Division, Pharmaceutical Discovery, Abbott Laboratories, Abbott Park, Illinois 60064

Received August 24, 1994[®]

Ligands which activate neuronal nicotinic acetylcholine receptors (nAChRs) represent a potential approach for the palliative treatment for the symptoms of memory loss associated with Alzheimer's disease (AD). Based upon this approach, a series of novel 3,5-disubstituted isoxazoles and isothiazoles were prepared and evaluated *in vitro* as cholinergic channel activators (ChCAs) of neuronal nAChRs. Many of the 3-substituted 5-(2-pyrrolidinyl)isoxazoles were found to have nanomolar binding affinities comparable to (*S*)-nicotine (**2a**) in a preparation of whole rat brain. However, in a paradigm measuring the evoked release of [³H]dopamine from a preparation of rat striatum, there were differences in the agonist potencies and efficacies of these analogues relative to **2a**. The differences in agonist potency observed between compounds of comparable binding potency may be due to differences in ligand interactions with various subtypes of neuronal nAChRs.

Introduction

Alzheimer's disease (AD) is a central nervous system neurodegenerative disorder affecting up to 10% of the population over the age of 65. Currently, it is the fourth leading cause of death in the developed world and it is anticipated that as the size of the elderly population increases, so will the number of people afflicted with AD. Although the etiology of AD is not yet known, neurochemical studies performed on autopsied brain tissue have shown that the disease adversely affects a number of brain neurochemical systems. Of all the abnormalities noted, the most pronounced and consistent change is a deficiency in markers of cholinergic tone in the neocortex and hippocampus. The degree of cholinergic dysfunction has been correlated with the severity of dementia and these collective findings have led to the cholinergic hypothesis of AD-related memory loss.^{1–3}

Pilot clinical studies have suggested that (*S*)-nicotine (**2a**) may be useful in the palliative treatment of deficits in attention and information processing associated with AD.^{4,5} Since **2a** also produces a number of adverse side effects including irritations of the gastrointestinal tract^{6,7} and negative effects upon the cardiovascular system,^{8–12} it does not represent an optimal entity for the safe and effective treatment of the symptoms of memory loss associated with AD. The shortcomings of **2a**, coupled with the existence of a diversity of mRNAs which code for the component protein subunits which form nAChRs^{13,14} prompted our efforts to prepare cholinergic channel activators (ChCAs) which might be differentiated from **2a**. By selectively interacting with the appropriate putative subtypes of neuronal nAChRs, it was felt that it might be possible to prepare ChCAs which retain the




Figure 1. The structures of **3b** and **4a** with the corresponding nornicotine (**1b**) and nicotine (**2a**) standards.


ability to produce the beneficial actions associated with neuronal nAChR stimulation by **2a** while also having a dramatically reduced propensity to elicit adverse side effects. Such ChCAs represent a potentially novel class of psychoactive drugs for the safe and effective treatment of AD.¹⁵

Recently, reports from these laboratories have described the synthesis, receptor binding properties, and rodent behavioral activities of **ABT-418** (**4a**) and **A-82695** (**3b**), two novel ChCAs in which a substituted isoxazole ring has been incorporated as a bioisosteric replacement for the pyridine ring found in (*S*)-nicotine (**2a**) and (*R*)-nornicotine (**1b**), respectively (Figure 1).^{16–18} From receptor binding studies, it was found that both **4a** and **3b** have binding potencies comparable to **2a**; however, their abilities to elicit various behavioral effects differed significantly. Beneficial cognitive enhancing activity as well as anxiolytic effects were observed in mice at doses of **4a** and **3b** which were 10–30-fold lower than the doses of **2a** required to produce the same effects. Evaluation of these compounds for adverse effects showed that the doses of **3b** or **4a** required to produce negative effects were at dose levels 6–100-fold higher than those at which **2a** showed significant liabilities.

The variety of behavioral effects elicited by **2a** *in vivo* may be pharmacologically attributed to its ability to modulate the release of several neurotransmitters including noradrenaline, dopamine, acetylcholine, and 5-hydroxytryptamine from several different brain re-

[†] Current Address: NitroMed Research Laboratories, 801 Albany St., Boston, MA 02118.

[®] Abstract published in *Advance ACS Abstracts*, December 1, 1994.

Scheme 1^a

^a Reagents: (a) *n*-BuLi, THF, -78°C , then NaHCO_3 (aq); (b) PhNCO , Et_3N , RCH_2NO_2 , benzene; or $\text{RCH}_2\text{C}(\text{N}=\text{OH})\text{Cl}$, base (c) $\text{TFA}-\text{CH}_2\text{Cl}_2$; (d) 37% (CH_2O) (aq), 88% HCO_2H , reflux.


gions.¹⁹ To develop a better understanding of the structure-activity relationships (SAR) associated with **4a** and **3b**, an expanded series of disubstituted isoxazoles and isothiazoles have been prepared and characterized *in vitro* with respect to their brain nAChR binding affinities at the sites labeled by [³H]cytisine as well as in a functional assay measuring the evoked release of [³H]dopamine.

Chemistry


The nornicotine enantiomers **1a** and **1b** were resolved via HPLC on a chiral column.¹⁶ The 3-substituted 5-(2-pyrrolidinyl)isoxazoles **3–19** were prepared as outlined in Scheme 1. The dibromovinyl derivative **23**²⁰ was converted to the terminal acetylene **24**. Reacting compound **24** with a nitrile oxide generated *in situ* from either a nitro or hydroximoyl chloride precursor afforded the protected isoxazole derivatives **25–34**. Nitrogen deprotection gave the product secondary amines **3, 6, 8, 10, 12, and 14**. Deprotection and the N-methylation under Eschweiler-Clarke conditions afforded the tertiary amines **4, 7, 9, 11, 13, 15–19**. The *N*-ethylpyrrolidine **5a** was prepared by acetylation of **3a** followed by LiAlH_4 reduction. As evidence that racemization did not occur during the synthetic sequence, the enantiomeric purities of **4a** and **4b** were determined to be >99% ee by HPLC assay on a chiral column.¹⁶

The 3-methyl-5-(2-pyrrolidinyl)isoxazole **20a** was prepared as outlined in Scheme 2. The oxime **35a** prepared from the protected (*S*)-prolinal was converted to the chloro oxime by treatment with *N*-chlorosuccinimide. Generation of the corresponding nitrile oxide and reaction with 2-bromopropene in the presence of excess triethylamine afforded **36a**. Deprotection of the nitrogen followed by N-methylation gave the desired product **20a**.

The isothiazoles **21** and **22** were prepared as outlined in Scheme 3. The (*S*)- and (*R*)-vinyl dibromides **23** were independently converted to the acetylide anions by the action of *n*-butyllithium and then quenched with acetaldehyde. Swern oxidation of the resulting alcohols produced the ketones **37**, which were then transformed into the protected isothiazoles **38** following the procedure of Lucchesini et al.²¹ Deprotection of the pyrrolidine nitrogens gave the secondary amines **21**, which were N-methylated to afford **22**.

Scheme 2^a

^a Reagents: (a) NCS, $\text{ClCH}_2\text{CH}_2\text{Cl}$, 0°C ; (b) 2-Bromopropene, Et_3N , $\text{ClCH}_2\text{CH}_2\text{Cl}$; (c) $\text{TFA}-\text{CH}_2\text{Cl}_2$, then 37% (CH_2O) (aq), 88% HCO_2H , reflux.

Scheme 3^a

^a Reagents: (a) *n*-BuLi, THF, -78°C , then CH_3CHO , then $(\text{COCl})_2$, DMSO, -60°C , Et_3N ; (b) $\text{H}_2\text{NOSO}_3\text{H}$, $\text{CH}_3\text{OH}-\text{H}_2\text{O}$, then NaSH , NaHCO_3 ; (c) HCl -dioxane; (d) 37% (CH_2O) (aq), 88% HCO_2H , reflux.

Pharmacology

Compounds were evaluated for their binding affinities to neuronal nAChRs by measuring the displacement of [³H]cytisine from a preparation of whole rat brain according to the procedure described by Arneric et al.¹⁷ The binding results of the test compounds have been normalized to their equilibrium dissociation constants (K_i).

Ligand intrinsic activity responses were determined by measuring the magnitude of the neuronal nAChR-evoked [³H]dopamine released from a prelabeled superfused preparation of rat striatal slices as described by Arneric et al.¹⁷ Studies from our laboratories utilizing this assay have shown that the evoked release response produced by the ChCA standard **2a** reaches a maximum which begins to plateau at a concentration of approximately $0.1\ \mu\text{M}$. Thus, all of the test compounds were evaluated at a single high dose concentration of $10\ \mu\text{M}$, and the results are expressed as a percentage of the response evoked by the agonist standard **2a** which is defined as producing a 100% response at a concentration of $0.1\ \mu\text{M}$. Release responses were blocked by the competitive nAChR antagonist dihydro- β -erythroidine ($10\ \mu\text{M}$). Abbreviated "three-point" dose-response curves




Figure 2. Abbreviated dose-response curves for the evoked release of [³H]dopamine from rat striatum for the N-unsubstituted pyrrolidines **1a**, **1b**, **3a**, and **3b**. Intrinsic activities are normalized to the response produced by **2a** at a concentration of 10^{-7} M, which is defined as 1.




Figure 3. Abbreviated dose-response curves for the evoked release of [³H]dopamine from rat striatum for the N-methylated pyrrolidines **2a**, **2b**, **4a**, and **4b**. Intrinsic activities are normalized to the response produced by **2a** at a concentration of 10^{-7} M, which is defined as 1.

were generated for the ChCA standards **1** and **2** as well as the isoxazoles **3** and **4** and are shown in Figures 2 and 3.

Results and Discussion

The binding and single concentration intrinsic activity profiles of the standards **1** and **2** along with the novel isoxazole and isothiazole analogues **3–22** are shown in Table 1. N-Methylation of **1** enhances the binding affinity of the compound in the (*S*)-enantiomeric series while for the (*R*)-enantiomeric form it has no effect. From the binding SAR of the 3-substituted 5-(2-pyrrolidinyl)isoxazoles (**3–19**), it is apparent that potent nanomolar affinity to brain nAChRs can be achieved with compounds having either the (*S*)- or the (*R*)-configuration. Interestingly, N-methylation of the pyrrolidine nitrogen produces divergent trends in binding potency between the enantiomeric series of isoxazole compounds. For the (*S*)-enantiomers, there is a consistent enhancement in the binding potency of at least 50-

Table 1. Neuronal nAChR *in Vitro* Profile

Standards					
no.	conf	compd	binding affinity ^a (<i>K_i</i> , nM)	% dopamine release ^{a,d}	
1a	<i>S</i>	nornicotine	22.0 ± 3.0	155 ± 2	
1b	<i>R</i>	nornicotine	17.0 ± 2.0	93 ± 6	
2a	<i>S</i>	nicotine	1.1 ± 0.4	126 ± 18 ^b	
2b	<i>R</i>	nicotine	16.0 ± 3.4	114 ± 11	

Analogues					
no.	conf	structure		binding affinity ^a (<i>K_i</i> , nM)	% dopamine release ^{a,d}
		R	R'		
X = N, Y = O					
3a	<i>S</i>	CH ₃	H	333 ± 35	43 ± 4
3b	<i>R</i>	CH ₃	H	7.4 ± 0.7	111 ± 4
4a	<i>S</i>	CH ₃	CH ₃	4.2 ± 0.6	95 ± 7
4b	<i>R</i>	CH ₃	CH ₃	52.9 ± 12.4	85 ± 10
5a	<i>S</i>	CH ₃	C ₂ H ₅	34.3 ± 4.0	64 ± 11
6a	<i>S</i>	C ₂ H ₅	H	552 ± 63	30 ± 5
6b	<i>R</i>	C ₂ H ₅	H	11.7 ± 0.1	79 ± 21
7a	<i>S</i>	C ₂ H ₅	CH ₃	7.6 ± 0.3	78 ± 8
7b	<i>R</i>	C ₂ H ₅	CH ₃	71.0 ± 6.3	46 ± 6
8a	<i>S</i>	<i>n</i> -C ₃ H ₇	H	>10000	
9a	<i>S</i>	<i>n</i> -C ₃ H ₇	CH ₃	14.0 ± 1.8	45 ± 4
10a	<i>S</i>	C ₆ H ₅	H	>10000	
11a	<i>S</i>	C ₆ H ₅	CH ₃	788 ± 160	
12a	<i>S</i>	CH ₂ C ₆ H ₅	H	3500 ^c	
13a	<i>S</i>	CH ₂ C ₆ H ₅	CH ₃	7.4 ± 0.7	30 ± 6
14a	<i>S</i>	<i>n</i> -C ₄ H ₉	H	5000 ^c	
15a	<i>S</i>	<i>n</i> -C ₄ H ₉	CH ₃	24.2 ± 1.4	38 ± 8
16a	<i>S</i>	CO ₂ C ₂ H ₅	CH ₃	890 ± 84	
17a	<i>S</i>	CH ₂ OH	CH ₃	126 ± 20	41 ± 10
18a	<i>S</i>	CH ₂ OCH ₃	CH ₃	21.9 ± 5.7	51 ± 4
19a	<i>S</i>	CF ₃	CH ₃	6.1 ± 2.0	92 ± 16
X = O, Y = N					
20a	<i>S</i>	CH ₃	CH ₃	1333 ± 333	
X = N, Y = S					
21a	<i>S</i>	CH ₃	H	51 ± 5	61 ± 6
21b	<i>R</i>	CH ₃	H	228 ± 23	56 ± 3
22a	<i>S</i>	CH ₃	CH ₃	222 ± 14	41 ± 5
22b	<i>R</i>	CH ₃	CH ₃	207 ± 30	22 ± 2

^a Values are the means ± SEM. ^b Value was determined at the analogue concentration of 10^{-6} M. ^c Mean value of two determinations. ^d Percent relative to **2a** at 10^{-7} M.

fold which appears to be independent of the substituent at the 3-position of the isoxazole. A methyl group on the pyrrolidine nitrogen is optimal, as there is a drop in the binding potency of the *N*-ethyl analogue **5a**. With the (*R*)-enantiomeric series, the unsubstituted pyrrolidines **3b** and **6b** have potencies which are approximately 7.5-fold higher than the corresponding *N*-methyl analogues **4b** and **7b**.

A variety of substituents have been incorporated at the 3-position of the isoxazole ring of compounds with the (*S*)-configuration. Lipophilic substituents of small to intermediate size are well-tolerated with respect to brain nAChR binding. Lower straight chain alkyls (**4a**, **7a**, **9a**, **15a**), the benzyl derivative **13a**, the ether **18a**, and the trifluoromethyl derivative **19a** all bind with *K_i* values of less than 25 nM. More polar substituents such as the ester **16a** and the alcohol **17a** bind with significantly lower affinity. The low binding potency of the phenyl-substituted derivative **11a** suggests that branching at the position adjacent to the heteroaromatic ring is not tolerated.

The substitution pattern as well as the heteroatoms contained within the 5-membered aromatic ring both appear to be very important. Isoxazole **20a**, which has the position of the nitrogen and oxygen inverted relative to **4a**, has only micromolar binding affinity. Replacement of the oxygen atom in the aromatic heterocycle of compounds **3** and **4** with a sulfur yields the corresponding isothiazoles **21** and **22**. With the exception of the pyrrolidine unsubstituted (*S*)-enantiomer **21a**, this modification resulted in analogues which have lower neuronal nAChR binding potencies than their isoxazole counterparts.

In contrast to the lack of effects that relatively small changes in the steric size of the lipophilic substituent at 3-position of the isoxazole has upon binding affinity, these modifications were found to have a much greater modulatory effect upon the maximal agonist response in the dopamine release assay. Substituents with minimal steric requirements such as the methyl group found in **3b** and **4a**, the ethyl group in **6b** and **7a**, and trifluoromethyl functionality in **19a** afforded analogues which produced a robust agonist response. In contrast, larger lipophilic substituents at this position yielded compounds which were at best only partial agonists in the functional assay. The three-point dose-response curves for 1–4 shown in Figures 2 and 3 further illustrate the potency differences of the agonists which were evaluated. With the exception of **1a**, all of the other analogues included in the Figures appear as full agonists, however, the maximal response achieved with **2a** is at a concentration at least 1 order of magnitude lower than any of the other compounds tested.

[³H]-(-)-Cytisine has been shown to bind with high affinity to the $\alpha 4\beta 2$ subtype of nAChR, the major subtype in rodent brain, accounting for greater than 90% of the (-)-nicotine binding sites.^{22,23} On the other hand, the stimulated release of dopamine from rat striatal slices has been suggested to involve $\alpha 3$ subunit activation.²⁴ Thus, the differences in the agonist potencies relative to the binding potencies upon comparison of the *in vitro* profiles of the standards with many of the isoxazole analogues may be the result of selective interactions of these ChCAs with different subtypes of neuronal nAChRs. Such a selective interaction of ChCAs with subtypes of nAChRs may also contribute to the potency differences observed *in vivo* with compounds which show a much greater therapeutic window with respect to the production of beneficial behavioral effects as opposed to the undesirable side effects also associated with nAChR activation.

In summary, a number of suitably substituted isoxazoles have been identified as potent bioisosteric replacements of the pyridine ring found in **2** when the ligands are evaluated at neuronal nAChRs. Many of the isoxazole ChCAs have binding potencies to the neuronal $\alpha 4\beta 2$ subtype which differ by less than 10-fold. In contrast, when their intrinsic activities are measured in a evoked dopamine release assay which is believed to involve the activation of a neuronal nAChR containing an $\alpha 3$ subunit, the variation in potency for the same compounds is greater than 1 order of magnitude. Selective interaction of ChCAs with subtypes of neuronal nAChRs may explain the potency differences observed between these two *in vitro* assays and may also play a part in the explaining the potency variations with

respect to positive as well as negative effects produced *in vivo*.

Experimental Section

Proton magnetic resonance spectra were obtained on a General Electric GN-300 (300 MHz) instrument or, where specified, a General Electric GN-500 (500 MHz) instrument. Chemical shifts are reported as δ values (ppm) relative to Me₄-Si as an internal standard unless otherwise indicated. Mass spectra were obtained with a Hewlett-Packard HP5965 spectrometer. These determinations were performed by the PPD Analytical Research Department, Abbott Laboratories. Elemental analyses were obtained from Robertson Microлит Laboratories, Inc. (Madison, NJ).

Flash chromatography was carried out using Merck silica gel 60 (200–400 mesh) and column chromatography was carried out using Merck silica gel 60 (70–230 mesh). Melting points are uncorrected and were determined on a Büchi melting point apparatus. Optical rotation data was obtained on a Perkin-Elmer Model 241 polarimeter. All reactions were performed under an inert atmosphere of nitrogen and with anhydrous solvents unless otherwise noted.

(2S)-2-Ethynyl-N-(tert-butyloxycarbonyl)pyrrolidine (24a). A solution of **23a** (27.1 g, 76.3 mmol) in THF (550 mL) was cooled to –75 °C using a dry ice bath, and then a 2.5 M solution of *n*-butyllithium in hexane (62.6 mL, 156.5 mmol) was added dropwise over a 15 min period. After stirring for 1 h, saturated aqueous sodium bicarbonate was added dropwise to the reaction flask. The dry ice bath was removed and an additional portion of saturated aqueous sodium bicarbonate was added. The mixture was extracted with EtOAc (3 \times), and the combined organic phases were dried over anhydrous sodium sulfate and concentrated *in vacuo*. The resulting residue was purified by flash chromatography (diethyl ether/hexane 1:6 \rightarrow 1:5) to give 11.5 g (77% yield) of the product as an oil: [α]_D²⁵ = –92.1° (c 2.20, MeOH); MS (CI) *m/e* 196 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.48 (s, 9 H) 1.85–2.25 (m, 5 H), 3.24–3.53 (m, 2 H), 4.45 (m, 1 H).

(2R)-2-Ethynyl-N-(tert-butyloxycarbonyl)pyrrolidine (24b): oil; [α]_D²⁵ = +113.0° (c 0.94, MeOH).

Method A. 3-Methyl-5-(N-(tert-butyloxycarbonyl)-2-(S)-pyrrolidinyl)isoxazole (25a). To a stirred solution of **24a** (1.45 g, 7.43 mmol) and phenyl isocyanate (1.45 mL, 13.37 mmol) in benzene (3.5 mL) was added a benzene (2 mL) solution of triethylamine (10 drops) and nitroethane (535 μ L, 7.43 mmol). A precipitate began to form about 2–3 min after addition was complete. The reaction mixture was stirred at ambient temperature for 2 h, heated at reflux for 1.5 h, and then allowed to cool to ambient temperature and stirred overnight. The reaction mixture was then filtered and the filter cake washed with benzene. The filtrate was concentrated *in vacuo* and the residue was purified using flash chromatography (EtOAc/hexane 1:8) to give, after concentration *in vacuo*, 1.02 g (55% yield) of the product as a viscous yellow oil: [α]_D²⁵ = –104.4° (c 0.90, MeOH); MS (DCI/NH₃) *m/e* 253 (M + H)⁺, 270 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆; 100 °C) δ 1.32 (s, 9 H), 1.80–1.90 (m, 3 H), 2.16 (s, 3 H), 2.19 (m, 1 H), 3.31–3.42 (m, 2 H), 4.87 (dd, 1 H), 6.04 (s, 1 H).

3-Methyl-5-(N-(tert-butyloxycarbonyl)-2(R)-pyrrolidinyl)isoxazole (25b) was prepared from **24b** (1.96 g, 10 mmol) according to method A in 60% yield. [α]_D²⁵ = +102.4° (c 0.70, MeOH).

Method B. 3-Methyl-5-(2(S)-pyrrolidinyl)isoxazole (3a). Compound **25a** (880 mg, 3.49 mmol) was dissolved in CH₂Cl₂ (7.5 mL) and cooled to 0 °C. Excess trifluoroacetic acid (7.5 mL) was added and the reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was concentrated *in vacuo*, leaving an amber oil. The oil was dissolved in saturated aqueous sodium bicarbonate solution and continuously extracted with CH₂Cl₂ for approximately 16 h. The solvent was evaporated and the residue was purified by flash chromatography (5% MeOH/CHCl₃ \rightarrow 10% MeOH/CHCl₃) to give 456 mg (86% yield) of the product: [α]_D²⁵ = –13.1° (c 0.9, MeOH); MS (DCI/NH₃) *m/e* 153 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.80–2.00

(m, 3 H), 1.99 (br s, 1 H), 2.17 (m, 1 H), 2.28 (s, 3 H), 2.96–3.16 (m, 2 H), 4.32 (dd, 1 H), 5.95 (s, 1 H).

3-Methyl-5-(2(R)-pyrrolidinyl)isoxazole (3b) was prepared from **25b** (1.41 g, 5.6 mmol) according to method B in 78% yield: $[\alpha]_D^{25} = +11.6^\circ$ (c 1.0, MeOH).

Method C. 3-Methyl-5-(2(S)-pyrrolidinyl)isoxazole Oxalate (3a-Ox). A solution of **3a** (20 mg, 0.188 mmol) in diethyl ether was prepared. To this solution was added a solution of oxalic acid (25 mg, 0.282 mmol) in diethyl ether in a dropwise fashion with rapid stirring. The resultant white precipitate was filtered and triturated with three portions of diethyl ether. The white solid was recrystallized from CH₃OH/Et₂O to give, after evaporating the residual solvent, 23.7 mg (52% yield) of the product: mp = 133–135 °C; MS (DCI/NH₃) *m/e* 253 (M + H)⁺, 270 (M + NH₄)⁺; ¹H NMR (D₂O) δ 2.11–2.33 (m, 3 H), 2.31 (s, 3 H), 2.55 (m, 1 H), 2.48 (dd, 2 H), 4.92 (t, 1 H), 6.52 (s, 1 H). Anal. (C₈H₁₂N₂O·C₂H₂O₄) C, H, N.

3-Methyl-5-(2(R)-pyrrolidinyl)isoxazole Benzoate (3b-Bz). To a stirred solution of **3b** (855 mg, 5.62 mmol) in diethyl ether was added benzoic acid (755 mg, 6.18 mmol). The reaction was allowed to stir for 1 h, after which time the ether was evaporated. The remaining solid was recrystallized from hot diethyl ether (2×) to give 601 mg (39% yield) of the product as pale-tan long needles: mp = 90.5–91.5 °C; $[\alpha]_D^{25} = +9.5^\circ$ (c 0.58, MeOH); MS (DCI/NH₃) *m/e* 153 (M + H)⁺, 170 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 1.95–2.23 (m, 3 H), 2.22 (s, 3 H), 2.31 (m, 1 H), 3.00–3.32 (m, 2 H), 4.62 (dd, *J* = 7.4 Hz, 5.9 Hz, 1 H), 6.12 (s, 1 H), 7.39–7.44 (m, 2 H), 7.51 (m, 1 H), 8.01–8.04 (m, 2 H), 8.14 (br s, 1 H). Anal. (C₈H₁₂N₂O·C₇H₆O₂) C, H, N.

Method D. 3-Methyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole (4a). A solution of **3a** (93.5 mg, 0.61 mmol), in a mixture of 37% aqueous formaldehyde (1.5 mL) and 88% aqueous formic acid (1.5 mL), was heated at reflux for 1 h. The reaction mixture was allowed to cool to ambient temperature and was then extracted with diethyl ether. The aqueous layer was made basic (pH ~ 10–11) by sequential addition of saturated aqueous sodium bicarbonate solution and solid potassium carbonate. The basic, aqueous solution was then extracted with three portions of CHCl₃ and combined with the remaining organic phase. The combined organic phases were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. The residue was purified by flash chromatography (EtOAc/hexane 1:1) to give 71 mg (70% yield) of the product as a colorless oil: $[\alpha]_D^{25} = -101^\circ$ (c 0.68, MeOH); MS (FAB) *m/e* 167 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.78–2.03 (m, 3 H), 2.17–2.42 (m, 2 H), 2.29 (s, 3 H), 2.34 (s, 3 H), 3.17 (m, 1 H), 3.43 (dd, 1 H), 5.99 (s, 1 H).

Method E. 3-Methyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole Hydrochloride (4a-HCl). Compound **4a** (1.04 g, 6.26 mmol) was dissolved in diethyl ether (100 mL) and cooled to 0 °C. While the solution was being stirred, an ethereal solution of HCl was added to the reaction, resulting in the formation of a white precipitate. The solvent was evaporated *in vacuo* and the remaining solid was recrystallized from MeOH/Et₂O to give 543 mg (86% yield) of the product as hygroscopic white needles: mp = 155–157 °C; $[\alpha]_D^{25} = -32.4^\circ$ (c 0.58, MeOH); MS (DCI/NH₃) *m/e* 167 (M + H)⁺, 184 (M + NH₄)⁺; ¹H NMR (D₂O) δ 2.23–2.48 (m, 3 H), 2.34 (s, 3 H), 2.61 (m, 1 H), 2.92 (br s, 3 H), 3.38 (m, 1 H), 3.77 (m, 1 H), 4.74–4.84 (partly buried in H₂O peak, 1 H), 6.65 (s, 1 H). Anal. (C₉H₁₄N₂O·HCl) C, H, N.

3-Methyl-5-(1-methyl-2(R)-pyrrolidinyl)isoxazole (4b) was prepared from **3b** (370 mg, 2.23 mmol) according to method D in 64% yield: $[\alpha]_D^{25} = +101.0^\circ$ (c 0.76, MeOH).

3-Methyl-5-(1-methyl-2(R)-pyrrolidinyl)isoxazole hydrochloride (4b-HCl) was prepared from **4b** (228 mg, 1.38 mmol) according to method E. The white solid obtained was recrystallized from MeOH/Et₂O to give 248 mg (89% yield) of the product as white needles: mp = 154–155 °C; $[\alpha]_D^{25} = +29.1^\circ$ (c 0.80, MeOH). Anal. (C₉H₁₄N₂O·HCl) C, H, N.

3-Methyl-5-(1-ethyl-2(S)-pyrrolidinyl)isoxazole (5a). Compound **3a** (90 mg, 0.59 mmol) and acetic anhydride (120 mg, 1.2 mmol) were combined in a 1,4-dioxane (1.5 mL) and refluxed for 1 h. The reaction was allowed to cool to ambient temperature and the solvent was evaporated *in vacuo*. The

crude product was purified by flash chromatography (2% MeOH/CHCl₃) to give 117 mg (100%) of the amide as a clear yellow oil: MS (DCI/NH₃) *m/e* 195 (M + H)⁺, 212 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 1.86–2.45 (m, 10 H), 3.47–3.74 (m, 2 H), minor conformational isomer 5.00 (d, *J* = 7.0 Hz, 1 H), major isomer 5.30 (dd, *J* = 5.9 Hz, 2.6 Hz, 1 H), minor conformational isomer 5.91 (s, 1 H), major isomer 5.96 (s, 1 H).

The amide (108 mg, 0.56 mmol) in THF (1 mL) was treated with a 1.0 M THF solution of LiAlH₄ (506 mL, 0.56 mmol). The reaction was allowed to stir at ambient temperature for 2 h and then worked up under standard conditions. The crude product was then purified using flash chromatography (EtOAc/hexane 1:1) to give 63 mg (63% yield) of the product as a clear oil: MS (DCI/NH₃) *m/e* 181 (M + H)⁺; ¹H NMR (CDCl₃) δ: 1.07 (t, *J* = 7.2 Hz, 3 H), 1.82–2.00 (m, 3 H), 2.14–2.47 (m, 3 H), 2.28 (s, 3 H), 2.72 (dq, *J* = 12.1 Hz, 7.4 Hz, 1 H), 3.23 (m, 1 H), 3.60 (dd, *J* = 8.3 Hz, 6.4 Hz, 1 H), 5.97 (s, 1 H).

3-Methyl-5-(1-ethyl-2(S)-pyrrolidinyl)isoxazole hydrochloride (5a-HCl) was prepared from **5a** (59 mg, 0.33 mmol) according to method E. The white solid obtained was recrystallized from CH₂Cl₂/hexane to give the product as fine white needles in 58% yield: mp = 166–168 °C; $[\alpha]_D^{25} = -33.3^\circ$ (c 0.33, MeOH); MS (DCI/NH₃) *m/e* 181 (M + H)⁺; ¹H NMR (D₂O) δ 1.27 (t, *J* = 7.4 Hz, 3 H), 2.18–2.41 (m, 4 H), 2.31 (s, 3 H), 2.56 (m, 1 H), 3.16 (m, 1 H), 3.28–3.33 (m, 2 H), 3.75 (m, 1 H), 6.61 (s, 1 H). Anal. (C₁₀H₁₆N₂O·HCl) C, H, N.

3-Ethyl-5-(N-(tert-butyloxycarbonyl)-2(S)-pyrrolidinyl)isoxazole (26a) was prepared from **24a** (885 mg, 4.45 mmol) and 1-nitropropane according to method A in 52% yield: MS (DCI/NH₃) *m/e* 267 (M + H)⁺, 284 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆, 100 °C) δ 1.19 (t, *J* = 7.5 Hz, 3 H), 1.34 (s, 9 H), 1.89–1.95 (m, 3 H), 2.25 (m, 1 H), 2.60 (q, *J* = 7.5 Hz, 2 H), 3.39–3.45 (m, 2 H), 4.91 (dd, *J* = 7.5 Hz, 2.5 Hz, 1 H), 6.10 (s, 1 H).

3-Ethyl-5-(2(S)-pyrrolidinyl)isoxazole (6a) was prepared from **26a** (610 mg, 2.29 mmol) according to method B in 64% yield: MS (DCI/NH₃) *m/e* 167 (M + H)⁺, 184 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 1.25 (t, 3 H), 1.80–1.95 (m, 3 H), 2.07 (br s, 1 H), 2.17 (m, 1 H), 2.66 (q, 2 H), 2.97–3.16 (m, 2 H), 4.32 (dd, 1 H), 5.99 (s, 1 H).

3-Ethyl-5-(2(S)-pyrrolidinyl)isoxazole oxalate (6a-Ox) was prepared from **6a** (51.2 mg, 0.35 mmol) according to method C. Recrystallization from MeOH/Et₂O afforded the product as white crystals in 86% yield: mp = 131–133 °C; MS (DCI/NH₃) *m/e* 167 (M + H)⁺, 184 (M + NH₄)⁺; ¹H NMR (CD₃OD) δ 1.27 (t, *J* = 7.5 Hz, 3 H), 2.17–2.31 (m, 3 H), 2.50 (m, 1 H), 2.73 (q, *J* = 7.5 Hz, 2 H), 3.42–3.47 (m, 2 H), 4.91 (buried in H₂O peak, 1 H), 6.56 (s, 1 H). Anal. (C₉H₁₄N₂O·C₂H₂O₄) C, H, N.

3-Ethyl-5-(2(R)-pyrrolidinyl)isoxazole oxalate (6b-Ox) was prepared from **6b** (84 mg, 0.51 mmol) according to method C. Recrystallization from MeOH/Et₂O afforded the product as white crystals in 67% yield: mp = 131–132 °C; $[\alpha]_D^{25} = +8.3^\circ$ (c 0.46, MeOH). Anal. (C₉H₁₄N₂O·C₂H₂O₄) C, H, N.

3-Ethyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole (7a) was prepared from **6a** (150 mg, 0.90 mmol) according to method D to afford the product as a clear colorless oil in 95% yield: ¹H NMR (CDCl₃) δ 1.26 (t, 3 H), 1.81–2.02 (m, 3 H), 2.23 (m, 1 H), 2.34 (s, 3 H), 2.67 (q, 2 H), 3.13–3.21 (m, 2 H), 3.43 (dd, 1 H), 6.03 (s, 1 H).

3-Ethyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole hydrochloride (7a-HCl) was prepared from **7a** (72 mg, 0.42 mmol) according to method E. The resultant white precipitate was triturated (4×) with diethyl ether and then the solvent evaporated *in vacuo* to give 72 mg (80% yield) of a hygroscopic white solid: mp = 135–136 °C; $[\alpha]_D^{25} = -28.6^\circ$ (c 0.42, MeOH); MS (DCI/NH₃) *m/e* 181 (M + H)⁺, 198 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆, 500 MHz) δ 1.21 (t, *J* = 7.8 Hz, 3 H), 2.05–2.28 (m, 4 H), 2.67 (q, *J* = 7.8 Hz, 2 H), 2.81 (br s, 3 H), 3.20 (m, 1 H), 3.67 (m, 1 H), 4.72 (m, 1 H), 6.85 (br s, 1 H). Anal. (C₁₀H₁₆N₂O·HCl) C, H, N.

3-Ethyl-5-(1-methyl-2(R)-pyrrolidinyl)isoxazole hydrochloride (7b-HCl) was prepared from **7b** (130 mg, 0.72 mmol) according to method E. The white precipitate obtained was recrystallized from MeOH/Et₂O to give the product as fine

white needles in 42% yield: mp = 134–135 °C; $[\alpha]_D^{25} = +33.2^\circ$ (c 0.44, MeOH). Anal. ($C_{10}H_{16}N_2O \cdot HCl$) C, H, N.

3-*n*-Propyl-5-(*N*-(*tert*-butyloxycarbonyl)-2(S)-pyrrolidinyl)isoxazole (27a) was prepared from **24a** (1.23 g, 6.30 mmol) and 1-nitrobutane according to method A in 61% yield: $[\alpha]_D^{25} = -51.4^\circ$ (c 0.80, MeOH); MS (DCI/NH₃) *m/e* 281 (M + H)⁺, 298 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆, 500 MHz, 100 °C) δ 0.93 (t, *J* = 7.5 Hz, 3 H), 1.34 (s, 9 H), 1.64 (qt, *J* = 7.5 Hz, 7.3 Hz, 2 H), 1.88–1.95 (m, 3 H), 2.24 (m, 1 H), 2.56 (t, *J* = 7.3 Hz, 2 H), 3.37–3.46 (m, 2 H), 4.92 (dd, *J* = 8.3 Hz, 2.6 Hz, 1 H), 6.08 (s, 1 H).

3-*n*-Propyl-5-(2(S)-pyrrolidinyl)isoxazole (8a) was prepared from **27a** (987 mg, 3.52 mmol) according to method B to afford a 92% yield of the product as an amber oil: $[\alpha]_D^{25} = -11.5^\circ$ (c 1.2, MeOH); MS (DCI/NH₃) *m/e* 181 (M + H)⁺, 198 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 0.97 (t, *J* = 7.4 Hz, 3 H), 1.68 (tq, *J* = 7.5 Hz, 7.4 Hz, 2 H), 1.80–1.97 (m, 3 H), 2.06 (br s, NH), 2.18 (m, 1 H), 2.61 (t, *J* = 7.5 Hz, 2 H), 2.98–3.16 (m, 2 H), 4.32 (dd, *J* = 7.7 Hz, 5.5 Hz, 1 H), 5.97 (s, 1 H).

3-*n*-Propyl-5-(2(S)-pyrrolidinyl)isoxazole hydrochloride (8a·HCl) was prepared from **8a** (174 mg, 0.97 mmol) according to method E. The white solid obtained was recrystallized from MeOH/Et₂O to give the product as fine white needles in 72% yield: $[\alpha]_D^{25} = -4.2^\circ$ (c 0.5, MeOH); mp = 79–81 °C; MS (DCI/NH₃) *m/e* 181 (M + H)⁺; ¹H NMR (D₂O) δ 0.92 (t, *J* = 7.5 Hz, 3 H), 1.63–1.75 (m, 2 H), 2.22–2.38 (m, 3 H), 2.55 (m, 1 H), 2.69 (t, *J* = 7.4 Hz, 2 H), 3.47–3.53 (m, 2 H), 4.95 (dd, *J* = 8.1 Hz, 7.4 Hz, 1 H), 6.59 (s, 1 H). Anal. ($C_{10}H_{16}N_2O \cdot HCl$) C, H, N.

3-*n*-Propyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole (9a) was prepared from **8a** (370 mg, 2.05 mmol) according to method D to afford the product as a clear yellow oil in 74% yield: $[\alpha]_D^{25} = -84.1^\circ$ (c 1.2, MeOH); MS (DCI/NH₃) *m/e* 195 (M + H)⁺, 212 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 0.98 (t, *J* = 7.4 Hz, 3 H), 1.69 (tq, *J* = 7.7 Hz, 7.4 Hz, 2 H), 1.80–2.20 (m, 3 H), 2.24 (m, 1 H), 2.37 (m, 1 H), 2.34 (br s, 3 H), 2.62 (t, *J* = 7.7 Hz, 2 H), 3.17 (m, 1 H), 3.43 (dd, *J* = 8.1 Hz, 7.4 Hz, 1 H), 6.01 (s, 1 H).

3-*n*-Propyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole hydrochloride (9a·HCl) was prepared from **9a** (250 mg, 1.29 mmol) according to method E. Evaporation of the solvent afforded the product as hygroscopic short white needles in 90% yield: mp = 112–114 °C; $[\alpha]_D^{25} = -27.2^\circ$ (c 0.66, MeOH); MS (DCI/NH₃) *m/e* 195 (M + H)⁺; ¹H NMR (D₂O) δ 0.92 (t, *J* = 7.4 Hz, 3 H), 1.70 (tq, *J* = 7.4 Hz, 7.4 Hz, 2 H), 2.22–2.48 (m, 3 H), 2.61 (m, 1 H), 2.70 (t, 7.4 Hz, 2 H), 2.93 (br s, 3 H), 3.42 (m, 1 H), 3.77 (m, 1 H), 4.77–4.87 (partly buried in H₂O peak, 1 H), 6.70 (s, 1 H). Anal. ($C_{10}H_{16}N_2O \cdot HCl$) C, H, N.

3-Phenyl-5-(*N*-(*tert*-butyloxycarbonyl)-2(S)-pyrrolidinyl)isoxazole (28a). Benzohydroxamoyl chloride (1.10 g, 6.96 mmol) was treated with sodium hydroxide to generate the benzonitrile oxide. The nitrile oxide in diethyl ether was immediately added to a stirring solution of **24a** (680 mg, 3.48 mmol) in diethyl ether (1.75 mL). The reaction was stirred for 2 h, then refluxed for 2 h, and finally allowed to stir overnight at room temperature. The solvent was evaporated *in vacuo* and the residue purified by flash chromatography (EtOAc/hexane 1:10) to give 579 mg (53% yield) of the product as a white solid: $[\alpha]_D^{25} = -100^\circ$ (c 0.7, MeOH); mp = 88–90 °C; MS (DCI/NH₃) *m/e* 315 (M + H)⁺; ¹H NMR (DMSO-*d*₆, 500 MHz, 100 °C) δ 1.35 (s, 9 H), 1.93–2.01 (m, 3 H), 2.31 (m, 1 H), 3.44 (m, 1 H), 3.50 (m, 1 H), 5.00 (dd, *J* = 8.1 Hz, 2.5 Hz, 1 H), 6.71 (s, 1 H), 7.47–7.51 (m, 3 H), 7.81–7.83 (m, 2 H).

3-Phenyl-5-(2(S)-pyrrolidinyl)isoxazole (10a) was prepared from **28a** (547 mg, 1.74 mmol) according to method B to afford the product in 46% yield: $[\alpha]_D^{25} = -13.0^\circ$ (c 0.7, MeOH); MS (DCI/NH₃) *m/e* 215 (M + H)⁺; ¹H NMR (DMSO-*d*₆) δ 1.71–1.88 (m, 3 H), 2.14 (m, 1 H), 2.92 (t, *J* = 6.8 Hz, 2 H), 4.34 (dd, *J* = 7.35 Hz, 5.15 Hz, 1 H), 6.87 (s, 1 H), 7.47–7.54 (m, 3 H), 7.83–7.87 (m, 2 H).

3-Phenyl-5-(2(S)-pyrrolidinyl)isoxazole oxalate (10a·Ox) was prepared from **10a** (86.5 mg, 0.38 mmol) according to method C and recrystallized from MeOH/Et₂O to afford the product as a white solid in 71% yield: mp = 176–178 °C; MS (DCI/NH₃) *m/e* 215 (M + H)⁺; ¹H NMR (CD₃OD) δ 2.17–2.43

(m, 3 H), 2.58 (m, 1 H), 3.50 (t, *J* = 7.0 Hz, 2 H), 5.00 (dd, *J* = 7.72 Hz, 7.72 Hz, 1 H), 7.11 (s, 1 H), 7.48–7.52 (m, 3 H), 7.85–7.89 (m, 2 H). Anal. ($C_{13}H_{14}N_2O \cdot C_2H_2O_4$) C, H, N.

3-Phenyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole (11a) was prepared from **10a** (99.2 mg, 0.46 mmol) according to method D to give the product as a viscous oil in 88% yield: MS (DCI/NH₃) *m/e* 229 (M + H)⁺; ¹H NMR (DMSO-*d*₆) δ 1.78–1.96 (m, 3 H), 2.17–2.39 (m, 2 H), 2.27 (s, 3 H), 3.07 (m, 1 H), 3.52 (m, 1 H), 6.94 (s, 1 H), 7.46–7.54 (m, 3 H), 7.85–7.89 (m, 2 H).

3-Phenyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole oxalate (11a·Ox) was prepared from **11a** (86.5 mg, 0.38 mmol) according to method C to give the product as a white solid in 73% yield: mp = 106–108 °C; MS (DCI/NH₃) *m/e* 229 (M + H)⁺; ¹H NMR (CD₃OD) δ 2.24–2.34 (m, 2 H), 2.48 (m, 1 H), 2.63 (m, 1 H), 2.93 (s, 3 H), 3.34–3.38 (m, partially buried in MeOH peak 1 H), 3.73 (m, 1 H), 4.76 (dd, *J* = 8.4 Hz, 8.4 Hz, 1 H), 7.21 (s, 1 H), 7.48–7.55 (m, 3 H), 7.86–7.91 (m, 2 H). Anal. ($C_{14}H_{16}N_2O \cdot C_2H_2O_4$) C, H, N.

3-Benzyl-5-(*N*-(*tert*-butyloxycarbonyl)-2(S)-pyrrolidinyl)isoxazole (29a) was prepared from **24a** and 2-phenyl-1-nitroethane²⁵ according to method A in 59% yield: MS (DCI/NH₃) *m/e* 329 (M + H)⁺, 346 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆, 100 °C) δ 1.27 (s, 9 H), 1.84–1.97 (m, 3 H), 2.24 (m, 1 H), 3.33–3.46 (m, 2 H), 3.95 (s, 2 H), 4.90 (dd, *J* = 7.8 Hz, 2.7 Hz, 1 H), 6.02 (s, 1 H), 7.20–7.34 (m, 5 H).

3-Benzyl-5-(2(S)-pyrrolidinyl)isoxazole (12a) was prepared from **29a** (600 mg, 1.83 mmol) according to method B to afford the product as a pale yellow oil in 63% yield: MS (DCI/NH₃) *m/e* 229 (M + H)⁺, 246 (M + NH₄)⁺; ¹H NMR (CDCl₃/D₂O exchange) δ 1.88–1.92 (m, 3 H), 2.16 (m, 1 H), 2.93–3.10 (m, 2 H), 3.97 (s, 2 H), 4.27 (dd, *J* = 7.5 Hz, 5.3 Hz, 1 H), 5.88 (s, 1 H), 7.21–7.37 (m, 5 H).

3-Benzyl-5-(2(S)-pyrrolidinyl)isoxazole oxalate (12a·Ox) was prepared from **12a** (60.0 mg, 0.263 mmol) according to method C and recrystallized from MeOH/Et₂O to afford the product as a white solid in 90% yield: mp = 155–157 °C; $[\alpha]_D^{25} = -12.0^\circ$ (c 0.7, MeOH); MS *m/e* 229 (M + H)⁺, 246 (M + NH₄)⁺; ¹H NMR (CD₃OD) δ 2.10–2.30 (m, 3 H), 2.47 (m, 1 H), 3.38–3.44 (m, 2 H), 4.03 (s, 2 H), 4.82–4.95 (partially buried under water peak (m, 1 H), 6.48 (s, 1 H), 7.20–7.34 (m, 5 H). Anal. ($C_{14}H_{16}N_2O \cdot C_2H_2O_4$) C, H, N.

3-Benzyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole (13a) was prepared from **12a** (191 mg, 0.84 mmol) according to method D to give the product as a colorless oil in 75% yield: MS (DCI/NH₃) *m/e* 243 (M + H)⁺, 260 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 1.79–2.00 (m, 3 H), 2.18 (m, 1 H), 2.34 (m, 1 H), 2.30 (s, 3 H), 3.14 (m, 1 H), 3.40 (m, 1 H), 3.99 (s, 2 H), 5.92 (s, 1 H), 7.21–7.36 (m, 5 H).

3-Benzyl-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole hydrochloride (13a·HCl) was prepared from **13a** (97 mg, 0.40 mmol) according to method E. Evaporation of the solvent gave the product as a clear colorless viscous oil in quantitative yield: $[\alpha]_D^{25} = -22.3^\circ$ (c 0.26, MeOH); MS (DCI/NH₃) *m/e* 243 (M + H)⁺; ¹H NMR (CD₃OD) δ 2.21–2.44 (m, 3 H), 2.57 (m, 1 H), 2.91 (br s, 3 H), 3.30–3.42 (partly buried in MeOH peak, 1 H), 3.78 (m, 1 H), 4.05 (s, 2 H), 4.76 (m, 1 H), 6.60 (s, 1 H), 7.21–7.32 (m, 5 H). Anal. ($C_{15}H_{18}N_2O \cdot HCl \cdot 0.8H_2O$) C, H, N.

3-*n*-Butyl-5-(*N*-(*tert*-butyloxycarbonyl)-2(S)-pyrrolidinyl)isoxazole (30a) was prepared from **24a** and 1-nitropentane according to method A in 61% yield: $[\alpha]_D^{25} = -90.0^\circ$ (c 0.60, MeOH); MS (DCI/NH₃) *m/e* 295 (M + H)⁺, 312 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆, 500 MHz, 100 °C) δ 0.90 (t, *J* = 7.7 Hz, 3 H), 1.12–1.40 (m, 2 H), 1.34 (s, 9 H), 1.60 (tt, *J* = 7.4 Hz, 2 H), 1.87–1.96 (m, 3 H), 2.25 (m, 1 H), 2.58 (t, *J* = 7.4 Hz, 2 H), 3.37–3.47 (m, 2 H), 4.91 (dd, *J* = 7.8, 2.9 Hz, 1 H), 6.08 (s, 1 H).

3-*n*-Butyl-5-(2(S)-pyrrolidinyl)isoxazole (14a) was prepared from **30a** (540 mg, 1.83 mmol) according to method B to afford the product as a pale yellow oil in 85% yield: MS (DCI/NH₃) *m/e* 195 (M + H)⁺, 212 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 0.93 (t, *J* = 7.4 Hz, 3 H), 1.32–1.44 (m, 2 H), 1.58–1.68 (m, 2 H), 1.80–1.97 (m, 3 H), 2.13–2.23 (m, 1 H), 2.63 (t, *J* = 7.5 Hz, 2 H), 2.97–3.15 (m, 2 H), 4.31 (dd, *J* = 7.4 Hz, 5.5 Hz, 1 H), 5.97 (s, 1 H).

3-*n*-Butyl-5-(2(*S*)-pyrrolidinyl)isoxazole hydrochloride (14a-HCl) was prepared from **14a** (71.6 mg, 0.37 mmol) according to method E. Evaporation of the solvent afforded the product as a hygroscopic solid in 85% yield: $[\alpha]_D^{25} = -6.2^\circ$ (c 0.38, MeOH); MS *m/e* 195 (M + H)⁺, 212 (M + NH₄)⁺; ¹H NMR (D₂O) 0.91 (t, *J* = 7.4 Hz, 3 H), 1.33 (qt, *J* = 7.7, 7.4 Hz, 2 H), 1.65 (tt, *J* = 7.7, 7.4 Hz, 2 H), 2.18–2.34 (m, 3 H), 2.55 (m, 1 H), 2.72 (t, *J* = 7.4 Hz, 2 H), 3.49 (t, *J* = 7.0 Hz, 2 H), 4.94 (dd, *J* = 8.1, 7.7 Hz, 1 H), 6.58 (s, 1 H). Anal. (C₁₁H₁₈N₂O·HCl·0.2H₂O) C, H, N.

3-*n*-Butyl-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole (15a) was prepared from **14a** (219 mg, 1.1 mmol) according to method D to give the product as a colorless oil in 63% yield: $[\alpha]_D^{25} = -54.4^\circ$ (c 0.59, MeOH); MS (DCI/NH₃) *m/e* 209 (M + H)⁺; ¹H NMR (CDCl₃) δ 0.93 (t, *J* = 7.4 Hz, 3 H), 1.38 (br tq, *J* = 7.7 Hz, 7.0 Hz, 2 H), 1.59–1.69 (m, 3 H), 1.80–2.02 (m, 2 H), 2.23 (m, 1 H), 2.36 (m, 1 H), 2.34 (s, 3 H), 2.65 (t, *J* = 7.7 Hz, 2 H), 3.17 (m, 1 H), 3.42 (br dd, *J* = 7.0, 7.0 Hz, 1 H), 6.00 (s, 1 H).

3-*n*-Butyl-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole hydrochloride (15a-HCl) was prepared from **15a** (99.0 mg, 0.48 mmol) according to method E. Evaporation of the solvent afforded the product as hygroscopic solid in 61% yield: mp = 100–102 °C; $[\alpha]_D^{25} = -25.2^\circ$ (c 0.40, MeOH); MS (DCI/NH₃) *m/e* 209 (M + H)⁺; ¹H NMR (D₂O) δ 0.91 (t, *J* = 7.4 Hz, 3 H), 1.33 (br tq, *J* = 7.7, 7.4 Hz, 2 H), 1.66 (tt, *J* = 7.7, 7.4 Hz, 2 H), 2.22–2.48 (m, 3 H), 2.63 (m, 1 H), 2.74 (t, *J* = 7.7 Hz, 2 H), 2.91 (br s, 3 H), 3.37 (m, 1 H), 3.76 (m, 1 H), 4.74–4.82 (partly buried in H₂O peak, 1 H), 6.69 (s, 1 H). Anal. (C₁₂H₂₀N₂O·HCl·0.4H₂O) C, H, N.

3-(Ethoxycarbonyl)-5-(*N*-(*tert*-butyloxycarbonyl)-2(*S*)-pyrrolidinyl)isoxazole (31a). Ethyl chlorooximidatoacetate (5.40 g, 35.4 mmol), **24a** (2.30 g, 11.8 mmol), and dried powdered 4A sieves (2.3 g) were combined in CH₂Cl₂ (8 mL) and allowed to stir at room temperature for 13 days.²⁶ The reaction was then filtered through Celite and the filtrate concentrated *in vacuo*. The residue was purified by flash chromatography (EtOAc/hexane 1:4) to give 4.88 g of a viscous yellow oil of ~85% purity: MS *m/e* 311 (M + H)⁺, 328 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆, 500 MHz, 100 °C) δ 1.30 (t, *J* = 7.0 Hz, 3 H), 1.34 (s, 9 H), 1.91–2.01 (m, 3 H), 2.30 (m, 1 H), 3.40–3.51 (m, 2 H), 4.32 (q, *J* = 7.0 Hz, 2 H), 5.02 (dd, *J* = 8.2, 3.5 Hz, 1 H), 6.60 (s, 1 H).

3-(Ethoxycarbonyl)-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole (16a). Compound **31a** (315 mg, 1.0 mmol) was deprotected according to method B to afford the secondary amine as an amber oil in 39% yield: MS (DCI/NH₃) *m/e* 211 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.41 (t, *J* = 7.2 Hz, 3 H), 1.84–2.01 (m, 3 H), 2.25 (m, 1 H), 3.05–3.17 (m, 2 H), 4.43 (q, *J* = 7.2 Hz, 2 H), 4.47 (m, 1 H), 6.57 (s, 1 H). The amine (75 mg, 0.36 mmol) was methylated according to method D to give the product as a colorless oil in 48% yield: MS (DCI/NH₃) *m/e* 225 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.41 (t, *J* = 7.2 Hz, 3 H), 1.85–2.03 (m, 3 H), 2.21–2.45 (m, 2 H), 2.34 (s, 3 H), 3.17 (m, 1 H), 3.54 (dd, *J* = 8.1, 6.99 Hz, 1 H), 4.43 (q, *J* = 7.2 Hz, 2 H), 6.59 (s, 1 H).

3-(Ethoxycarbonyl)-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole (16a-Ox) was prepared from **16a** (35.6 mg, 0.16 mmol) according to method C. Evaporation of the solvent gave the product as a semisolid in quantitative yield: MS (DCI/NH₃) *m/e* 225 (M + H)⁺; ¹H NMR (D₂O, 500 MHz) δ 1.39 (t, *J* = 7.3 Hz, 3 H), 2.24–2.36 (m, 2 H), 2.48 (m, 1 H), 2.68 (m, 1 H), 2.98 (br s, 3 H), 3.39 (m, 1 H), 3.88 (m, 1 H), 4.47 (q, *J* = 7.3 Hz, 2 H), 4.88 (m, 1 H), 7.19 (s, 1 H). Anal. (C₁₁H₁₈N₂O₃·1.1C₂H₅O₄) C, H, N.

3-(Hydroxymethyl)-5-(*N*-(*tert*-butyloxycarbonyl)-2(*S*)-pyrrolidinyl)isoxazole (32a). To a solution of semipurified **31a** (4 g) in EtOH/H₂O (70 mL) was added KOH (1.35 g) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was then acidified with 2 N HCl and extracted with CHCl₃ (3×). The organic phase was dried over Na₂SO₄ and concentrated *in vacuo*, and the residue was purified by flash chromatography (10% MeOH/CHCl₃ → 0.5% HOAc/10% MeOH/CHCl₃) to afford, after evaporation of the solvent and azeotroping with toluene, the acid (3.09 g) as a yellow foam.

The acid (387 mg, 1.37 mmol) and 1.0 M borane–THF complex (4.80 mL, 4.80 mmol) were combined at room temperature in THF (4.5 mL) and then heated to reflux for 4 h. The reaction was cooled to room temperature and then saturated NaHCO₃ solution was added. The mixture was stirred for 1 h and then diluted with ethyl acetate. The two phases which formed were separated. The aqueous phase was extracted with CHCl₃ (1×) and the organic extract was combined with the original organic phase. The mixture was then dried over Na₂SO₄, concentrated *in vacuo*, and purified by flash chromatography (2% MeOH/CHCl₃) to give 266 mg (72% yield) of a clear viscous oil: MS (CI) *m/e* 269 (M + H)⁺; ¹H NMR (DMSO-*d*₆, 500 MHz, 100 °C) δ 1.35 (s, 9 H), 1.87–1.96 (m, 3 H), 2.25 (m, 1 H), 3.37–3.46 (m, 2 H), 4.47 (s, 2 H), 4.94 (dd, *J* = 8.4, 2.8 Hz, 1 H), 6.17 (s, 1 H).

3-(Hydroxymethyl)-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole (17a). Compound **32a** (266 mg, 1.0 mmol) was treated with 4 N HCl in 1,4-dioxane to give crude N-deprotected material. MS (DCI/NH₃) *m/e* 169 (M + H)⁺. The crude secondary amine was then methylated according to method D and the resulting crude product was purified by flash chromatography (1% MeOH/CHCl₃ → 2% MeOH/CHCl₃) to give 104 mg (58% yield) of a clear oil: MS (DCI/NH₃) *m/e* 183 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.81–2.03 (m, 3 H), 2.25 (m, 1 H), 2.34 (s, 3 H), 2.39 (m, 1 H), 3.18 (m, 1 H), 3.49 (m, 1 H), 4.74 (s, 2 H), 6.24 (s, 1 H).

3-(Hydroxymethyl)-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole hydrochloride (17a-HCl) was prepared from **17a** (97.5 mg, 0.54 mmol) according to method E. Evaporation of the solvent and titration with Et₂O afforded the product as a solid in 85% yield: MS (DCI/NH₃) *m/e* 183 (M + H)⁺; ¹H NMR (D₂O) δ 2.24–2.51 (m, 3 H), 2.65 (m, 1 H), 2.93 (s, 3 H), 3.39 (m, 1 H), 3.78 (m, 1 H), 4.76 (s, 2 H), 4.81 (m, 1 H), 6.80 (s, 1 H). Anal. (C₉H₁₄N₂O₂·HCl·0.2H₂O) C, H, N.

3-(Methoxymethyl)-5-(*N*-(*tert*-butyloxycarbonyl)-2(*S*)-pyrrolidinyl)isoxazole (33a). Compound **32a** (274 mg, 1.02 mmol), in THF (1 mL), was added to a stirring slurry of 80% sodium hydride (31 mg, 1.02 mmol), in THF (1 mL). The reaction was stirred for 15 min at ambient temperature and then iodomethane (190 mL, 3.06 mmol) was added. After an additional 15 min of stirring, the reaction was poured over a two phase solution of EtOAc/saturated aqueous NH₄Cl. The organic phase was separated, dried over Na₂SO₄, and concentrated *in vacuo*, and the residue was purified by flash chromatography (EtOAc/hexane 1:3) to afford 234 mg (81% yield) of a clear yellow oil: MS (CI) *m/e* (M + H)⁺ 283; ¹H NMR (DMSO-*d*₆, 500 MHz, 100 °C) δ 1.34 (s, 9 H), 1.87–1.96 (m, 3 H), 2.26 (m, 1 H), 3.32 (s, 3 H), 3.36–3.47 (m, 2 H), 4.43 (s, 2 H), 4.94 (dd, *J* = 7.8, 2.6 Hz, 1 H), 6.20 (s, 1 H).

3-(Methoxymethyl)-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole (18a). N-Deprotection of **33a** with 4 N HCl–dioxane followed by methylation according to method D afforded the product in 93% yield: MS (CI) *m/e* (M + H)⁺ 197; ¹H NMR (CDCl₃) δ 1.80–2.03 (m, 3 H), 2.24 (m, 1 H), 2.34 (s, 3 H), 2.36 (m, 1 H), 3.17 (m, 1 H), 3.39 (s, 3 H), 3.47 (dd, *J* = 8.1, 7.4 Hz, 1 H), 4.51 (s, 2 H), 6.20 (s, 1 H).

3-(Methoxymethyl)-5-(1-methyl-2(*S*)-pyrrolidinyl)isoxazole Fumarate (18a-Fu). Compound **18a** (34 mg, 0.17 mmol) and fumaric acid (20 mg, 0.17 mmol) were combined in MeOH (1 mL) and stirred for 30 min. The solvent was evaporated *in vacuo* and the remaining viscous oil was left on the high-vacuum line overnight to give 37 mg of the product as a clear viscous oil (60% yield): MS (CI) *m/e* (M + H)⁺ 197; ¹H NMR (MeOD) δ 2.24–2.03 (m, 3 H), 2.42 (m, 1 H), 2.58 (s, 3 H), 6.72 (s, 2 H), 2.82 (ddd, *J* = 10.7, 10.3, 8.5, 1 H), 3.39 (s, 3 H), 3.40 (m, 1 H), 4.08 (dd, *J* = 8.1, 8.1 Hz, 1 H), 4.52 (s, 2 H), 6.54 (s, 1 H). Anal. (C₁₀H₁₆N₂O₂·1.4C₄H₄O₄·0.2H₂O) C, H, N.

3-(Trifluoromethyl)-5-(*N*-(*tert*-butyloxycarbonyl)-2(*S*)-pyrrolidinyl)isoxazole (34a). To a solution of **24a** (200 mg, 1.02 mmol) in toluene (10 mL) was added solid K₂CO₃ (414 mg, 3.00 mmol) followed by freshly prepared (trifluoroacetyl)-hydroxymoyl chloride²⁷ (295 mg, 2.00 mmol), and the reaction mixture brought to reflux. After refluxing the mixture for ~20 h second aliquots of K₂CO₃ (450 mg) and (trifluoroacetyl)-hydroxymoyl chloride (592 mg, 4.01 mmol) were added, and

refluxing was continued for an additional 7 h. The reaction mixture was then diluted with Et₂O (50 mL) and washed with 20 mL portions of saturated aqueous NaHCO₃, 10% aqueous citric acid, and brine, dried (MgSO₄), and concentrated to afford the crude product as an oil. Flash chromatographic purification (EtOAc/hexane 1:6) afforded 130 mg (41% yield) of the pure product as a pale yellow oil: MS (CI) *m/e* 307 (M + H)⁺, 324 (M + NH₄)⁺; ¹H NMR (CDCl₃) δ 1.34 (s, 9 H), 1.94–2.43 (m, 4 H), 3.37–3.66 (m, 2), 4.99 (br s, 1 H) 5.11 (br s, 1 H), 6.29 (br s, 1 H), 6.35 (br s, 1 H).

3-(Trifluoromethyl)-5-(1-methyl-2(S)-pyrrolidinyl)isoxazole Hydrochloride (19a·HCl). Nitrogen deprotection of **34a** (37 mg, 0.121 mmol) according to method B followed by methylation according to method D and salt formation according to method E gave, after evaporation of the solvent, the product as a hygroscopic solid in 73% yield: MS (CI) *m/e* 221 (M + H)⁺, 238 (M + NH₄)⁺; ¹H NMR (D₂O) δ 2.28–2.43 (m, 2 H), 2.53 (m, 1 H), 2.74 (m, 1 H), 3.02 (m, 4 H), 3.47 (br s, 1 H), 3.85 (br s, 1 H), 7.28 (s, 1 H). Anal. (C₉H₁₁N₂OF₃·HCl·0.3H₂O) C, H, N.

N-(tert-Butyloxycarbonyl)pyrrolidine 2(S)-Aldoxime (35a). To a stirred 0 °C solution of hydroxylamine hydrochloride (488 mg, 7.03 mmol) in pyridine (7 mL) was added N-(tert-butyloxycarbonyl)-(S)-proline (1.40 g, 7.03 mmol). The reaction mixture was stirred for 16 h while gradually warming to room temperature. The reaction mixture was concentrated *in vacuo* and the residue partitioned between aqueous buffer (pH = 4) and EtOAc/diethyl ether (1:1). The organic phase was dried (MgSO₄), filtered, and evaporated *in vacuo*, to give 1.40 g (93% crude yield) of a oil which was carried forward without purification: MS (DCI/NH₃) *m/e* 215 (M + H)⁺.

5-Methyl-3-(N-(tert-butyloxycarbonyl)-2(S)-pyrrolidinyl)isoxazole (36a). N-Chlorosuccinimide (1.23 g, 9.09 mmol) was added to an ice-cooled solution of **35a** (1.85 g, 8.65 mmol) in dichloroethane (20 mL). The reaction was stirred for 0.75 h at 0 °C, followed by 0.5 h at room temperature. The reaction mixture was then cooled to –5 °C and an excess of 2-bromopropene was added, followed by the gradual addition of triethylamine (1.19 mL, 8.56 mmol). After stirring at 0 °C for an additional 2.5 h, excess triethylamine was added to the mixture and stirring was continued for 16 h. The reaction mixture was concentrated *in vacuo* and the solid residue dissolved in CHCl₃ (40 mL). The organic phase was washed with 10% KHSO₄, followed by saturated NaHCO₃ and brine. The organic phase was then dried (MgSO₄), filtered, and concentrated *in vacuo*. The residue was purified by flash chromatography (EtOAc/hexane 1:1) to yield 607.2 mg (27%) of the pure product: MS (DCI/NH₃) *m/e* 253 (M + H)⁺, 270 (M + NH₄)⁺; ¹H NMR (DMSO-*d*₆, 100 °C) δ 1.34 (s, 9 H), 1.86–1.97 (m, 3 H), 2.22 (m, 1 H), 2.98 (s, 3 H), 3.32–3.47 (m, 2 H), 4.84 (dd, *J* = 8.0 Hz, *J* = 3.0 Hz, 1 H), 6.03 (s, 1 H).

5-Methyl-3-(1-methyl-2(S)-pyrrolidinyl)isoxazole (20a). Deprotection of **36a** (607.2 mg, 2.41 mmol) according to method B followed by N-methylation according to method D and chromatographic purification (5% MeOH/CH₂Cl₂) afforded 108 mg (27% yield) of the product as a colorless oil: MS (DCI/NH₃) *m/e* 167 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.80–2.04 (m, 3 H), 2.15–2.37 (m, 2 H), 2.27 (s, 3 H), 2.40 (s, 3 H), 3.20 (t, *J* = 8.5 Hz, 1 H), 3.35 (t, *J* = 8.0 Hz, 1 H), 6.00 (d, *J* = 0.7 Hz, 1 H).

5-Methyl-3-(1-methyl-2(S)-pyrrolidinyl)isoxazole oxalate (20a·Ox) was prepared from **20a** (31.3 mg, 0.19 mmol) according to method C. Evaporation of the solvent afforded 24 mg (50% yield) of an oil: MS (DCI/NH₃) *m/e* 167 (M + H)⁺; ¹H NMR (D₂O) δ 2.18–2.34 (m, 3 H), 2.47 (s, 3 H), 2.65 (m, 1 H), 2.96 (s, 3 H), 3.33 (m, 1 H), 3.89 (m, 1 H), 4.60 (m, 1 H), 6.36 (s, 1 H). Anal. (C₉H₁₄N₂O·1.3C₂H₂O₄) C, H, N.

3-Keto-1-[N-(tert-butyloxycarbonyl)-2(S)-pyrrolidinyl]-1-butyne (37a). To a solution of **23a** (2.0 g, 5.63 mmol) in THF (10 mL) cooled to –75 °C was added *n*-BuLi (4.6 mL, 11.54 mmol of a 2.5 M solution in hexane) dropwise over a period of 10 min. This solution was stirred for 20 min before adding acetaldehyde (377 μL, 6.75 mmol). This mixture was allowed to warm slowly to ambient temperature over several hours. The reaction was then quenched by adding aqueous saturated sodium bicarbonate solution. The aqueous mixture was extracted with two portions of EtOAc and the organic

phase was dried over anhydrous sodium sulfate and then concentrated *in vacuo* to an orange oil. The oil was purified by column chromatography (EtOAc/hexane 1:2) to give 1.27 g (91% yield) of the product as a colorless oil; MS (DCI/NH₃) *m/e* 240 (M + H)⁺; ¹H NMR (CDCl₃) δ 1.43 (d, *J* = 6.6 Hz, 3 H), 1.48 (s, 9 H), 1.4–2.1 (m, 4 H), 3.2–3.5 (m, 3 H), 4.45 (br s, 1 H), 4.53 (q, *J* = 6.6 Hz, 1 H).

To a solution of oxalyl chloride (1.28 mL, 14.7 mmol) in CH₂Cl₂ (30 mL) at –60 °C was added DMSO (1.12 mL, 15.8 mmol). The reaction mixture was stirred for 10 min at –60 °C, and then a solution of the alcohol (1.26 g, 5.26 mmol) in CH₂Cl₂ (5 mL) was slowly added over a 2 min period. This mixture was stirred for 15 min at –60 °C before diisopropylethylamine (5.5 mL, 31.6 mmol) was added. After an additional 10 min of stirring at –60 °C, the reaction mixture was warmed to 0 °C and quenched with aqueous saturated ammonium chloride. The aqueous phase was extracted with CH₂Cl₂ and the organic extract was combined with organic phase from the original reaction mixture. The combined organic phases were then dried over anhydrous sodium sulfate and concentrated *in vacuo*. The residue was purified by column chromatography (EtOAc/hexane 1:1) to give 824 mg (66% yield) of the product as a colorless oil: [α]_D²⁵ = –142.3° (c 1.4, CH₂Cl₂); MS (DCI/NH₃) *m/e* 238 (M + H)⁺; ¹H NMR (DMSO-*d*₆) δ 1.3–1.5 (m, 9 H), 1.80–2.29 (m, 3 H), 2.30 (s, 2 H), 3.20–3.40 (m, 4 H), 4.60 (br s, 1 H).

3-Keto-1-[N-(tert-butyloxycarbonyl)-2(R)-pyrrolidinyl]-1-butyne (37b) was prepared in a manner similar to that described for **37a**: [α]_D²⁵ = +143.6° (c 1.6, CH₂Cl₂).

3-Methyl-5-(N-(tert-butyloxycarbonyl)-2(S)-pyrrolidinyl)isothiazole (38a) was prepared using a modification of the procedure described by Lucchesini *et al.*²¹ A solution of **37a** in 50% aqueous methanol (8 mL) was cooled to 0 °C. (Hydroxylamino)sulfonic acid (338 mg, 3 mmol) was added to the solution and the reaction mixture was stirred for 45 min. After the 45 min of mixing, solid sodium bicarbonate (250 mg, 3 mmol) was added to the reaction mixture, followed by the addition of 2.3 mL of a 1.4 M aqueous solution of sodium hydrosulfide (3.3 mmol). The reaction mixture was then stirred at ambient temperature for 6.5 h. The reaction mixture was diluted with brine and extracted with two portions of EtOAc. The aqueous phase was made basic by the addition of excess sodium bicarbonate and additional sodium hydrosulfide (800 μL of 1.4 M solution) was added to the aqueous reaction mixture which was then stirred overnight. The aqueous phase was again extracted with EtOAc and the organic extract was combined with the organic extracts from the previous day. The combined organic phases were dried over anhydrous sodium sulfate and concentrated *in vacuo*. The residue was purified by column chromatography (EtOAc/hexane 1:1) to yield 135 mg (17% yield) of the product as a yellow oil: [α]_D²⁵ = –90.9° (c 1.28, CH₂Cl₂); MS (DCI/NH₃) *m/e* 269 (M + H)⁺; ¹H NMR (DMSO-*d*₆) δ 1.36 (s, 9 H), 1.83–1.98 (m, 3 H), 2.31 (s, 1 H), 2.37 (s, 3 H), 3.40 (dd, *J* = 7.5, 6.1 Hz, 2 H), 5.14 (dd, *J* = 9.8, 2.4 Hz, 1 H), 6.96 (s, 1 H).

3-Methyl-5-(N-(tert-butyloxycarbonyl)-2(R)-pyrrolidinyl)isothiazole (38b) was prepared in a manner similar to that described for **38a**: [α]_D²⁵ = +107.7°.

3-Methyl-5-(2(S)-pyrrolidinyl)isothiazole Hydrochloride (21a·HCl). A saturated solution of hydrogen chloride in dioxane (2 mL) was added to **38a** (115 mg, 0.43 mmol). The reaction mixture was left at ambient temperature for 30 min before it was concentrated *in vacuo*. The residue was dissolved in CH₂Cl₂ and washed with saturated aqueous sodium bicarbonate. The organic phase was dried over anhydrous sodium sulfate and concentrated *in vacuo* to afford 40 mg of a yellow oil. A sample of this oil (9.5 mg) was purified by column chromatography on silica gel (10% EtOH/EtOAc) to afford 9 mg of a colorless oil. This amine was dissolved in ethanol (1 drop) and diethyl ether (~1.5 mL). A saturated solution of hydrogen chloride in diethyl ether was then added to this solution. A precipitate formed and was collected by centrifugation. The collected precipitate was washed with diethyl ether and dried *in vacuo* to yield the product as a white powder: mp = 129–130 °C; [α]_D²⁵ = +13.4° (c 0.8, MeOH); MS (DCI/NH₃) *m/e* 169 (M + H)⁺; ¹H NMR (D₂O) δ 2.15–2.37

(m, 3 H), 2.49 (s, 3 H), 2.65 (m, 1 H), 3.47–3.56 (m, 2 H), 5.07 (dd, $J = 8.5, 6.6$ Hz, 1 H), 7.31 (s, 1 H). Anal. ($C_8H_{12}N_2S \cdot 1.2HCl \cdot 0.1Et_2O$) C, H, N.

3-Methyl-5-(2(R)-pyrrolidinyl)isothiazole hydrochloride (21b·HCl) was prepared from **38b** (957.2 mg, 4.03 mmol) in a similar fashion as that described for **21a·HCl** to give 514 mg (62% yield) of the product as a beige solid: mp = 132–133 °C; $[\alpha]_D^{23} = -14.8^\circ$ (c 0.7, MeOH). Anal. ($C_8H_{11}N_2S \cdot HCl$) C, H, N.

3-Methyl-5-(1-methyl-2(S)-pyrrolidinyl)isothiazole Hydrochloride (22a). Methylation of **21a** (30 mg, 0.15 mmol) according to method D and salt formation according to method E gave, after evaporation of the solvent and recrystallization from EtOH/Et₂O, the product as a white powder in 30% yield: mp = 153–154 °C; MS (DCI/NH₃) m/e 183 ($M + H$)⁺, 270 ($M + NH_4$)⁺; ¹H NMR (D₂O) δ 2.21–2.39 (m, 3 H), 2.51 (s, 3 H), 2.72 (m, 1 H), 2.84 (br s, 3 H), 3.35 (m, 1 H), 3.78 (m, 1 H), 4.79 (m, 1 H), 7.39 (s, 1 H). Anal. ($C_9H_{14}N_2S \cdot HCl \cdot 0.25 H_2O$) C, H, N.

3-Methyl-5-(1-methyl-2(R)-pyrrolidinyl)isothiazole Hydrochloride (22b). Methylation of **21b** (150 mg, 0.89 mmol) according to method D and salt formation according to method E gave, after evaporation of the solvent and recrystallization from EtOH/Et₂O, the product as a white powder in 30% yield: mp = 147–149 °C. Anal. ($C_9H_{14}N_2S \cdot HCl \cdot 0.4H_2O$) C, H, N.

References

- Perry, E. K. The Cholinergic Hypothesis—Ten Years On. *Br. Med. Bull.* **1986**, *42*, 63–69.
- Bartus, R. T.; Dean, R. L.; Fisher, S. K. Cholinergic Treatment for Age-Related Memory Disturbances: Dead or Barely Coming of Age? In *Treatment Development Strategies for Alzheimer's Disease*; Crook, T., Bartus, R. T., Ferris, S., Gershon, S., Eds.; Mark Powley Assoc.: Madison, CN, 1986; pp 421–450.
- Coyle, J. T.; Price, D. L.; DeLong, M. R. Alzheimer's Disease: A Disorder of Cortical Cholinergic Innervation. *Science* **1983**, *219*, 1184–1190.
- Newhouse, P. A.; Sunderland, Y.; Tariot, P. N.; Blumhardt, C. L.; Weingartner, H.; Mellow, A. Intravenous Nicotine in Alzheimer's Disease: A Pilot Study. *Psychopharmacol.* **1988**, *95*, 171–175.
- Sahakian, B.; Jones, G.; Levy, R.; Gray, J.; Warburton, D. The Effects of Nicotine on Attention, Information Processing, and Short-Term Memory in Patients with Dementia of the Alzheimer Type. *Br. J. Psych.* **1989**, *154*, 797–800.
- Wormsley, K. G. Smoking and Duodenal Ulcer. *Gastroenterology* **1978**, *75*, 139–152.
- Eastwood, G. L. The Role of Smoking in Peptic Ulcer Disease. *J. Clin. Gastroenterol.* **1988**, *10*, 519–523.
- Gebber, G. L. Neurogenic Basis for the Rise in Blood Pressure Evoked by Nicotine in the Cat. *J. Pharmacol. Exp. Ther.* **1969**, *166*, 255–263.
- Larson, P. S.; Silvette, H. *Tobacco: Experimental and Clinical Studies*; Williams & Wilkins: Baltimore, 1968; Supplement I.
- Larson, P. S.; Silvette, H. *Tobacco: Experimental and Clinical Studies*; Williams & Wilkins: Baltimore, 1975; Supplement III.
- Quick, C. R. G.; Cotton, L. T. The Measured Effect of Stopping Smoking on Intermittent Claudication. *Br. J. Surg.* **1982**, *69*, 524–526.
- Lusby, R. J.; Bauminger, B.; Woodcock, J. P.; Skidmore, R.; Baird, R. N. Cigarette Smoking: Acute Main and Small Vessel Hemodynamic Responses in Patients with Arterial Disease. *Am. J. Surg.* **1981**, *142*, 169–173.
- Deneris, E. S.; Connolly, J.; Rogers, S. W.; DuVoisin, R. Pharmacological and Functional Diversity of Neuronal Nicotinic Acetylcholine Receptors. *Trends Pharmacol. Sci.* **1991**, *12*, 34–40.
- Sargent, P. B. The Diversity of Neuronal Nicotinic Acetylcholine Receptors. *Annu. Rev. Neurosci.* **1993**, *16*, 403–443.
- Arneric, S. P.; Williams, M. Nicotinic Agonists in Alzheimer's Disease: Does The Molecular Diversity of Nicotine Receptors Offer the Opportunity for Developing CNS Selective Cholinergic Channel Activators? In *Recent Advances in the Treatment of Neurodegenerative Disorders and Cognitive Function*; Racagni, G., Brunello, N., Langer, S. Z., Eds.; Karger: Basel 1994; Vol. 7, pp 58–70.
- Garvey, D. S.; Wasicak, J. T.; Decker, M. W.; Brioni, J. D.; Buckley, M. J.; Sullivan, J. P.; Carrera, G. M.; Holladay, M. W.; Arneric, S. P.; Williams, M. Novel Isoxazoles which Interact with Brain Cholinergic Channel Receptors Have Intrinsic Cognitive Enhancing and Anxiolytic Activities. *J. Med. Chem.* **1994**, *37*, 1055–1059.
- Arneric, S. P.; Sullivan, J. P.; Briggs, C. A.; Donnelly-Roberts, D.; Anderson, D. J.; Raszkievicz, J. L.; Hughes, M.; Cadman, E. D.; Adams, P.; Garvey, D. S.; Wasicak, J.; Williams, M. (S)-3-Methyl-5-(1-Methyl-2-Pyrrolidinyl) Isoxazole (ABT-418): A Novel Cholinergic Ligand with Cognition-Enhancing and Anxiolytic Activity: I. In Vitro Characterization. *J. Pharmacol. Exp. Ther.* **1994**, *270*, 310–318.
- Decker, M. W.; Brioni, J. P.; Sullivan, J. P.; Buckley, M. J.; Sullivan, J. P.; Radek, R. J.; Raszkievicz, J. L.; Kang, C. H.; Kim, D. J. B.; Giardina, W. J.; Wasicak, J.; Garvey, D. S.; Williams, M.; Arneric, S. P. (S)-3-Methyl-5-(1-Methyl-2-Pyrrolidinyl) Isoxazole (ABT-418): A Novel Cholinergic Ligand with Cognition-Enhancing and Anxiolytic Activity: II. In Vivo Characterization. *J. Pharmacol. Exp. Ther.* **1994**, *270*, 319–328.
- Rowell, P. P. Current Concepts on the Effects of Nicotine in the Central Nervous System. In *Tobacco Smoking and Health*; Martin W. R. et al., Eds.; Plenum: New York, 1987; pp 191–207.
- Garvey, D. S.; Wasicak, J. T.; Chung, J. Y.-L.; Shue, Y. K.; Carrera, G. M.; May, P. D.; McKinney, M. M.; Anderson, D.; Cadman, E.; Vella-Rountree, L.; Nadzan, A. M.; Williams, M. Synthesis and In Vitro Characterization of Novel Amino Terminally Modified Oxotremorine Derivatives for Brain Muscarinic Receptors. *J. Med. Chem.* **1992**, *35*, 1550–1557.
- Luchenisini, F.; Picci, N.; Pocci, M.; De Munno, A.; Bertini, V. Synthesis of 4-Unsubstituted Isothiazoles. *Heterocycles* **1989**, *29*, 97–102.
- Flores, C. M.; Rogers, S. W.; Pabreza, L. A.; Wolfe, B.; Kellar, K. J. A Subtype of Nicotinic Cholinergic Receptor is Composed of $\alpha 4$ and $\beta 2$ Subunits and is Upregulated by Chronic Nicotine Treatment. *Mol. Pharm.* **1992**, *41*, 31–37.
- Whiting, P.; Schoepfer, R.; Lindstrom, J.; Priestly, T. Structural and Pharmacological Characterization of the Major Brain Nicotinic Acetylcholine Receptor Subtype Stably Expressed in Mouse Fibroblasts. *Mol. Pharm.* **1992**, *40*, 463–472.
- Grady, S.; Marks, M. J.; Wonnacott, S.; Collins, A. C. Characterization of Nicotinic Receptor-Mediated [³H]Dopamine Release from Synaptosomes Prepared from Mouse Striatum. *J. Neurochem.* **1992**, *59*, 848–856.
- Sinhababu, A. K.; Borchart, R. T. Silica Gel-Assisted Reduction of Nitrostyrenes to 2-Aryl-1-Nitroalkanes with Sodium Borohydride. *Tetrahedron Lett.* **1983**, *24*, 227–230.
- Kim, J. N.; Ryu, E. K. 1,3-Dipolar Cycloaddition: Molecular Sieves Assisted Generation of Nitrile Oxides from Hydroximoyl Chlorides. *Heterocycles* **1990**, *31*, 1693–1697.
- Middleton, W. J. Trifluoroacetonitrile Oxide. *J. Org. Chem.* **1984**, *49*, 919–922.