

# ChemComm

## Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. Okumura, Y. Takeda, K. Kiyokawa and S. Minakata, *Chem. Commun.*, 2013, DOI: 10.1039/C3CC45469J.



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE


# Hypervalent Iodine(III)-Induced Oxidative [4+2] Annulation of *o*-Phenylenediamines and Electron-Deficient Alkynes: Direct Synthesis of Quinoxalines from Alkyne Substrates under Metal-Free Conditions

Sota Okumura,<sup>a</sup> Youhei Takeda,<sup>\*a,b</sup> Kensuke Kiyokawa<sup>a</sup> and Satoshi Minakata<sup>\*a</sup>

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX  
DOI: 10.1039/b000000x

**Hypervalent iodine(III)-induced oxidative [4+2] annulation of *o*-phenylenediamines and electron-deficient alkynes under metal-free conditions has been developed. The reaction allows for direct access to quinoxalines bearing two electron-withdrawing groups in an efficient manner.**

Quinoxaline derivatives not only constitute an important class of biologically active agents,<sup>1</sup> but also find tremendous applications in materials science such as luminescent materials<sup>2</sup> and low-band-gap polymers.<sup>3</sup> Of the reported methods,<sup>4</sup> the most widely used approach involves condensation of *o*-phenylenediamines with 1,2-dicarbonyl compounds bearing electron-rich or neutral substituents, which are generally prepared by oxidation of upstream alkynes (Scheme 1a). On the other hand, the synthetic methods of the quinoxalines bearing electron-withdrawing groups (EWGs, *e.g.*, -COR, -CO<sub>2</sub>R, -SO<sub>2</sub>R) have been poorly explored,<sup>5</sup> although such compounds can serve as promising candidates for (opto)electronic materials<sup>5a,6</sup> and as versatile synthetic intermediates. Herein we present a hypervalent iodine(III) reagent-induced oxidative [4+2] annulation of *o*-phenylenediamines and electron-deficient alkynes (Scheme 1b), which allows for direct access to electron-deficient quinoxalines from alkynes instead of diketo substrates in an efficient manner.





**Scheme 1** Synthetic approaches to quinoxalines

Recently, we have reported an oxidative dimerization of anilines through the agency of a unique and powerful iodinating reagent, *tert*-butyl hypoiodite (*t*-BuOI), leading to aromatic azo compounds in an efficient and selective manner.<sup>7</sup> The key to the success is an efficient two-fold iodination of nitrogen-center, forming ArNI<sub>2</sub> species which then serves as an electrophile to form N–N bond. Based on the findings, we envisioned that a tandem process consisting of i) the Michael-addition of *o*-phenylenediamine to an electron-deficient alkyne and ii) a

subsequent nucleophilic attack on the highly electrophilic N-center (NI<sub>2</sub>) by the resulting enamine could form dihydro quinoxaline skeleton. The following elimination of HI would produce quinoxalines. At the outset, we attempted the oxidative annulation of *o*-phenylenediamine (**1a**) and DMAD (**2a**) as a model reaction (Table 1). However, contrary to our preliminary expectation, the results were disappointing: the treatment of an equimolar mixture of **1a** and **2a** with *t*-BuOI (4 equiv) at -20 °C gave *cis,cis*-mucononitrile (**5**) as a major product, which should be formed through oxidative dearomatization and the following C–C bond cleavage of the benzene core (entries 1 and 2).<sup>8</sup> These results clearly suggested that *t*-BuOI is not appropriate oxidant for the aimed transformation, probably due to the rapid H–I exchange and dearomatization processes of **1a** prior to Michael-addition. After extensive screening of iodine-containing reagents, we were delighted to find that the employment of phenyliodine diacetate (PIDA) was highly effective for the progression of the desired annulation (entries 3–6). It should be noted that protecting group-free phenylenediamines, which are usually labile to oxidation reactions, were applicable to the annulation. Hypervalent iodine(III) compounds have been emerging as powerful reagents in organic synthesis due to their diverse reactivity as well as to high-availability and environmentally-benign character.<sup>9</sup> Specifically, PIDA and its derivatives have been utilized to develop privileged oxidative C–N bond forming reactions.<sup>10,11</sup> Nonetheless, to the best of our knowledge, hypervalent iodine(III)-mediated oxidative annulation reaction that leads to quinoxaline, has not been reported to date.<sup>12</sup> Intriguingly, a significant solvent effect was observed (entries 3–6): as the polarity of solvent increased, the yield of **3a** was enhanced while by-product **4a**<sup>13</sup> decreased.<sup>14</sup> Other representative hypervalent iodine(III) reagents were found ineffective for the annulation (entries 7–10), and PIDA was indispensable for the annulation reaction (entry 11).

Having optimized the reaction conditions, the scope of the annulation was investigated (Table 2). A wide variety of diamines bearing an electron-rich, -neutral, and -deficient substituent reacted with DMAD to give corresponding quinoxalines **3b–3j** in high yields. Multiple-substituted diamines afforded **3k** and **3l**. Furthermore, sterically demanding diamine gave the corresponding product **3m** in excellent yield. Although the reaction of naphthalene-2,3-diamine with DMAD required


**Table 1** Summary of the screening of the reaction conditions<sup>a</sup>


Entry	Oxidant (equiv)	Solvent	T [°C]	Yield [%] <sup>b</sup>		
				3a	4a	5
1	<i>t</i> -BuOI (4)	THF	-20	12	0	64
2	<i>t</i> -BuOI (4)	DME	-20	2	0	33
3	PhI(OAc) <sub>2</sub> (2)	CH <sub>2</sub> Cl <sub>2</sub>	-20	5	40	0
4	PhI(OAc) <sub>2</sub> (2)	THF	-20	63	18	0
5	PhI(OAc) <sub>2</sub> (2)	DME	-20	60 <sup>c</sup>	4 <sup>c</sup>	0
6	PhI(OAc) <sub>2</sub> (2)	DMF	-20	92 <sup>c</sup>	4 <sup>c</sup>	0
7	PhI(OCOCF <sub>3</sub> ) <sub>2</sub> (2)	DMF	-20	3	–	0
8	PhI=O (2)	DMF	-20	0	–	0
9	 (2)	DMF	-20	0	–	0
10	PhI(OH)OTs (2)	DMF	-20	0	–	0
11	–	DMF	-20	0	21	0


<sup>a</sup> Reaction conditions: **1a** (0.25 mmol), **2a** (0.25 mmol), and iodine-containing oxidant (0.50–1.0 mmol) were mixed in a solvent (3 mL) at the temperature in the column and stirred for 24 h. <sup>b</sup> <sup>1</sup>H NMR yields. <sup>c</sup> Isolated yield.

prolonged time, N-heteroacene **3n**, which constitutes a family of electron-transporting materials,<sup>15</sup> was obtained in 56% yield. Using the method, biquinoxaline **3o** was prepared in good yield. In respect to alkyne substrates, dibutyl acetylenedicarboxylate was successfully applied to the reaction conditions to afford **3p** and **3q** in 77% and 44% yield, respectively. In addition, an unsymmetrical alkyne having an ester and a sulfonyl groups also successfully underwent the annulation to give **3r** in good yield.

Taking advantage of the ester functionality, **3a** was diversely derivatized into functionalized quinoxalines (Scheme 2). For example, diester of **3a** easily underwent hydrolysis to give dicarboxylic acid **6** in high yield, which was further efficiently converted to **7** by dehydration. Moreover, anhydride **7** was successfully transformed into imide-fused quinoxaline **8** by the condensation with *p*-toluidine, which is an N-analogue of triboluminescent material.<sup>16</sup> It is noted that such compounds are quite difficult to prepare by traditional condensation method.<sup>17</sup>


**Scheme 2** Derivatization of **3a**

To investigate the reaction pathways, several experiments were conducted as follows: enamine **9**, which was readily prepared by the Michael-addition of *N*-Boc-protected *o*-phenylenediamine to DMAD,<sup>13</sup> was treated with PIDA in the presence of trifluoroacetic acid (Scheme 3).<sup>18</sup> At -20 °C, **9** underwent oxidative cyclization to give *N*-Boc dihydro quinoxaline **10** in 45% yield,<sup>19</sup> while **9** was quantitatively recovered in the absence

**Table 2** Scope of the oxidative [4+2] annulation<sup>a</sup>


Product	Yield [%]
<b>3a</b>	92%
<b>3b</b>	85%
<b>3c</b>	97%
<b>3d</b>	94%
<b>3e</b>	87%
<b>3f</b>	85%
<b>3g</b>	91%
<b>3h</b>	80%
<b>3i</b>	95%
<b>3j</b>	89%
<b>3k</b>	83%
<b>3l</b>	76%
<b>3m</b>	98%
<b>3n</b>	56% <sup>[d]</sup>
<b>3o</b>	71% <sup>[d]</sup>
<b>3p</b>	77%
<b>3q</b>	44% <sup>[d]</sup>
<b>3r</b>	65%


<sup>a</sup> Reaction conditions: **1** (0.25 mmol), **2** (0.25 mmol), and PhI(OAc)<sub>2</sub> (0.50 mmol) were mixed in DMF (3 mL) at -20 °C and stirred for 24 h. <sup>b</sup> The values in parentheses indicate the yields of quinoxaline products. <sup>c</sup> Reaction time: 48 h. [d] [1,1'-biphenyl]-3,3',4,4'-tetraamine (**1o**) (0.25 mmol), **2a** (0.50 mmol), PhI(OAc)<sub>2</sub> (1.0 mmol) were employed.

**Scheme 3** Oxidative cyclization of enamine **9**

of PIDA. In contrast, at room temperature, quinoxaline **3a** was obtained in 54% yield. In conjunction with the fact that DMAD does not react with PIDA in the absence of *o*-phenylenediamine, the most likely intermediate of the annulation would be the deprotected counterpart of the Michael-adduct **9** as preliminary assumed.

On the basis of the experimental results and accumulated literature knowledge about hypervalent iodine(III)-mediated oxidative C–N bond forming reactions using enamine substrates,<sup>20–22</sup> conceivable reaction pathways are illustrated in Scheme 4. The reaction would start with Michael addition of *o*-phenylenediamine to DMAD, forming enamine intermediate **A**, which has three possible reactive points when react with PIDA, namely, β-carbon of enamine (**a**),<sup>20</sup> enamine nitrogen (**b**),<sup>21</sup> and nitrogen on the benzene moiety (**c**).<sup>22</sup> Accordingly, three species should be extrapolated as intermediates prior to cyclization: i) α-iodo(III) imine **B** (*route a*); ii, iii) enamines **C** and **D** (*route b and c, respectively*). Successive cyclizative nucleophilic substitution on the iodine-attached *sp*<sup>3</sup>-carbon (from **B**), on the enamine carbon in a pseudo-S<sub>N</sub>2' manner (from **C**), or on the electrophilic N-center (from **D**) would provide common intermediate **E**. Oxidative aromatization of **E** with another equivalent of PIDA should lead to quinoxaline **F**. Ma and Lei

reported an oxidative dimerization of aromatic amines using PIDA to give azobenzene.<sup>22</sup> No azo compounds were detected in our system, suggesting that the pathway via intermediacy of **D** (route *c*) might be excluded. On one hand, according to the reactivity of carbonyl-conjugated enamines (*i.e.*, enaminones),<sup>23</sup> electrophilic reagents, including iodine electrophiles such as BTMA·ICl<sub>2</sub>,<sup>24a</sup> I(Py)<sub>2</sub>BF<sub>4</sub>,<sup>24b</sup> and CF<sub>3</sub>CH<sub>2</sub>I(OH)(OTs),<sup>24c</sup> react exclusively at the enamine β-carbon. Taken together, we believe that the most likely reaction pathway is route *a*, although routes *b*, *c*, and other possible pathways cannot be excluded completely.<sup>25</sup>



Scheme 4 Conceivable reaction pathways

## Conclusions

In summary, a simple, efficient and metal-free synthesis of electron-deficient quinoxalines through oxidative annulation of *o*-phenylenediamines and alkynes has been developed. Further investigations into mechanism and application to the construction of functional materials are currently underway in our laboratory.

## Notes and references

<sup>a</sup> Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Yamaoka 2-1, Suita, Osaka 565-0871, Japan. Fax: +81-6-6879-7402; Tel: +81-6-6879-7400; E-mail: minakata@chem.eng.osaka-u.ac.jp

<sup>b</sup> Frontier Research Base for Global Young Researchers, Graduate School of Engineering, Osaka University, Yamadaoka 2-1, Suita, Osaka 565-0871, Japan. E-mail: takeda@chem.eng.osaka-u.ac.jp

† Electronic Supplementary Information (ESI) available: Procedure for the synthesis and experimental data for quinoxalines, NMR spectra. See DOI: 10.1039/b000000x/

‡ This research was partly supported by a research Grant from the Ogasawara Foundation for the Promotion of Science & Engineering (to Y.T.), by a Grant-in-Aid for Scientific Research (B) from the JSPS, Japan (No. 25288047, to S.M.), and by a research Grant from the Nagase Science and Technology Foundation (to S.M.). Also, Y.T. acknowledges all support from the Frontier Research Base for Global Young Researchers, Osaka University, from the Program of MEXT.

1 (a) L. Yan, F.-W. Liu, G.-F. Dai and H.-M. Liu, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 609; (b) L. E. Seitz, W. J. Sulung and R. C. Reynolds, *J. Med. Chem.*, 2002, **45**, 5604.

2 S. Achelle, C. Baudequin and N. Plé, *Dyes Pigm.*, 2013, **98**, 575.

- 3 Y. Zhang, J. Zou, H.-L. Yip, K.-S. Chen, D. F. Zeigler, Y. Sun and A. K.-Y. Jen, *Chem. Mater.*, 2011, **23**, 2289 and references therein.
- 4 For reviews of synthetic methods of quinoxalines, see: (a) D. F. Saifina and V. A. Mamedov, *Russ. Chem. Rev.*, 2010, **79**, 351; (b) G. Sakata, K. Makino and Y. Kurasawa, *Heterocycles*, 1988, **27**, 2481.
- 5 (a) P. Gawrys, T. Marszalek, E. Bartnik, M. Kucinska, J. Ulanski and M. Zagorska, *Org. Lett.*, 2011, **13**, 6090; (b) T. M. V. D. Pinho e Melo, C. S. J. Lopes, A. M. d'A. Rocha Gonsalves, A. M. Beja, J. A. Paixão, M. R. Silva and L. A. da Veiga, *J. Org. Chem.*, 2001, **67**, 66; (c) H. W. Rothkopf, D. Wöhrle, R. Müller and G. Koßmehl, *Chem. Ber.*, 1975, **108**, 875.
- 6 Q. Tang, Z. Liang, J. Liu and J. Xu, Q. Miao, *Chem. Commun.*, 2010, **46**, 2977.
- 7 (a) Y. Takeda, S. Okumura and S. Minakata, *Synthesis*, 2013, **45**, 1029; (b) Y. Takeda, S. Okumura and S. Minakata, *Angew. Chem., Int. Ed.*, 2012, **51**, 7804.
- 8 (a) V. N. Telvelar and H. M. Bachhav, *Synlett*, 2009, 2059; (b) V. N. Telvekar and B. S. Takale, *Tetrahedron Lett.*, 2010, **51**, 3940.
- 9 (a) T. Dohi and Y. Kita, *Chem. Commun.*, 2009, 2073; (b) V. V. Zhdankin and P. J. Stang, *Chem. Rev.*, 2008, **108**, 5299; (c) R. M. Moriarty, *J. Org. Chem.*, 2005, **70**, 2893; (d) T. Wirth, *Angew. Chem., Int. Ed.*, 2005, **44**, 3656; (e) V. V. Zhdankin and P. J. Stang, *Chem. Rev.*, 2002, **102**, 2523.
- 10 For a review, see: M. A. Ciufolini, N. A. Braun, S. Canesi, M. Ousmer, J. Chang and D. Chai, *Synthesis*, 2007, 3759.
- 11 For recent examples, see: (a) J. A. Souto, D. Zian and K. Muñiz, *J. Am. Chem. Soc.*, 2012, **134**, 7242; (b) C. Röben, J. A. Souto, Y. González, A. Lishchynskiy and K. Muñiz, *Angew. Chem., Int. Ed.*, 2011, **50**, 9478; (c) A. A. Kantak, S. Potavathri, R. A. Barham, K. M. Romano and B. DeBoef, *J. Am. Chem. Soc.*, 2011, **133**, 19960; (d) S. Hwan, J. Yoon and S. Chang, *J. Am. Chem. Soc.*, 2011, **133**, 5996; (e) A. P. Antonchick, R. Samanta, K. Kulikov and J. Lategahn, *Angew. Chem., Int. Ed.*, 2011, **50**, 8605; (f) Y. Du, R. Liu, G. Linn and K. Zhao, *Org. Lett.*, 2006, **8**, 5919; (g) M. Ousmer, N. A. Braun, C. Bavoux, M. Perrin and M. A. Ciufolini, *J. Am. Chem. Soc.*, 2001, **123**, 7534.
- 12 Two-step synthesis of quinoxalines through hypervalent iodine(III)-mediated oxidation of alkynes and condensation of the resultant diketones with diamines has been reported: M. Tingoli, M. Mazzella, B. Panunzi and A. Tuzi, *Eur. J. Org. Chem.*, 2011, 399.
- 13 G. Choudhary and R. K. Peddinti, *Green Chem.*, 2011, **13**, 3290.
- 14 For the detailed explanation, see the ESI.
- 15 U. H. F. Bunz, J. U. Engelhart, B. D. Lindner and M. Schaffroth, *Angew. Chem., Int. Ed.*, 2013, **52**, 3810.
- 16 H. Nakayama, J. Nishida, N. Takada, H. Sato and Y. Yamashita, *Chem. Mater.*, 2012, **24**, 671.
- 17 L. Hanaineh-Abdelnour, S. Bayyuk and R. Theodorie, *Tetrahedron*, 1999, **55**, 11859.
- 18 Although we have tried preparation of the Michael-adduct starting from **1a** (deprotection form of **9**) by a similar method reported in ref 13, it failed only to produce **4a** alone. Trifluoroacetic acid was added for the purpose of detaching the Boc group of resulting intermediates.
- 19 The deprotected counterpart of **9** was not formed at all.
- 20 P. Gao, J. Liu and Y. Wei, *Org. Lett.*, 2013, **15**, 2872.
- 21 (a) W. Liu, C. Chen and Q. Zhang, *Org. Biomol. Chem.*, 2011, **9**, 6484; (b) W. Liu, H. Jiang and L. Huang, *Org. Lett.*, 2010, **12**, 312; (c) J.-Y. Wang, S.-P. Liu and W. Yu, *Synlett*, 2009, 2529; (d) X. Li, Y. Du, Z. Liang, X. Li, Y. Pan and K. Zhao, *Org. Lett.*, 2009, **11**, 2643; (e) W. Yu, Y. Du and K. Zhao, *Org. Lett.*, 2009, **11**, 2417.
- 22 H. Ma, W. Li, J. Wang, G. Xiao, Y. Gong, C. Qi, Y. Feng, X. Li, Z. Bao, Q. Cao, Q. Sun, C. Veaceslav, F. Wang and Z. Lei, *Tetrahedron*, 2012, **68**, 8358.
- 23 (a) G. Negri, C. Kascheres and A. J. Kascheres, *J. Heterocyclic Chem.*, 2004, **41**, 461; (b) A.-Z. A. Elassar and A. A. El-Khair, *Tetrahedron*, 2003, **59**, 8463; (c) J. V. Greenhill, *Chem. Soc. Rev.*, 1977, **6**, 277.
- 24 (a) C. P. Kordik and A. B. Reitz, *J. Org. Chem.*, 1996, **61**, 5644; (b) P. J. Campos, J. Arranz and M. A. Rodriguez, *Tetrahedron Lett.*, 1997, **38**, 8397; (c) I. Papoutsis, S. Spyroudis, A. Varvoglis, J. A. Callies and V. V. Zhdankin, *Tetrahedron Lett.*, 1997, **38**, 8401.

- 25 Diels-Alder reaction of oxidatively generated 1,2-diimines and DMAD might be possible, although the matching of the frontier orbitals of these substrates are less likely to be favorable.