


ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: B. wan, Y. Hu, R. Yi, X. Yu, X. Xin and C. Wang, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC05715A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

COMMUNICATION

O-Transfer-facilitated Cyclizations of Propargylamides with TMSN₃: Selective Synthesis of Tetrazoles and Dihydroimidazoles†

Received 00th January 20xx,
Accepted 00th January 20xxYancheng Hu, Ruxia Yi, Xinzhang Yu, Xiaoyi Xin, Chunxiang Wang,* and Boshun Wan*^a

DOI: 10.1039/x0xx00000x


www.rsc.org/

An unprecedented formal [3+2] annulation of propargylamides with TMSN₃ to deliver functionalized tetrazoles is developed. Oxygen-atom transfer (OAT) from amide group to C≡C bond was realized via a NIS-triggered-cyclization/ring-opening cascade pathway. The OAT process enables the amide to serve as a two-atom unit in the reactions. Notably, in situ umpolung of azide occurred when terminal propargylamides were employed in this reaction, providing an array of diiodomethylated dihydroimidazoles.

Cyclization of alkynes containing adjacent nucleophilic centers has emerged as an exceptionally mild and practical approach for the synthesis of functionalized carbo- and heterocycles.^[1] Among them, cyclizations of ketone-, amine *N*-oxide-, nitro-, sulfoxide-, and nitron-alkynes involving an intramolecular oxygen transfer process have been investigated.^[2,3] For example, alkyne-carbonyl metathesis of ketone-alkynes to produce oxygen transferred carbocycles through a [2+2] pathway has been reported (Scheme 1a).^[2] Transition-metal-catalyzed oxygen transfer reactions of alkynes possessing the polar Z⁺-O⁻ groups have been proposed to generate reactive α-oxo carbenoids, allowing the generation of various carbonyl compounds (Scheme 1b).^[3] However, despite these achievements, oxygen-atom transfer (OAT) from amide into alkynes has never been reported before.

Recently, propargylamides have been well studied as versatile precursors for the synthesis of 5- or 6-membered oxaheterocycles.^[4,5] As our continued efforts to explore new reaction patterns of *N*-tosyl propargylamides,^[5] we envisaged that whether oxygen-atom transfer could facilitate the cyclizations of propargylamides with nucleophilic coupling partners to deliver some unexpected aza-heterocycles (Scheme 1c). Indeed, by employing TMSN₃ as the coupling partner, the iodocyclization of

internal *N*-Ts propargylamides furnished various tetrazoles **2**, whereas that of terminal substrates gave dihydroimidazoles **4** (Scheme 1d). Intramolecular OAT from the amide groups to carbon-carbon triple bonds was observed in both cyclizations. The OAT process enables the amide to serve as a two-atom unit in the reactions. Thus, we have realized the O-transfer-facilitated cyclizations of propargylamides with TMSN₃ to selectively construct functionalized tetrazoles^[6,7] and diiodomethylated dihydroimidazoles^[8] for the first time. Herein, we report the preliminary results of our investigations.



Scheme 1. An overview of oxygen transfer reactions of functionalized alkynes


Our initial studies were carried out using internal propargylamide **1a** as substrate, NIS as iodine source, and TMSN₃ as coupling partner. When the reaction was performed in DCE at 60 °C, 1,5-disubstituted tetrazole **2a** was obtained. The structure of **2a** was unambiguously confirmed by X-ray crystallography.^[9] It is noteworthy that the most known synthesis of tetrazoles involves the cycloaddition reactions between azides and nitriles.^[7] Nonetheless, our method provides an appealing alternative for the

^a Dalian Institute of Chemical Physics, Chinese Academy of Sciences, 457 Zhongshan Road, Dalian 116023, China. E-mail: bswan@dicp.ac.cn; cxwang@dicp.ac.cn; Fax: +86-411-84379223; Tel: +86-411-84379260

† Electronic Supplementary Information (ESI) available: General experimental procedures, characterization data, ¹H and ¹³C NMR spectra and X-ray crystallographic analysis of compounds **2a** (CCDC 1026405) and **4a** (CCDC 1026409). For ESI and crystallographic data in CIF or other electronic format, see DOI: 10.1039/x0xx00000x

COMMUNICATION


synthesis of functionalized tetrazoles from amides and azides,^[10] which prompted us to screen the reaction conditions for this process (see the Supporting Information). Gratifyingly, the reaction of **1a** with NIS (2.4 equiv), TMSN₃ (3.6 equiv) and H₂O (1.2 equiv) in DCE at 80 °C delivered tetrazole **2a** in 81% yield (Scheme 2). The scope of this transformation was further examined (Scheme 2). Propargylic amides possessing a broad variety of aromatic motifs in R¹, R² and R³ groups underwent this reaction smoothly, providing the corresponding tetrazoles (**2b–2k**) in good yields. The electronic effects and the positions of the substituents on the phenyl ring had less influence on the results. However, when aliphatic moiety was employed as R³ (e.g. R³ = Me) or R¹ (e.g. R¹ = *i*Pr) group, no reactions occurred. Butyl substituent in R² group was also tolerated, albeit with lower yield (**2l**). Moreover, NBS could also trigger this cyclization(**2m**), whereas NCS gave no desired product.




Scheme 2. Direct transformation of internal propargylic amides into tetrazoles. Reaction conditions: **1** (0.20 mmol), NIS (0.48 mmol), TMSN₃ (0.72 mmol), H₂O (0.24 mmol), DCE (3.0 mL), at RT for 10 min, then 80 °C for 2–8 h; isolated yields are given.

Having established the transformations of internal propargylamides, we wondered whether the chemo- and regioselectivity of the reactions can be tuned by changing the substituents on the amides. To our delight, when terminal propargylamide **3a** was subjected to the same conditions, diiodomethylated dihydroimidazole **4a** was obtained.^[9] Since dihalogen-derived compounds have been shown to possess useful biological activity,^[11] this reaction might be applicable to prepare analogous diiodomethylated heterocycles.^[12] After careful screening of the reaction conditions (see the Supporting Information), we observed that the desired product **4a** could be obtained in 80% yield at 60 °C (Scheme 3). Variations on the acyl moiety (R³) were first examined. Both electron-withdrawing and electron-donating substituents (R) on the phenyl ring were compatible with this transformation, giving the corresponding dihydroimidazo-4-ols **4b–4g** in moderate to good yields. It is noteworthy that when *p*-OCH₃ substituted propargylamide **3g** was submitted to the standard conditions, no desired product was formed. Pleasingly, switching the solvent to 1,4-dioxane could provide dihydroimidazole **4g** in 38% yield. It turns out that 1,4-dioxane is much more effective than DCE for the cyclization of heteroaryl- and alkyl-derived substrates. For example, 2-furyl,

methyl and cyclopentyl substituents in R³ group (**3h–3j**) were all well tolerated with the reaction. Unexpectedly, the R¹ group significantly influences the reactivity. When phenyl group was employed as R¹, dihydroimidazole **4k** was only obtained in 15% yield. Gratifyingly, the protocol can be further extended to oxygen-linked propargylic carboxylates bearing various aromatic moieties (**3l–3n**), furnishing the products in moderate yields. It is noted that neither NBS nor NCS could promote the cyclization of **3a**.



Scheme 3. Substrate scope. Reaction conditions: **3** (0.20 mmol), NIS (0.48 mmol), TMSN₃ (0.72 mmol), H₂O (0.40 mmol), DCE (3.0 mL), 60 °C, 8–18 h; isolated yields are given. ^[a] 1,4-dioxane (3.0 mL) was used as solvent.




Scheme 4. Synthesis of heterocyclic carbaldehydes. Reaction conditions: **4** (0.20 mmol), Martin's sulfuran dehydrating reagent (0.40 mmol), CHCl₃ (2.0 mL), at room temperature for 12–18 h; isolated yields are given.

To illustrate the synthetic potential of the products, additional investigations were performed on their further transformations. Pleasingly, we found that the resulting dihydroimidazo-4-ols and dihydrooxazo-4-ols could be efficiently converted into the corresponding carbaldehydes in the presence of Martin's sulfuran dehydrating agent in CHCl₃ (Scheme 4).^[13] The elimination of H₂O leads to diiodomethylimidazoles or diiodomethylloxazoles, followed by nucleophilic substitution and elimination of HI to furnish the final products in 55–89% yields. This general protocol can be readily applied for the construction of diversified heterocyclic carbaldehydes under mild conditions.

To probe the reaction mechanism, isotopic labelling experiments were first conducted. When ¹⁸O was incorporated into the carbonyl group of propargylamide **1a** [Eq. (1)] or **3a** [Eq. (2)]


respectively, ^{18}O -labeled products were observed for both cyclizations. Besides, when H_2^{18}O was employed in the reactions, only normal products **2a** and **4a** were obtained (see the Supporting Information). These results indicate that intramolecular OAT from amide group to carbon-carbon triple bond occurs in the reaction process. The deuterated substrate **3a-d** underwent this reaction efficiently to afford the desired product **4a-d** without any loss of deuterium (see the Supporting Information), which implies that the terminal C–H bond of alkyne is not cleaved in the reaction and the proton of hydroxyl group originates exclusively from H_2O [Eq. (3)].



Notably, only one equivalent of azide or nitrogen atom is ultimately incorporated into the product, while much excess amount of TMSN₃ (3.6 equiv) is required to promote the reactions. This phenomenon indicates that the role of TMSN₃ is not only limited to be a nucleophile (or coupling partner), but might be a counteranion.^[14] Meanwhile, in situ umpolung of azide^[15] may also occur in the reaction. To confirm this hypothesis, control experiments were then carried out [Eq. (4), entries A–F]. Indeed, the reaction of **3a** with 1 equivalent of TMSN₃ only gave trace amount of **4a** [Eq. (4), entry A]. In contrast, with the assistance of other anions (e.g. I[–], Br[–] and PhSO_2Na), non-negligible amount of product could be produced [Eq. (4), entries B–D]. It is noteworthy that the using of NaN_3 as additive could also promote this reaction [Eq. (4), entry E], albeit with much lower yield than its TMSN₃ analogue [Eq. (4), entry F].

On the basis of these preliminary mechanistic studies, a plausible mechanism is proposed (Scheme 5). In the case of internal substrate **1**, an attack of the carbonyl oxygen atom onto the I⁺-activated carbon-carbon triple bond via a 6-*endo-dig* cyclization mode produces intermediate **B**. The iminium ion of **B** can be trapped by an azide anion formed in situ from TMSN₃ and H_2O , followed by the elimination of 4-methylbenzenesulfonic acid^[5a] and retroelectrocyclization to give O-transfer intermediate **E**. Finally, an

intramolecular cyclization affords tetrazole **2**. In contrast, a 5-*exo-dig* cyclization of terminal substrate **3** occurs, resulting in the formation of intermediate **H**. The activation of exo-cyclic double bond by I⁺ and intramolecular OAT lead to the ring opening product **J**. The in situ formed azidocarbenium ion^[14] is likely to be surrounded by azide anion through a contact ion pair. Subsequently, an excess of azide anion acts as another molecule of nucleophile to promote the cyclization and furnish intermediate **K**. Ultimately, the succinimide anion traps the azide group of **K** (when X = N₃), releasing *N*-azide succinimide by an umpolung step, followed by protonation to generate the desired product **4**. However, attempts to isolate or detect the byproduct *N*-azide succinimide did not succeed, which was presumably due to its lability in the reactions. Fortunately, the molecular weight of *N*-phenylsulfonyl succinimide (239.0) was detected by LC-MS when PhSO_2Na was employed in the reaction [Eq. (4), entry D], thus providing an indirect evidence to support the proposed mechanism.



Scheme 5. Proposed mechanism

In conclusion, we have developed the O-transfer-facilitated cyclizations of propargylamides with TMSN₃ to

COMMUNICATION

Journal Name

selectively afford functionalized tetrazoles and diiodomethylated dihydroimidazoles. OAT from amide group to C=C bond was realized via a NIS-triggered-cyclization/ring-opening cascade pathway. This process enables the amide to serve as a two-atom unit in the cyclizations. The resulting products can be further transformed into the corresponding heterocyclic carbaldehydes. We expect our findings can be applicable to develop new reaction patterns of amides.

This work was supported by the NSFC (21172218 and 21572225). We also thank one reviewer for helpful suggestions on revising the manuscript.

Notes and references

- For some leading reviews, see: a) A. V. Gulevich, A. S. Dudnik, N. Chernyak and V. Gevorgyan, *Chem. Rev.*, 2013, **113**, 3084; b) P. T. Parvatkar, P. S. Parameswaran and S. G. Tilve, *Chem.-Eur. J.*, 2012, **18**, 5460; c) B. Godoi, R. F. Schumacher and G. Zeni, *Chem. Rev.*, 2011, **111**, 2937; d) A. S. K. Hashmi and G. J. Hutchings, *Angew. Chem. Int. Ed.*, 2006, **45**, 7896.
- a) L.-P. Liu, D. Malhotra, R. S. Paton, K. N. Houk and G. B. Hammond, *Angew. Chem. Int. Ed.*, 2010, **49**, 9132; b) L.-P. Liu, D. Malhotra, Z. Jin, R. S. Paton, K. N. Houk and G. B. Hammond, *Chem.-Eur. J.*, 2011, **17**, 10690; c) T. Jin, F. Yang, C. Liu and Y. Yamamoto, *Chem. Commun.*, 2009, 3533; d) J. U. Rhee and M. J. Krische, *Org. Lett.*, 2005, **7**, 2493; e) K. C. M. Kurtz, R. P. Hsung and Y. Zhang, *Org. Lett.*, 2005, **8**, 231; f) T. Jin and Y. Yamamoto, *Org. Lett.*, 2007, **9**, 5259; g) T. Jin and Y. Yamamoto, *Org. Lett.*, 2008, **10**, 3137; h) C. González-Rodríguez, L. Escalante, J. A. Varela, L. Castedo and C. Saá, *Org. Lett.*, 2009, **11**, 1531.
- For leading reviews, see: a) J. Xiao and X. Li, *Angew. Chem. Int. Ed.*, 2011, **50**, 7226; b) L. Zhang, *Acc. Chem. Res.*, 2014, **47**, 877; c) H.-S. Yeom and S. Shin, *Acc. Chem. Res.*, 2014, **47**, 966; For O-atom transfer of nitro-alkynes, see: d) X. Li, C. D. Incarvito, T. Vogel and R. H. Crabtree, *Organometallics*, 2005, **24**, 3066; e) C. V. Ramana, P. Patel, K. Vanka, B. Miao and A. Degterev, *Eur. J. Org. Chem.*, 2010, 5955; f) I. Cikotiene, *Eur. J. Org. Chem.*, 2012, 2766; For O-atom transfer of sulfoxide- and nitro-alkynes, see: g) K. Pati and R.-S. Liu, *Chem. Commun.*, 2009, 5233; h) G. Song, D. Chen, Y. Su, K. Han, C.-L. Pan, A. Jia and X. Li, *Angew. Chem. Int. Ed.*, 2011, **50**, 7791; i) D. Chen, G. Song, A. Jia and X. Li, *J. Org. Chem.*, 2011, **76**, 8488; j) B. Lu, Y. Li, Y. Wang, D. H. Aue, Y. Luo and L. Zhang, *J. Am. Chem. Soc.*, 2013, **135**, 8512; k) Y. Wang, Z. Zheng and L. Zhang, *Angew. Chem. Int. Ed.*, 2014, **53**, 9572; l) X. Zhang, Z. Qi and X. Li, *Angew. Chem. Int. Ed.*, 2014, **53**, 10794; m) R. B. Dateer and S. Chang, *J. Am. Chem. Soc.*, 2015, **137**, 4908; n) L. Kong, F. Xie, S. Yu, Z. Qi and X. Li, *Chin. J. Catal.*, 2015, **36**, 925; o) H. Yan, H. Wang, X. Li, X. Xin, C. Wang and B. Wan, *Angew. Chem. Int. Ed.*, 2015, **54**, DOI: 10.1002/anie.201503997; For O-atom transfer of pyridine-N-oxides, see: p) T. Wang, S. Shi, M. Rudolph and A. S. K. Hashmi, *Adv. Synth. Catal.*, 2014, **356**, 2337; q) T. Wang, L. Huang, S. Shi, M. Rudolph and A. S. K. Hashmi, *Chem.-Eur. J.*, 2014, **20**, 14868.
- For a recent review, see: a) Y. Hu, X. Xin and B. Wan, *Tetrahedron Lett.*, 2015, **56**, 32; For non-transition-metal-mediated cyclizations, see: b) P.-Y. Coqueron, C. Didier and M. A. Ciufolini, *Angew. Chem. Int. Ed.*, 2003, **42**, 1411; c) P. Wipf, Y. Aoyama and T. E. Benedum, *Org. Lett.*, 2004, **6**, 3593; d) E. Merkul and T. J. J. Muller, *Chem. Commun.*, 2006, 4817; e) Y. M. Pan, F. J. Zheng, H. X. Lin and Z. P. Zhan, *J. Org. Chem.*, 2009, **74**, 3148; f) A. Saito, A. Matsumoto and Y. Hanzawa, *Tetrahedron Lett.*, 2010, **51**, 2247; For transition-metal-catalyzed cyclizations, see: g) A. Arcadi, S. Cacchi, L. Cascia, G. Fabrizi and F. Marinelli, *Org. Lett.*, 2001, **3**, 2501; h) A. Baccchi, M. Costa, B. Gabriele, G. Pelizzi and G. Salerno, *J. Org. Chem.*, 2002, **67**, 4450; i) A. S. K. Hashmi, J. P. Weyrauch, W. Frey and J. W. Bats, *Org. Lett.*, 2004, **6**, 4391; j) M. D. Milton, Y. Inada, Y. Nishibayashi and S. Uemura, *Chem. Commun.*, 2004, 2712; k) R. Robles-Machín, J. Adrio and J. C. Carretero, *J. Org. Chem.*, 2006, **71**, 5023; l) E. M. Beccalli, E. Borsini, G. Broggini, G. Palmisano and S. Sottocornola, *J. Org. Chem.*, 2008, **73**, 4746; m) A. S. K. Hashmi, A. M. Schuster and F. Rominger, *Angew. Chem. Int. Ed.*, 2009, **48**, 8247; n) J. P. Weyrauch, A. S. K. Hashmi, A. Schuster, T. Hengst, S. Schetter, A. Littmann, M. Rudolph, M. Hamzic, J. Visus, F. Rominger, W. Frey and J. W. Bats, *Chem.-Eur. J.*, 2010, **16**, 956; o) A. S. K. Hashmi, A. M. Schuster, M. Schmuck and F. Rominger, *Eur. J. Org. Chem.*, 2011, 4595; p) X. Meng and S. Kim, *Org. Biomol. Chem.*, 2011, **9**, 4429; q) G. C. Senadi, W.-P. Hu, J.-S. Hsiao, J. K. Vandavasi, C.-Y. Chen and J.-J. Wang, *Org. Lett.*, 2012, **14**, 4478; r) H. v. Wachenfeldt, P. Röse, F. Paulsen, N. Loganathan and D. Strand, *Chem.-Eur. J.*, 2013, **19**, 7982.
- a) X. Xin, D. Wang, X. Li and B. Wan, *Angew. Chem. Int. Ed.*, 2012, **51**, 1693; b) Y. Hu, C. Wang, D. Wang, F. Wu and B. Wan, *Org. Lett.*, 2013, **15**, 3146; c) X. Yu, X. Xin, B. Wan and X. Li, *J. Org. Chem.*, 2013, **78**, 4895; d) Y. Hu, R. Yi, F. Wu and B. Wan, *J. Org. Chem.*, 2013, **78**, 7714; e) Y. Hu, R. Yi, C. Wang, X. Xin, F. Wu and B. Wan, *J. Org. Chem.*, 2014, **79**, 3052.
- For the potential applications of tetrazoles in medicinal, agrochemical and material sciences, see: a) L. L. Dai, S. F. Cui, G. L. V. Damu and C. H. Zhou, *Chin. J. Org. Chem.*, 2013, **33**, 224; b) J. Roh, K. Vávrová and A. Hrabálek, *Eur. J. Org. Chem.*, 2012, 6101; c) T. M. Klapötke and C. M. Sabaté, *Chem. Mater.*, 2008, **20**, 3629; d) A. Hammerl, G. Holl, T. M. Klapötke, H. Nöth and M. Warchhold, *Propell. Explos. Pyrotech.*, 2003, **28**, 165.
- Synthesis of tetrazoles involves cycloadditions between nitriles and NaN₃, see: a) J. S. Mihina and R. M. Herbst, *J. Org. Chem.*, 1950, **15**, 1082; b) W. G. Finnegan, R. A. Henry and R. Lofquist, *J. Am. Chem. Soc.*, 1958, **80**, 3908; c) Z. P. Demko and K. B. Sharpless, *J. Org. Chem.*, 2001, **66**, 7945; d) F. Himzo, Z. P. Demko, L. Noodleman and K. B. Sharpless, *J. Am. Chem. Soc.*, 2002, **124**, 12210; e) Z. P. Demko and K. B. Sharpless, *Angew. Chem. Int. Ed.*, 2002, **41**, 2110; f) Z. P. Demko and K. B. Sharpless, *Angew. Chem. Int. Ed.*, 2002, **41**, 2113; g) F. Himzo, Z. P. Demko, L. Noodleman and K. B. Sharpless, *J. Am. Chem. Soc.*, 2003, **125**, 9983; For cycloadditions between nitriles and TMSN₃, see: h) S. J. Wittenberger and B. G. Donner, *J. Org. Chem.*, 1993, **58**, 4139; i) D. Amantini, R. Beleggia, F. Fringuelli, F. Pizzo and L. Vaccaro, *J. Org. Chem.*, 2004, **69**, 2896; j) T. Jin, F. Kitahara, S. Kamijo and Y. Yamamoto, *Tetrahedron Lett.*, 2008, **49**, 2824; k) T. Jin, F. Kitahara, S. Kamijo and Y. Yamamoto, *Chem.-Asian J.*, 2008, **3**, 1575; l) J. Bonnamour and C. Bolm, *Chem.-Eur. J.*, 2009, **15**, 4543; m) F. Chen, C. Qin, Y. Cui and N. Jiao, *Angew. Chem. Int. Ed.*, 2011, **50**, 11487; n) M. Gaydou and A. M. Echavarren, *Angew. Chem. Int. Ed.*, 2013, **52**, 13468; o) X.-R. Song, Y.-P. Han, Y.-F. Qiu, Z.-H. Qiu, X.-Y. Liu, P.-F. Xu and Y.-M. Liang, *Chem.-Eur. J.*, 2014, **20**, 12046; p) B. V. Rokade, K. Gadde and K. R. Prabhu, *Adv. Synth. Catal.*, 2014, **356**, 946.
- For the synthesis of dihydroimidazoles, see: a) R. J. Ferm and J. L. Riebsomer, *Chem. Rev.*, 1954, **54**, 593; b) G. Neef, U. Eder and G. Sauer, *J. Org. Chem.*, 1981, **46**, 2824; c) D. H. Huh, H. Ryu and Y. G. Kim, *Tetrahedron*, 2004, **60**, 9857; d) S.-L. You and J. W. Kelly, *Org. Lett.*, 2004, **6**, 1681.
- CCDC 1026405 (**2a**) and CCDC 1026409 (**4a**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- Only two reports on the reactions of TMSN₃ with amides, see: a) J. V. Duncia, M. E. Pierce and J. B. Santella, *J. Org. Chem.*, 1991, **56**, 2395; b) E. W. Thomas, *Synthesis*, 1993, 767.
- A. S. Mayhoub, M. Khaliq, C. Botting, Z. Li, R. J. Kuhn and M. Cushman, *Bioorg. Med. Chem.*, 2011, **19**, 3845;
- For the synthesis of diiodo-substituted compounds, see: a) A. Guirado, B. Martiz, R. Andreu and D. Bautista, *Tetrahedron*, 2011, **67**, 5811; b) J. Liu, W. Li, C. Wang, Y. Li and Z. Li, *Tetrahedron Lett.*, 2011, **52**, 4320; c) A. Zall, D. Bensinger and B. Schmidt, *Eur. J. Org. Chem.*, 2012, 1439; d) J. Wang, H. Li, D. Zhang, P. Huang, Z. Wang, R. Zhang, Y. Liang and D. Dong, *Eur. J. Org. Chem.*, 2013, 5376; e) J. M. D'Oyley, A. E. Aliev and T. D. Sheppard, *Angew. Chem. Int. Ed.*, 2014, **53**, 10747.
- For the synthesis of 4-oxazolecarbaldehydes, see: a) E. J. Zylstra, M. W. L. She, W. A. Salamant and J. W. Leahy, *Synlett*, 2007, 623; b) S. R. Shah, S. S. Navathe, A. G. Dikundwar, T. N. Guru Row and A. T. Vasella, *Eur. J. Org. Chem.*, 2013, 264.
- a) S. Pramanik and P. Ghorai, *Org. Lett.*, 2014, **16**, 2104; b) J. P. Richard, T. L. Amyes and M. M. Toteva, *Acc. Chem. Res.*, 2001, **34**, 981.
- a) M. V. Vita and J. Waser, *Org. Lett.*, 2013, **15**, 3246; b) T. Harschneck, S. Hummel, S. F. Kirsch and P. Klahn, *Chem.-Eur. J.*, 2012, **18**, 1187.