RSC Advances

PAPER

Cite this: RSC Adv., 2014, 4, 27780

Received 14th April 2014 Accepted 10th June 2014

DOI: 10.1039/c4ra03394a

www.rsc.org/advances

Introduction

Polymer-supported catalysts have attracted much attention in recent decades due to their inherent advantages in synthetic chemistry, e.g., simplification of reaction procedures including easy recovery of the catalyst by filtration, application to automated systems, and recycling of the catalyst.1 Polymer-supported organobases have attracted much attention in recent years. For example, polymer-supported bicyclic guanidine, 1,5,7-triazabicyclo[4,4,0]dec-5-ene polystyrene (PS-TBD), acts as a good catalyst for the Henry reaction,² ring-openings of epoxides,^{3,4} aldol-type condensations and Michael additions.³ Polymer-supported formamides, phosphoramide, diazaphosphorine (PS-BEMP) and phosphine (PS-PPh₃) also act as good catalysts for carbon-carbon bond forming reactions including Michael additions,5 ring-openings of epoxides,6 cyanosilylation,⁷ allylation of aldehydes with allyltrichlorosilane and aldol reactions with trichlorosilyl enol ethers.8,9 Recently, we have also reported cyanosilylation of aldehydes, ketones, imines and aziridines catalyzed by PS-TBD.10 In an effort to apply polymer-supported organobases to other useful reactions, Mukaiyama aldol reactions of aldehydes and imines were examined.

The aldol reaction has become a fundamental method for forming carbon–carbon bonds in modern organic synthesis. One of the most commonly used and best understood is the Mukaiyama aldol reaction (addition of silyl enol ethers to carbonyl compounds).^{11,12} This method has many advantages over the previous aldol reactions, such as mild reaction conditions, nonreversibility and lower production of side products. Therefore, several efficient catalysts have been developed in the last 40 years to realize high yields and selectivities of this reaction.¹³ This reaction can be catalyzed either by a Lewis acid

Polymer-supported PPh₃ as a reusable organocatalyst for the Mukaiyama aldol and Mannich reaction[†]

Satoru Matsukawa,* Kazuki Fukazawa and Junya Kimura

An easily accessible and user-friendly polymer-supported phosphine PS-PPh₃ catalyzes the aldol reaction of aldehydes and imines. A broad range of aldehydes and imines could be applied under mild conditions using 5–10 mol% PS-PPh₃. PS-PPh₃ was easily recovered and reused with minimal loss of activity.

via activation of the electrophiles or by a Lewis base *via* activation of the nucleophiles. For example, the reaction could be catalyzed by fluoride,¹⁴ phosphines,¹⁵ lithium amides,¹⁶ lithium acetate,¹⁷ lithium alkoxides,¹⁸ quaternary ammonium dendrimers,¹⁹ ammonium phenoxide,²⁰ *N*-oxides,²¹ *N*-methyl imidazole,²² sodium phenoxide–phosphine oxides,²³ *N*-heterocyclic carbenes,²⁴ Verkade superbases²⁵ and dendritic HMPA.²⁶ Herein, we report that easily accessible and user-friendly polymer-supported phosphine PS-PPh₃ acts as an efficient and reusable catalyst for the Mukaiyama aldol and Mannich reactions.^{27,28}

Results and discussion

First, we examined the catalytic activity of various commercially available polymer-supported bases (Fig. 1) in the reaction of 4methoxybenzaldehyde with a trimethylsilyl enol ether derived from methyl isobutyrate. The reaction was carried out by adding the trimethylsilyl enol ether and aldehyde to DMF containing 5 mol% of the polymer-supported base at room temperature. The reaction was monitored by TLC. After the reaction was completed, EtOAc was added to the mixture and the polymersupported base was removed by filtration. The filtrate was concentrated under vacuum and purified by column chromatography to afford the pure product. The results are



Fig. 1 Polymer-supported bases



View Article Online

View Journal | View Issue

Department of Science Education, Faculty of Education, Ibaraki University, Ibaraki, 310-8512 Japan. E-mail: smatsuka@mx.ibaraki.ac.jp; Fax: +81 29 228 8234 † electronic supplementary information (ESI) available: Copies of ¹H and ¹³C NMR Spectra of products **3a-3l**, **5a-5b**, **7a-7f**. See DOI: 10.1039/c4ra03394a

summarized in Table 1. Good product yields were obtained in the presence of a catalytic amount of the PS-PPh₃. Interestingly, the product was obtained in lower yields when PS-TBD and PS-BEMP, which have higher basicity than PS-PPh₃, were used as catalysts. PS-DMAP was also used as a catalyst, however chemical yields were low. The reactions performed in THF, MeCN, CH₂Cl₂ and toluene were inferior when compared to that performed in DMF. Solvent free conditions (SFC) were also effective. However, a longer reaction time was required to obtain the product in a good yield.

The recovery and reuse of PS-PPh3 are illustrated in Scheme 1 for the reaction of 4-methoxybenzaldehyde with a trimethylsilyl enol ether. After the reaction was completed, ethyl acetate was added to the reaction mixture and the catalyst was recovered by filtration. The recovered catalyst was washed (H₂O, NH₃, acetone), dried and then reused. The catalyst maintained its catalytic activity after three runs.

In order to clarify the scope of this reaction, several aldehydes were examined in the presence of 10 mol% PS-PPh₃ 3 (Table 2). Aromatic aldehydes with either electron-donating or



^b PS-BEMP: ^a PS-PPh₃: diphenylphosphino-polystyrene. 2-tertbutylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphorine polystyrene. ^c PS-DMAP: N-(methyl polystyrene)-4-(methylamino) pyridine. ^d SFC: solvent free condition. ^e Reaction time: 24 h.



Scheme 1 Reuse of recovered PS-PPh₃.

Paper

Table 2 PS-PPh₃ catalyzed Mukaiyama aldol reaction of various aldehydes



	I 2a-21	38-31			
Entry	Aldehyde	Product	Time	Yield $(\%)^a$	
1	о сн ₃ о Н (2а)	3a	8 h	94	
2	О Н (2b)	3b	8 h	91	
3	о сі Н (2с)	3c	4 h	95	
4	о сн ₃ Н (2d)	3d	6 h	88	
5	NO ₂ H (2e)	3e	8 h	85	
6	Me ₂ N (2f)	3f	8 h	90	
7	(2g)	3g	8 h	86	
8	• (2h)	3h	4 h	95	
9	о Н (2i)	3i	8 h	93	
10	о Н (2j)	3j	24 h	87	
11	C_8H_{17} H (2k)	3k	24 h	91	
12		31	4 h	84	

^a Isolated yield.

electron-withdrawing groups afforded the corresponding products in high yields. Compared with the aromatic aldehyde, the aliphatic aldehyde required a longer reaction time, which was similar to that for other Lewis base catalysts. Heteroaromatic aldehydes also worked well.

This PS-PPh₃ catalyzed Mukaiyama aldol reaction was also examined by using other silyl enol ethers (Table 3). Good product yields were obtained trimethylsilyl enol ether derived from ethyl acetate. In this case, the reaction also proceeded smoothly when 1 mol% of PS-PPh₃ was used. Moderate *syn*diastereoselectivity was observed irrespective of the geometry of the silyl enol ether derived from methyl propionate. This trend is the same as previous reports. Unfortunately, we had no success with ketone silyl enol ethers under these conditions.

Furthermore, we have extended the present catalytic system to the reaction with imines (Mannich reaction).^{29–31} The corresponding β -amino esters were obtained in moderate to good yields in the presence of 10 mol% PS-PPh₃ (Table 4). In this case, the reaction rates were influenced by the type of substituent on the benzene ring. When aromatic aldimines having electron-donating groups were used, the reaction proceeded more slowly compared with those using aldimines having electron-withdrawing groups. The product was obtained in lower yields when other polymer-supported organobases such as PS-TBD and PS-BEMP were used instead of PS-PPh₃ (Table 4, entries 1 *vs.* 2, 3). We also examined three component Mannich reaction between silyl enol ether 1, benzaldehyde (2b) and tosylamine, unfortunately the product was obtained as a mixture of aldol product 3b and Mannich product 7b.

A possible mechanism is illustrated in Scheme 2. First, phosphorus atom of PS-PPh₃ coordinates the silicon atom of the silyl enol ether to form activated hexa coordinated silicon species **A** or penta coordinated silicon species **A'** and the O–Si bond is activated. Next, the nucleophilicity of the enol ether is enhanced and readily reacts with an electrophile to produce the alkoxide or amino ion **B** and silylphosphonium salt. Finally, immediate silylation occurs to give the silylated adduct **C** with regeneration of PS-PPh₃. This mechanism is similar to that of previously reported Lewis base catalyzed aldol reactions.^{13a,16-25,31}

Experimental

All reactions were performed under an argon atmosphere using oven-dried glassware. Flash column chromatography was performed using silica gel Wakogel C-200. Preparative thin-layer chromatography was carried out on silica gel Wakogel B-5F. Dehydrate DMF, THF, toluene and CH₃CN were purchased from Wako Chemical. Other commercially available reagents was used as received without further purification. Yields refer to isolated compounds estimated to be >95% pure, as determined



^{*a*} Isolated yield. ^{*b*} In this case, silyl ether product was unstable, then product was obtained as OH form after hydrolysis. ^{*c*} 1 mol% of PS-PPh₃ was used.

Table 4	PS-PPh ₃ catalyzed	aldol reaction	of various	Imines
---------	-------------------------------	----------------	------------	--------



 a Isolated yield. b PS-TBD was used instead of PS-PPh3, c PS-BEMP was used instead of PS-PPh3.



Scheme 2 Proposed mechanism.

by ¹H NMR spectroscopy. IR spectra were recorded on a JUSCO FT/IR-430 spectrometer.¹H and ¹³C NMR spectra were determined for solutions in CDCl₃ with Me₄Si as internal standard on a Bruker Avance III instrument. HRMS data were measured on a JEOL JMS-700 mass spectrometer.

General procedure for PS-PPh₃ catalyzed Mukaiyama aldol reaction of aldehyde with silyl ketene acetals

To a suspension of PS-PPh₃ (0.05 mmol) in DMF (1 mL) was added aldehyde (1.0 mmol) and silyl ketene acetal (1.25 mmol) at room temperature. After the reaction was complete (as determined by TLC), EtOAc (5 ml) was added to the mixture and PS-PPh₃ was separated by filtration. The filtrate was concentrated under vacuum and purified by column chromatography on silica gel (EtOAc : hexane = 1 : 3) to give the corresponding product. The recovered catalyst is reusable after washing (acetone, NH₄OH and water) and drying *in vacuo*.

General procedure for PS-PPh₃ catalyzed Mannich reaction of imine with silyl ketene acetals

To a suspension of PS-PPh₃ (0.10 mmol) in DMF (1 mL) was added imine (1.0 mmol) and silyl ketene acetal (1.25 mmol) at room temperature or 30 °C. After the reaction was complete (as determined by TLC), EtOAc (15 ml) was added to the mixture and PS-PPh₃ was separated by filtration. The filtrate was washed with brine and water and dried over anhydrous Na₂SO₄, then evaporated. The crude mixture was purified by preparative TLC (EtOAc : hexane = 1 : 1) to afford the corresponding product.

Methyl 3-(4-methoxyphenyl)-2,2-dimethyl-3-trimethylsilyloxy-propionate (3a).³² Colorless oil; yield: 303 mg (94%); ¹H NMR (500 MHz, CDCl3): δ 0.09 (s, 9H), 0.93 (s, 3H), 1.06 (s, 3H), 3.63 (s, 3H), 3.75 (s, 3H), 4.88 (s, 1H), 6.77 (d, J = 8.6 Hz, 5H), 7.13 (d, J = 8.6 Hz, 5H); ¹³C NMR (125 MHz, CDCl3) $\delta = 0.0$, 19.1, 21.7, 49.1, 51.6, 55.2, 78.8, 112.8, 128.8, 133.0, 158.9, 177.4; HRMS (FAB): m/z: calcd for C₁₆H₂₇O₄Si: 323.1679; found: 323.1961 [M + H]⁺.

Methyl 2,2-dimethyl-3-phenyl-3-trimethylsilyloxypropionate (3b).^{17c} Colorless oil; yield: 255 mg (91%); ¹H NMR (500 MHz, CDCl3): δ 0.04 (s, 9H), 0.97 (s, 3H), 1.11 (s, 3H), 3.66 (s, 3H), 4.96 (s, 1H), 4.92 (s, 1H), 7.22–7.32 (m, 5H); ¹³C NMR (125 MHz, CDCl3) $\delta = 0.0$, 19.0, 22.0, 49.1, 51.7, 79.2, 127.4, 127.8, 140.7, 177.3; HRMS (FAB): *m/z*: calcd for C₁₅H₂₅O₃Si: 281.1573; found: 281.1568 [M + H]⁺.

Methyl 3-(4-chlorophenyl)-2,2-dimethyl-3-trimethylsilyloxypropionate (3c).^{17e} White solid; m.p. 53 °C; yield: 299 mg (95%); ¹H NMR (500 MHz, CDCl3): δ 0.06 (s, 9H), 0.94 (s, 3H), 1.08 (s, 3H), 3.64 (s, 3H), 7.18 (d, *J* = 8.4 Hz, 5H), 7.23 (d, *J* = 8.6 Hz, 5H); ¹³C NMR (125 MHz, CDCl3) δ = 0.0, 19.2, 21.4, 48.9, 51.7, 78.5, 127.6, 129.2, 133.1, 139.5, 177.0; HRMS (FAB): *m/z*: calcd for C₁₅H₂₄ClO₃Si: 315.1183; found: 315.1175 [M + H]⁺.

Methyl 3-(4-methylphenyl)-2,2-dimethyl-3-trimethylsilyloxypropionate (3d).³³ Colorless oil; yield: 259 mg (88%); ¹H NMR (500 MHz, CDCl3): δ 0.07 (s, 9H), 0.96 (s, 3H), 1.09 (s, 3H), 2.30 (s, 3H), 3.65 (s, 3H), 4.91 (s, 1H), 7.05 (d, *J* = 8.0 Hz, 5H), 7.13 (d, *J* = 8.0 Hz, 5H); ¹³C NMR (125 MHz, CDCl3) δ = 0.0, 19.1, 21.1, 21.7, 49.0, 51.6, 79.0, 127.7, 128.1, 136.9, 137.8, 177.5; HRMS (FAB): m/z: calcd for C₁₆H₂₇O₃Si: 295.1729; found: 295.1727 [M + H]⁺.

Methyl 2,2-dimethyl-3-(4-nitromethoxyphenyl)-3-trimethylsilyloxypropionate (3e).³³ White solid; m.p. 116 °C; yield: 275 mg (85%); ¹H NMR (500 MHz, CDCl3): δ 0.04 (s, 9H), 0.97 (s, 3H), 1.11 (s, 3H), 3.65 (s, 3H), 5.03 (s, 1H), 7.43 (d, *J* = 8.5 Hz, 5H), 8.13 (d, *J* = 8.5 Hz, 5H); ¹³C NMR (125 MHz, CDCl3) δ = 0.0, 19.7, 21.3, 49.1, 52.0, 78.3, 122.8, 128.5, 147.4, 148.7, 176.5.; HRMS (FAB): *m/z*: calcd for C₁₅H₂₄NO₅Si: 326.1424; found: 326.1417 [M + H]⁺.

Methyl 2,2-dimethyl-3-(naphthalene-1-yl)-3-trimethylsilyloxy-propionate (3g).³⁴ Colorless oil; yield: 285 mg (86%); ¹H NMR (500 MHz, CDCl3): δ -0.11 (s, 9H), 0.93 (s, 3H), 1.21 (s, 3H), 3.64 (s, 3H), 6.00 (s, 1H), 7.40-7.51 (m, 3H), 7.69 (d, *J* = 7.4 Hz, 1H), 7.76 (d, *J* = 8.2 Hz, 1H), 7.82 (d, *J* = 7.4 Hz, 1H), 8.15 (d, *J* = 8.2 Hz, 1H); ¹³C NMR (125 MHz, CDCl3) δ = -0.1, 18.5, 23.0, 50.0, 51.8, 72.9, 123.2, 124.8, 125.0, 125.7, 127.0, 127.9, 128.8, 131.6, 133.2, 136.8, 177.5; HRMS (FAB): *m/z*: calcd for C₁₉H₂₇O₃Si: 331.1729; found: 331.1736 [M + H]⁺.

Methyl 2,2-dimethyl-3-(naphthalene-2-yl)-3-trimethylsilyloxy-propionate (3h).³⁵ White solid; m.p. 70 °C; yield: 315 mg (95%); ¹H NMR (500 MHz, CDCl3): δ –0.05 (s, 9H), 1.02 (s, 3H), 1.15 (s, 3H), 3.67 (s, 3H), 5.13 (s, 1H), 7.40–7.47 (m, 3H), 7.67 (d, J = 4.4 Hz, 1H), 7.74 (d, J = 8.5 Hz, 1H), 7.81 (d, J = 7.4 Hz, 2H); ¹³C NMR (125 MHz, CDCl3) δ = 0.0, 19.2, 21.8, 49.3, 51.7, 79.3, 125.7, 125.9, 126.1, 126.6, 127.0, 127.6, 128.0, 132.8, 133.0, 138.5, 177.3; HRMS (FAB): *m/z*: calcd for C₁₉H₂₇O₃Si: 331.1729; found: 331.1715 [M + H]⁺.

Methyl (4*E*)-2,2-dimethyl-5-phenyl-3-trimethylsilyloxypent-4enoate (3i).³³ Colorless oil; yield: 285 mg (93%); ¹H NMR (500 MHz, CDCl3): δ 0.00 (s, 9H), 1.02 (s, 3H), 1.11 (s, 3H), 3.59 (s, 3H), 4.40 (d, *J* = 7.0 Hz, 1H), 6.03 (dd, *J* = 7.0, 16.0 Hz, 1H), 6.41 (d, *J* = 16.0 Hz, 1H), 7.16 (d, *J* = 7.4 Hz, 1H), 7.24 (dd, *J* = 7.2, 7.5 Hz, 2H), 7.28 (d, *J* = 7.2 Hz, 2H); ¹³C NMR (125 MHz, CDCl3) δ = 0.3, 19.8, 21.2, 48.5, 51.8, 78.5, 126.4, 127.6, 128.6, 128.8, 132.0, 136.8, 177.1; HRMS (FAB): *m/z*: calcd for C₁₇H₂₇O₃Si: 307.1729; found: 307.1740 [M + H]⁺.

Methyl 2,2-dimethyl-5-phenyl-3-trimethylsilyloxypentanoate (3j).³³ Colorless oil; yield: 268 mg (87%); ¹H NMR (500 MHz, CDCl3): δ 0.14 (s, 9H), 1.08 (s, 3H), 1.16 (s, 3H), 1.61–1.66 (m, 2H), 2.40–2.55 (m, 2H), 3.64 (s, 3H), 3.97 (d, J = 12.0 Hz, 1H), 7.16–7.22 (m, 2H), 7.24–7.29 (m, 2H); ¹³C NMR (125 MHz, CDCl3) $\delta = 0.9$, 20.3, 21.6, 33.5, 35.2, 48.4, 51.8, 77.5, 125.8, 128.3, 128.4, 142.2, 177.6; HRMS (FAB): m/z: calcd for $C_{17}H_{29}O_3$ Si: 309.1886; found: 309.1876 [M]⁺.

Methyl 2,2-dimethyl3-trimethylsilyloxy-dodecanoate (3k).^{17c} Colorless oil; yield: 288 mg (91%); ¹H NMR (500 MHz, CDCl3): δ 0.09 (s, 9H), 0.85 (t, J = 8.1 Hz, 3H), 1.04 (s, 3H), 1.11 (s, 3H), 1.11–1.45 (m, 12H), 1.53–1.64 (m, 2H), 3.64 (s, 3H), 3.83 (d, J =

9.1 Hz, 1H); ¹³C NMR (125 MHz, CDCl3) $\delta = 0.7$, 14.1, 20.1, 21.6, 22.6, 27.0, 29.2, 29.5, 29.7, 31.8, 32.8, 48.2, 51.5, 77.7, 177.8; HRMS (FAB): m/z: calcd for $C_{17}H_{37}O_3Si$: 317.2512; found: 317.2523 $[M + H]^+$.

Methyl 3-(2-furyl)-2,2-dimethyl-3-trimethylsilyloxy-propionate (3l).^{17c} Colorless oil; yield: 242 mg (84%); ¹H NMR (500 MHz, CDCl3): δ 0.00 (s, 9H), 1.04 (s, 3H), 1.23 (s, 3H), 3.68 (s, 3H), 5.00 (s, 1H), 6.20 (d, J = 3.2 Hz, 1H), 6.31 (d, J = 3.2 Hz, 1H), 7.32–7.34 (m, 1H); ¹³C NMR (125 MHz, CDCl3) δ = -0.3, 19.6, 21.2, 48.6, 51.8, 73.5, 107.9, 109.9, 141.5, 154.4, 176.8; HRMS (FAB): m/z: calcd for C₁₃H₂₃O₄Si: 271.1366; found: 271.1363 [M + H]⁺.

Ethyl 3-3-hydroxy-3-phenylpropionate (5a).³³ Colorless oil; yield: 165 mg (85%); ¹H NMR (500 MHz, CDCl3): δ 1.26 (t, J = 7.2 Hz, 3H), 2.70 (dd, J = 3.9, 16.4 Hz, 1H), 2.76 (dd, J = 9.0, 16.4 Hz, 1H), 3.30–3.34 (brs, 3H), 4.18 (q, J = 7.2 Hz, 2H), 5.13 (dd, J = 2.8, 8.8 Hz, 1H), 7.28–7.31 (m, 1H), 7.35–7.39 (m, 4H); ¹³C NMR (125 MHz, CDCl3) δ = 14.1, 43.2, 60.9, 70.3, 125.6, 127.8, 128.5, 142.4, 172.4; HRMS (FAB): m/z: calcd for C₁₁H₁₅O₃: 195.1021; found: 195.1010 [M + H]⁺.

Methyl 3-3-hydroxy-2-methyl-3-phenylpropionate (5b).^{17c} syn-anti mixture: colorless oil; yield: 175 mg (90%); HRMS (FAB): *m*/z: calcd for C₁₁H₁₅O₃: 195.1021; found: 195.1033 [M + H]⁺. syn Isomer: ¹H NMR (500 MHz, CDCl3): δ 1.12 (d, J = 7.2 Hz, 3H), 2.78 (dq, J = 4.1, 7.2 Hz, 1H), 2.93–2.98 (brs, 3H), 3.67 (s, 3H), 5.10 (d, J = 4.0 Hz, 1H), 7.23–7.30 (m, 1H), 7.31–7.39 (m, 4H); ¹³C NMR (125 MHz, CDCl3) δ = 10.6, 46.3, 51.8, 73.5, 125.9, 127.5, 128.2, 141.3, 176.2.; anti Isomer: ¹H NMR (500 MHz, CDCl3): δ 1.00 (d, J = 7.2 Hz, 3H), 2.83 (dq, J = 7.2, 8.4 Hz, 1H), 2.90–3.05 (brs, 3H), 3.72 (s, 3H), 4.74 (d, J = 8.4 Hz, 1H), 7.25–7.30 (m, 1H), 7.31–7.42 (m, 4H); ¹³C NMR (125 MHz, CDCl3) δ = 14.4, 47.1, 51.8, 76.4, 126.6, 128.1, 128.5, 141.5, 176.3.

Methyl 4-chlorophenyl-2,2-Dimethyl-3-(tosylamino)-propionate (7a).^{31e} White solid; m.p. 145 °C; yield: 332 mg (84%); ¹H NMR (500 MHz, CDCl3): δ 1.03 (s, 3H), 1.31 (s, 3H), 2.30 (s, 3H), 3.58 (s, 3H), 4.27 (d, J = 9.5 Hz, 1H), 6.13 (d, J = 9.5 Hz, 1H), 6.82 (d, J = 8.5 Hz, 2H), 6.94–7.00 (m, 4H), 7.35 (d, J = 8.5 Hz, 2H); ¹³C NMR (125 MHz, CDCl3) $\delta = 21.3$, 22.5, 24.9, 46.9, 52.2, 64.3, 126.8, 128.0, 129.1, 129.3, 135.7, 137.5, 143.0, 176.4; HRMS (FAB): m/z: calcd for C₁₉H₂₃ClNO₄S: 396.1036; found: 396.1022 [M + H]⁺.

Methyl 2,2-dimethyl-3-phenyl-3-(tosylamino)propionate (7b).^{31e} White solid; m.p. 130 °C; yield: 285 mg (82%); ¹H NMR (500 MHz, CDCl3): δ 1.08 (s, 3H), 1.26 (s, 3H), 2.24 (s, 3H), 3.60 (s, 3H), 4.38 (d, J = 9.9 Hz, 1H), 6.30 (d, J = 9.9 Hz, 1H), 6.88 (d, J = 7.0 Hz, 2H), 6.93 (d, J = 8.0 Hz, 2H), 9.68–7.10 (m, 5H), 7.39 (d, J = 8.0 Hz, 2H); ¹³C NMR (125 MHz, CDCl3) δ = 21.2, 221, 24.2, 41.3, 52.0, 64.5, 126.7, 127.2, 127.7, 127.8, 128.9, 136.9, 137.4, 142.5, 176.4; HRMS (FAB): m/z: calcd for C₁₉H₂₄NO₄S: 362.1426; found: 362.1429 [M + H]⁺.

Methyl 2,2-dimethyl-3-(4-nitrophenyl)-3-(tosylamino)-propionate (7c).^{31e} White solid; m.p. 180 °C; yield: 365 mg (90%); ¹H NMR (500 MHz, CDCl3): δ 1.05 (s, 3H), 1.33 (s, 3H), 2.26 (s, 3H), 3.59 (s, 3H), 4.39 (d, J = 9.3 Hz, 1H), 6.30 (d, J = 9.3 Hz, 1H), 6.98 (d, J = 8.0 Hz, 2H), 7.12 (d, J = 8.5 Hz, 2H), 7.39 (d, J = 8.0 Hz, 2H), 7.88 (d, J = 8.5 Hz, 2H); ¹³C NMR (125 MHz, CDCl3) δ =

21.4, 22.5, 24.9, 46.8, 52.4, 64.2, 123.0, 126.8, 129.0, 129.2, 137.3, 143.4, 144.7, 147.1, 176.1; HRMS (FAB): m/z: calcd for $C_{19}H_{23}N_2O_6S$: 407.1277; found: 407.1273 $[M + H]^+$.

Methyl 3-(4-methoxyphenyl)-2,2-dimethyl-3-(tosylamino)propionate (7d).^{31e} White solid; m.p. 114 °C; yield: 289 mg (64%); ¹H NMR (500 MHz, CDCl3): δ 1.04 (s, 3H), 1.25 (s, 3H), 2.25 (s, 3H), 3.57 (s, 3H), 3.68 (s, 3H), 4.29 (d, J = 9.2 Hz, 1H), 6.12 (d, J = 9.2 Hz, 1H), 6.53 (d, J = 8.8 Hz, 2H), 6.78 (d, J = 8.5 Hz, 2H), 7.36 (d, J = 8.8 Hz, 2H), 7.81 (dd, J = 8.5, 8.8 Hz, 2H); ¹³C NMR (125 MHz, CDCl3) δ = 21.5, 22.5, 24.6, 47.6, 52.2, 55.4, 64.3, 113.1, 126.9, 129.0, 129.1, 129.4, 137.7, 142.4, 158.8, 176.4; HRMS (FAB): m/z: calcd for C₂₀H₂₆NO₅S: 392.1532; found: 392.1540 [M + H]⁺.

Methyl 3-[4-(dimethylamino)phenyl]-2,2-dimethyl-3-(tosylamino)-propionate (7e).^{31e} White solid; m.p. 155 °C; yield: 226 mg (56%); ¹H NMR (500 MHz, CDCl3): δ 1.04 (s, 3H), 1.25 (s, 3H), 2.23 (s, 3H), 2.82 (s, 6H), 3.57 (s, 3H), 4.23 (d, J = 9.6 Hz, 1H), 5.99 (d, J = 9.6 Hz, 1H), 6.34 (d, J = 7.1 Hz, 2H), 6.68 (d, J = 8.6 Hz, 2H), 6.91 (d, J = 8.0 Hz, 2H), 7.34 (d, J = 8.6 Hz, 2H); ¹³C NMR (125 MHz, CDCl3) δ = 21.3, 22.2, 24.5, 40.4, 47.3, 51.9, 64.4, 111.9, 126.9, 128.6, 128.8, 142.0, 149.7, 162.5, 176.8; HRMS (FAB): m/z: calcd for C₂₀H₂₆NO₅S: 392.1532; found: 392.1540 [M + H]⁺.

Methyl 2,2-dimethyl-3-phenyl-3-(phenylamino)propionate (7f).^{31e} Colorless oil; yield: 234 mg (82%); ¹H NMR (500 MHz, CDCl3): δ 1.13 (s, 3H), 1.24 (s, 3H), 2.23 (s, 3H), 3.62 (s, 3H), 4.46 (s, 1H), 4.70 (br, 1H), 6.46 (d, *J* = 8.5 Hz, 2H), 6.56 (dd, *J* = 8.5, 8.8 Hz, 1H), 7.00 (d, *J* = 8.5, 8.8 Hz, 2H), 7.20–7.28 (m, 5H). ¹³C NMR (125 MHz, CDCl3) δ = 20.7, 24.5, 47.0, 52.0, 64.3, 113.3, 117.2, 127.4, 128.0, 128.2, 129.0, 139.2, 146.9, 177.0; HRMS (FAB): *m/z*: calcd for C₁₈H₂₂NO₂: 284.1651; found: 286.1648 [M + H]⁺.

Conclusions

In conclusion, we demonstrated the easily accessible and userfriendly polymer-supported phosphine PS-PPh₃ catalyzed Mukaiyama aldol and Mannich reactions. A broad range of aldehydes and imines could be applied under mild conditions using 5–10 mol% PS-PPh₃. Furthermore, PS-PPh₃ was easily recovered and reused with minimal loss of activity after three runs. These reactions provide a simple and environmentally friendly method for the synthesis of β -hydroxy esters and β -amino esters.

Notes and references

 Recent reviews, see: (a) C. A. McNamara, M. J. Dixon and M. Bradley, Chem. Rev., 2002, 102, 3275-3300; (b) M. Benaglia, A. Puglisi and F. Cozzi, Chem. Rev., 2003, 103, 3401-3429; (c) F. Cozzi, Adv. Synth. Catal., 2006, 348, 1367-1390; (d) W. J. Sommer and M. Weck, Cood. Chem. Rev., 2007, 251, 860-873; (e) C. K. Kwong, R. Huang, M. Zhang and P. H. Toy, Chem.-Eur. J., 2007, 13, 2369-2376; (f) M. R. Buchmeiser, Chem. Rev., 2009, 109, 303-321; (g) Z. Wang, G. Chen and K. Ding, Chem. Rev., 2009, 109, 322-359; (h) S. Ikegami and H. Hamamoto, Chem. Rev., 2009, 109, 583-593; (i) J. Lu and P. H. Toy, Chem. Rev., 2009, 109, 815–838; (j) T. E. Kristensen and T. Hansen, *Eur. J. Org. Chem.*, 2010, **17**, 3179–3204; (k) Y. Zhang and S. N. Riduan, *Chem. Soc. Rev.*, 2012, **41**, 2093–2094.

- 2 D. Simoni, R. Rondanin, M. Morini, R. Baruchello and F. P. Invidiata, *Tetrahedron Lett.*, 2000, **41**, 1607–1610.
- 3 F. Fringuelli, F. Pizzo, C. Vittoriani and L. Vaccaro, *Chem. Commun.*, 2004, 2756–2757.
- 4 (a) F. Fringuelli, F. Pizzo, C. Vittoriani and L. Vaccaro, *Eur. J. Org. Chem.*, 2006, 1231–1236; (b) D. Lanari, R. Balini, A. Palmieri, F. Pizzo and L. Vaccaro, *Eur. J. Org. Chem.*, 2011, 2874–2884.
- 5 (a) D. Bensa, T. Constantieux and J. Rodriguez, Synthesis, 2004, 923–927; (b) A. W. Pilling, J. Boehmer and D. J. Dixon, Angew. Chem., Int. Ed, 2007, 46, 5428–5430; (c) A. W. Pilling, J. Boehmer and D. J. Dixon, Chem. Commun., 2008, 832–834; (d) R. Ballini, L. Barboni, L. Castrica, F. Fringuelli, D. Lanari, F. Pizzo and L. Vaccaro, Adv. Synth. Catal., 2008, 350, 1218–1224; (e) S. Bonollo, D. Lanari, J. M. Longo and L. Vaccaro, Green Chem., 2012, 14, 164–169.
- 6 T. Angelini, F. Fringuelli, D. Lanari, F. Pizzo and L. Vaccaro, *Tetrehedron Lett.*, 2010, **51**, 1566–1569.
- 7 G. Strappaveccia, D. Lanari, D. Gelman, F. Pizzo, O. Rosati, M. Curini and L. Vaccaro, *Green Chem.*, 2013, 15, 199–204.
- 8 C. Ogawa, M. Sugiura and S. Kobayashi, *Chem. Commun.*, 2003, 191–193.
- 9 R. A. Flowers II, X. Liu, C. Timmons and G. Li, *Eur. J. Org. Chem.*, 2004, 2988–2990.
- 10 (a) S. Matsukawa and S. Fujikawa, *Tetrahedron Lett.*, 2012, 53, 1075–1077; (b) S. Matsukawa, T. Harada and S. Yasuda, *Org. Biomol. Chem.*, 2012, 10, 4886–4890; (c) S. Matsukawa, K. Tsukamoto, S. Yasuda and T. Harada, *Synthesis*, 2013, 45, 2959–2965.
- (*a*) T. Mukaiyama, K. Narasaka and K. Banno, *Chem. Lett.*, 1973, 1011–1114; (*b*) T. Mukaiyama, K. Banno and K. Narasaka, *J. Am. Chem. Soc.*, 1974, **96**, 7503–7509; (*c*) T. Mukaiyama, T. Izawa and K. Saigo, *Chem. Lett.*, 1974, 323–326.
- 12 For selected reviews, see: (a) T. Mukaiyama, Org. React., 1982, 28, 203–331; (b) T. Mukaiyama and S. Kobayashi, Org. React., 1994, 46, 1–103; (c) S. G. Nelson, Tetrahedron: Asymmetry, 1998, 9, 357–389; (d) H. Groger, E. M. Vogl and M. Shibasaki, Chem.-Eur. J., 1998, 4, 1137–1141; (e) R. Mahrwald, Chem. Rev., 1999, 99, 1095–1120; (f) T. Mukaiyama, Tetrahedron, 1999, 55, 8609–8670; (g) T. Mukaiyama, Angew. Chem., Int. Ed., 2004, 43, 5590–5614; (h) Modern Aldol Reactions, ed. R. Mahrwald, Wiley-VCH, Weinheim, 2004, Vol. 1, p. 2; (i) E. M. Carreira, A. Fettes and C. Marti, Org. React., 2006, 67, 1–216; (j) M. Shibasaki and S. Matsunaga, Chem. Soc. Rev., 2006, 35, 269–279; (k) J. Gawronski, N. Wascinska and J. Gajewy, Chem. Rev., 2008, 108, 5227–5252; (l) T. Kitanosono and S. Kobayashi, Adv. Synth. Catal., 2013, 355, 3095–3118.
- 13 (a) G. L. Beutner and S. E. Denmark, Angew. Chem., Int. Ed., 2013, 52, 9086–9096; (b) S. B. J. Kan, K. K. H. Ng and I. Paterson, Angew. Chem., Int. Ed., 2013, 52, 9097–9108; (c) J. Matsuo and M. Murakami, Angew. Chem., Int. Ed., 2013, 52, 9109–9118.

- 14 (a) R. Noyori, I. Nishida and J. Sakata, J. Am. Chem. Soc., 1983, 105, 1598–1608; (b) E. Nakamura, M. Shimizu, I. Kuwajima, J. Sakata, K. Yokoyama and R. Noyori, J. Org. Chem., 1983, 48, 932–945.
- 15 S. Matsukawa, N. Okano and T. Imamoto, *Tetrahedron Lett.*, 2000, **41**, 103–107.
- 16 T. Mukaiyama, H. Fujisawa and T. Nakagawa, *Helv. Chim.* Acta, 2002, **85**, 4518.
- 17 (a) T. Nakagawa, H. Fujisawa and T. Mukaiyama, *Chem. Lett.*, 2003, 32, 696–697; (b) H. Fujisawa, E. Takahashi and T. Mukaiyama, *Eur. J. Chem.*, 2006, 12, 5082–5093; (c) T. Nakagawa, H. Fujisawa, Y. Nagata and T. Mukaiyama, *Bull. Chem. Soc. Jpn.*, 2004, 77, 1555–1567.
- 18 H. Fujisawa, T. Nakagawa and T. Mukaiyama, *Adv. Synth. Catal.*, 2004, **346**, 1241–1246.
- 19 T. Mizugaki, C. E. Hetrick, M. Murata, K. Ebitani, M. D. Amiridis and K. Kaneta, *Chem. Lett.*, 2005, 420–421.
- 20 H. Fujisawa, Y. Nagata, Y. Sato and T. Mukaiyama, *Chem. Lett.*, 2005, **34**, 842–843.
- 21 H. Hagiwara, H. Inoguchi, M. Fukushima, T. Hoshi and T. Suzuki, *Synlett*, 2005, 2388.
- 22 H. Hagiwara, H. Inoguchi, M. Fukushima, T. Hoshi and T. Suzuki, *Tetrahedron Lett.*, 2006, **47**, 5371.
- 23 M. Hatano, E. Takagi and K. Ishihara, *Org. Lett.*, 2007, 9, 4527–4530.
- 24 J. J. Song, Z. Tan, J. T. Reeves, N. K. Yee and C. H. Senanayake, *Org. Lett.*, 2007, 9, 1013–1016.
- 25 V. R. Chintareddy, K. Wadhwa and J. G. Verkade, J. Org. Chem., 2009, 74, 8118–8132.
- 26 F. Mummy and R. Haag, Synlett, 2012, 2672-2676.
- 27 For some recent PPh₃ catalyzed reactions, see: (a)
 D. B. Ramachary, C. Venkaiah and P. M. Krishna, Org. Lett., 2013, 15, 4714–4717; (b) L. Nicolas, P. Angibaud,
 I. Stansfield, L. Meerpoel, S. Reymond and J. Cossy, Tetrahedron Lett., 2014, 55, 849–852; (c) W. Li and J. Zhang, Org. Lett., 2014, 16, 162–165; (d) E. Li, P. Jia, L. Liang and
 Y. Huang, ACS Catal., 2014, 4, 600–603; (e) W. Dong, P. Hu,
 J. Hu and X. Tong, Tetrahedron Lett., 2014, 55, 1682–1685.

- 28 Silica-supported TBD catalyzed Mukaiyama aldol reactions have also been reported. However, a high temperature is required and chemical yield is not high. See: R. Srivastava, *J. Mol. Catal. A: Chem.*, 2007, **264**, 146–152.
- 29 (a) I. Ojima, S. Inaba and K. Yoshida, *Tetrahedron Lett.*, 1977, 3643–3646; (b) I. Ojima, S. Inaba and M. Nagai, *Synthesis*, 1991, 545–547.
- 30 Reviews, see: (a) M. Arend, B. Westermann and N. Risch, Angew. Chem., Int. Ed, 1998, 37, 1044–1070; (b)
 S. Kobayashi and H. Ishitani, Chem. Rev., 1999, 99, 1069– 1094; (c) A. Cordova, Acc. Chem. Res., 2004, 37, 102–112; (d)
 G. K. Friestad and A. K. Mathies, Tetrahedron, 2007, 63, 2541–2569; (e) D. Ferraris, Tetrahedron, 2007, 63, 9581– 9597; (f) B. Yin, Y. Zhang and L. W. Xu, Synthesis, 2010, 3583–3595; (g) M. Hatano and K. Ishihara, Synthesis, 2010, 3785–3801; (h) S. Kobayashi, Y. Mori, Y. J. S. Fossey and M. M. Salter, Chem. Rev., 2011, 111, 2626–2704; (i)
 P. Merino and T. Tejero, Synlett, 2011, 1965; (j)
 P. S. Bhadury and H. Li, Synlett, 2012, 1108–1131.
- 31 Lewis base catalyzed reactions, see: (a) H. Fujisawa, E. Takahashi, T. Nakagawa and T. Mukaiyama, *Chem. Lett.*, 2003, 32, 1036–1037; (b) E. Takahashi, H. Fujisawa and T. Mukaiyama, *Chem. Lett.*, 2004, 33, 936–937; (c) S. Matsukawa and K. Obu, *Chem. Lett.*, 2004, 33, 1626–1627; (d) E. Takahashi, H. Fujisawa and T. Mukaiyama, *Chem. Lett.*, 2005, 34, 84–85; (e) H. Fujisawa, E. Takahashi and T. Mukaiyama, *Chem.-Eur. J.*, 2006, 12, 5082–5093.
- 32 N. Giuseppone, P. Van de Weghe and M. Mellah, J. Collin Tetrahedron, 1998, 54, 13129–13148.
- 33 G. Onodera, T. Toeda, N. Toda, D. Shibagishi and R. Takeuchi, *Tetrahedron*, 2010, 66, 9021–9031.
- 34 K. Gedrich, M. Heitbaum, A. Notzon, I. Senkovska, R. Fröhlich, J. Getzschmann, U. Mueller, F. Glorius and S. Kaskel, *Chem.-Eur. J.*, 2011, 17, 2099–2106.
- 35 M. Abe, M. Ikeda, Y. Shirodai and M. Nojima, *Tetrahedron Lett.*, 1996, **37**, 5901–5904.