


RSC Advances



This article can be cited before page numbers have been issued, to do this please use: J. Yang, H. Wang, J. Wang, X. Guo and Y. Zhang, *RSC Adv.*, 2015, DOI: 10.1039/C5RA20772J.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Light-Modulated Aggregation Behavior of Some Unsubstituted Cinnamate-Based Ionic Liquids in Aqueous Solutions

View Article Online
DOI: 10.1039/C5RA20772J

Jie Yang,^{ac} Huiyong Wang,^{ab} Jianji Wang,^{*ab} Xiaojia Guo^{ab} and Yue Zhang^{ab}

^a Henan Key Laboratory of Green Chemical Media and Reactions, School of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang, Henan 453007, P. R. China

^b Collaborative Innovation Center of Henan Province for Green Manufacturing of Fine Chemicals, Key Laboratory of Green Chemical Media and Reactions, Ministry of Education, Henan Normal University, Xinxiang, Henan 453007, P. R. China

^c School of Basic Medicine, Xinxiang Medical University, Xinxiang, Henan 453007, P. R. China

ABSTRACT: Light modulation of the self-assembly behavior of ionic liquids in aqueous solution is of great importance from viewpoint of fundamental and technology. In this work, a new class of light-responsive ionic liquids composed of 1-alkyl-3-methylimidazolium cations, $[C_n\text{mim}]^+$ ($n = 4, 6, 8, 10, 12$), and trans-cinnamic acid ($[CA]$) anion was synthesized and characterized. These compounds were used to study the modulation action of UV light on the aggregation behavior of ionic liquids in aqueous solution. For this purpose, critical aggregation concentration (CAC), ionization degree of the aggregates (α), standard Gibbs energy of aggregation (ΔG_m°) and aggregate size of the ionic liquids were determined before and after UV light irradiation by conductivity and dynamic light scattering techniques. The percentage of cis-isomer of these ionic liquids was also reported at photoisomerism equilibrium from HPLC measurements. It was found that the aggregation behavior of $[C_n\text{mim}][CA]$ ($n = 8, 10, 12$) could be efficiently modulated by UV light in aqueous solution. After UV light irradiation, the values of CAC , α and ΔG_m° increased, while the size of aggregates decreased. These results have been rationalized from photo-isomerization of $[CA]$ anion, hydrophilicity and structural feature of the cis- isomer of $[CA]$ anion.

Introduction

In the past two decades, ionic liquids have been developed rapidly due to their

attractive properties such as negligible volatility, low melting point, high thermal stability, and strong solubility capacity for many materials.^[1-4] These liquid materials have drawn increasing interest as alternative media in a variety of chemical synthesis, separation, electrochemical and catalytic processes.^[5-14] Because of the amphiphilic property, ionic liquids are able to form aggregates in solutions.^[15-19] In fact, many applications of ionic liquids, such as environmental pollution control, chromatographic application, material preparation, and separation, are often closely dependent on their aggregation behavior. Therefore, the study of aggregation behavior in solutions is highly necessary for understanding the role of ionic liquids playing in practical applications.

At present, many investigations have been focused on the modulation of aggregation behavior of ionic liquids in aqueous solutions by altering the length of alkyl chain in cation^[20,21], the structure of cation, the type of anions^[21,22], pH value^[23,24] and temperature^[25] of the system, and by adding inorganic or organic additive in solutions as well.^[26-28] As an external stimuli, light trigger is of great importance and superiority, since no “invasion” can be introduced into the system during the stimuli-responsive process, and the optical signal is easy to obtain with stability and reliability. Furthermore, light can be directed at a precise spot with a resolution of a few micrometers, which is of particularly value in nanoscience and nanotechnology applications. Thus photo-responsive ionic liquids have been created in recent years. To this end, Asaka and co-workers^[29] synthesized a photo-responsive ionic liquid, 3-butyl-1-methyl-2-phenylazoimidazolium bis(trifluoro-methanesulfonyl) amide, and investigated its isomerization mechanism and solvent viscosity effect in E-Z photoisomerization. Branco *et al*^[30] prepared some methyl orange based photochromic ionic liquids and determined rate constants of their thermal recovery. Tamura *et al*^[31] reported some photo-responsive stilbene ionic liquids and their emissive properties and melting points. Some azobenzene-based photo-responsive ionic liquids were prepared by Zhang *et al*,^[32] and reversible modulation of ionic conductivity in some organic solvents was examined under UV/visible light irradiation. In a previous work,^[33] we reported another series of cinnamate-based ionic

liquids composed of 1-alkyl-3-methylimidazolium cations $[C_n\text{mim}]^+$ and trans-ortho-methoxycinnamic acid anion ([OMCA]). Highly efficient conductivity modulation of these light-responsive ionic liquids was studied in aqueous solutions. However, to the best of our knowledge, there was only one report^[34] concerning with light modulation of the self-assembly behavior of ionic liquids in aqueous solution up to now. In that work, Shen *et al.*^[34] synthesized a new surface active ionic liquid 4-butylazobenzene-4'-hexyloxy-trimethyl-ammoniumtrifluoroacetate. It was shown that reversible micelle-vesicle transformation was achieved in aqueous solution by alternative UV/vis light irradiation.

Considering the fact that physical and chemical properties of ionic liquids can be finely tuned by a judicious design of the cations and anions structures, we are interested in the creation of novel ionic liquids whose aggregation behavior in solution can be modulated by light irradiation. In this report, we designed and synthesized five kinds of light responsive ionic liquids composed of 1-alkyl-3-methylimidazolium cations, $[C_n\text{mim}]^+$ ($n = 4, 6, 8, 10, 12$), and trans-cinnamic acid ([CA]) anion which can be photoisomerized from trans- to cis-isomer upon exposure to UV light. Because its photoisomerization is not reversible,^[35] the irradiated samples will not undergo any changes when stored under ambient conditions for a long time, and it is easy to conduct subsequent tests using appropriate methods. The modulation of aggregation behavior of these ionic liquids in water by UV light irradiation has been investigated by conductivity and dynamic light scattering techniques. The results have been used to understand the effect of light-induced transformation of [CA] anion from trans- to cis-isomers on the self-assembly of $[C_n\text{mim}][\text{CA}]$ ionic liquids in aqueous solutions.

Materials and methods

Chemicals

1-Methylimidazole (99%), 1,1,1-trichloroethane (AR), ethyl acetate (AR) and petroleum ether (bp 60-90 °C) was acquired from Shanghai Chem. Co.; 1-bromobutane (99%), 1-bromohexane (99%), 1-bromooctane (99%), 1-bromodecane

(99%), 1-bromododecane (99%), trans-cinnamic acid (98%) and Ambersep 900(OH) anion exchange resin were purchased from Alfa Aesar. These chemicals were used without further purification.

Synthesis of the ionic liquids

[C_nmim]Br was prepared and purified by using the procedures described in literature.^[36] Briefly, the reaction of 1-methylimidazole (0.10 mol) with an excess of 1-brominated alkanes (0.13 mol) was performed in 1,1,1-trichloroethane (100 mL) at 70 °C for 48 h to obtain [C_nmim]Br. Among the products prepared, [C₄mim]Br, [C₆mim]Br, [C₈mim]Br and [C₁₀mim]Br were washed with 1,1,1-trichloroethane, and the residual solvents were removed by heating at 70 °C under vacuum, while [C₁₂mim]Br was recrystallized three times from ethyl acetate and ethyl acetate/acetonitrile (3:2 by volume), respectively.

An aqueous solution of [C_nmim]Br was allowed to pass through a column filled with Ambersep 900(OH) anion exchange resin to obtain [C_nmim][OH]. The aqueous [C_nmim][OH] solution was then neutralized with equal molar trans-cinnamic acid (H[CA]). After removing water by evaporation under reduced pressure, [C₄mim][CA] was thoroughly washed with diethyl ether or petroleum ether, and the product was obtained as slightly brown viscous liquid. The other ionic liquids were recrystallized three times from ethyl acetate, and finally dried under vacuum for 72 h at 60 °C to obtain products as white solids (faint yellow solid for [C₆mim][CA]).

The chemical structures of these ionic liquids were confirmed by ¹H NMR, ¹³C NMR (Bruker Avance-400 spectroscopy) and HR-mass spectroscopy (Bruker MicroTOF II spectrometer). The detailed data were given in ESI. Their glass transition temperature (*T_g*), melting temperature (*T_m*) and thermal decomposition temperature (*T_d*) were determined by differential scanning calorimetry (Mettler Toledo 822e) and thermogravimetric analysis (NETZSCH STA 449 C), respectively. Bromide content in these ionic liquids was determined by means of a Br⁻ selective electrode (Shanghai Precision & Scientific Instrument Co. Ltd) coupled with a saturated calomel electrode (Shanghai Precision & Scientific Instrument Co. Ltd).

UV-light irradiation and UV-vis spectroscopy measurements

Aqueous [C_nmim][CA] solutions were irradiated with UV light from a 300W mercury pressure short arc lamp with a filter (<400 nm). Each sample (30 mL) was placed in a quartz tube with a cover and then irradiated for specific duration under stirring. In order to avoid overheating from the UV irradiation, the lamp was put in a cell with a water circulating jacket, and the temperature was kept to be 298.2 ± 0.5 K.

UV-vis spectra for the solutions of ionic liquids were determined before and after UV-irradiation by a UV-4100 spectrophotometer at room temperature. Deionized water was utilized as blank in the experiments.

Conductivity measurements

Conductivity measurements were performed with a Shanghai DJS-1 electrode at 298.15 K by using a Wayne-Kerr 6430B Auto Balance Bridge with a resolution of 1×10⁻⁵ μS·cm⁻¹. The conductance cell was equipped with a water circulating jacket, and the temperature was controlled within ± 0.01°C with a HAAKE V26 thermostat (Thermo Electron, Germany). The cell was calibrated with aqueous KCl solutions at different concentrations, and a cell constant of 1.0127 cm⁻¹ was determined.

Dynamic Light Scattering (DLS) measurements

Dynamic light scattering (DLS) measurements were carried out at 298.2 K by using a laser light scattering photometer (Nano-ZS90, Malvern, U. K). Light of λ= 633 nm from a solid-state He-Ne laser (4.0 mW) was used as the incident beam. All sample solutions were filtered through a 0.22 μm hydrophilic PVDF membrane filter. All measurements were made at 90° scattering angle. At least three measurements were taken for each solution, and the reproducibility of aggregate sizes from DLS data was found to be within ±3%.

High Performance Liquid Chromatography (HPLC) measurements

To determine the percentage of the cis-isomer of [C_nmim][CA], the irradiated solutions were analyzed by using HPLC (Agilent Eclipse XDB-C18 column, 5μm, 4.6 mm×150 mm) at a flow rate of 0.8 mL/min. The eluting solvent was an aqueous methanol solution (45:55 by volume) containing 1.0% acetic acid, and the column

temperature was at 30 °C. The peaks were detected using UV absorption at 254 nm, and the cis-trans ratio was estimated by their integrated peak area.

Results and discussion

Table 1 shows the melting temperature (T_m), glass transition temperature (T_g), and thermal decomposition temperature (T_d) for the investigated ionic liquids. It can be seen that [C₄mim][CA] showed only glass transition temperature at about -30 °C, and the other compounds possess a melting point below 51 °C. This suggests that all these compounds are ionic liquids. In addition, all these ionic liquids have good thermal stability (234–248 °C) under an inert atmosphere as estimated from their T_d values.

Table 1. Thermal properties of [C_nmim][CA] (n = 4, 6, 8, 10, 12)

IL	T_m / °C	T_g / °C	T_d / °C
[C ₄ mim][CA]	-	-30	248
[C ₆ mim][CA]	50	-	242
[C ₈ mim][CA]	46	-	246
[C ₁₀ mim][CA]	37	-	243
[C ₁₂ mim][CA]	39	-	234

It has been reported that impurity in ionic liquids has an important effect on their physico-chemical properties.^[37] Considering the fact that in the synthesis process of our ionic liquids, the main impurity was bromide, therefore, the bromide content in these ionic liquids was determined by a Br⁻ selective electrode, and the data were given in Table S1 (ESI[†]). It can be seen that the bromide content was less than 0.0079 mol kg⁻¹. These results suggest that the quality of the ionic liquids prepared in this work is reliable to investigate the aggregation behavior of ionic liquids in aqueous solutions.

Photo-isomerization of [C_nmim][CA] ionic liquids in aqueous solution

As a typical example, UV-vis spectra for [C₁₂mim][CA] and trans- cinnamic acid (H[CA]) at the concentration of 1 mM were shown in Fig. 1 and S1, respectively, where a slightly excess of base was added to aqueous solutions to increase solubility

of trans- cinnamic acid in water. It can be seen that the maximal absorption wavelength of $[C_n\text{mim}][\text{CA}]$ was at around 267 nm before irradiation and at 254 nm after irradiation (at photostationary state) in aqueous solutions, and the absorbance peak heights were also significantly reduced after irradiation. This blue shift of the maximal absorption wavelength is an indicative of trans to cis photo-isomerization,^[35] which is exactly similar to what has been observed for trans- cinnamic acid (see Fig.S1). This indicates that the spectral features of these ionic liquids were surely originated from their $[\text{CA}]$ anion, and almost have nothing to do with their cations and alkyl chain length. Thus the photoisomerization of $[C_n\text{mim}][\text{CA}]$ ionic liquids can be represented by the reaction shown in Fig.2. By HPLC measurements, it was confirmed that percentage of cis-isomer of $[C_n\text{mim}][\text{CA}]$ ($n = 8, 10, 12$) corresponding to photoisomerism equilibrium was ca. 83% at 298.2 K.




Fig. 1. UV-vis spectra of aqueous $[C_{12}\text{mim}][\text{CA}]$ solution at the concentration of 1 mM before and after irradiation.




Fig. 2. Photoisomerism of $[C_n\text{mim}][\text{CA}]$ ionic liquids.

Aggregation parameters for the ionic liquids in water before irradiation

The experimental conductivities for aqueous solutions of $[C_n\text{mim}][\text{CA}]$ ($n = 4, 6, 8, 10, 12$) before irradiation were shown in Fig. 3 as a function of ionic liquid

concentration at 298.15 K. It can be seen that a characteristic shape of the curves was observed in these $[C_n\text{mim}][\text{CA}]$ ($n = 8, 10, 12$) solutions, except for $[C_4\text{mim}][\text{CA}]$ and $[C_6\text{mim}][\text{CA}]$. These curves exhibited typical behavior with two linear fragments, and the concentration at which the two linear fragments intersect was assigned to the critical aggregate concentration (CAC). The degree of ionization (α) of the aggregates was taken to be the ratio of the values of dk/dC above and below the CAC , and can be calculated from the slope of the two straight lines in these regions. Actually, α value indicates the ability of anion bound on the aggregates. An increase of α value suggests a decrease in the ability of the anions bound to the aggregates.^[38] The CAC and α values of $[C_n\text{mim}][\text{CA}]$ ionic liquids calculated from a linear least-squares analysis were presented in Table 2.




Fig. 3. Plots of the conductivity of aqueous $[C_n\text{mim}][\text{CA}]$ solutions before irradiation as a function of the ILs concentration at 298.15 K.

Table 2. CAC , α and ΔG_m° values of aqueous $[C_n\text{mim}][\text{CA}]$ ($n = 8, 10, 12$) before and after irradiation (at photostationary equilibrium) at 298.15 K

IL	Before irradiation			Photostationary equilibrium		
	$CAC / \text{mol kg}^{-1}$	α	$\Delta G_m^\circ / \text{kJ mol}^{-1}$	$CAC / \text{mol kg}^{-1}$	α	$\Delta G_m^\circ / \text{kJ mol}^{-1}$
$[C_{12}\text{mim}][\text{CA}]$	0.00320	0.0462	-47.3	0.00545	0.154	-42.2
$[C_{10}\text{mim}][\text{CA}]$	0.0118	0.0570	-40.7	0.0194	0.175	-36.0
$[C_8\text{mim}][\text{CA}]$	0.0454	0.0625	-34.2	0.0767	0.180	-29.7

Obviously, the CAC values decreased with the increase of carbon atom number in alkyl chain (N_c) of cations of the ionic liquids, just similar to what have been

observed for conventional ionic surfactants. Interestingly, a linear correlation exists between $\log(CAC)$ and N_c as shown in Fig. 4. This linear correlation can be described by equation (1) with a correlation coefficient of 0.9998.

$$\log(CAC) = 0.957 - 0.288N_c \quad (1)$$

The slope of the straight line was -0.29, which is close to -0.28 obtained from $[C_n\text{mim}]\text{Br}$ ($n = 10, 12, 14, 16$) ionic liquids,^[39] and is in the range reported for conventional ionic surfactants (-0.27~ -0.33).^[40] This result is consistent with the findings that the effect of alkyl chain length of the cation on the CAC does not depend markedly on the structure of the cations and the type of counterions.^[22]




Fig. 4. Logarithmic plot of CAC at 298.15K against carbon atom number in alkyl chain, N_c , of $[C_n\text{mim}][CA]$ ($n = 8, 10, 12$).

It is worth to note that the CAC values of $[C_n\text{mim}][CA]$ are far lower than that of other imidazolium-based ionic liquids with the same carbon atom number (see Table 3). This means that the surface activity of $[C_n\text{mim}][CA]$ is advantageous to other imidazolium-based ionic liquids with the same carbon atom number, and these ionic liquids are more easier to form aggregates. Furthermore, it can be seen that the α values of $[C_n\text{mim}][CA]$ was decreased with the increase of alkyl chain length of the cations. Similar phenomenon was reported for some surfactants in aqueous solutions.^[41] On the other hand, it was found that the α values of $[C_n\text{mim}][CA]$ ($n=8, 10, 12$) were much less than those of other $[C_n\text{mim}]X$ ($n=8, 10, 12$) ionic liquids with different anions such as $[\text{CH}_3\text{COO}]$, Cl , Br , $[\text{NO}_3]$ and $[\text{CF}_3\text{COO}]$ (see Table 3 for

example). This indicates that the interactions between imidazolium head group and [CA] anion is much stronger than those between imidazolium head group and [CH₃COO]⁻, Cl⁻, Br⁻, [NO₃]⁻ or [CF₃COO]⁻. This stronger association was resulted from the stronger hydrophobicity of [CA] anion.

Table 3. Comparison of *CAC*, α and ΔG_m° values of [C₈mim][X] ionic liquids at 298.15 K

IL	<i>CAC</i> / (mol kg ⁻¹)	α	ΔG_m° / kJ mol ⁻¹
[C ₈ mim][CH ₃ COO] ^a	0.22	0.42	-21.5
[C ₈ mim]Cl ^a	0.21	0.37	-22.4
[C ₈ mim]Br ^a	0.16	0.34	-23.6
[C ₈ mim][NO ₃] ^a	0.15	0.29	-25.1
[C ₈ mim][CF ₃ COO] ^a	0.12	0.19	-27.2
[C ₈ mim][CA]	0.045	0.063	-34.2

^a reported in ref.[22]

On the basis of the pseudophase model of micellization, the standard Gibbs energy of aggregation can be calculated from the following equation:^[42]

$$\Delta G_m^\circ = (2 - \alpha)RT \ln x_{CAC} \quad (2)$$

where x_{CAC} is the critical aggregation concentration expressed in mole fraction scale. The values of ΔG_m° calculated for [C_nmim][CA] before irradiation were also included in Table 2. It was found from Table 2 that the standard Gibbs energy values for [C_nmim][CA] aggregation became more negative with increasing alkyl chain length of the ionic liquids. This is similar to the case of other types of ionic liquids, indicating that the aggregation was driven mainly by hydrocarbon-hydrocarbon interactions.^[22] In addition, the values of standard Gibbs energy for the aggregation of [C₈mim][CA] was unusually more negative than those of the other [C₈mim]X (X = [CH₃COO], Cl, Br, [NO₃], [CF₃COO]) ionic liquids, which suggests that the aggregation of [C₈mim][CA] was more favorable than the ionic liquids mentioned. In general, aggregate formation was determined by a balance between the repulsive headgroup interactions and the attractive forces arising from a need to minimize the exposure of the hydrophobic core to water. The anion acts by altering such a balance.

As shown in Table 3, the more negative values of ΔG_m° for $[C_8mim][CA]$ was resulted from the stronger absorption of $[CA]$ anion on the aggregates (i.e, lower α), which decreased the electrostatic repulsion between the imidazolium headgroups, and facilitated the formation of aggregates in aqueous solutions.

Photo-induced modulation of aggregation behavior of the ionic liquids

Fig.5. Shows the variation of conductivity for $[C_nmim][CA]$ ($n = 8, 10, 12$) in aqueous solutions as a function of the ionic liquids concentration before and after irradiation. By using the same procedure described before, we calculated the values of CAC , α and ΔG_m° at 298.15 K for $[C_nmim][CA]$ ($n = 8, 10, 12$) ionic liquids after irradiation (at photostationary equilibrium). For the sake of comparison, these values were also included in Table 2.




Fig. 5. Plots of the conductivity of $[C_nmim][CA]$ in aqueous solutions as a function of the ionic liquids concentration before irradiation and after irradiation. (a), $[C_8mim][CA]$; (b), $[C_{10}mim][CA]$; (c), $[C_{12}mim][CA]$; and (d), $[C_{12}mim][CA]$ at different photoisomerism percentage.

It is clear from Fig.5 and Table 2 that UV irradiation significantly increased the conductivity, CAC , α and ΔG_m° values of these aqueous ionic liquids solutions. The increase in α value means that the association of cis-CA anion with imidazolium head groups became weak after UV irradiation, which was not beneficial to the formation of cationic aggregates and resulted in the increase of CAC and ΔG_m° values and the reduction of aggregation ability of the ionic liquids. Actually, the CAC values of aqueous $[C_n\text{mim}][\text{CA}]$ solution at photostationary equilibrium were roughly doubled. In addition, it is evident from Fig.5d that the extent of photo-induced isomerization of the anion from trans- to cis-isomers had an important affect on both α and CAC values. The higher the percentage of cis-isomers of the anion is, the higher the α and CAC values are. Therefore, UV irradiation can be conveniently used to modulate aggregation behavior of ionic liquids in aqueous solutions.

From the above discussion, it is clear that the difference in the interaction between counter ion and headgroup before and after light irradiation is responsible for light-induced aggregation behavior modulation of the ionic liquids in water. Before irradiation, trans-CA anions are the major existing form of the counter ion, and their hydrophobic part (aromatic ring) is relatively "distant" from the hydrophilic part (carboxylate group). This structure feature makes it convenient for the counter ions to intercalate its hydrophobic part into the micelle interior and remain its hydrophilic part outside water. That is to say, trans-CA anions can tightly associate with cationic aggregates, thus promoting the growth of the aggregates. Under these circumstance, CAC , α and ΔG_m° values of $[C_n\text{mim}][\text{CA}]$ were particularly small. On the other hand, when trans-CA anions were transformed into cis-CA anions upon UV light irradiation, molecular skeleton of the anion was curved, and the distance (D) between the hydrophobic benzene ring and the hydrophilic carboxylate group was greatly shortened (see Fig.6). Thus, benzene ring was located close to carboxylate group, resulting in the difficulty for the benzene ring to orient inward to the $[C_n\text{mim}]^+$ aggregates interface and the carboxylate group outward. Furthermore, the difference in the relative hydrophobicity of trans- and cis-CA also led to the difference in the interactions of two isomers with $[C_n\text{mim}]^+$ aggregates. It has been noted for a variety

of compounds that the trans isomer is more hydrophobic than its cis isomer, this difference in hydrophobicity has been ascribed to the lower net dipole moment of the trans isomer compared to the cis isomer.^[35] This implies that trans-CA would readily bind to $[C_n\text{mim}]^+$ aggregates, whereas the more hydrophilic cis-CA could be more likely to remain in solution and this is not desirable for the growth of aggregates. As a result, CAC , α and ΔG_m^0 values of $[C_n\text{mim}][CA]$ significantly increased after light irradiation.




Fig.6. Alteration of distance (D) between hydrophobic benzene ring and hydrophilic carboxylate group after irradiation

Photo-induced modulation of the aggregates size of $[C_n\text{mim}][CA]$ ionic liquids

In order to further investigate the influence of UV light irradiation on the aggregates size, DLS measurement was employed to detect the changes of aggregates size of $[C_n\text{mim}][CA]$ ($n = 8, 10, 12$) before and after UV irradiation in aqueous solution. Fig. 7 shows the size distribution for the aggregates from DLS measurements at 0.1 mol/kg of $[C_n\text{mim}][CA]$ before and after light irradiation for 3 h at 298.2 K. It can be seen that at the same concentration, the aggregate size increased with increasing alkyl chain length of the ILs before irradiation. For example, the aggregate size grew from 5.2 to 6.4 nm when the alkyl chain length increased from C8 to C10. In the meantime, size distribution for $[C_{12}\text{mim}][CA]$ aggregate showed obvious polydispersity with two peaks at 3.56 and 77.5 nm, indicating the formation of larger sized aggregates. It can also be seen from Fig. 7 that after irradiation for 3 h, the aggregate size was changed from 5.2, 6.4, and 3.6 nm to 3.5, 3.4 and 2.5 nm for $[C_8\text{mim}][CA]$, $[C_{10}\text{mim}][CA]$ and $[C_{12}\text{mim}][CA]$, respectively, and the aggregate with a diameter of 77.5 nm disappeared for $[C_{12}\text{mim}][CA]$. These results indicate that the aggregate size of the ionic liquids became small upon UV irradiation, proving that UV irradiation is not conducive to the growth of the aggregates. This is highly consistent with the

conclusion deduced from conductivity measurements.

View Article Online
DOI: 10.1039/C5RA20772J




Fig. 7. The size distribution for the aggregates from DLS measurements at 0.1 mol/kg of [C_nmim][CA] (n = 8, 10, 12) before and after UV irradiation for 3 h at 298.2 K.

Conclusions

In summary, a new class of light-responsive ionic liquids [C_nmim][CA] (n = 4, 6, 8, 10, 12) was synthesized and characterized for the first time, and their aggregation behavior was investigated before and after UV light irradiation. It was found that after UV irradiation, the critical aggregation concentration, ionization degree of the aggregates, and standard Gibbs energy of aggregation of the ionic liquids increased, while the size of aggregates decreased. This could be attributed to the photoisomerization of anion of [C_nmim][CA] from trans- to cis- isomers upon exposure to UV light. The cis-[CA] has unfavorable geometry and weaker hydrophobicity, which makes the association interactions of the counter ions with [C_nmim]⁺ aggregates to weaken. These findings are expected to be useful in light controlled drug delivery, microscale photo-control devices and sensors.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 21133009, 21273062 and 21203057).

View Article Online
DOI: 10.1039/C5RA20772J

Electronic Supplementary Information Available: Synthesis and characterization of the ionic liquids, bromide content in the ionic liquids and UV-vis spectra of trans-cinnamic acid before and after UV light irradiation. See DOI: 10.1039/x0xx00000x

References


- 1 A. W. T. King, J. Asikkala, I. Mutikainen, P. Jarvi and I. Kilpelainen, *Angew. Chem., Int. Ed.*, 2011, **50**, 6301.
- 2 X. Han and D. W. Armstrong, *Acc. Chem. Res.*, 2007, **40**, 1079.
- 3 D. Xiong, G. Cui, J. Wang, H. Wang, Z. Li, K. Yao, and S. Zhang, *Angew. Chem.* 2015, **127**, 7373-7377.
- 4 Y. Lu, K. Korf, Yu Kambe, Z. Tu and L. A. Archer, *Angewandte Chemie*, 2014, **126**, 498.
- 5 S. Zhang, J. Sun, X. Zhang, J. Xin, Q. Miao and J. Wang, *Chem. Soc. Rev.*, 2014, **43**, 7838.
- 6 T. Erdmenger, C. Guerrero-Sanchez, J. Vitz, R. Hoogenboom and U. S. Schubert, *Chem. Soc. Rev.*, 2010, **39**, 3317.
- 7 S. N. Riduan, Y. Zhang, *Chem. Soc. Rev.*, 2013, **42**, 9055.
- 8 X. Zhang, X. Zhang, H. Dong, Z. Zhao, S. Zhang and Y. Huang, *Energy Environ. Sci.*, 2012, **5**, 6668.
- 9 J. Chen, X. Chen, M. Su, J. Ye, J. Hong, Z. Yang, *Chem. Eng. J.*, 2015, **279**, 136.
- 10 F. J. Hernández-Fernández, A. P. De los Ríos, F. Mateo-Ramirez, C. Godinez, L. J. Lozano-Blanco, J. I. Moreno, F. Tomas-Alonso, *Chem. Eng. J.*, 2015, **279**, 115.
- 11 N. Liu, Z. Ma, *Biosensors and Bioelectronics*, 2014, **51**, 184.
- 12 A. F. Cláudio, M. Neves, K. Shimizu, J. N. C. Lopes, M. G. Freire and J. A. P. Coutinho, *Green Chem.*, 2015, **17**, 3948.
- 13 J. P. Hallett, T. Welton, *Chem. Rev.*, 2011, **111**, 3508.
- 14 A. W. T. King, J. Asikkala, I. Mutikainen, P. J. Distillable, *Angew. Chem.* 2011, **123**, 6425.

- 15 J. Bowers, C. P. Butts, P. J. Martin, M. C. Vergara-Gutierrez, *Langmuir*, 2004, **20**, 2191. View Article Online
DOI: 10.1039/C5RA20772J
- 16 T. L. Greaves, C. J. Drummond, *Chem. Soc. Rev.*, 2013, **42**, 1096.
- 17 K. Shimizu, C. E. S. Bernardes, *J. Phys. Chem. B*, 2014, **118**, 567.
- 18 H. Wang, L. Zhang, J. Wang, Z. Li and S. Zhang, *Chem. Commun.*, 2013, **49**, 5222.
- 19 W. Xu, T. Wang, N. Cheng, Q. Hu, Y. Bi, Y. Gong and L. Yu, *Langmuir*, 2015, **31**, 1272.
- 20 J. Wang, H. Wang, S. Zhang, H. Zhang and Y. Zhao, *J. Phys. Chem. B*, 2007, **111**, 6181.
- 21 T. Singh and A. Kumar, *J. Phys. Chem. B*, 2007, **111**, 7843.
- 22 H. Wang, J. Wang, S. Zhang and X. Xuan, *J. Phys. Chem. B*, 2008, **112**, 16682.
- 23 H. Wang, B. Tan, H. Zhang and J. Wang, *RSC Adv.*, 2015, **5**, 65583.
- 24 H. Wang, B. Tan, J. Wang, Z. Li and S. Zhang, *Langmuir*, 2014, **30**, 3971.
- 25 B. Dong, X. Xing, R. Wang, B. Wang, X. Zhou, C. Wang, L. Yu, Z. Wu and Y. Gao, *Chem. Commun.*, 2015, **51**, 11119.
- 26 J. Wang, L. Zhang, H. Wang, and C. Wu, *J. Phys. Chem. B*, 2011, **115**, 4955.
- 27 V. Pino, C. Yao and J. L. Anderson, *J. Colloid. Interface Sci.*, 2009, **333**, 548.
- 28 H. Wang, Q. Feng, J. Wang and H. Zhang, *J. Phys. Chem. B*, 2010, **114**, 1380.
- 29 T. Asaka, Akai, A. Kawai, K. Shibuya, *J. Photochem. Photobiology A*, 2010, **209**, 12.
- 30 L. C. Branco, F. Pina, *Chem. Commun.*, 2009, 6204.
- 31 H. Tamura, Y. Shinohara T. Arai, *Chem. Lett.*, 2010, **39**, 240.
- 32 S. G. Zhang, S. M. Liu, Q. H. Zhang, Y. Q. Deng, *Chem. Commun.*, 2011, **47**, 6641.
- 33 J. Yang, H. Wang, J. Wang, Y. Zhang and Z. Guo, *Chem. Commun.*, 2014, **50**, 14979-14982.
- 34 S. Shi, T. Yin, X. Tao, W. Shen, *RSC Adv.*, 2015, **5**, 75806.
- 35 H. Sakai, Y. Orihara, H. Kodashima, A. Matsumura, T. Ohkubo, K. Tsuchiya and M. Abe, *J. Am. Chem. Soc.*, 2005, **127**, 13454.
- 36 P. D. María, *Angew. Chem. Int. Ed.*, 2008, **47**, 6960.
- 37 K. R. Seddon, A. Stark and M. Torres, *Pure Appl. Chem.*, 2000, **72**, 2275.
- 38 A. González-Pérez, J. L. del Castillo, J. Czapkiewicz and J. R. Rodríguez, *J. Phys. Chem. B*, 2001, **105**, 1720.
- 39 T. Inoue, H. Ebina, B. Dong and L. Zheng, *J. Colloid. Interface Sci.*, 2007, **314**, 236.
- 40 M. J. Rosen, *Surfactants and Interfacial Phenomena*, second ed., Wiley, New York, 1989.

- 41 T. Asakawa, H. Kubode, T. Ozawa, A. Ohta, S. Miyagishi, *J. Oleo Sci.* 2005, **54**, 545. [View Article Online](#)
10.1039/C5RA20772J
- 42 S. K. Metha, K. K. Bhasin, A. Kamar and S. Dham, *Colloids Surf. A*, 2006, **278**, 17.

Table of content entry

View Article Online
DOI: 10.1039/C5RA20772J



The aggregation behavior of a new class light-reponseive ionic liquids was modulated efficiently by UV light irradiation in aqueous solutions.