

View Article Online View Journal

# ChemComm

## Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: P. Gilormini, C. Lion, D. Vicogne, C. mariller, S. Potelle, T. Levade, Y. Guerardel, C. Biot and F. Foulquier, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC08838K.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

# YAL SOCIETY CHEMISTRY

### Journal Name

#### COMMUNICATION



#### distinct tools to unravel sialic acids metabolic pathways<sup>++</sup> Received 00th January 20xx, P.A. Gilormini,<sup>a, b</sup> C. Lion,<sup>a, b</sup> D. Vicogne,<sup>a, b</sup> T. Levade,<sup>c</sup> S. Potelle,<sup>a, b</sup> C. Mariller,<sup>a, b</sup> Y. Guérardel,<sup>a, b</sup> C. Biot,<sup>a, b</sup> \*† and F. Foulguier<sup>a, b</sup> \*†

Sequential Bioorthogonal Dual Strategy: ManNAI and SiaNAI as

Accepted 00th January 20xx

DOI: 10 1039/x0xx00000x

www.rsc.org/

Recent methodological developments metabolic in oligosaccharide engineering (MOE) pave the way for tremendous advances in glycobiology. Herein, we propose a Sequential Bioorthogonal Dual Strategy (SBDS) combining the use of two unprotected alkyne-tagged monosaccharide reporters (ManNAI and SiaNAI) with the bioligation of fluorescent probes by coppercatalysed azide-alkyne cycloaddition (CuAAC). With SBDS, we are able to shed light on trafficking and cellular uptake mechanisms of sialic acid. Using their corresponding analogues, we visualized that SiaNAI enters via endocytosis, whereas its biosynthetic intermediate ManNAI uptake is mediated by a yet unknown but specific plasma membrane transporter. Sialin, a lysosomal protein, is shown to be crucial for the export of exogenous sialic acid from lysosomes to the cytosol. Metabolic labeling with alkyne-tagged derivatives of N-acetylneuraminic acid (Neu5Ac) or Nacetylmannosamine (ManNAc) could thus be used to follow endocytosis in physiological vs pathological conditions.

With the advent of bioorthogonal click chemistry, the emergence of metabolic oligosaccharide engineering (MOE) has opened a completely new field of investigation in the past few years.<sup>1</sup> Indeed, a traditional limitation in the field of glycosciences used to be the molecular imaging of mono-, oligo- or polysaccharides in terms of subcellular localization and quantification. Metabolic inclusion of a synthetically modified monosaccharide reporter in living cells or organisms, followed by bioorthogonal ligation of a probe, now provides a

very elegant strategy to visualize and quantify various glycoconjugate families.<sup>2-6</sup> Several methodologies have specifically been developed for the study of sialylated glycoconjugates.7-12 Sialic acids are a diverse family of monosaccharides that consistently occupy the non-reducing termini of glycan chains in glycoproteins and glycolipids.<sup>13-14</sup> The sialic acid metabolism occurs in different compartments of the cell. Its synthesis starts in the cytosol from UDP-GlcNAc, which is epimerized and phosphorylated into ManNAc-6P by UDP-GlcNAc 2-epimerase (GNE/MNK). ManNAc-6P is then condensed with PEP by the Neu5Ac-9-Phosphate synthase (NANS) to give rise to Neu5Ac-9P. After dephosphorylation by N-acetylneuraminic acid phosphatase (NANP), the synthesis then continues in the nucleus where the Neu5Ac is activated into CMP-Neu5Ac by the CMP-Neu5Ac synthase (CSS). Once activated, the CMP-Neu5Ac is exported from the nucleus into the cytosol then to the Golgi apparatus via the SLC35A1 transporter, before its incorporation into glycoconjugates via sialyltransferases.15

However, biological insights into the metabolism of glycoconjugates have yet to emerge from such chemical reporter strategies. To our knowledge, all reported studies but two 16-17 have so far taken an interest in observing glycoconjugates incorporated at the cellular membrane exclusively.

The available chemical toolbox for such investigations mainly consists of peracetylated saccharide derivatives bearing a reporter group, typically an azide or a terminal alkyne. As those unnatural monosaccharides are presumed not to be membrane permeable when unprotected, they have so far been systematically peracetylated in order to increase their hydrophobicity and thus facilitate their cellular entry via passive diffusion.<sup>18</sup> While powerful, these tools do not allow the fine study of cellular uptake and metabolic compartmentation.

In order to highlight such processes, we have successfully developed a Sequential Bioorthogonal Dual Strategy (SBDS) based on the distinct metabolic incorporation of two unprotected monosaccharide reporters, namely N-(4-

<sup>&</sup>lt;sup>a.</sup> Univ. Lille. UMR 8576 – UGSF – Unité de Glycobiologie Structurale et Fonctionnelle, F- 59000 Lille, France

E-mail: christophe.biot@univ-lille1.fr, francois.foulquier@univ-lille1.fr

<sup>&</sup>lt;sup>b.</sup> CNRS. UMR 8576, F-59000 Lille, France.

<sup>&</sup>lt;sup>c.</sup> Laboratoire de Biochimie Métabolique, IFB, CHU Purpan, INSERM UMR 1037, CRCT, Université Paul Sabatier Toulouse III, Toulouse (France).

<sup>++</sup> This study was funded by grants from Université de Lille1 (Bonus Qualité Recherche) to C.B. and Y.G., from Agence Nationale de la Recherche (ANR-SOLV\_CDG), from ERANet for Research Programmes on Rare Diseases Joint Transnational Call 2011 (ERARE 11-135 (EURO-CDG)) to F.F., and from the Ministère de l'Enseignement Supérieur to P.A.G. and S.P.

<sup>+</sup> These authors contributed equally

Electronic Supplementary Information (ESI) available: [details of any information supplementary available should be included here]. See DOI: 10.1039/x0xx00000x

pathway.

#### COMMUNICATION

staining methodology.<sup>20</sup>

Published on 16 December 2015. Downloaded by Central Michigan University on 26/12/2015 10:02:17.

Depending on the used chemical tool (ManNAl or SiaNAl), the dynamics of incorporation into glycoconjugates are strikingly different in accordance with the known sialic acids metabolic

While sialylated glycoconjugates can be observed in the Golgi region after only 3h of cellular SiaNAI metabolic labeling, the same Golgi pattern is only observed with ManNAI after 7h of labeling.

In order to confirm the sialin-mediated uptake pathway proposed by Bardor et al.,<sup>21</sup> we applied our SBDS in sialin deficient patients' cells. Indeed, sialin (Q9NRA2) is an anion transporter known to be a membrane protein that exports sialic acids out of lysosomes into the cytosol.<sup>22</sup> To highlight the potential role of sialin in cellular SiaNAI/ManNAI incorporation, control and sialin-deficient patients' cells were metabolically labeled for 8h with either SiaNAl or ManNAl. After incorporation of ManNAl, labeled sialylated glycoconjugates were mainly observed in the perinuclear Golgi like region of both control and sialin deficient patients' cells (Fig. 2). In contrast, the sialin deficient cells metabolically labeled with SiaNAI displayed no staining. The sialin deficient patients' cells have the capacity to transform ManNAl into CMP-SiaNAl, which can then be incorporated into the newly synthesized glycoconjugates. These result show the inability of SiaNAI to reach the cytosol and be converted to CMP-SiaNAI in sialin deficient cells. In support of the pathway suggested previously for Neu5Gc,<sup>21</sup> the present study not only confirms the crucial role of sialin in SiaNAI metabolism but also the potential of this metabolic labeling methodology to decipher deficiencies in sialic acid pathways.



7h Figure 1. A. Structure of the two chemical reporters, SiaNAI and ManNAI. B. Fibroblasts from healthy individuals were metabolically labelled with 500  $\mu$ M of SiaNAl or ManNAl for 2. 5. and 7 h respectively. They were then stained with AzidoFluor545 fluorescent probe (alkyne tagged sialic acid linked to glycoconjugates in red). Staining was

metabolically labelled with 500 %M of ManNAl or SiaNAl for 8 h and stained with AzidoFluor 545 fluorescent probe. Cells were then examined using confocal microscopy. Scale bar: 50 um

visualized using confocal microscopy. Scale bar: 50 µm



pentynoyl)neuraminic acid (SiaNAI) and N-(4-pentynoyl)

mannosamine (ManNAI) (Fig. 1). The visualization of tagged

glycoconjugates was then obtained by chemical ligation of an

azido probe via copper-catalyzed azide-alkyne cycloaddition

In order to study the dynamics of incorporation of our

chemical reporters into glycoconjugates, we have successfully

used SBDS with the two key players ManNAI and SiaNAI after

optimization of their synthesis and of our previously described

Human fibroblast cells from healthy individuals were

metabolically labeled at different time points (2, 5 and 7h,

respectively) with SiaNAI or ManNAI. Figure 1 shows the

distinctive staining patterns observed as a function of labeling

time and alkyne-tagged monosaccharide. When comparing the

incorporation of SiaNAI with that of ManNAI, two pools of

labeled sialylated glycoconjugates can easily be differentiated:

one is perfectly localized in the Golgi compartment while the

other is localized in early endosomes (respectively shown by

(CuAAC),<sup>19</sup> and imaging by confocal fluorescence microscopy.

DOI: 10.1039/C5CC08838K

Journal Name

Published on 16 December 2015. Downloaded by Central Michigan University on 26/12/2015 10:02:17.

Journal Name

View Article Online DOI: 10.1039/C5CC08838K COMMUNICATION



As no staining was visible in sialin deficient cells after SiaNAI labeling, we wanted to discriminate between an endocytosis

Figure 3. Quantification by HPLC after DMB derivatization (A) SiaNAI, both free and linked to glycoconjugates into Sialin deficient patients cells versus control fibroblast cells. (B) Free and linked Neu5Ac into  $\Delta$ Sialin cells versus control fibroblast cells. Bars represent arithmetic means +/- SEM of two independent DMB experiments.

or a lysosomal export defect of sialic acids mediated by sialin. Control and sialin deficient cells were labeled with SiaNAI for 8h and the amount of free and linked SiaNAI was guantified by reverse phase fluorescence HPLC after DMB (1,2-diamino-4,5methylenedioxybenzene dihydrochloride) derivatization (see SI). As shown in Fig. 3, a similar amount of total SiaNAI is observed in sialin deficient cells and control cells. However, while 15% of the observed SiaNAI is incorporated in glycoconjugates in control cells, SiaNAI is exclusively observed as the unbound free monosaccharide in sialin deficient cells. This result demonstrates that SiaNal entry into sialin deficient cells is not prevented. Once in the cell, SiaNAI is blocked into cellular compartments (most likely endosomes and lysosomes). preventing its downstream metabolic incorporation into glycoconjugates. Interestingly, a 60-fold increase in the amount of unbound Neu5Ac is found in sialin deficient cells compared to control cells. Opposite to the aforementioned SiaNAI results, the amount of linked Neu5Ac is unchanged, which indicates that the biosynthetic pathway of sialoglycoconjugates is not affected. Altogether, these data clearly demonstrate that sialin deficiency does not prevent SiaNAI entry, likely achieved via endocytosis, but its export from lysosomes into cytosol.

Subsequently, we investigated the specificity of ManNAI cellular entry mechanism in healthy fibroblasts, as no membrane transporters have been reported for ManNAc to date. Competitive uptake experiments were performed with the relevant mannose-like carbohydrates whose transporters are known<sup>23</sup> (Fig. S2). As expected, the incorporation level into glycoconjugates indicated that N-acetylmannosamine strongly compete with ManNAI for uptake. In contrast, mannosamine and mannose did not. Interestingly we showed that in contrast to ManNAI and SiaNAI, no signal is observed when cells were metabolically labelled with unprotected GalNAI and GlcNAI, then arguing for a specific ManNAI entry mechanism (data not shown). Although we cannot completely exclude a passive diffusion through the plasma membrane for ManNAl, our results re-enforce the presence of a specific ManNAc transporter at the plasma membrane as previously suggested by Varki's group<sup>21</sup>.

In summary, the present report based on the use of unprotected monosaccharides demonstrates the power of these two different metabolic strategies. SiaNAI and ManNAI are shown to be suitable tools to track sialylated glycoconjugate metabolism and particularly to dissect the sialic acids metabolic routes. Indeed, our data bring evidences that both synthetic monosaccharides are up-taken by differential cellular mechanisms.

Then, we demonstrated that this Sequential Bioorthogonal Dual Strategy is suitable to elucidate fundamental intracellular glycobiological questions. In this instance, we confirmed that sialin is crucial to the export of exogenous sialic acids from lysosomes to cytosol. SiaNAI could be a major tool to identify lysosomal diseases or endocytosis deficiencies. Such labeling strategies could also prove valuable in the search of therapeutic molecules to restore sialin functions. Complementary to SiaNAI metabolic labeling, ManNAI is an alternative tool to visualize the ManNAc downstream pathway. For example, in case of GNE deficiency,<sup>24</sup> this methodology could simply be used to assess treatment efficacy.<sup>25</sup>

#### References

- H. Kayser, R. Zeitler, C. Kannicht, D. Grunow, R. Nuck and W. Reutter, J. Biol. Chem., 1992, 267, 16934; L. K. Mahal, K. J. Yarema and C. R. Bertozzi, Science, 1997, 276, 1125; U. Aich and K. J. Yarema, in Glycoscience, Springer, 2008, pp. 2133.
- 2 D. J. Vocadlo, H. C. Hang, E.-J. Kim, J. A. Hanover and C. R. Bertozzi, Proc. Natl. Acad. Sci., 2003, 100, 9116.
- 3 B. W. Zaro, Y.-Y. Yang, H. C. Hang and M. R. Pratt, Proc. Natl. Acad. Sci., 2011, 108, 8146.
- 4 T. Zheng, H. Jiang, M. Gros, D. Soriano del Amo, S. Sundaram, G. Lauvau, F. Marlow, Y. Liu, P. Stanley and P. Wu, Angew. Chem. Int. Ed., 2011, 50, 4113.
- 5 J. Mas Pons, A. Dumont, G. Sautejeau, E. Fugier, A. Baron, S. Dukan and B. Vauzeilles, Angew. Chem. Int. Ed., 2014, 53, 1275.
- 6 K. N. Chuh and M. R. Pratt, Glycoconj. J., 2015 (in press).
- 7 N. J. Agard, J. A. Prescher and C. R. Bertozzi, J. Am. Chem. Soc., 2004, 126, 15046.
- 8 Homann, R. Qamar, S. Serim, P. Dersch and J. Seibel, *Beilstein J. Org. Chem.*, 2010, **6**.
- 9 D. Soriano del Amo, W. Wang, H. Jiang, C. Besanceney, A. C. Yan, M. Levy, Y. Liu, F. L. Marlow and P. Wu, J. Am. Chem. Soc., 2010, 132, 16893.
- 10 A. Niederwieser, A.-K. Späte, L. D. Nguyen, C. Jüngst, W. Reutter and V. Wittmann, Angew. Chem. Int. Ed., 2013, 52, 4265.
- 11 D. M. Patterson, L. A. Nazarova and J. A. Prescher, ACS Chem. Biol., 2014, 9, 592.
- 12 L. Rong, L.-H. Liu, S. Chen, H. Cheng, C.-S. Chen, Z.-Y. Li, S.-Y. Qin and X.-Z. Zhang, Chem Commun, 2014, 50, 667.
- 13 T. Angata and A. Varki, Chem. Rev., 2002, 102, 439.
- 14 C. Traving and R. Schauer, Cell. Mol. Life Sci. CMLS, 1998, 54, 1330.
- 15 R. Schauer and J. P. Kamerling, in Glycoproteins II, Elsevier., 1997, vol. 29b, pp. 243–402.
- 16 J. Vanbeselaere, D. Vicogne, G. Matthijs, C. Biot, F. Foulquier and Y. Guerardel, Chem. Commun., 2013, 49, 11293.

This journal is C The Royal Society of Chemistry 20xx

Published on 16 December 2015. Downloaded by Central Michigan University on 26/12/2015 10:02:17.

- 17 N. E. Mbua, H. Flanagan-Steet, S. Johnson, M. A. Wolfert, G.-J. Boons and R. Steet, Proc. Natl. Acad. Sci., 2013, 110, 10207.
- 18 M. B. Jones, H. Teng, J. K. Rhee, N. Lahar, G. Baskaran and K. J. Yarema, Biotechnol. Bioeng., 2004, 85, 394.
- 19 V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, Angew. Chem. Int. Ed., 2002, 41, 2596; W. Tornøe, C. Christensen and M. Meldal, J. Org. Chem., 2002, 67, 3057.
- 20 R. Péanne, J. Vanbeselaere, D. Vicogne, A.-M. Mir, C. Biot, G. Matthijs, Y. Guérardel and F. Foulquier, in Methods in Cell Biology, Elsevier, 2013, vol. 118, pp. 157.
- 21 M. Bardor, D. H. Nguyen, S. Diaz and A. Varki, J. Biol. Chem., 2005, 280, 4228.
- 22 P. Morin, C. Sagné and B. Gasnier, EMBO J., 2004, 23, 4560.
- 23 K. Panneerselvam and H. H. Freeze, J. Biol. Chem., 1996, 271, 9417.
- 24 C. Oetke, S. Hinderlich, W. Reutter and M. Pawlita, Biochem. Biophys. Res. Commun., 2003, 308, 892-898.
- 25 S.-K. Wang and C.-M. Cheng, Chem Commun, 2015 (in press).

4 | J. Name., 2012, 00, 1-3

Page 4 of 5



Published on 16 December 2015. Downloaded by Central Michigan University on 26/12/2015 10:02:17.

Journal Name

e ChemCömmnargins

View Article Online DOI: 10.1039/C5CC08838K COMMUNICATION



Figure 4. A working hypothesis of cellular uptake and metabolization of ManNAI and SiaNAI. SiaNAI enters via endocytosis and is recycled from lysosomes via Sialin. Once inside the cytosol, SiaNAI is converted into CMP-SiaNAI in the nucleus lumen and used at the Golgi level by the sialyltransferases. The newly sialylated glycoconjugates are then exported either to the plasma membrane or at the early endosomes. We hypothesize that ManNAI enters via a facilitate transporter uncharacterized yet. The cytosolic ManNAI is converted into SiaNAI by the different enzymes (GNE, NANS, NANP). In sialin deficient patients and after ManNAI labelling. In conclusion, SiaNAI labelling can be extremely to identify endosomal/lysosomal dysfunctions while ManNAI labelling allow the sialylated glycoconjugates monitoring.