

RSC Advances



This article can be cited before page numbers have been issued, to do this please use: G. Burgé, A. L. Flourat, B. Pollet, H. E. Spinnler and F. Allais, *RSC Adv.*, 2015, DOI: 10.1039/C5RA18274C.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

3-Hydroxypropionaldehyde (3-HPA) quantification by HPLC using synthetic acrolein-free 3-hydroxypropionaldehyde system as analytical standard

G. Burgé^{§1,2,3}, A.L. Flourat^{§1,4,5}, B. Pollet^{2,3}, H.E. Spinnler^{2,3}, F. Allais^{1,2,3*}

¹ Chaire Agro-Biotechnologies Industrielles (ABI) - AgroParisTech, F-51100 Reims, France

² AgroParisTech, UMR 782 Génie et Microbiologie des Procédés Alimentaires (GMPA), F-78850 Thiverval-Grignon, France

³ INRA, UMR 782 Génie et Microbiologie des Procédés Alimentaires (GMPA), F-78850 Thiverval-Grignon, France

⁴ AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), F-78026 Versailles Cedex, France

⁵ INRA, Institut Jean-Pierre Bourgin (IJPB), F-78026 Versailles Cedex, France

[§] These authors contributed equally to this work.

ABSTRACT HPLC-based quantification of 3-HPA using a synthetic acrolein-free 3-HPA standard obtained from commercially available 1,2,4-butanetriol through a straightforward and easy synthetic process has the advantages over previous colorimetric methods of easier and safer implementation, and greater specificity. This HPLC method is very simple to implement in a lab, does not need any extra handling of the sample to be analyzed, and is suitable even in the presence of other aldehydes and 3-HPA derivatives, provided that the latter do not have similar retention times.

Keywords

3-Hydroxypropionaldehyde, reuterin, chemical synthesis, NMR, HPLC, HESI-MS, GC-MS

Introduction

3-Hydroxypropionaldehyde (aka 3-HPA, β -hydroxypropionaldehyde, reuterin) was originally discovered in 1910 by Voisenet.¹ Its formation was observed during bacterial spoilage of wine by *Bacillus amaracrylus* (now classified as *Paenibacillus polymixa*). Voisenet assumed that 3-HPA was produced by dehydration of glycerol, which was later confirmed by Abeles.² 3-HPA is a viscous and colorless liquid with a slightly acrid smell. Nowadays, the name reuterin is used as a synonym for 3-HPA. This molecule exhibits powerful antimicrobial activity against many pathogenic microorganisms (Gram-positive and Gram-negative, yeast, protozoa),³ particularly into the gastrointestinal tract.⁴ Minimal inhibitory (MIC) and minimal bactericidal concentrations (MBC) were determined for several intestinal bacteria, showing that they were very sensitive to this molecule.⁵ 3-HPA therefore participates, at least in part, on the *L. reuteri* probiotic behavior,⁶ which was confirmed by randomized, double-blinded, and placebo-controlled clinical studies.⁷ 3-HPA may also cause oxidative stress in cells, by reacting with thiol groups of proteins, enzymes and small molecules. However, the mechanisms responsible for this toxicity have not been identified yet.⁸ 3-HPA finds also use as a food preservative and as a therapeutic auxiliary agent in the pharmaceutical industry.⁹

In addition, 3-HPA is a platform chemical of considerable industrial interest as it can be relatively easily converted into a number of commodity chemicals. It can be dehydrated to acrolein then oxidized to acrylic acid. 3-HPA can also be reduced to 1,3-propanediol (1,3-PDO) through the Degussa and Shell processes,¹⁰ or oxidized to 3-hydroxypropionic acid (3-HP) to produce malonic acid. All these molecules are starting materials for polymers and therefore of great interest for the industrial applications. Given the potential of this building block, biotechnological production of 3-HPA from glycerol is therefore being intensively studied¹¹ and commercial production is expected to start in the near future. The biotechnological production of 3-HPA by *Lactobacillus reuteri* is initiated during the stationary growth phase, in the presence of glycerol as substrate.¹²

In water, 3-HPA system is made of several monomeric and oligomeric components that exist in a dynamic equilibrium, and depends upon the set conditions (Fig. 1). For example, studies on the structure of 3-HPA showed that 3-HPA in water is part of a concentration- and pH-dependent dynamic system and revealed that at high concentrations (4.9 M), the 3-HPA system was predominantly composed of 3-HPA dimer along with acrolein and unspecified HPA derivatives.¹⁰




Fig. 1 3-HPA system: (A) 3-HPA monomer, (B) 3-HPA hydrate (aka 1,1,3-trihydroxypropane, 1,1,3-propanetriol or 1,1,3-PPT), (C) 3-HPA cyclic dimer (2-(2-hydroxyethyl)-4-hydroxy-1,4-dioxane), (D) 3-HPA open dimer, (E) and (F) 3-HPA trimers

Given the great interest of 3-HPA, analytical methods for quantification of 3-HPA were developed using MIC (Minimum Inhibitory Concentration, calculated from the highest dilution showing a complete inhibition of the tested strain),⁸ colorimetry and HPLC. Although being one of the most used methods, MIC assay has a major drawback as it only provides relative 3-HPA concentrations expressed in arbitrary units (units 3-HPA.mL⁻¹). Indeed, there is a definite advantage for using methods allowing the determination of defined molar 3-HPA concentrations. To access the latters, colorimetric methods can be used.¹³⁻¹⁵ They all consist in transforming the complex 3-HPA system (monomers and oligomers) into acrolein by dehydrating 3-HPA, thus preventing the formation of the dynamic system. Acrolein is then further converted to UV absorbing compounds allowing its quantification through

colorimetry. Though they allow the determination of the total amount of 3-HPA (mol.L^{-1}), these methods have two main drawbacks. Not only they require extra handling of the samples and the use of harsh conditions (toxic or harmful reagents), but they also do not distinguish between 3-HPA and eventual other aldehydes - such as acrolein - originally present in the samples, thus leading to inaccurate values. HPLC-based analytical methods were also reported,³ however pure 3-HPA (or 3-HPA system) analytical standards being not readily available from common chemical suppliers, such method requires preparing them before performing the analyses. Up to now, pure 3-HPA aqueous samples were obtained through bioconversion of glycerol using *Lactobacillus reuteri*,³ followed by purification of the resulting fermentation broth involving semi-preparative HPLC^{3a} or flash chromatography on silica gel¹⁰ to remove contaminants (e.g., glycerol, 1,3-PDO, acrolein). Though effective, semi-preparative HPLC and silica gel chromatography provide 3-HPA in small quantity or with relatively low yield (45%), respectively. In view of these considerations, the preparation of pure, acrolein- and water-free 3-HPA standard in high yield through a scalable chemical synthesis under mild conditions appeared as a simpler and faster alternative. The main objectives of the present work were therefore to (i) design a straightforward and efficient chemical synthesis of 3-HPA that could be either easily implemented in an analytical chemistry laboratory, or used by a chemical supplier to produce commercial 3-HPA standards, (ii) fully characterize pure synthetic compound using ¹³C NMR, HPLC, GC-MS, LC-MS and FT-IR, as well as (iii) use it to develop a reliable quantitative analytical method using HPLC.

EXPERIMENTAL SECTION

Materials and methods.

Chemicals (1,2,4-butanetriol, sodium periodate, acrolein (>99.0% analytical standard, 0.2% hydroquinone as stabilizer) were purchased from Aldrich and used as received. Reagent grade solvents were purchased from ThermoFisher Scientific. Deuterated chloroform (CDCl_3) was purchased from Euriso-top.

Instrumentation.

Evaporations were conducted under reduced pressure at temperature below 25 °C to avoid 3-HPA evaporation. Column chromatography was carried out with an automated flash chromatography system (PuriFlash 4100, Interchim) and pre-packed INTERCHIM PF-30SI-HP columns (30 μm silica gel).

IR and UV-Visible analyses were performed on Cary 60 UV-VIS and Cary 630 FT-IR from Agilent technologies, respectively.

HESI-MS analyses were performed on a Q-Exactive (Thermo-Fischer Scientific) at a flow rate of 5 $\mu\text{L}/\text{min}$ in positive mode. The electrospray voltage was set to 3.70 kV, the S lens RF level was set to 55 V. The sheath auxiliary and sweep gas (nitrogen) flow rates were set to 30, 10 and 0 arbitrary units, respectively. The capillary and heater temperatures were set to 320

and 50 °C, respectively. Mass spectra were recorded from m/z 50 to m/z 300 at a resolution of 70 000.

GC-MS analyses were performed on a Varian 1200 (quadrupole) equipped with a DB5-MS capillary column 30 x 0.25 mm, 0.25 μ m film thickness (J and W scientific), operated in the temperature program (from 40 °C held 1 min to 80 °C at +30 °C/min, then 80 to 250 °C at +3 °C/min), with helium as carrier gas (constant flow 1.2 mL/min), injector at 260 °C operating in splitless mode and a transfer line to MS at 280 °C. The mass spectral analyses were run with a quadrupole in positive mode (electronic impact, 70 eV).

NMR analyses were performed on a Bruker Fourier 300. ^1H NMR spectra of samples were recorded in CDCl_3 at 300 MHz, chemicals shifts were reported in parts per million relative to the internal standard tetramethylsilane (TMS, $\delta = 0.00$ ppm). ^{13}C NMR spectra of samples were recorded at 75 MHz (CDCl_3 residual signal at $\delta = 77.16$ ppm).

HPLC analyses¹² were performed on a Biorad column (300 mm x 7.8 mm; Biorad, Richmond, USA) equipped with an Aminex HPX-87H Guard column (30 mm x 4.6 mm; Biorad) at 50 °C eluted with aqueous H_2SO_4 (0.005 M) at a flow rate of 0.6 mL.min⁻¹ (Waters 717 plus autosampler; Waters Associates, Millipore, Molsheim, France). Detection was performed by a refractometer (Waters 2414 refractive index) and a UV spectrophotometer (Waters 2489 UV) at 210 nm. Results were processed by Empower software (Waters Associates).

Procedure for the synthesis of synthetic 3-HPA system.

1,2,4-Butanetriol (1.0 g, 9.4 mmol, 1 equiv) was dissolved in acetone, THF or 1,4-dioxane (190 mL, C=0.05 M). Water (4.7 mL) and sodium periodate (8.0 g, 37.6 mmol, 4 equiv) were then added, and the mixture was magnetically stirred at 1100 rpm at room temperature. Reaction was followed by TLC (9/1 Ethyl acetate/methanol) until completion. The crude mixture was then filtered on Celite[®] and the latter washed with 1,4-dioxane. The solvent was removed under reduced pressure, and the crude oil was purified by flash chromatography (100% Ethyl acetate) to provide 3-HPA as a colorless viscous oil.

Heated Electrospray ionization mass spectrometry (HESI-MS) analysis.

The characterization of the 3-HPA system was determined by heated electrospray ionization mass spectrometry (HESI-MS). The sample was dissolved in pestipur grade acetonitrile to obtain a solution with a concentration about 10 ng/ μ L, then filtered through a 0.22 μ m pore-size filter in nylon (Millipore). The sample was infused into the HESI source of the mass spectrometer.

Gas chromatography mass spectrometry (GC-MS) analysis.

Analyses of the 3-HPA system were performed on a gas chromatograph coupled to an electronic impact mass spectrometer and fitted with a 30 m x 0.25 mm HP5 capillary column (film thickness: 0.25 μ m). Extracted sample product was dissolved in pestipur grade dichloromethane to obtain a solution with a concentration of about 1 μ g/ μ L, then dried over

anhydrous sodium sulfate. 5 μL of dried sample solution were silylated in presence of 50 μL of *N,O*-bis-trimethyl-trifluoroacetamide and 5 μL of pyridine (one night at room temperature) before injection onto the GC-MS system.

¹³C Nuclear magnetic resonance (NMR) analysis.

20 mg of synthetic 3-HPA system, after purification, was dissolved in CDCl_3 .

High performance liquid chromatography (HPLC) analysis.

To obtain calibration curves, synthetic 3-HPA was diluted in water at different concentrations. Samples were filtered through a 0.22 μm pore-size filter in nylon (Millipore). Citric acid was used as internal standard and added at 50% (v/v) to the samples before HPLC analysis (20 μL aliquot was injected).

3-HPA system quantification by colorimetric method.

The assay for 3-HPA content was based on the colorimetric method of Circle *et al.* (1945),¹³ modified by Lüthi-Peng *et al.* (2002)¹⁴. To obtain standard curves, 5.6–280 mg.L^{-1} of acrolein (>99.0% analytical standard, 0.2% hydroquinone as stabilizer, Sigma-Aldrich) in water and 3.9–780 mg.L^{-1} of freshly diluted 3-HPA system in water were supplemented with 0.75 mL of DL-tryptophan (Sigma-Aldrich) solution (0.01 M solution in 0.05 M HCl) and 3.0 mL of 37% HCl were added immediately. For 3-HPA quantification, a 1 mL sample was mixed with 0.75 mL of DL-tryptophan (Sigma-Aldrich) solution and 3 mL of HCl 37%. Mixtures containing samples and standards were incubated for 20 min at 37 °C and the optical density was measured at 560 nm (OD_{560}). If necessary, samples were diluted with the same medium to ensure an $\text{OD}_{560} < 1.8$.

RESULTS AND DISCUSSION

Production and purification of synthetic 3-HPA




Fig. 2 Oxidative cleavage of 1,2,4-butanetriol into 3-hydroxypropionaldehyde.

The most commonly used chemical method for producing 3-HPA is the catalytic hydration of acrolein under strong acidic conditions.¹⁶ However, it results in modest yields with many contaminants. Herein, we report the one-step synthesis and study of 3-HPA from commercially available biobased 1,2,4-butanetriol. This precursor is chemically produced via hydrogenation of malate¹⁷ or can be synthesized by bioconversion¹⁸ starting from xylose (*E. coli*) or arabinose (*Pseudomonas fragi*). These microbial syntheses relied on the creation of biosynthetic pathways that do not exist in nature. The aim of this study was to convert 1,2,4-

butanetriol to 3-HPA by oxidative cleavage of the 1,2-diol in presence of sodium periodate (Fig. 2). The reaction was carried out at room temperature and did not need the use of an inert atmosphere. Different solvents were tested in order to evaluate their impact on reaction time (determined by TLC) and final yield (Table 1). To limit formation of oligomers, reaction media was very dilute ($C=0.05M$). Solubility of sodium periodate proved to be a critical factor for the kinetic of oxidation. The more soluble sodium periodate is, the faster the reaction (i.e., 2.5, 4 and 6 h for THF, acetone and 1,4-dioxane, respectively) (Table 1). Even though it required longer reaction time, 1,4-dioxane was the solvent of choice as large amount of salts were still present for reaction performed in acetone and THF, despite many filtration/evaporation cycles. It is noteworthy to mention that, due to its low molecular weight, 3-HPA is easily evaporated during sample concentration. To prevent this phenomenon, the water bath of rotavapor needs to be maintained at 25 °C (or below).

Table 1 Impact of solvent on reaction yield.

Solvent	Reaction time (h)	Yield (%)
Acetone	4	30
THF	2.5	28
1,4-Dioxane	6	70

Crude 3-PA was then purified by flash chromatography on silica gel. Ethyl acetate was used as eluent because of its high polarity, low toxicity, and its inactivity towards aldehydes. Isolated spot of 3-HPA ($R_f = 0.58$ with 9/1 ethyl acetate/methanol) was actually composed of monomeric, dimeric and few trimeric forms, as identified by NMR spectroscopy and GC-MS. To summarize, these quick and simple synthetic and purification procedures allowed 3-HPA to be obtained in very good yield (70%) through a one-step synthetic pathway without any temperature or atmosphere regulation. This method can be easily scaled-up if larger amounts of 3-HPA are needed. It is also noteworthy to mention that, sealed under nitrogen and stored in a freezer (4 °C), this synthetic 3-HPA can be used over a long period of time (ca. 6 months). In addition, in water for concentrations between 0.5 and 10 g/L and without citric acid, no change in signal area and calibration curve coefficients has been observed for both IR and UV detections over several weeks, demonstrating that no deterioration of synthetic 3-HPA system was observed for several weeks.

Synthetic 3-HPA system characterization by HESI-MS

Analysis of 10 ng/ μL 3-HPA solution in acetonitrile with HESI-MS (Fig. 3) showed signals for compounds with a m/z 74 corresponding to 3-HPA monomers, dimers, trimers, as well as their Na^+ and K^+ adducts. It is noteworthy that 3-HPA hydrate could not be detected with this method. Table 2 gives the mass to charge (m/z) ratio of the components of the 3-HPA system. Under these conditions, the 3-HPA dimer was the most ionized form of 3-HPA as shown by peaks with the highest intensity with m/z of 171 corresponding to the sodium adduct of 3-HPA dimer ($[C_6H_{12}O_4Na]^+$). Smaller amounts of monomeric 3-HPA (m/z 74 + 23 = 97 and

m/z 74 + 39 = 113), 3-HPA trimer (m/z 222 + 23 = 245) were also found in the mixture and observed in the spectrum. Conversely, acrolein ($M = 56 \text{ g.mol}^{-1}$, m/z 56 + 23 = 79) was not detected in the synthesized 3-HPA system.




Fig. 3 HESI-MS spectrum of purified 3-HPA (100 ng/ μ L) in acetonitrile. m/z values of 23, 39, 55, 71, 87, 119, and 141 correspond to 3-HPA and its derivatives and their Na^+ , K^+ , $\text{CH}_3\text{CN}+\text{Na}^+$, and $\text{CH}_3\text{CN}+\text{K}^+$ adducts. A detailed description of the m/z values is given in Table 2.

Table 2 m/z values of synthesized 3-HPA system (100 ng/ μ L) in acetonitrile measured by HESI-MS.

3-HPA	3-HPA dimer (C)	3-HPA trimer (E)	3-HPA trimer (F)
74 [C ₃ H ₆ O ₂]	148 [C ₆ H ₁₂ O ₄]	204 [C ₁₂ H ₂₄ O ₈]	222 [C ₉ H ₁₈ O ₆]
97 [+Na] ⁺	171 [+Na] ⁺	227 [+Na] ⁺	245 [+Na] ⁺
113 [+K] ⁺	187 [+K] ⁺	243 [+K] ⁺	261 [+K] ⁺
138 [+CH ₃ CN+Na] ⁺	212 [+CH ₃ CN+Na] ⁺	268 [+CH ₃ CN+Na] ⁺	286 [+CH ₃ CN+Na] ⁺
154 [+CH ₃ CN+Na] ⁺	228 [+CH ₃ CN+K] ⁺	284 [+CH ₃ CN+K] ⁺	302 [+CH ₃ CN+K] ⁺

Synthetic 3-HPA system characterization by GC-MS

In the total ion current (Fig. 4), several forms of 3-HPA (monomeric 3-HPA at 6.1 min, 3-HPA dimers between 18.9 and 20.3 min, 3-HPA trimers between 25.8 and 26.2 min and 3-HPA tetramers at 38.3 min) can be observed. Under these conditions, the 3-HPA dimers were the most abundant. The spectra corresponding to the different species are shown in the

additional data. Results thus confirm that dimers are the major components of the 3-HPA system.




Fig. 4 GC-MS chromatogram of 3-HPA system. Identification of each peak has been made thanks to the mass spectrum of each trimethylsilyl derivative (see Supporting Information). The peaks with retention times around 6, 20, 25 and 38 minutes correspond to 3-HPA monomer, 3-HPA dimers, 3-HPA trimers and 3-HPA tetramers, respectively.

3-HPA system characterization by FT-IR

In infrared spectra (Fig. 5), in addition to the aldehyde band (1714 cm^{-1}), bands in the ether area confirmed a dynamic system rather than pure 3-HPA as these bands correspond to the acetals and hemiacetals belonging to dimeric and trimeric forms (Fig. 5). Furthermore, no alkene band (around 1650 cm^{-1}) has been detected, confirming the absence of acrolein.




Fig. 5 FT-IR analysis of 3-HPA system after purification by flash chromatography on silica gel. The sample purity was tested by HESI-MS and GC-MS analyses before FT-IR analysis. The transmittance of the major functional groups of 3-HPA system are noted.

Table 3 Identification of characteristic vibration bands

Wavelength (cm ⁻¹)	Groups	Bonding	Vibration type
3355.5	Alcohol	Hydrogen bonded OH	Stretch
2962.9-2878.6	Alkanes	CH ₂	Stretch
1714.5	Aldehyde	C=O	Stretch
1247.7-1368.7-1424.3	Alcohol	O-H	Bend
947.8-1031.8-1118.9	ether (hemiacetal and acetal)	C-O	Stretch

Synthetic 3-HPA system characterization by ¹H & ¹³C NMR

Monomer analysis:

¹H NMR (ppm, CDCl₃, 300 MHz) δ = 2.75 (td, J = 5.4 Hz and 0.9 Hz, H₂), 3.95 (m, H₃), 9.86 (t, J = 0.9 Hz, H₁)

It is noteworthy to mention that the chemical shifts obtained here are in accordance with those reported by Kachele *et al.*²⁰ (¹H NMR in D₂O at 400 MHz : 2.78, 3.96 and 9.74 respectively).

¹³C NMR (ppm, CDCl₃, 75 MHz) δ = 46.1 (t, C₂), 56.6 (t, C₃), 202.4 (d, C₁)

Dimers and trimers analysis:

¹³C NMR (ppm, CDCl₃, 75 MHz): Between 88-100 ppm: C belonging to acetal and hemiacetal; between 57-68 ppm: CH₂-OH; between 29-37 ppm: HO-CH₂-CH₂-C.

In the acetal and hemiacetal areas (88-100 ppm, Fig. 6), various signals can be distinguished. They correspond to 1,1,3-PPT (**B**), the two diastereomers of the cyclic dimer (**C**), the opened dimer (**D**) and the trimers (**E**) and (**F**) (Fig. 1).




Fig. 6 ^{13}C NMR spectrum of pure synthetic 3-HPA system in CDCl_3 .

In addition, the major signal in the aldehyde area (see ^1H NMR in Supporting Information) corresponds to monomeric 3-HPA as proved by the correlations in 2D NMR spectra (see Supporting Information). In accordance with the literature,^{14,19} the four most intense signals observed in the acetal area (89.1, 92.7, 95.1 and 99.2 ppm) belong to the two diastereomers of cyclic dimer (**C**), demonstrating that cyclic dimer (**C**) is the most abundant form in our conditions. The other signals probably belong to 1,1,3-PPT (**B**), the open dimer (**D**) and trimers (**E**) and (**F**). No signal corresponding to the double bond of acrolein (around 140 ppm) was detected. This irrefutably demonstrates that no dehydration of 3-HPA into acrolein occurs during this chemical synthesis.

Synthetic 3-HPA system characterization by HPLC and its use as analytical standard for the quantification of 3-HPA in fermentation broth using HPLC

As the efficient separation and analysis of glucose, glycerol, formate, lactate, ethanol, 1,3-PDO, 3-HP and 3-HPA by HPLC using RI detection was already reported in the literature,¹² the conditions published were used to assess the suitability of the synthetic 3-HPA system as analytical standard. Under these conditions, retention times of 3-HPA system and citric acid (internal standard) were: citric acid (8.01 and 8.38 min by UV and RI detectors, respectively), 3-HPA system (15.03 and 15.07 min with UV and RI detectors, respectively).

On the RI chromatogram (Fig. 7 Top), only one peak (retention time = 15.07 min) was obtained. On the contrary, on the UV chromatogram (Fig. 7 Bottom), at least two peaks were obtained (retention times = 15.03 and 27.34 min), indicating that at least two forms of 3-HPA were detected. The unexpected asymmetric shape of the peak is probably due to the dynamic equilibrium between the different forms of 3-HPA system. Unfortunately, despite many attempts (modification of column temperature and/or mobile phase flow), we were not able to improve further the resolution of the peaks in UV detection.

The linearity of the method was assessed by analytical curves of 3-HPA system with citric acid as internal standard for both detections (RI and UV). The calibration curves obtained showed linear regression with R^2 coefficients >0.999 in the concentration range tested (Fig. 8). Repeatability was evaluated by measuring intra-day and inter-day variations of one sample 5 times in one day and in triplicate for three consecutive days. The overall intra- and inter-day relative standard deviations were less than 1.3% and 1.8%, respectively. This analytical method is therefore appropriate for the detection and quantification of 3-HPA system.




Fig. 7 Analysis of 3-HPA by HPLC (distilled water supplemented with 4 g/L of synthetic 3-HPA system). Chromatogram obtained by (Top) Refractive Index detection and (Bottom) UV detection at 210 nm. Citric acid solution ($C = 5$ mM) was used as internal standard and added at 50% (v/v) to the sample before HPLC analysis.




Fig. 8 Quantification using HPLC of synthetic 3-HPA system by (Top) UV detection at 210 nm and (Bottom) Refractive Index detection.

To demonstrate that synthetic 3-HPA system can be used as analytical standard to quantify efficiently and effectively 3-HPA using HPLC, we realized the following experiments (Fig. 9). Using Lüthi-Peng's method,¹⁴ a calibration curve of the optical density at 560 nm ($OD_{(560 \text{ nm})}$) for different concentration of the synthetic 3-HPA system in water (mg.L^{-1}) was obtained and provided the following equation :

$$OD_{(560 \text{ nm})} = 0.0029 [\text{3-HPA system}] + 0.6712 \text{ (Eq. 1)}$$

The same methodology was then applied on aqueous acrolein solutions (mmol.L^{-1}) at different concentrations and gave the following equation :

$$OD_{(560 \text{ nm})} = 0.1434 [\text{acrolein}] + 0.6674 \text{ (Eq. 2)}$$




Fig. 9 Calibration curves obtained for (Top) aqueous acrolein solutions and (Bottom) aqueous synthetic 3-HPA system, using Lüthi-Peng's method.

Using the two equations above, the concentration of the 3-HPA system ([3-HPA system] in mg.L^{-1}) can now be correlated to the actual quantity of 3-HPA in the synthetic 3-HPA system ([3-HPA] in mmol.L^{-1}) knowing that $[\text{acrolein}] = [\text{3-HPA}]$ and with the approximation that the two curves have the same intercept:

$$[\text{3-HPA}] = (0.0029/0.1434) ([\text{3-HPA system}]) \text{ (Eq. 3)}$$

$$[\text{3-HPA}] = 0.0202 [\text{3-HPA system}] \text{ (Eq. 4)}$$

$$[\text{3-HPA}] = [\text{3-HPA system}]/49.4483 \text{ (Eq. 5)}$$

It is noteworthy to mention that $49.4483 \text{ g.mol}^{-1}$ is lower than the molecular weight of monomeric 3-HPA (i.e., $74.0790 \text{ g.mol}^{-1}$) which is consistent with the fact 3-HPA is in the form of a mixture of oligomers.

Using equation 5 and the above calibration curves (Fig. 8 & 9), two different fermentation broths containing 3-HPA have been analyzed using both methods (i.e., HPLC and Lüthi-Peng's). Results are reported in Table 4.

Table 4 Quantification of 3-HPA in two fermentation broths

Sample	[3-HPA] ^a using HPLC				[3-HPA] ^a using colorimetry		Difference (%)
	UV		RI		mmol.L ⁻¹	g.L ⁻¹	
	mmol.L ⁻¹	g.L ⁻¹	mmol.L ⁻¹	g.L ⁻¹			
Broth 1	165.42	12.25	161.98	12.00	169.06	12.52	4.2% (RI) - 2.2% (UV)
Broth 2	191.48	14.18	172.50	12.78	175.73	13.02	1.8% (RI) – 9.0 % (UV)

^a [3-HPA] in g.L⁻¹ is obtained by multiplying [3-HPA] in mol.L⁻¹ by 74.08 g.mol⁻¹ (M_(3-HPA))

In summary, even though synthetic 3-HPA system is a complex mixture, it can be used as analytical standard to reliably quantify 3-HPA using HPLC. However, because of the less resolved peak in UV detection, more reliable data are obtained with RI detection. This HPLC method based on the use of synthetic 3-HPA has been recently applied to monitor 3-HPA in a study dedicated to the production capabilities of diverse *Lactobacillus reuteri* strains.²¹

CONCLUSION

Herein, we propose an efficient and straightforward procedure to synthesize and purify an acrolein- and water-free 3-HPA system in high yield and in only one-step from commercially available 1,2,4-butanetriol under mild conditions (e.g., room temperature, atmospheric pressure). NMR, HPLC and IR analyses confirmed that acrolein was not produced during the course of the chemical reaction. Analyses also indicated that the dimeric form (C) was predominant in the 3-HPA system among several multimeric forms which is consistent given that the composition of the 3-HPA systems is concentration dependent.

This pure synthetic 3-HPA system was then successfully used as analytical standard and allowed a reliable quantitative analysis of 3-HPA using HPLC. Unlike the HPLC and colorimetric methods used so far to quantify 3-HPA, this HPLC method based on synthetic 3-HPA is very simple to implement in a lab, does not need any extra handling of the sample to be analyzed, and is suitable even in the presence of other aldehydes and 3-HPA derivatives, provided that the latter do not have similar retention times.

ASSOCIATED CONTENT

Support information

The Supporting Information is available free of charge on the RSC Publications website at DOI/10.1021/

AUTHOR INFORMATION

Corresponding author

*E-mail: florent.allais@agroparistech.fr Phone: + 33 326364368

Author Contributions

The manuscript was written through contributions of all named authors. All authors have given approval to the final version of the manuscript.

Notes

The Authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors are grateful to the Region Champagne-Ardenne, the Conseil Général de la Marne and Reims Métropole for their financial support.

REFERENCES

1. Voisenet, M. E. *CR Acad. Sci.* **1910**, *150*, 1614-1616.
2. Abeles, R. H.; Brownstein, A. M.; Randles, C. H. *Biochim. Biophys.* **1960**, *41*, 530-531.
3. (a) Talarico, T. L.; Casas, I. A.; Chung, T. C.; Dobrogosz, W. J. *Antimicrob. Agents Chemother.* **1988**, *32*, 1854-1858; (b) Spinler, J. K.; Taweechotipatr, M.; Rognerud, C. L.; Ou, C. N.; Tumwasorn, S.; Versalovic, J. *Anaerobe* **2008**, *14*, 166-171.
4. Dobrogosz, W. J.; Casas, I. A.; Pagano, G. A.; Sjöberg, B. M.; Talarico, T. L.; Karlsson, M. (1989) *Lactobacillus reuteri* and the enteric microbiota in the regulatory and protective role of the normal microflora (R. Grubb, T. Midtvedt, and E. Norin, eds.) Macmillan Press Ltd., London.
5. Cleusix, V.; Lacroix, C.; Vollenweider, S.; Duboux, M.; Le Blay, G. *BMC Microbiol.* **2007**, *7*:101.
6. Casas, I. A.; Dobrogosz, W. J. *Microb. Ecol. Health Dis.* **2000**, *12*, 247-285.
7. Valeur, N.; Engel, P.; Carbajal, N.; Connolly, E.; Ladefoged, K. *Appl. Environ. Microbiol.* **2004**, *70*, 1176-1181.
8. Schaefer, L.; Auchtung, T. A.; Hermans, K. E.; Whitehead, D.; Borhan, B.; Britton, R. A. *Microbiol.* **2010**, *156*, 1589-1599.
9. Sung, H. W.; Chen, C. N.; Liang, H. F.; Hong, M. H. *Biomaterials* **2003**, *24*, 1335-1347.
10. Vollenweider, S.; Grassi, G.; König, I.; Puhon, Z. *J. Agric. Food Chem.* **2003**, *51*, 3287-3293.
11. (a) Lüthi-Peng, Q.; Dileme, F. B.; Puhon, Z. *Appl. Microbiol. Biotechnol.* **2002**, *59*, 289-296; (b) Lüthi-Peng, Q.; Schärer, S.; Puhon, Z. *Appl. Microbiol. Biotechnol.* **2002**,

- 60, 73-80; (c) Ulmer, C.; Zeng, A. P. *Chem. Biochem. Eng.* **2007**, *21*, 321-326; (d) Krauter, H.; Willke Vorlop, K. D. *New Biotechnol.* **2012**, *29*, 211-217.
12. El-Ziney, M. G.; Arneborg, N.; Uyttendaele, M.; Debevere, J.; Jakobsen, M. *Biotechnol. Lett.* **1998**, *20*, 913-916.
 13. Circle, S. J.; Stone, L.; Boruff, C. S. *Ind. Eng. Chem.* **1945**, *17*, 259-262.
 14. Lüthi-Peng, Q.; Schärer, S.; Puhán, Z. *Appl. Microbiol. Biotechnol.* **2002**, *60*, 73-80.
 15. Vollenweider, S.; Evers, S.; Zubriggen, K.; Lacroix, C. *J. Agric. Food Chem.* **2010**, *58*, 10315-10322.
 16. Hall, R. H.; Stern, E. S. *J. Chem. Soc.* **1950**, 490-498.
 17. Valdehuesa, K. N. G.; Liu, H.; Nisola, G. M.; Chung, W. J.; Lee, S. H.; Park, S. J. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 3309-3321.
 18. Niu, W.; Molefe, M. N.; Frost, J. W. *J. Am. Chem. Soc.* **2003**, *125*, 12998-12999.
 19. Nielsen, A. T.; Moore, D. W.; Schuetze, A., Jr. *Pol. J. Chem.* **1981**, *55*, 1393-1403.
 20. Kächele, M.; Monakhova, Y. B.; Kuballa, T.; Lachenmeier, D. W. *Anal. Chim. Acta* **2014**, *820*, 112-118.
 21. Burgé, G.; Saulou-Bérion, C.; Moussa, M.; Pollet, B.; Flourat, A. L.; Allais, F.; Athès, V.; Spinnler, H. E. *Appl. Biochem. Biotechnol.* **2015**, *177*, 923-939.