Synthesis of β -analogues of *C*-mannosyltryptophan, a novel *C*-glycosylamino acid found in proteins[†]

Toshio Nishikawa,*^{a,b} Yuya Koide,^a Akira Kanakubo,^a Hiroshi Yoshimura^a and Minoru Isobe^a

Received 16th November 2005, Accepted 27th January 2006 First published as an Advance Article on the web 16th February 2006 DOI: 10.1039/b516282c

 α -*C*-Mannosyltryptophan (α -C-Man-Trp) has been found to be a novel post-translational modification of tryptophan found from some biologically important glycoproteins. In order to analyze the biological functions of α -C-Man-Trp, we have developed an efficient synthetic strategy for α -C-Man-Trp and its glucose and galactose analogues, starting from α -*C*-glycosidation of the corresponding hexapyranoside derivatives with tinacetylene. According to the synthetic routes, we describe here syntheses of β -anomers of C-Man-Trp, and its glucose and galactose analogues from the corresponding β -*C*-glycosylacetylenes. During this study, we have developed a highly stereocontrolled synthesis of β -*C*-mannosylacetylene that is required for the synthesis of β -C-Man-Trp, while the precedented method gave an anomeric mixture of the *C*-mannosylacetylene. The synthetic C-Man-Trp and its analogues were analyzed by HPLC.

Introduction

 α -C-Mannosyltryptophan (α -C-Man-Trp (1) in Fig. 1) was first found as a novel linkage "C-glycoside" between a protein and carbohydrate from human ribonuclease 2 (RNase 2).1 Interestingly, the mannose moiety of α -C-Man-Trp adopts a ${}^{1}C_{4}$ conformation as a major conformer due to a bulky tryptophan moiety occupying the α -position of the anomeric center and a lack of anomeric effect. Since its discovery, C-Man-Trp has been identified in several biologically important proteins such as interleukin-12,² the terminal four components of a complement system (C6, C7, C8α, β and C9),³ properdin,⁴ thrombospondin,⁵ F-spondin,⁶ mucins (MUC5AC and MUC5B),⁷ and erythropoietin receptors.⁸ Extensive studies on the biosynthetic pathway have revealed that "C-mannosyltransferase," still not identified as an enzyme for installation of mannose to tryptophan, recognizes the amino acid sequence W-X-X-W to modify the first Trp of this motif.9 Since the recognition sequence is included in conserved sequences such as TSP-1 (W-X-X-W-X-X-W-X-X-C) and the WS motif (W-S-X-W-S) in thrombospondin type 1 repeats (TSRs),¹⁰ it is likely that other proteins with these motifs may be modified by Cmannosylation. On the other hand, the biological functions of α -C-Man-Trp have not been clarified, although several possibilities have been studied.11 In order to identify a-C-Man-Trp from proteins, peptide sequencing utilizing Edman degradation, the MS/MS method,12 and specific antibodies against α-C-Man-Trp^{13–15} have been employed; however, such methods have not been able to exclude the possibility of other stereoisomers having a different carbohydrate moiety with the same molecular weight as that of α-C-Man-Trp. Although NMR spectroscopy is the only



Fig. 1 Structure and preferable conformation of α-C-Man-Trp (1).

way to discriminate these isomers, a sufficient amount of the sample containing the modification is difficult to obtain. Thus, a new analytical method to differentiate the isomers of α -C-Man-Trp has been highly desired.^{3,6,8} To supply the α -C-Man-Trp for analyzing its biological functions, we have previously synthesized α -C-Man-Trp, and its glucose and galactose analogues,^{16,17} and its probes for use in biochemical studies.¹⁸ The corresponding β -isomers, however, have been reported to be obtained in only minute amounts by heating L-tryptophan with D-mannose, galactose, and glucose in the presence of acid.¹⁹ We therefore describe herein stereocontrolled syntheses of the β series of C-Man-Trp and its glucose analogues, which can be employed as authentic samples for their trace analyses.

Synthetic plan for β-C-glycosyltryptophan

In our previous papers, we established the synthesis of α -C-Man-Trp and its glucose and galactose analogues as outlined in Scheme 1. Thus, *C*-glycosidation of glycosyl-1-acetate **2** with tinacetylene in the presence of TMSOTf as a Lewis acid exclusively gave the corresponding α -*C*-glycosylacetylene **3**,²⁰ which was coupled with *N*-Ts-*o*-iodoanilide **4** by palladium catalyst to *o*ethylaniline derivatives **5**. Copper(1)-mediated indole formation was followed by deprotection of the Ts group to give α -*C*glycosylindole **6**, which reacted with L-serine-derived aziridine carboxylate **7** in the presence of Sc(ClO₄)₃ as a Lewis acid²¹ to afford a fully protected α -C-Gly-Trp **8**. Finally, two-step

^aLaboratory of Organic Chemistry, Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya, 464-8601, Japan. E-mail: nisikawa@agr.nagoya-u.ac.jp; Fax: (+)81-52-789-4111; Tel: (+)81-52-789-4115

^bPRESTO, Japan Science and Technology Agency, 4-1-8 Honcho, Kawaguchi, Saitama, 332-0012 Japan

[†] Electronic supplementary information (ESI) available: NMR data. See DOI: 10.1039/b516282c



Scheme 1 Synthesis of α-C-Man-Trp and its analogues.

deprotection furnished α -*C*-mannosyltryptophan (1, α -C-Man-Trp), α -*C*-glucosyltryptophan (9, α -C-Glc-Trp), and α -*C*-galactosyltryptophan (10, α -C-Gal-Trp). Based on this synthesis, we planned to synthesize the β -series of the analogues starting from β -*C*-glycosylacetylenes.

Highly stereoselective synthesis of β-C-mannosylacetylene

The β -*C*-glycosylacetylenes, including glucose, galactose, and mannose, were required as the starting materials. Meldal and Vasella *et al.*²² have reported the syntheses of these β -sugar acetylenes as tetrabenzyl ether according to Kishi's protocol,²³ a widely employed and reliable synthetic method for β -*C*-hexapyranosides (Scheme 2); addition of lithium acetylide to the corresponding sugar lactones **A** followed by reduction of the

resulting ketal **B** with triethylsilane in the presence of a Lewis acid.²⁴ The synthetic sequence enabled us to synthesize β -*C*-glucosylacetylene **11** β and β -*C*-galactosylacetylene **12** β in highly stereoselective manner.²² However, when the same procedure was applied to D-mannopyranolactone, reduction of the corresponding ketal proceeded in a low stereoselective manner, giving a 1 : 2 mixture of α - and β -*C*-mannosylacetylenes **13** (see Table 1). Kishi has also reported that the reduction of an allyl-substituted ketal of mannose gave a 1 : 1 mixture of the anomeric isomer. Many efforts have been reported to achieve stereoselective synthesis of β -*C*-mannosylacetylene has been reported to date.²⁷ We thus initiated our studies from the perspective of improving the stereoselectivity in the synthesis of β -*C*-mannosylacetylene **13** β , a starting material for β -C-Man-Trp.

The aforementioned results imply that the β -benzyloxy substituent at the C-2 position of the mannose derivative induces a different conformation of the oxocarbenium cation from those of the corresponding glucose and galactose derivatives. This assumption might be supported by a recent study of Shuto and co-workers,28 from which it was reported that conformationally restricted ketal derived from a 4,6-benzylidene acetal-protected mannopyranolactone undertook a highly stereoselective reduction with Et₃SiH and TMSOTf to exclusively afford β-C-mannoside. We supposed that even if the transition state conformation is different from those of the corresponding glucose and galactose intermediates, the α -face of the oxocarbenium cation should be less hindered and therefore sterically hindered silanes would improve the stereoselectivity of the reduction. We thus examined the influence of sterically hindered silanes in the reduction of ketal 16, which was prepared from 2,3,4,6-tetra-O-benzylmannose (14) *via* the corresponding lactone 15 (Scheme 3 and Table 1).²⁹ To our delight, tris(trimethylsilyl)silane (TMS)₃SiH, one of the most hindered silanes, reduced the ketal 16 to afford β -Cmannosylacetylene 13β as a single isomer (entry 2). Further examination led us to find that triisopropylsilane (i-Pr₃SiH) exhibits a high stereoselectivity, giving 13β in 67% yield along with a trace amount of 13α .³⁰ Since the coupling constant between



Scheme 2 General synthetic route for β -C-glycosylacetylenes.

Table 1 Stereoselectivities of the reduction of ketal 16 with silanes

	Conditions for 1) ^a			Products		
Entry	Silane	Temp.	Conditions for 2)	Yield ^b	$13\alpha:13\beta^c$	
1 2 3	Et ₃ SiH (TMS) ₃ SiH i-Pr ₃ SiH	-40 °C 0 °C 0 °C	TBAF TBAF K2CO3, MeOH	56% (from 14) 76 (from 16) 67 (from 16)	1:2 0:100 1:99	

^{*a*} All reactions were performed with BF₃·OEt₂ as a Lewis acid in CH₃CN–CH₂Cl₂ (85 : 15). ^{*b*} The yields were isolated yields. ^{*c*} The ratios were determined by ¹H NMR.



Scheme 3 Synthesis β -C-mannosylacetylene 13 β .

the protons at the C-1 and C-2 positions of 13a and 13 β are very close (J = 2.0 Hz for 13a, J = 1.0 Hz for 13 β), the α -configuration of 13 β was confirmed by the NOESY spectrum of tetraacetate 17 obtained by acetolysis of 13 β^{22} (Scheme 4).



Scheme 4 Proof of the configuration of 13β

Synthesis of fully protected β-C-glycosyltryptophan

With three β -*C*-glycosylacetylenes (**11** β , **12** β and **13** β) from glucose, galactose, and mannose in hand, we next transformed these products to β -*C*-glycosylindoles, as shown in Table 2, according to our previous studies. The details are including in the supporting information[†]. It is worth noting that in the case of mannosylacetylene **13** β , the cross coupling between **13** β and **4**

gave a mixture of **20** and **23** in a *ca.* $3.5 : 1 \text{ ratio},^{31}$ which could be directly transformed into **26** by treatment with TBAF in refluxing THF.³²

Coupling of β -*C*-glycosylindoles **24–26** with aziridine carboxylate **7** was then carried out as shown in Table 3 to afford β -C-Man-Trp and its glucose and galactose analogues in moderate yields.

Table 3 $Sc({\rm ClO}_4)_3$ -promoted coupling of glycosylindole (24–26) with the aziridine 7



Table 2 Pd-catalyzed coupling between glycosylacetylenes (11 β -13 β) and N-Ts-o-iodoanilide 4, and synthesis of β -C-glycosylindoles (24-26)

					TBAF/THF, refl	ux	Ţ	
	BnO**	H OBn Pd(OAc Et ₃ N, 60	NHTS)2, Ph ₃ P BnO ^{ve}	n H O OBn OBn	NHTs Cul Et ₃ N-E 80 %	DMF BnO st	H NR OBn Bn	
	0.011							
	11β-1 3 β			18-20	T	BAF/THF 21 eflux 24	-23 R = Ts -26 R = H	
	11β-13β		Pd coupling	18-20 g with 4	T r Synthesis of	BAF/THF 21 eflux 24	-23 R = Ts -26 R = H Deprotectio	n of Ts group
Entry	11β-13β Glycosylacetylene		Pd coupling Products	18-20 g with 4 Yield (%)	T Synthesis of Products	BAF/THF 21 eflux 24 f indole Yield (%)	-23 R = Ts $-26 R = H$ $Deprotectio$ $Products$	n of Ts group Yield (%)
Entry 1	11β-13β Glycosylacetylene	Glucose	Pd coupling Products 18	18-20 g with 4 Yield (%) 86	Synthesis of Products 21	BAF/THF 21 eflux 24 f indole Yield (%) 91	-23 R = Ts -26 R = H Deprotectio Products 24	n of Ts group Yield (%) 94
Entry 1 2	11β-13β Glycosylacetylene 11β 12β	Glucose Galactose	Pd coupling Products 18 19	18-20 g with 4 Yield (%) 86 93	Synthesis of Products 21 22	BAF/THF 21 eflux 24 f indole Yield (%) 91 90	-23 R = Ts -26 R = H Deprotectio Products 24 25	n of Ts group Yield (%) 94 92

Deprotection

Deprotection of the methyl ester and benzyl groups of 27, 28 and 29 would then be carried out according to the synthesis of analogues of α -C-Man-Trp (1). However, many modifications of the conditions were necessary for satisfactory yields of the final products 33, 34 and 35.

First, methyl esters of the fully protected C–Gly–Trp (27– 29) were hydrolyzed with aqueous lithium hydroxide (Table 4). In the syntheses of analogues of α -C-Man-Trp (1), 2-propanol was employed as a solvent for the hydrolysis, however, the corresponding β -analogues 27–29 were found to be not sufficiently soluble in 2-propanol. After some experiments, we found a 1 : 1 mixture of acetonitrile and methanol to be a suitable solvent.³³ Under the improved conditions, the esters of 29, 28, and 27 were hydrolyzed to give 30, 31, and 32 in good to moderate yields (entries 1, 2, and 4).

Benzyl groups of tetrabenzyl β-C-Man-Trp 30 were then removed under hydrogenolytic conditions with 5% Pd-C in methanol (Table 4). Because of our initial concern regarding a reductive opening of the pyranose ring that we encountered in the synthesis of α -C-Man-Trp, we were reluctant to add hydrochloric acid. However, debenzylation of 30 was very sluggish in the absence of the acid. Fortunately, the debenzylation proceeded in the presence of concentrated hydrochloric acid to give β-C-Man-Trp (33) in moderate yield (entry 1). Under these conditions, the ring-opening by-product was not observed. When deprotection of tetrabenzyl C-Gal-Trp (31) was conducted with 5% Pd-C in methanol in the presence of 1 N HCl for 42 hours, β -C-Gal-Trp (34) was obtained in 54% yield along with the corresponding Nmethyl C-Gal-Trp in 9% yield (entry 2). However, we found that a shorter time (18.5 hours) suppressed the side reaction to give a 69% yield of the desired product 34 with a trace amount of the *N*-methyl product (entry 3). In the deprotection of tetrabenzyl β -C-Glc-Trp (32), we found that both the substrate 32 and the product 35 were not sufficiently soluble in methanol, resulting in lower yields. A mixture of dioxane and water was found to be a better solvent,¹⁹ and β -C-Glc-Trp 35 was thus obtained in good yield (entry 4). The NMR spectra of these synthetic materials are in good agreement with those of the literature.¹⁸

Table 4 Deprotection of esters and benzyl groups

HPLC analysis of the synthetic analogues of C-Man-Trp

With all six possible isomers (1, 9, 10, 33, 34, and 35) of C-Man-Trp in hand, we next analyzed the synthetic compounds by HPLC with an ODS column attached to UV and fluorescence detectors. Fortunately, all six isomers were clearly separated under conventional conditions as shown in Fig. 2. Interestingly, only β -C-Man-Trp (33) was eluted at a completely different retention time from those of other analogues.

Conclusion

The synthesis of β -C-Man-Trp and its Glc, Gal analogues was carried out in a highly stereoselective manner. The synthesized materials should be useful as authentic samples for developing new analytical methods.

Experimental

General

Optical rotations were measured on a JASCO DIP-370 digital polarimeter. Infrared spectra (IR) were recorded on a JASCO FT/IR-8300 spectrophotometer and are reported in wavenumbers (cm⁻¹). Proton nuclear magnetic resonance (¹H NMR) spectra were recorded on a Bruker AMX-600 (600 MHz), a Bruker ARX-400 (400 MHz), a Bruker AVANCE-400 (400 MHz) or a Varian Gemini-2000 (300 MHz) spectrometer. Data are reported as follows; chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, br = broadened, m = multiplet), couplingconstant and assignment. Carbon nuclear magnetic resonance (13C NMR) spectra were recorded on a Bruker AMX-600 (150 MHz), a Bruker ARX-400 (100 MHz), a Bruker AVANCE-400 (100 MHz) or a Varian Gemini-2000 (75 MHz) spectrometer. High resolution mass spectra (HRMS) were recorded on a JEOL JMS-700 or a LC-MATE spectrometer and are reported in m/z. Elemental analyses were performed by the Analytical Laboratory at the Graduate School of Bioagricultural Sciences, Nagoya University. Reactions were monitored by thin-layer chromatography (TLC) on 0.25 mm silica gel coated glass plates 60 F_{254} (Merck. #1.05715).

	CbzHN, COOMe OBn H N H Bno ^{rr} OBn CH ₃ CI			IN LiOH 3CN-MeOH rt	CbzHN, C OBn H BnO ^{rr} O OBn	$H_2N_{,,}COOH$ $H_2N_{,,}COOH$ $H_2N_{,,}COOH$ $H_2N_{,,}COOH$ H_2				
		27-29			30-32			33-35		
	Deprotection of ester			Deprotection of benzyl groups						
Entry	Substrate		Product	Yield	Cat.	Additive	Solvent	Time/h	Product	Yield (%)
1	29	Mannose	30	86%	5% Pd–C	12 N HCl	MeOH	25	33	43
2	28	Galactose	31	64	5% Pd–C	1 N HCl	MeOH	42	34 ^{<i>a</i>}	54
3					5% Pd–C	1 N HCl	MeOH	18.5	34	69
5								-		_



Fig. 2 HPLC chromatogram of synthetic α -C-Man-Trp and its analogues. Conditions: column ODS-5 Develosil, size; 4.6 \times 250 mm, mobile phase: 10% MeOH–H₂O, 0.1% TFA, flow rate: 0.5 ml min⁻¹.

Cica-reagent silica gel 60 (particle size 0.063–0.2 mm ASTM) and Silica Gel 60 N (spherical, neutral) were used for opencolumn chromatography. Preparative thin-layer chromatographic separations were carried out on 0.5 mm silica gel plates 60 F_{254} (Merck. #1.05774). Unless otherwise noted, non-aqueous reactions were carried out in oven-dried (120 °C) or flame-dried glassware under a nitrogen atmosphere. Dry THF was purchased from Wako Pure Chemical Industries, Ltd. Dry CH₂Cl₂, CH₃CN and Et₃N were distilled from CaH₂ under a nitrogen atmosphere. Sc(ClO₄)₃ was prepared according to the literature. All other commercially available reagents were used as received.

1-Trimethylsilylethynyl-2,3,4,6-tetra-O-benzyl-D-mannopyranose (16). (1) A solution of mannolactol 14 (6.29 g, 11.6 mmol) in DMSO (36 ml) and Ac₂O (24 ml) was stirred at rt for 7 h, and the reaction mixture was quenched with H₂O at 0 °C. The resulting mixture was extracted with AcOEt ($\times 2$). The combined organic extracts were washed with water $(\times 2)$ and brine $(\times 2)$, dried over anhydrous Na₂SO₄, and concentrated. The residue was dissolved in Et₂O, and passed through a short column packed with neutral silica gel, and the eluate was concentrated to afford lactone 15. This material 15 (5.82 g) was used for the next step without further purification. (2) Trimethylsilylethyne (3.05 ml, 21.6 mmol) was dissolved in dry THF (60 ml) and the solution was cooled to -78 °C, and stirred at -78 °C for 20 min. To this solution was added n-BuLi (10.3 ml, 16.2 mmol, 1.57 M in hexane). After stirring at -78 °C for 40 min, the solution was warmed to 0 °C and stirred at 0 °C for 40 min. This solution was again cooled to -78 °C and stirred at -78 °C for 20 min. To this solution

was added a THF (56 ml) solution of the lactone **15** (5.82 g, 10.8 mmol, dried azeotropically with toluene). After stirring for 3.5 h, the reaction was quenched with saturated NH₄Cl solution and extracted with AcOEt (×3). The combined organic extracts were washed with H₂O (×2) and brine (×2), dried over anhydrous Na₂SO₄, and concentrated. The residue (6.87 g) was purified by silica gel column chromatography (150 g, AcOEt–hexane = 1 : 6) to afford ketal **16** (5.52 g, 75% in 2 steps).

(2,3,4,6-Tetra-O-benzyl- β -D-mannopyranosyl)ethyne (13 β). Reduction with $(TMS)_3SiH$ (entry 2 in Table 1). (1) The ketal 16 (3.25 g, 5.11 mmol) was dried azeotropically from toluene, and dissolved in dry CH₃CN (83 ml) and CH₂Cl₂ (15 ml). To this solution cooled at 0 °C were added (TMS)₃SiH (7.90 ml, 25.6 mmol) and BF₃·Et₂O (1.94 µl, 15.3 mmol). After 1.5 h, the reaction was quenched with saturated NaHCO₃ solution and extracted with AcOEt $(\times 3)$. The combined organic extracts were washed with $H_2O(\times 2)$ and brine ($\times 2$), dried over anhydrous Na_2SO_4 , and concentrated. The residue (8.39 g) was purified by silica gel column chromatography (150 g, AcOEt-hexane = 1: $15 \rightarrow 1: 10 \rightarrow 1: 7$) to afford β -C-trimethylsilylmannosylacetylene (2.58 g, 81%). (2) The β -C-trimethylsilylmannosylacetylene (2.58 g, 4.16 mmol) was dissolved in THF (74 ml)-H₂O (4 ml), and TBAF (4.16 ml, 4.16 mmol, 1 M in THF) was added. After stirring at rt for 1.5 h, the reaction was quenched with aqueous NH₄Cl solution and extracted with AcOEt (×3). The combined organic extracts were washed with $H_2O(\times 2)$ and brine (\times 2), dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by silica gel column chromatography (120 g,

AcOEt-hexane = 1 : 6) to afford β -C-mannosylacetylene 13 β (2.13 mg, 93%) as a white solid: $[a]_{10}^{30}$ -31.4 (c 1.05, CHCl₃); IR (KBr) v_{max} 3284, 3032, 2866, 2127, 1954, 1877, 1811, 1606 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.47 (1H, d, J = 2 Hz, C \equiv CH), 3.47 (1H, ddd, J = 9.5, 5, 2 Hz, H-5), 3.55 (1H, dd, J = 9.5, 3 Hz, H-3), 3.68-3.80 (2H, m, H-6), 3.92 (1H, t, J = 9.5 Hz, H-4), 3.97 (1H, dd, J = 3, 1 Hz, H-2), 4.14 (1H, dd, J = 2, 1 Hz, H-1), 4.53 (1H, d, J = 11 Hz, CH_AH_BPh), 4.56 (1H, d, J = 13 Hz, $CH_{\rm C}H_{\rm D}$ Ph), 4.60 (1H, d, J = 11.5 Hz, $CH_{\rm E}H_{\rm F}$ Ph), 4.63 (1H, d, J = 13 Hz, CH_CH_DPh), 4.65 (1H, d, J = 11.5 Hz, CH_EH_FPh), 4.86 (1H, d, J = 11 Hz, CH_AH_BPh), 4.98 (2H, s, CH₂Ph), 7.12–7.50 (20H, m, aromatic); ¹³C NMR (75 MHz, $CDCl_3$) δ 69.0, 69.3, 72.0, 73.4, 74.3, 74.5, 74.6, 75.2, 75.8, 79.9, 80.1, 83.4, 127.5, 127.6, 127.7, 127.7, 128.0, 128.1, 128.1, 128.2, 128.3, 128.4, 128.5, 138.1, 138.2, 138.3, 138.5; Anal. Calcd for C₃₆H₃₆O₅: C, 78.81; H, 6.61. Found: C, 78.81; H, 6.70.

Reduction with $(i-Pr)_3SiH$ (entry 3 in Table 1). The ketal 16 (55 mg, 0.086 mmol) was dried azeotropically with toluene and dissolved in dry CH₃CN-CH₂Cl₂ (1.40 ml-0.25 ml). To this solution was added (i-Pr)₃SiH (88 µl, 0.43 mmol). After the solution was stirring at 0 °C for 25 min, BF₃·OEt₂ (33 µl, 0.26 mmol) was added. After 15 min, the reaction was quenched with saturated NaHCO₃ solution and extracted with AcOEt (\times 3). The combined organic extracts were washed with $H_2O(\times 2)$ and brine (\times 2), dried over anhydrous Na₂SO₄, and concentrated. The residue (52 mg) was dissolved in MeOH (1.0 ml), and K₂CO₃ (52 mg, 0.38 mmol) was added. After stirring at room temperature for 20 min, the reaction was guenched with saturated NH₄Cl solution and extracted with AcOEt $(\times 3)$. The combined organic extracts were washed with $H_2O(\times 2)$ and brine ($\times 2$), dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by column chromatography (silica gel 1.6 g, AcOEt-hexane = 1 : $7 \rightarrow 1:5$) to afford β -C-mannosylacetylene **13** β (32 mg, 67% in 2 steps).

(2,3,4,6-Tetra-O-acetyl- β -D-mannopyranosyl)ethyne (17). To an ice-cold solution of β -C-mannosylacetylene 13 β (54 mg, 0.098 mmol) in Ac₂O (1.6 ml) was added TMSOTf (0.13 ml, 0.63 mmol). After stirring at rt for 38 h 45 min, the reaction was quenched with saturated NaHCO₃ solution and extracted with AcOEt (×3). The combined organic extracts were washed with saturated NaHCO₃ solution (×2), H₂O (×2) and brine (×2), dried over anhydrous Na₂SO₄, and concentrated. The residue (98 mg) was purified by silica gel column chromatography (10 g, Et₂O-hexane = 1 : 2 \rightarrow 1 : 1 \rightarrow 2 : 1) to afford β -Ctetraacetylmannosylacetylene 17 (22 mg, 62%) as a yellow oil. The NMR spectra were identical to those of the literature, diagnostic NOESY correlations were observed between H-1 (δ 4.46, dd, J = 2, 1 Hz) and H-3 (δ 5.06, dd, J = 10, 3.5 Hz) as well as H-1 and H-5 (δ 3.67, 1H, ddd, J = 10, 5.5, 2 Hz).

1-(2,3,4,6-Tetra-*O***-benzyl-**β**-D-glucopyranosyl)**-2-*o*-(*p***-toluensulfoamidyl)phenylethyne (18).** A two-necked round-bottomed flask was charged with β-*C*-glucosylacetylene **11**β (278 mg, 0.507 mmol), *N*-Ts-*o*-iodoanilide **4** (376 mg, 1.01 mmol) and PPh₃ (13.3 mg, 0.0507 mmol), and connected to a vacuum/argon line. The flask was evacuated and then filled with argon. This evacuation–filling cycle was repeated three times. Et₃N (8.3 ml) was added and then the mixture was heated to 60 °C. After these reagents were completely dissolved, Pd(OAc)₂ (5.6 mg, 0.025 mmol) was added. After stirring at 60 °C for 2 h 45 min, the mixture was cooled to room temperature. The reaction was then quenched with saturated NH₄Cl solution and extracted with AcOEt (\times 3). The combined organic extracts were washed with saturated NH₄Cl solution (\times 2), H₂O (\times 2) and brine (\times 2), dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by silica gel column chromatography (30 g, $CH_2Cl_2 \rightarrow$ Et_2O -hexane = 1 : 1) to afford glucosyl- β -1-ethynylaniline 18 (346 mg, 86%) as yellow oil: $[a]_{D}^{27}$ +20 (c 0.98, CHCl₃); IR (KBr) v_{max} 3031, 2868, 2343, 1495, 1454, 1400, 1342, 1167, 1091, 1028 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 2.26 (3H, s, CH₃ of Ts), 3.50–3.83 (6H, m, H-2, H-3, H-4, H-5, H-6), 4.24 (1H, d, J = 9 Hz, H-1), 4.57 (1H, d, J = 12 Hz, CH_AH_BPh), 4.59 (1H, d, J =11 Hz, $CH_{C}H_{D}Ph$), 4.65 (1H, d, J = 12 Hz, $CH_{A}H_{B}Ph$), 4.80 (1H, d, J = 11 Hz, CH_EH_FPh), 4.83 (1H, d, J = 14 Hz, CH_GH_HPh), 4.85 (1H, d, J = 11 Hz, $CH_{c}H_{p}Ph$), 4.90 (1H, d, J = 14 Hz, CH_GH_HPh), 4.92 (1H, d, J = 11 Hz, CH_EH_FPh), 6.98 (1H, t, J =7 Hz, aromatic), 7.11 (2H, d, J = 8 Hz, H-3" of Ts), 7.15–7.38 (23H, m, aromatic, NH), 7.59 (1H, d, J = 8 Hz, aromatic), 7.68 (2H, d, J = 8 Hz, H-2" of Ts); ¹³C NMR (CDCl₃, 100 MHz) δ 21.4, 68.9, 70.2, 73.6, 75.1, 75.5, 75.7, 77.6, 79.2, 80.7, 82.2, 86.1, 93.5, 113.2, 119.8, 124.2, 127.5, 127.7, 127.7, 127.8, 127.9, 127.9, 127.9, 128.0, 128.4, 129.6, 129.9, 132.1, 136.1, 137.7, 137.9, 138.0, 138.2, 138.4, 143.9; Anal. Calcd for C₄₉H₄₇NO₇S: C, 74.12; H, 5.97; N, 1.76. Found: C, 74.05; H, 6.14; N, 1.62.

1-(2,3,4,6-Tetra-O-benzyl-β-D-galactopyranosyl)-2-o-(p-toluensulfoamidyl)phenylethyne (19). Following the procedure for 18, 19 (2.50 g, 93%) was obtained as a yellow oil from $\beta\text{-}C\text{-galactosylacetylene}$ 12 β (1.85 g, 3.38 mmol) and N-Ts-oiodoanilide 4 (2.51 g, 6.76 mmol) after column chromatography (silica gel 120 g, CH₂Cl₂ \rightarrow AcOEt-hexane = 1:3): $[a]_{D}^{27}$ -9.0 (c 1.5, CHCl₃); IR (KBr) v_{max} 3032, 2868, 1496, 1456, 1341, 1167, 1093, cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 2.22 (3H, s, CH₃ of Ts), 3.57 (1H, dd, J = 9.5, 3 Hz, H-3), 3.61-3.71 (3H, m, H-5, H-6), 4.00 (1H, t, J = 9.5 Hz, H-2), 4.02 (1H, dd, J = 3, 0.5 Hz, H-4), 4.19 (1H, d, J = 9.5 Hz, H-1), 4.46 (1H, d, J = 12 Hz, $CH_{A}H_{B}Ph$), 4.54 (1H, d, J = 12 Hz, $CH_{A}H_{B}Ph$), 4.65 (1H, d, J = 11.5 Hz, $CH_{c}H_{D}Ph$), 4.73 (1H, d, J = 12 Hz, $CH_{E}H_{F}Ph$), 4.78 (1H, d, J = 12 Hz, CH_EH_FPh), 4.79 (1H, d, J = 11 Hz, $CH_{G}H_{H}Ph$), 4.86 (1H, d, J = 11 Hz, $CH_{G}H_{H}Ph$), 4.98 (1H, d, J = 11.5 Hz, CH_cH_DPh), 6.98 (1H, dt, J = 7.5 1 Hz, aromatic), 7.06 (2H, br d, J = 8 Hz, aromatic), 7.17–7.41 (23H, m, aromatic, NH), 7.59 (1H, br d, J = 8 Hz, aromatic), 7.64 (2H, br d, J =8 Hz, aromatic); ¹³C NMR (CDCl₃, 100 MHz) δ 21.4, 68.7, 70.6, 72.7, 73.7, 74.0, 75.0, 75.7, 77.5, 78.8, 80.1, 83.5, 93.7, 120.3, 124.3, 127.5, 127.6, 127.8, 127.9, 128.1, 128.1, 128.3, 128.4, 128.5, 129.6, 129.8, 132.1, 138.1, 138.1; Anal. Calcd for C₄₉H₄₇NO₇S: C, 74.12; H, 5.97; N, 1.76. Found: C, 74.12; H, 5.79; N, 1.66.

1-(2,3,4,6-Tetra-*O***-benzyl-β-D-mannopyranosyl)-2-***o***-(***p***-toluen-sulfoamidyl)phenylethyne (20).** Following the procedure for **18**, **20** (2.05 g, 67%) was obtained as a yellow oil along with mannosyl-β-1-tosylindole **23** (579 mg, 19%) from β-*C*-mannosylacetylene **13β** (2.13 g, 3.89 mmol) and *N*-Ts-*o*-iodoanilide **4** (2.89 g, 7.77 mmol) after column chromatography (silica gel 120 g, CH₂Cl₂ → AcOEt-hexane = 1 : 4 → 1 : 2): $[a]_D^{30}$ –19.1 (*c* 0.24, CHCl₃); IR (KBr) ν_{max} 3063, 3032, 2865, 1954, 1811, 1653, 1598, 1496, 1093 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 2.26 (3H, s, CH₃ of Ts), 3.55 (1H, ddd, *J* = 9, 5 Hz, H-5), 3.64 (1H, dd, *J* = 9, 3 Hz,

H-3), 3.73–3.85 (2H, m, H-6 × 2), 3.99 (1H, t, J = 9 Hz, H-4), 4.01 (1H, dd, J = 3, 1 Hz, H-2), 4.36 (1H, d, J = 1 Hz, H-1), 4.57 (1H, d, J = 10.5 Hz, CH_AH_BPh), 4.58 (1H, d, J = 12 Hz, CH_CH_DPh), 4.65 (1H, d, J = 12 Hz, CH_CH_DPh), 4.68 (1H, d, J = 12 Hz, CH_EH_FPh), 4.72 (1H, d, J = 12 Hz, CH_EH_FPh), 4.90 (1H, d, J = 10.5 Hz, CH_AH_BPh), 4.91 (1H, d, J = 12 Hz, CH_GH_HPh), 5.01 (1H, d, J = 12 Hz, CH_GH_HPh), 6.98 (1H, td, J = 3, 1 Hz, aromatic), 7.10–7.42 (26H, m, aromatic), 7.54 (1H, dd, J = 8, 1 Hz, aromatic), 7.68 (1H, dt, J = 8.5, 2 Hz, aromatic);¹³C NMR (CDCl₃, 100 MHz) δ 21.4, 69.5, 69.8, 72.4, 73.5, 74.6, 74.6, 75.3, 76.0, 79.9, 80.7, 83.6, 92.8, 113.2, 119.6, 124.1, 127.5, 127.5, 127.6, 127.7, 127.8, 128.0, 128.1, 128.1, 128.3, 128.4, 128.5, 129.6, 129.8, 132.1, 136.1, 138.0, 138.2, 138.3, 143.8; Anal. Calcd for $C_{49}H_{47}NO_7S$: C, 74.12; H, 5.97; N, 1.76. Found: C, 74.11; H, 6.05; N, 1.53.

2-(2,3,4,6-Tetra-O-benzyl-β-D-glucopyranosyl)-1-(p-toluenesulfonyl)indole (21). Glucosyl-β-1-ethynylaniline 18 (493 mg, 0.623 mmol) was dissolved in Et₃N (9.9 ml) and DMF (4.9 ml), and CuI (23 mg, 0.13 mmol) was added. This solution was stirred at 80 °C for 2.5 h. The reaction was quenched with saturated NH₄Cl solution and extracted with AcOEt (\times 3). The combined organic extracts were washed with saturated NH₄Cl solution (\times 2), H₂O (\times 2) and brine (\times 2), dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by silica gel column chromatography (15 g, AcOEt–hexane = 1:5) to afford glucosyl- β -1-tosylindole **21** (447 mg, 91%) as a yellow oil: $[a]_{D}^{27}$ -46.5 (c 1.03, CHCl₃); IR (KBr) v_{max} 3031, 2865, 1598, 1497, 1453, 1368, 1176, 1092 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 2.20 (3H, s, CH₃ of Ts), 3.68–4.04 (6H, m, H-2, H-3, H-4, H-5, H-6), 4.47 (1H, d, J = 12 Hz, CH_AH_BPh), 4.52 (1H, d, J = 11 Hz, $CH_{C}H_{D}Ph$), 4.58 (1H, d, J = 12 Hz, $CH_{A}H_{B}Ph$), 4.64 (1H, d, J =11 Hz, CH_EH_FPh), 4.77 (1H, d, J = 11 Hz, CH_CH_DPh), 4.88 (1H, d, J = 11 Hz, CH_EH_FPh), 4.94 (1H, d, J = 11.5 Hz, CH_GH_HPh), 4.98 (1H, d, J = 11.5 Hz, CH_GH_HPh), 5.39 (1H, br d, J = 9 Hz, H-1), 6.68 (1H, s, H-3'), 6.92 (2H, br d, J = 8 Hz, aromatic), 6.97 (2H, br dd, J = 8, 1 Hz, aromatic), 7.07–7.36 (20H, m, aromatic) 7.39 (1H, br d, J = 7.5 Hz, aromatic), 7.78 (2H, d, J = 8 Hz aromatic), 8.08 (1H, d, J = 9 Hz, aromatic); ¹³C NMR (CDCl₃, 75 MHz) δ 21.4, 69.1, 73.3, 74.3, 74.9, 75.7, 78.3, 79.0, 81.2, 87.6, 115.2, 121.3, 123.6, 125.0, 127.0, 127.5, 127.6, 127.7, 127.8, 127.9, 128.1, 128.3, 128.5, 128.5, 129.1, 129.5, 135.6, 137.0, 137.9, 138.1, 138.3, 138.6, 144.4; Anal. Calcd for C₄₉H₄₇NO₇S: C, 74.12; H, 5.97; N, 1.76. Found: C, 74.07; H, 6.00; N, 1.64.

2-(2,3,4,6-Tetra-O-benzyl-β-D-galactopyranosyl)-1-(p-toluenesulfonyl)indole (22). Following the procedure for 21, 22 (2.24 g, 90%) was obtained as a yellow oil from galactosyl-β-1ethynylaniline 19 (2.50 g, 3.16 mmol) after column chromatography (120 g, AcOEt-hexane = 1 : 5): $[a]_{D}^{27}$ -49.7 (c 0.35, CHCl₃); IR (KBr) v_{max} 3031, 2869, 1598, 1497, 1497, 1454, 1367, 1092 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 2.18 (3H, s, CH₃ of Ts), 3.54–4.14 (5H, m, H-3, H-4, H-5, H-6), 4.32 (1H, t, J = 9.5 Hz, H-2),4.42 (1H, d, J = 12 Hz, CH_AH_BPh), 4.46 (1H, d, J = 12 Hz, CH_AH_BPh), 4.58 (1H, d, J = 11 Hz, CH_CH_DPh), 4.68 (1H, d, J = 11 Hz, $CH_{\rm E}H_{\rm F}Ph$), 4.74 (1H, d, J = 11.5 Hz, $CH_{\rm G}H_{\rm H}Ph$), 4.82 (1H, d, J = 11.5 Hz, CH_GH_HPh), 4.89 (1H, d, J = 11 Hz, $CH_{c}H_{p}Ph$), 5.00 (1H, d, J = 11 Hz, $CH_{E}H_{F}Ph$), 5.41 (1H, d, J = 9.5 Hz, H-1), 6.71 (1H, s, H-3'), 6.88 (2H, br d, J = 8.5 Hz, aromatic), 7.00 (2H, dd, J = 8, 1 Hz, aromatic), 7.07–7.42 (21H, m, aromatic), 7.73 (2H, d, J = 8.5 Hz, aromatic), 8.06 (1H, d,

 $J = 8.5 \text{ Hz, aromatic}; {}^{13}\text{C NMR} (\text{CDCl}_3, 100 \text{ MHz}) \delta 21.4, 68.6, 72.5, 73.2, 73.4, 73.9, 74.7, 77.1, 77.2, 77.7, 85.5, 111.0, 115.1, 121.3, 123.5, 124.8, 126.9, 127.4, 127.5, 127.7, 127.9, 128.1, 128.2, 128.4, 128.5, 129.2, 129.3, 135.6, 136.9, 137.9, 138.2, 138.3, 138.9, 139.0, 144.2; Anal. Calcd for <math>C_{49}H_{47}NO_7S$: C, 74.12; H, 5.97; N, 1.76. Found: C, 74.13; H, 6.06; N, 1.78.

2-(2,3,4,6-Tetra-O-benzyl-β-D-mannopyranosyl)-1-(p-toluenesulfonyl)indole (23). Following the procedure for 21, 23 (17.2 mg, 87%) was obtained as a yellow oil from mannosyl-β-1-ethynylaniline 20 (19.8 mg, 0.025 mmol) after column chromatography (5 g, AcOEt-hexane = 1 : 4): $[a]_{D}^{30}$ -94.6 (c 0.76, CHCl₃); IR (KBr) v_{max} 3309, 3064, 3031, 2864, 1952, 1597, 1497, 1454 cm⁻¹; ¹H NMR $(CDCl_3, 400 \text{ MHz}) \delta 2.28 (3H, s, CH_3 \text{ of Ts}), 3.68-3.74 (1H, m, m)$ H-5), 3.79–3.86 (2H, m, H-6), 3.88 (1H, dd, J = 9.5, 2.5 Hz, H-3), 4.01 (1H, t, J = 9.5 Hz, H-4), 4.25 (1H, d, J = 11.5 Hz, CH_AH_BPh),4.50 (1H, d, J = 11.5 Hz, CH_A H_B Ph), 4.53 (1H, br d, J = 2.5 Hz, H-2), 4.60 (1H, d, J = 12 Hz, $CH_{c}H_{D}Ph$), 4.61 (1H, d, J = 11 Hz, $CH_{\rm E}H_{\rm F}Ph$), 4.65 (1H, d, J = 12 Hz, $CH_{\rm G}H_{\rm H}Ph$), 4.70 (1H, d, J = 12 Hz, CH_CH_DPh), 4.75 (1H, d, J = 12 Hz, CH_GH_HPh), 4.95 $(1H, d, J = 11 Hz, CH_EH_FPh), 5.13 (1H, s, H-1), 6.89-7.45 (26H,$ m, aromatic, NH), 7.55 (2H, br d, J = 8 Hz, aromatic), 8.08 (1H, d, J = 8 Hz, aromatic); ¹³C NMR (CDCl₃, 100 MHz) δ 21.5, 69.9, 71.8, 73.4, 74.3, 74.9, 75.1, 75.6, 77.2, 80.3, 84.3, 112.4, 115.1, 120.9, 122.4, 123.9, 124.4, 126.3, 126.8, 127.3, 127.4, 127.5, 127.5, 127.6, 127.8, 127.9, 128.0, 128.0, 128.1, 128.3, 128.3, 128.4, 129.5, 129.6, 129.8, 130.1, 135.1, 137.1, 137.9, 138.1, 138.2, 138.5, 138.5, 139.1, 144.9; Anal. Calcd for C₄₉H₄₇NO₇S: C, 74.12; H, 5.97; N, 1.76. Found: C, 74.13; H, 6.06; N, 1.83.

2-(2,3,4,6-Tetra-O-benzyl-β-D-glucopyranosyl)-1H-indole (24). To a solution of glucosyl-β-1-tosylindole 21 (377 mg, 0.476 mmol) in THF (11 ml) was added TBAF (2.3 ml, 2.4 mmol, 1 M in THF). This solution was heated at reflux temperature with stirring for 2 h. The reaction was quenched with saturated NH₄Cl solution and extracted with AcOEt (×3). The combined organic extracts were washed with $H_2O(\times 2)$ and brine ($\times 2$), dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by silica gel column chromatography (5 g, CH2Cl2) to afford glucosyl-β-1-indole 24 (287 mg, 94%) as a yellow oil: $[a]_{D}^{27}$ -14.8 (c 1.01, CHCl₃); IR (KBr) v_{max} 3406, 3032, 2902, 2865, 1455, 1359, 1135, 1062 cm^{-1} ; ¹H NMR (CDCl₃, 400 MHz) δ 3.55–3.85 (6H, m, H-2, H-3, H-4, H-5, H-6), 4.02 (1H, s, J = 10.5 Hz, H-1) 4.50 (1H, d, J = 10 Hz, CH_AH_BPh), 4.56 (1H, d, J = 11.5 Hz, CH_CH_DPh), 4.59 (1H, d, J = 10 Hz, CH_AH_BPh), 4.62 (1H, d, J = 11.5 Hz, $CH_{C}H_{D}Ph$), 4.63 (1H, d, J = 11 Hz, $CH_{E}H_{F}Ph$), 4.88 (1H, d, J =11 Hz, CH_EH_FPh), 4.91 (1H, d, J = 10.5 Hz, CH_GH_HPh), 4.98 $(1H, d, J = 10.5 \text{ Hz}, CH_G H_H Ph), 6.58 (1H, s, indole), 7.00-7.40$ (23H, m, aromatic), 7.60 (1H, s, aromatic), 8.48 (1H, s, NH); ¹³C NMR (CDCl₃, 75 MHz) δ 68.9, 73.5, 75.0, 75.1, 75.7, 75.7, 77.9, 79.1, 82.5, 86.5, 101.6, 111.1, 119.8, 120.6, 121.9, 127.7, 127.8, 127.9, 127.9, 128.0, 128.2, 128.4, 128.4, 128.5, 135.7, 136.1, 137.5, 138.0, 138.1, 138.6; Anal. Calcd for C₄₂H₄₁NO₅: C, 78.85; H, 6.46; N, 2.19. Found: C, 78.86; H, 6.61; N, 2.19.

2-(2,3,4,6-Tetra-O-benzyl-β-D-galactopyranosyl)-1*H***-indole (25).** Following the procedure for **24, 25** (1.67 g, 92%) was obtained as a yellow solid from galactosyl-β-1-tosylindole **22** (2.24 g, 2.83 mmol) after column chromatography (silica gel 90 g, AcOEt–hexane = 1 : 4): $[a]_{D}^{2D}$ –20.8 (*c* 1.05, CHCl₃); IR (KBr) v_{max} 3423, 3032, 2869, 1497, 1455, 1364, 1294, 1101 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 3.64 (2H, dd, J = 6.5, 1.5 Hz, H-6), 3.68–3.78 (2H, m, H-3, H-5), 3.98 (1H, t, J = 9.5 Hz, H-2), 4.06–4.10 (1H, m, H-4), 4.09 (1H, d, J = 10 Hz, CH_AH_BPh), 4.45 (1H, d, J = 11.5 Hz, CH_CH_DPh), 4.47 (1H, d, J = 9.5 Hz, H-1), 4.50 (1H, d, J =11.5 Hz, CH_CH_DPh), 4.56 (1H, d, J = 10 Hz, CH_AH_BPh), 4.64 (1H, d, J = 11 Hz, CH_EH_FPh), 4.79 (2H, s, CH_2Ph), 5.00 (1H, d, J = 11 Hz, CH_EH_FPh), 6.55 (1H, br d, J = 8 Hz, aromatic), 8.59 (1H, s, NH); ¹³C NMR (CDCl₃, 75 MHz) δ 68.7, 72.7, 73.5, 74.3, 74.9, 75.2, 76.1, 77.4, 78.9, 84.0, 101.6, 111.1, 119.6, 120.6, 121.8, 127.6, 127.7, 127.8, 127.9, 128.0, 128.1, 128.3, 128.3, 128.4, 128.5, 135.9, 136.0, 137.9, 138.4, 138.7; Anal. Calcd for $C_{42}H_{41}NO_5$: C, 78.85; H, 6.46; N, 2.19. Found: C, 78.86; H, 6.35; N, 2.23.

2-(2,3,4,6-Tetra-O-benzyl-β-D-mannopyranosyl)-1H-indole (26) from 23. Following the procedure for 24, 26 (10 mg, quant.) was obtained as a yellow solid from mannosyl-\beta-1-tosylindole 23 (12 mg, 0.015 mmol) after column chromatography (silica gel 5 g, AcOEt-hexane = 1 : 4): $[a]_{D}^{30}$ -7.8 (c 1.30, CHCl₃); IR (KBr) v_{max} 3440, 3062, 2863, 1952, 1586, 1455, 1101 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 3.61 (1H, m, H-5), 3.76–3.81 (2H, m, H-6), 3.80 (1H, dd, J = 9.5, 2.5 Hz, H-3), 4.02 (1H, dd, J = 2.5, 1 Hz, H-2), 4.10 $(1H, t, J = 9.5 \text{ Hz}, H-4), 4.23 (1H, d, J = 11 \text{ Hz}, CH_AH_BPh),$ 4.54 (1H, d, J = 12 Hz, $CH_{C}H_{D}Ph$), 4.63 (1H, d, J = 10.5 Hz, $CH_{E}H_{F}Ph$), 4.64 (1H, d, J = 12 Hz, $CH_{C}H_{D}Ph$), 4.68 (1H, d, J = 1 Hz, H-1), 4.73 (1H, d, J = 12 Hz, $CH_{G}H_{H}Ph$), 4.77 (1H, d, J = 12 Hz, CH_GH_HPh), 4.78 (1H, d, J = 11 Hz, CH_AH_BPh), 4.94 (1H, d, J = 10.5 Hz, CH_EH_FPh), 6.34 (1H, d, J = 1.5 Hz, indole), 7.02–7.40 (23H, m, aromatic), 7.57 (1H, br d, J = 8 Hz, aromatic), 8.84–8.90 (1H, br, NH); ¹³C NMR (CDCl₃, 100 MHz) δ 69.4, 72.4, 73.5, 74.6, 74.8, 75.0, 75.3, 78.5, 79.7, 84.4, 99.5, 111.0, 119.5, 120.4, 121.7, 127.4, 127.5, 127.6, 127.7, 127.7, 127.8, 128.1, 128.2, 128.3, 128.3, 128.5, 129.6, 135.6, 135.8, 138.0, 138.2, 138.3, 138.3; Anal. Calcd for C42H41NO5: C, 78.85; H, 6.46; N, 2.19. Found: C, 78.83; H, 6.61; N, 2.15.

2-(2,3,4,6-Tetra-*O***-benzyl-**β**-**D**-mannopyranosyl)-1***H***-indole (26)** from 23. A mixture of β-1-ethynylmannose 20 (2.05 g, 2.59 mmol) and mannosyl-β-1-tosylindole 23 (579 mg, 0.73 mmol) was dissolved in THF (79 ml), and TBAF (10 ml, 10 mmol, 1 M in THF) was added. This solution was stirred at reflux temperature for 6 h. The reaction was quenched with saturated NH₄Cl solution and extracted with AcOEt (×3). The combined organic extracts were washed with H₂O (×2) and brine (×2), dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by column chromatography (silica gel 110 g, AcOEt–hexane = 1 : 4) to afford mannosyl-β-1-indole **26** (1.57 g, 74%) as a yellow solid.

2-(2,3,4,6-Tetra-*O***-benzyl-β-D-glucopyranosyl)-L-(***N***-carbobenzy-loxyl)-tryptophan methyl ester (27).** Sc(ClO₄)₃ (221 mg, 0.645 mmol) placed in a reaction vessel was freeze-dried with benzene for 1 h, then cooled to 0 °C. To this flask was added MS 5A (300 mg) and the vessel was connected to a vacuum/argon line. The flask was evacuated and then filled with argon. This evacuation–filling cycle was repeated three times. In a separate flask, glucosylindole 24 (206 mg, 0.322 mmol) and aziridine 7 (152 mg, 0.645 mmol) were dried azeotropically with benzene, and dissolved in dry CH₂Cl₂ (6 ml). The solution was added to the reaction vessel *via* cannular tubing. The reaction mixture was

stirred at the same temperature for 3 h, and directly subjected to silica gel column chromatography (15 g, AcOEt-hexane = 1: $4 \to 1$: 3) to give 27 (176 mg, 62%) as a brown oil: $[a]_{D}^{27}$ +8.4 (c 0.22, CHCl₃); IR (KBr) v_{max} 3311, 3032, 2922, 1954, 1718, 1455, 1210, 1062 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 3.19 (1H, dd, J = 15, 4 Hz, $CH_AH_BCHCOOMe$), 3.53 (1H, dd, J = 15, 6.5 Hz, CH_AH_BCHCOOMe), 3.53–3.58 (1H, m, H-5), 3.60 (3H, s, $COOCH_3$), 3.62 (1H, t, J = 9 Hz, H-2), 3.69 (1H, dd, J = 10.5, 2 Hz, H-6), 3.76 (1H, dd, J = 10.5, 3 Hz, H-6), 3.75–3.85 (2H, m, H-3, H-4), 4.15 (1H, d, J = 11 Hz, CH_AH_BPh), 4.43 (1H, d, J = 11.5 Hz, $CH_{c}H_{D}Ph$), 4.45 (1H, d, J = 11 Hz, $CH_{E}H_{F}Ph$), 4.52 (1H, d, J = 11 Hz, CH_AH_BPh), 4.56 (1H, d, J = 9 Hz, H-1), 4.58 (1H, d, J = 11.5 Hz, $CH_{c}H_{p}Ph$), 4.65–4.72 (1H, m, CHCOOCH₃), 4.74 (1H, d, J = 11 Hz, $CH_{G}H_{H}Ph$), 4.76 (1H, d, J = 11 Hz, CH_EH_FPh), 4.80 (1H, d, J = 11 Hz, CH_GH_HPh), 5.08 (1H, d, J = 12 Hz, CH_1H_3Ph), 5.12 (1H, d, J = 12 Hz, $CH_{I}H_{J}Ph$), 6.64 (1H, d, J = 8.5 Hz, aromatic), 6.90 (2H, d, J =7 Hz, aromatic), 7.06 (2H, br t, J = 7.5 Hz, aromatic), 7.09–7.39 (24H, m, aromatic), 7.53 (1H, d, J = 8 Hz aromatic), 8.41 (1H, s, NH of indole); ¹³C NMR (100 MHz, CDCl₃) δ 26.9, 52.2, 54.9, 66.9, 68.4, 73.5, 74.2, 74.5, 75.0, 75.5, 77.6, 79.0, 81.0, 86.9, 108.9, 111.2, 118.8, 119.5, 122.4, 127.6, 127.7, 127.8, 128.0, 128.1, 128.2, 128.3, 128.4, 128.4, 128.5, 132.5, 135.6, 137.0, 137.7, 138.2, 138.5, 156.3, 172.6; HRMS (FAB) Calcd for C₅₄H₅₅N₂O₉ (M + H): 875.3908, Found: 875.3887.

2-(2,3,4,6-Tetra-O-benzyl-β-D-galactopyranosyl)-L-(N-carbobenzyloxyl)-tryptophan methyl ester (28). Following the procedure for 27, 28 (60 mg, 39%) was obtained as a yellow oil from galactosylindole 25 (112 mg, 0.175 mmol) and aziridine 7 (82 mg, 0.35 mmol) after column chromatography (12 g, AcOEt-hexane = 1 : 4): $[a]_{D}^{27}$ +10.1 (c 0.36, CHCl₃); IR (KBr) v_{max} 3298, 3032, 2871, 1721, 1455, 1212, 1071 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 3.19 (1H, dd, J = 15, 4 Hz, $CH_AH_BCHCOOMe$), 3.53 (1H, dd, J = 15, 6.5 Hz, $CH_AH_BCHCOOMe$), 3.53–3.58 (1H, m, H-5), 3.60 (3H, s, COOC H_3), 3.62 (1H, t, J = 9 Hz, H-2), 3.69 (1H, dd, J = 10.5, 2 Hz, H-6), 3.76 (1H, dd, J = 10.5, 3 Hz, H-6), 3.75-3.85 (2H, m, H-3, H-4), 4.15 (1H, d, J = 11 Hz, CH_AH_BPh), 4.43 (1H, d, J = 11.5 Hz, $CH_{c}H_{D}Ph$), 4.45 (1H, d, J = 11 Hz, $CH_{E}H_{F}Ph$), 4.52 (1H, d, J = 11 Hz, $CH_{A}H_{B}Ph$), 4.56 (1H, d, J =9 Hz, H-1), 4.58 (1H, d, J = 11.5 Hz, CH_CH_DPh), 4.65–4.72 (1H, m, CHCOOCH₃), 4.74 (1H, d, J = 11 Hz, CH_GH_HPh), 4.76 (1H, d, J = 11 Hz, CH_EH_FPh), 4.80 (1H, d, J = 11 Hz, CH_GH_HPh), 5.08 (1H, d, J = 12 Hz, CH_1H_3Ph), 5.12 (1H, d, J = 12 Hz, CH_IH_JPh), 6.64 (1H, d, J = 8.5 Hz, aromatic), 6.90 (2H, d, J =7 Hz, aromatic), 7.06 (2H, br t, J = 7.5 Hz, aromatic), 7.09–7.39 (24H, m, aromatic), 7.53 (1H, d, J = 8 Hz aromatic), 8.41 (1H, s, NH of indole); ¹³C NMR (100 MHz, CDCl₃) δ 26.9, 52.1, 54.7, 66.7, 68.5, 73.5, 74.3, 74.4, 74.9, 77.2, 77.5, 84.7, 108.8, 111.2, 118.9, 119.5, 122.3, 127.5, 127.7, 127.7, 127.8, 127.9, 128.1, 128.2, 128.2, 128.3, 128.4, 128.4, 128.7, 128.8, 132.9, 135.5, 136.6, 137.1, 137.8, 138.6, 172.5; HRMS (FAB) Calcd for $C_{54}H_{55}N_2O_9$ (M + H): 875.3908, Found: 875.3887.

2-(2,3,4,6-Tetra-O-benzyl-β-D-mannopyranosyl)-L-(N-carbobenzyloxyl)-tryptophan methyl ester (29). Following the procedure for **27**, **29** (140 mg, 49%) was obtained as a yellow oil from mannosylindole **26** (210 mg, 0.329 mmol) and aziridine **7** (154 mg, 0.657 mmol) after column chromatography (15 g, AcOEt– hexane = 1 : 4 \rightarrow 1 : 3): $[a]_{D}^{27}$ +5.7 (*c* 0.35, CHCl₃); IR (KBr)

 v_{max} 3429, 3032, 2868, 1721, 1497, 1455, 1215, 1098 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 3.18 (1H, dd, J = 15, 6.5 Hz, $CH_{A}H_{B}CHCOOMe$), 3.25 (1H, dd, J = 15, 5.5 Hz, CH_AH_BCHCOOMe), 3.61 (3H, s, COOCH₃), 3.59–3.66 (1H, m, H-5), 3.69-3.80 (2H, m, H-6), 3.82 (1H, m, H-3), 4.05 (1H, m, H-2), 4.13 (1H, t, J = 9.5 Hz, H-4), 4.23 (1H, d, J = 10.5 Hz, $CH_{A}H_{B}Ph$), 4.42 (1H, d, J = 12 Hz, $CH_{C}H_{D}Ph$), 4.53–4.62 (1H, m, CHCOOCH₃), 4.56 (1H, d, J = 12 Hz, CH_CH_DPh), 4.61 (1H, d, J = 10.5 Hz, CH_EH_FPh), 4.68–4.78 (3H, m, H-1, CH_2Ph), 4.86 (1H, d, J = 10.5 Hz, CH_AH_BPh), 4.93 (1H, d, J = 10.5 Hz, CH_EH_FPh), 5.02 (1H, d, J = 12 Hz, CH_IH_JPh), 5.06 (1H, d, J =12 Hz, CH_1H_JPh), 5.44 (1H, d, J = 7 Hz, NH-Cbz), 7.03–7.39 (28H, m, aromatic), 7.49 (1H, d, J = 7.5 Hz aromatic), 8.96 (1H, s, NH of indole); ¹³C NMR (100 MHz, CDCl₃) δ 27.1, 52.4, 54.9, 66.9, 69.3, 72.0, 72.5, 73.4, 74.7, 75.1, 75.2, 77.2, 77.7, 79.8, 84.7, 106.4, 111.1, 118.7, 119.4, 122.2, 127.4, 127.6, 127.6, 127.7, 127.9, 128.0, 128.0, 128.1, 128.2, 128.3, 128.4, 128.4, 128.5, 133.0, 135.3, 136.2, 138.0, 138.2, 138.4, 155.8, 172.4; HRMS (FAB) Calcd for C₅₄H₅₅N₂O₉ (M + H): 875.3908, Found: 875.3887.

2-β-D-Mannopyranosyl-L-tryptophan (33). (1) To a solution of 29(130 mg, 0.149 mmol) in CH₃CN–MeOH (3 ml/3 ml) was added 1 N LiOH solution (0.30 ml, 0.30 mmol). After stirring at rt for 14 h, sat. NH₄Cl solution was added. The pH of the mixture was adjusted to 2 with 1 N HCl and then extracted with AcOEt (\times 3). The combined organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by column chromatography (silica gel 10 g, MeOH- $CH_2Cl_2 = 1:20$) to give **30** (110 mg, 86%). (2) A two-necked flask was charged with 5% Pd-C (31 mg) and connected to an inlet adaptor. The flask was evacuated and then filled with nitrogen. A solution of **30** (31 mg, 0.037 mmol) in MeOH (0.94 ml) and 1 N HCl (14 µl) were added. The flask was then evacuated and then filled with hydrogen. After vigorous stirring for 25 h, the mixture was filtered through a pad of Hyflo Super-Cel, and the precipitate was washed with MeOH and H₂O. The combined filtrate was concentrated. The residue (15.2 mg) was purified by preparative TLC (CHCl₃-MeOH-H₂O = 65 : 65 : 15) to give 33, which was further purified by reversed phase column chromatography (Cosmosil 75C₁₈, H_2O as an eluate) to give 33 (5.8 mg, 43%) as a white solid: $[a]_{D}^{22}$ +14.1 (*c* 0.17, H₂O); ¹H NMR (D₂O, 600 MHz) δ 3.26 (1H, dd, J = 15, 9.5 Hz, $CH_A H_B CHCOOH$), 3.61 (1H, dd, J = 15, 4.5 Hz, CH_A H_{B} CHCOOH), 3.66 (1H, ddd, J = 9.5, 6.5,2 Hz, H-5), 3.76 (1H, t, J = 9.5 Hz, H-4), 3.81 (1H, dd, J = 12, 6.5 Hz, H-6), 3.88 (1H, dd, J = 9.5, 3 Hz, H-3), 4.01 (1H, dd, J = 9.5, 4.5 Hz, CHCOOH), 4.04 (1H, dd, J = 12, 2 Hz, H-6), 4.29 (1H, d, J = 3 Hz, H-2), 5.08 (1H, s, H-1), 7.23 (1H, dd, J = 8, 7, Hz,indole), 7.31 (1H, dd, J = 8, 7, Hz, indole), 7.53 (1H, d, J = 8 Hz, indole), 7.75 (1H, d, J = 8 Hz, indole); ¹³C NMR (120 MHz, D₂O) δ 28.6, 58.0, 63.9, 69.7, 74.3, 76.5, 76.6, 82.9, 109.3, 114.6, 121.2, 122.5, 125.3, 129.8, 136.1, 138.3, 177.3; HRMS (FAB) Calcd for $C_{17}H_{23}N_2O_7$ (M + H): 367.1505, Found: 367.1500.

2-β-D-Galactopyranosyl-L-tryptophan (34). (1) To a solution of **28** (60 mg, 0.069 mmol) in CH₃CN–MeOH (2.7 ml/2.7 ml) was added 1 N LiOH solution (0.14 ml, 0.14 mmol). After stirring at rt for 6 h 40 min, sat. NH₄Cl solution was added. The mixture was adjusted to pH 2 with 1 N HCl and then extracted with AcOEt (\times 3). The combined organic layer was washed with water

and brine, dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by preparative TLC (10% MeOH-CH₂Cl₂) to give 31 (38 mg, 64%). (2) A two-necked flask was charged with 5% Pd-C (11.6 mg) and connected to an inlet adaptor. The flask was evacuated and then filled with nitrogen. A solution of 31 (11.6 mg, 0.013 mmol) in MeOH (0.34 ml) and 1 N HCl (4 µl) were added. The flask was then evacuated and then filled with hydrogen. After vigorous stirring for 18.5 h, the mixture was filtered through a pad of Hyflo Super-Cel, and the precipitate was washed with MeOH and H₂O. The combined filtrate was concentrated. The residue (4 mg) was purified by preparative TLC (CHCl₃-MeOH- $H_2O = 65: 65: 15$) to give 35, which was further purified by reversed phase column chromatography (Cosmosil 75C₁₈, H₂O as an eluate) to give **34** (3.3 mg, 69%) as a white solid: $[a]_{D}^{21}$ +3.0 $(c \ 0.17, \ H_2O); \ ^1H \ NMR \ (D_2O, \ 600 \ MHz) \ \delta \ 3.30 \ (1H, \ dd, \ J =$ 15, 9.5 Hz, $CH_AH_BCHCOOH$), 3.60 (1H, dd, J = 15, 4.5 Hz, $CH_AH_BCHCOOH$), 3.80 (1H, dd, J = 14, 12 Hz, H-6), 3.81 (1H, s, H-6), 3.85 (1H, dd, J = 9.5, 3 Hz, H-3), 3.94-3.97 (1H, m, H-5), 4.09 (1H, dd, J = 9.5, 4.5 Hz, CHCOOH), 4.11 (1H, t, J = 9.5 Hz, H-2), 4.11 (1H, br s, H-4), 4.68 (1H, d, J = 9.5 Hz, H-1), 7.23 (1H, ddd, J = 8, 7, 1 Hz, indole), 7.33 (1H, ddd, J = 8, 7, 1 Hz, indole), 7.55 (1H, d, J = 8 Hz, indole), 7.77 (1H, d, J = 8 Hz, indole); ¹³C NMR (120 MHz, D_2O) δ 28.4, 58.0, 64.2, 72.0, 72.8, 76.6, 76.8, 81.8, 111.0, 114.7, 121.6, 122.5, 125.8, 129.6, 136.2, 138.8, 177.1; HRMS (FAB) Calcd for $C_{17}H_{23}N_2O_7$ (M + H): 367.1505, Found: 367.1532.

2-β-D-Glucopyranosyl-L-tryptophan (35). (1) To a solution of 27 (172 mg, 0.197 mmol) in CH₃CN-MeOH (7.9 ml/7.9 ml) was added 1 N LiOH solution (0.39 ml, 0.39 mmol). After stirring at rt for 15 h, sat. NH₄Cl solution was added. The pH of the mixture was adjusted to 2 with 1 N HCl and then extracted with AcOEt (×3). The combined organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, and concentrated. The residue was purified by silica gel column chromatography (15 g, MeOH-CH₂Cl₂ = 1 : 20) to give **32** (109 mg, 64%). (2) A twonecked flask was charged with 10%Pd-C (6.4 mg) and connected to an inlet adaptor. The flask was evacuated and then filled with nitrogen. A solution of 32 (6.4 mg, 0.0074 mmol) in dioxane-H₂O (0.19 ml : 0.032 ml) was added. The flask was then evacuated and then filled with hydrogen. After vigorous stirring for 36 h, 1 N HCl $(2 \mu l)$ was added and stirring was continued for 7 h. The mixture was filtered through a pad of Hyflo Super-Cel, and the precipitate was rinsed with MeOH-CH₃Cl-H₂O (65:65:15). The combined filtrate was concentrated. The residue was washed with MeOH and the precipitate was further washed with CHCl₃-MeOH-H₂O (65:65:15) to give **35** (3.1 mg, quant.) as a white solid: $[a]_{D}^{27}$ +2.6 (c 0.16, H₂O); ¹H NMR (D₂O, 600 MHz) δ 2.88 (1H, dd, J = 15, 9 Hz, $CH_AH_BCHCOOH$), 3.15 (1H, dd, J = 15, 4.5 Hz, CH_A H_B CHCOOH), 3.34 (1H, t, J = 9.5 Hz, H-4), 3.41 (1H, t, *J* = 9.5 Hz, H-3), 3.41 (1H, ddd, *J* = 9.5, 5, 2 Hz, H-5), 3.53 (1H, dd, J = 12.5, 5 Hz H-6), 3.54 (1H, t, J = 9.5 Hz, H-2), 3.53–3.57 (1H, m, CHCOOH), 3.64 (1H, dd, J = 12.5, 2 Hz, H-6), 4.47 (1H, d, J = 9.5 Hz, H-1), 6.92 (1H, t, J = 7.5 Hz, indole), 7.02 (1H, t, J = 7.5 Hz, indole), 7.23 (1H, d, J = 8 Hz, indole), 7.48 (1H, d, J = 8 Hz, indole); ¹³C NMR (120 MHz, CDCl₃) δ 29.8, 58.6, 63.2, 72.1, 75.4, 76.1, 79.6, 82.3, 112.2, 114.2, 121.5, 122.1, 125.4, 129.6, 135.2, 138.6; HRMS (FAB) Calcd for $C_{17}H_{23}N_2O_7$ (M + H): 367.1505, Found: 367.1524.

Acknowledgements

NMR measurements of 600 MHz and elemental analyses were performed by Mr Koga and Mr M. Kitamura in the analytical division of the faculty, whom we thank. This work was financially supported by PRESTO, JST and Scientific research on priority area "Functional Glycomics" from MEXT.

References

- (a) J. Hofsteenge, D. R. Müller, T. Beer, A. Löffler, W. J. Richter and J. Vliegenthart, *Biochemistry*, 1994, 33, 13524–13530; (b) T. Beer, J. Vliegenthart, A. Löffler and J. Hofsteenge, *Biochemistry*, 1995, 34, 11785–11789.
- 2 M.-A. Doucey, D. Hess, M. J. J. Blommers and J. Hofsteenge, *Glycobiology*, 1999, 9, 435–441.
- 3 J. Hofsteenge, M. Blommers, D. Hess, A. Furmanek and O. Miroshnichenko, J. Biol. Chem., 1999, 274, 32786–32794.
- 4 S. Hartmann and J. Hofsteenge, J. Biol. Chem., 2000, 275, 28569-28574.
- 5 J. Hofsteenge, K. G. Huwiler, B. Macek, D. Hess, J. Lawler, D. F. Mosher and J. Peter-Katalinic, J. Biol. Chem., 2001, 276, 6485–6498.
- 6 A. Gonzalez de Peredo, D. Klein, M. Macek, D. Hess, J. Peter-Katalinic and J. Hofsteenge, *Mol. Cell. Proteomics*, 2002, 1, 11–18.
- 7 J. Perez-Vilar, S. H. Randell and R. C. Boucher, *Glycobiology*, 2004, 14, 325–337.
- 8 A. Furmanek, D. Hess, H. Rogniaux and J. Hofsteenge, *Biochemistry*, 2003, 42, 8452–8458.
- 9 J. Krieg, S. Hartmann, A. Vicentini, W. Gläsner, D. Hess and J. Hofsteenge, *Mol. Biol. Cell*, 1998, 9, 301–309.
- 10 (a) J. C. Adams and J. Lawler, *Curr. Biol.*, 1993, **3**, 188–190; (b) J. C. Adams and R. P. Tucker, *Dev. Dyn.*, 2000, **218**, 280–299.
- 11 (a) J. F. G. Vligenthart and F. Casset, *Curr. Opin. Struct. Biol.*, 1998, 8, 565–571; (b) A. Furmanek and J. Hofsteenge, *Acta Biochim. Pol.*, 2000, 47, 781–789.
- 12 J. Hofsteenge, A. Löffler, D. R. Müller, W. J. Richter, T. de Beer and J. F. G. Vligenthart, in *Techniques in protein Chemistry*, VII, ed. D. R. Marshak, Academic Press, New York, 1996, pp. 163–171.
- 13 J. Krieg, W. Gläsner, A. Viventini, M.-A. Douncy, A. Lffler, D. Hess and J. Hofsteenge, J. Biol. Chem., 1997, 272, 26687–26692.
- 14 H. Fujise, H. Horiuchi, K. Adachi, H. Sano and K. Suzuki, patent, WO99/0411; Chem. Abstr. 1999, 130.
- 15 Y. Ihara, S. Manabe, M. Kaneda, H. Kawano, T. Nakayama, I. Sekine, T. Kondo and Y. Ito, *Glycobiology*, 2005, **15**, 383–392.
- 16 (a) T. Nishikawa, M. Ishikawa and M. Isobe, Synlett, 1999, 123–125; (b) T. Nishikawa, M. Ishikawa, K. Wada and M. Isobe, Synlett, 2001, 945–947; (c) T. Nishikawa, Y. Koide, S. Kajii, K. Wada, M. Ishikawa and M. Isobe, Org. Biomol. Chem., 2005, 3, 687–700.
- 17 For the synthesis of C-Man-Trp from the other groups, see: (a) S. Manabe and Y. Ito, J. Am. Chem. Soc., 1999, **121**, 9754–9755; (b) S. Manabe, Y. Marui and Y. Ito, Chem.-Eur. J., 2003, **9**, 1435–1447.

- 18 T. Nishikawa, S. Kajii, C. Sato, Z. Yasukawa, K. Kitajima and M. Isobe, *Bioorg. Med. Chem.*, 2004, **12**, 2343–2348.
- 19 B. Gutsche, C. Grun, D. Scheutzow and M. Herderich, *Biochem. J.*, 1999, **343**, 11–19.
- 20 For accounts of the chemistry of sugar acetylene developed in this laboratory, see: (a) M. Isobe, R. Nishizawa, S. Hosokawa and T. Nishikawa, *Chem. Commun.*, 1998, 2665–2676; (b) M. Isobe and K. Kira, J. Synth. Org. Chem., Jpn., 2000, **58**, 23–30; (c) M. Isobe and K. Kira, J. Synth. Org. Chem., Jpn., 2000, **58**, 99–107.
- 21 T. Nishikawa, S. Kajii, K. Wada, M. Ishikawa and M. Isobe, *Synthesis*, 2002, 1658–1662.
- 22 T. Lowary, M. Meldal, A. Helmboldt, A. Vasella and K. Bock, J. Org. Chem., 1998, 63, 9657–9668.
- 23 M. D. Lewis, J. K. Cha and Y. Kishi, J. Am. Chem. Soc., 1982, 104, 4976–4978.
- 24 For the first application of Kishi's procedure to the synthesis of β-C-glycosylacetylene, see: J.-M. Lancelin, P. H. A. Zollo and P. Sinay, *Tetrahedron Lett.*, 1983, 24, 4833–4836.
- 25 The stereoselective synthesis of β-*C*-mannosides has been a general challenging subject as the synthesis of β-*O*-mannosides in carbohydrate chemistry. For a recent review on β-*O*-mannoside synthesis, see: S. C. Ennis and M. I. Osborn, in *Carbohydrate*, ed. H. M. I. Osborn, Academic Press, Oxford, 2003, pp. 239–276.
- 26 For leading references, see: (a) J. Beignet, J. Tiernan, C. H. Woo, B. M. Kariuki and L. R. Cox, J. Org. Chem., 2004, 69, 6341–6356; (b) G. Stork and J. J. L. Clair, J. Am. Chem. Soc., 1996, 118, 247–248; (c) T. Skrydstrup, D. Mazeas, M. Elmouchir, G. Doisnneau, C. Riche, A. Chiaroni and J.-M. Beau, Chem.–Eur. J., 1997, 3, 1342–1356; (d) J. P. Praly, G.-R. Chen, J. Gola and G. Hetzer, Eur. J. Org. Chem., 2000, 2831–2838.
- 27 Dondoni reported the synthesis of various sugar acetylenes: (a) A. Dondoni, G. Mariotti and A. Marra, *Tetrahedron Lett.*, 2000, 41, 3483–3486; (b) A. Dondoni, G. Moriotti and A. Marra, *J. Org. Chem.*, 2002, 67, 4475–4486.
- 28 M. Terauchi, H. Abe, A. Matsuda and S. Shuto, Org. Lett., 2004, 6, 3751–3754.
- 29 For examples of the reduction of ketals with silanes other than Et₃SiH, see: (a) Y. Wang, S. A. Babirad and Y. Kishi, J. Org. Chem., 1992, 57, 468–481; (b) S. A. Babirad, Y. Wang and Y. Kishi, J. Org. Chem., 1987, 52, 1372–1375; (c) A. Dondoni and M.-C. Scherrmann, Tetrahedron Lett., 1993, 34, 7319–7322; (d) E. Alvarez, R. Perez, M. Rico, R. M. Rodriguez and J. D. Martin, J. Org. Chem., 1996, 61, 3003–3016. For examples of the reduction of cyclic ketals, see: K. C. Nicolaou, C.-K. Hwang and D. A. Nugiel, J. Am. Chem. Soc., 1989, 111, 4136–4137.
- 30 Generality of the reduction with (TMS)₃SiH and i-Pr₃SiH is currently underway.
- 31 Similar cyclization was observed in the Sonogashira coupling (Pd and CuI catalyst) in the synthesis of α-C-Man-Trp, its glucose and galactose analogues, see ref. 16.
- 32 A. Yasuhara, Y. Kanamori, M. Kaneko, A. Numata, Y. Kondo and T. Sakamoto, J. Chem. Soc., Perkin Trans. 1, 1999, 529–534.
- 33 L. Dong and M. J. Miller, J. Org. Chem., 2002, 67, 4759-4770.