

View Article Online

Dalton Transactions

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. I. Nicasio, F. Montilla, E. Álvarez, R. Colodrero and A. Galindo, *Dalton Trans.*, 2016, DOI: 10.1039/C6DT03712G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the **author guidelines**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the ethical guidelines, outlined in our <u>author and reviewer resource centre</u>, still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/dalton

Synthesis and structural characterization of homochiral 2D coordination polymers of zinc and copper with conformationally flexible ditopic imidazolium-based dicarboxylate ligands

Antonio I. Nicasio,^a Francisco Montilla,^{* a} Eleuterio Álvarez,^b Rosario P. Colodrero^{a †} and Agustín Galindo^{* a}

Different novel coordination polymers containing zinc, 1-4, and copper, 5-8, metals, connected via chiral imidazolium-based dicarboxylate ligands, $[L^R]^-$, were isolated by reaction between zinc acetate or copper acetate with enantiomerically pure HL^R compounds. They were characterised and structurally identified by X-ray diffraction methods (single crystal and powder). These compounds are two-dimensional homochiral coordination polymers, $[M(L^R)_2]_n$, in which the metal ions are coordinated by the two carboxylate groups of $[L^R]^$ anions in a general bridging monodentate $\mu^2 - \kappa^1 - O^1 \cdot \kappa^1 - O^3$ fashion that afforded tetrahedral metal coordination environments for zinc, 1-4, and square planar for copper 5-8, complexes. In all the compounds the 3D supramolecular architecture is constructed by non-covalent interactions between the hydrophobic parts (R groups) of the homochiral 2D coordination polymers and, in some cases, by weak C-H···O non-classical hydrogen bonds that provided, in general, a dense crystal packing. DFT calculations on the $[L^R]^-$ anions confirmed their conformational flexibility as ditopic linkers and this fact makes possible the formation of different coordination polymers for four-coordinated metal centers. Preliminary studies in the Zn-catalyzed synthesis of chiral α -aminophosphonates were carried out and, unfortunately, no enantioselectivity was observed in these reactions.

^a Departamento de Química Inorgánica, Facultad de Química, Universidad de Sevilla, Aptdo 1203, 41071 Sevilla, Spain.

^b Instituto de Investigaciones Químicas, CSIC-Universidad de Sevilla, Avda. Américo Vespucio 49, 41092 Sevilla, Spain.

[†] Permanent address: Departamento de Química Inorgánica, Universidad de Málaga, Campus Teatinos s/n, 29071-Málaga, Spain.

Electronic Supplementary Information (ESI) available: Crystal data, supramolecular packing, Rietveld plots, XRPDs selected structural data of 1-3, TGA data, comparison between selected experimental and theoretical structural parameters, DFT optimised structures, energies and MOs. See DOI: 10.1039/x0xx00000x

Introduction

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

The interest in imidazolium-based carboxylic acids has increased markedly.¹ They are easily prepared, based on the Debus-Radziszewski synthesis,² and, in the last several years, they are employed as ionic liquids precursors,³ acid-base bifunctional catalysts,⁴ precursors of N-doped porous carbon materials,⁵ NHC-carbene precursors,⁶ and as ditopic linkers for the construction of coordination polymers^{7,8} or metal–organic frameworks.^{9,10} Moreover, the use of natural aminoacids in the Debus-Radziszewski reaction allowed the straightforward synthesis of chiral imidazolium-based dicarboxylic acids (Scheme 1).^{11,12,13} These compounds have also been employed as precursors of chiral NHC-carbene ligands¹⁴ and as chiral linkers in the construction of coordination polymers. In particular for the latter, compounds of manganese¹⁵ and zinc¹⁶ with imidazolium-based dicarboxylate ligands derived from L-alanine $(R = Me \text{ in Scheme 1, } HL^{Me})$ have been described. The possibility of forming enantiomerically pure metal coordination polymers or metal-molecular frameworks is particularly interesting in the emerging field of asymmetric catalysis,^{17,18} and, with these precedents, we become interested in the development of the coordination chemistry of chiral imidazolium-based dicarboxylic acids (HL^{R} in Scheme 1). Here, we described the synthesis and structural characterization of novel homochiral coordination polymers of zinc, 1-4, and copper, 5-8, metals with the ditopic 1,3-bis(1-carboxylate-1-alkyl)imidazolium linkers (HL^R, R = Me, ⁱPr, CH₂Ph). Complexes containing $[L^{iPr}]^{-1}$ and $[L^{CH2Ph}]^{-1}$ ligands are the first structurally characterised examples of coordination polymers with such a ligands.



Scheme 1 Synthesis of chiral HL^R compounds

Experimental

General

All preparations and other operations were carried out under aerobic conditions. Solvents were purified appropriately prior to use, using standard procedures. Chemicals were obtained from commercial sources and used as supplied. The compounds (S,S)-HL^R were prepared according to the literature procedures.¹³ Infrared spectra were recorded on a Perkin-Elmer FT-IR Spectrum Two spectrophotometer (pressed KBr pellets). NMR spectra were recorded at the Centro de Investigaciones, Tecnología e Innovación (CITIUS) of the University of Sevilla by using Avance III spectrometers with ${}^{13}C{}^{1}H$ and ${}^{1}H$ shifts referenced to the residual solvent signals. All data are reported in ppm downfield from $Si(CH_3)_4$. Polarimetry was carried out using a JASCO P-2000 Digital Polarimeter and the measurements were made at room temperature (concentration of ca. 10 mg/mL). Thermogravimetric Analyses (TGA) were carried out by the CITIUS of the University of Sevilla on a Q600 SDT TA instrument. Elemental analyses (C, H, N) and X-ray Powder Diffraction (XRPD) studies were conducted by the CITIUS of the University of Sevilla on an Elemental LECO CHNS 93 analyser and on a Bruker D8 advance A25 diffractometer, respectively. The HPLC chromatograms were performed on an Agilent 1260 Infinity instrument with a Chiralpak IA column at a flow rate of 0.5 mL/min with heptane/isopropanol = 9/1 (v/v) and using a UV detector at 254 nm.

Syntheses

[**Zn**((*S*,*S*)-**L**^{Me})₂]_n (1). This complex¹⁶ was prepared by a new experimental procedure: Over a solution of (*S*,*S*)-HL^{Me} (0.2122 g, 1 mmol) in a 3:1 mixture of EtOH/H₂O was added a solution of zinc(II) acetate dihydrated (0.1098 g, 0.5 mmol) in EtOH. The resulting mixture was stirred at 70 °C during 1 hour. The resulting white-yellowish solid was isolated by filtration and washed with EtOH and Et₂O and dried. Crystallization from a 1:3 mixture of H₂O/DMF afforded uncoloured crystals of compound **1** (0.145 g, 59 % yield). IR (KBr, cm⁻¹): 3442 (w,br), 3139 (w), 1632 (vs,br), 1560 (m), 1464 (m), 1391 (s), 1363 (s), 1248 (w), 1172 (s), 1111 (w), 1075 (w), 1025 (w), 974 (w), 881 (w), 741 (w), 712 (w), 682 (w), 525 (w), 510 (w). ¹H NMR (D₂O, 300 Hz): δ 1.73 (d, ³*J*_{HH} = 7.5 Hz, 6H, C*H*₃), 4.96 (q, ³*J*_{HH} = 7.5, 2H, C*H*CH₃), 7.48 (s, 2H, NC*H*CHN), 8.83 (s, 1H, NC*H*N). ¹³C{¹H} NMR (D₂O, 75.47 Hz): δ 17.5 (s, CH₃), 60.2 (s, CHCH₃), 123.5 (s, NCHCHN), 134.7 (s, NCHN), 175.5 (COO). Elemental Anal. Calc. for ZnC₁₈H₂₂N₄O₈: C, 44.32; H, 4.49; N, 11.35. Found: C, 44.53; H, 4.55; N, 11.49 %.

 $[Zn((S,S)-L^{iPr})_2 \cdot H_2O]_n$ (2) and $[Zn((R,R)-L^{iPr})_2 \cdot H_2O]_n$ (3). Over a solution of (S,S)-HL^{iPr} (0.2683 g, 1 mmol) in a 3:1 mixture of EtOH/H₂O was added a solution of zinc(II) acetate dihydrated (0.1097 g, 0.5 mmol) in EtOH. The resulting mixture was stirred at 60 °C during 1 hour. The solvent was evaporated and the resulting oil was dissolved in a 1:3 mixture of H₂O/DMF. Slow evaporation of this solution afforded uncoloured crystals of compound $[Zn((S,S)-L^{iPr})_2 H_2O]_n$ (2) (0.960 g, 31 % yield). IR (KBr, cm⁻¹): 3494 (m, br), 3138 (m), 2966 (s), 2877 (m), 1639 (vs), 1560 (m), 1470 (m), 1423 (m), 1375 (vs), 1242 (w), 1184 (w), 1160 (m), 1108 (w), 1021 (w), 982 (w), 920 (w), 843 (w), 818 (w), 754 (m), 717 (w), 650 (w), 501 (w), 452 (w), 401 (w). $[\alpha]_D^{25} = +105.1$ (H₂O). ¹H NMR (D₂O, 300 Hz): δ 0.91 (d, ³J_{HH} = 7.5 Hz, 6H, CHCH₃), 1.01 (d, ${}^{3}J_{HH} = 7.5$ Hz, 6H, CHCH₃), 2.39 (o, ${}^{3}J_{HH} = 7.5$ Hz, 2H, CHCH₃), 4.53 (d, ${}^{3}J_{HH} = 7.5$ Hz, 2H, CHCHCH₃), 7.52 (s, 2H, NCHCHN), 8.90 (s, 1H, NCHN). ¹³C{¹H} NMR (D₂O, 75.47 Hz): δ 17.5 (s, CHCH₃), 18.6 (s, CHCH₃), 31.1 (s, CHCH₃), 71.5 (s, CHCHCH₃), 122.0 (s, NCHCHN), 135.7 (s, NCHN), 173.8 (s, COO). Elemental Anal. Calc. for ZnC₂₆H₃₈N₄O₈·H₂O: C, 50.53; H, 6.52; N, 9.07. Found: C, 50.39; H, 6.33; N, 8.97 %. Following the same procedure, starting from (R,R)-HL^{iPr} (1 mmol), was obtained complex $[Zn((R,R)-L^{iPr})_2 \cdot H_2O]_n$ (3) (0.944 g, 59 % yield). IR (KBr, cm⁻¹): 3440 (vs, br), 3129 (m), 2966 (s), 2878 (w), 1643 (vs), 1563 (w), 1470 (w), 1424 (m), 1384 (vs), 1346 (m), 1247 (w), 1155 (s), 1117 (w), 1024 (w), 985 (w), 922 (w), 844 (w), 819 (w), 753 (s), 718 (w), 649 (w), 512 (vw, br). $[\alpha]_D^{24} = -103.8$ (H₂O). ¹H NMR (D₂O, 300 Hz): δ 0.77 (d, ³J_{HH} = 7.5 Hz, 6H, CH_3), 0.88 (d, ${}^{3}J_{HH} = 7.5$ Hz, 6H, CH_3), 2.38 (o, ${}^{3}J_{HH} = 7.5$ Hz, 2H, $CHCH_3$), 4.51 (d, ${}^{3}J_{HH} =$ 7.5 Hz, 2H, CHCHCH₃), 7.50 (s, 2H, NCHCHN), 8.89 (s, 1H, NCHN). ¹³C{¹H} NMR (D₂O, 75.47 Hz): δ 17.4 (s, CHCH₃), 18.6 (s, CHCH₃), 31.1 (s, CHCH₃), 71.4 (s, CHCHCH₃), 122.0 (s, NCHCHN), 135.7 (s, NCHN), 173.8 (s, COO). Elemental Anal. Calc. for ZnC₂₆H₃₈N₄O₈·2H₂O: C, 49.10; H, 6.66; N, 8.81. Found: C, 49.21; H, 6.40; N, 8.94 %.

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

 $[Zn((S,S)-L^{CH2Ph})_2 \cdot 2DMF \cdot H_2O]_n$ (4). Over a solution of HL^{CH2Ph} (0.1822 g, 0.5 mmol) in MeOH (5 mL) was added a solution of zinc(II) acetate dihydrated (0.0549 g, 0.25 mmol) in MeOH. The resulting mixture was stirred at 80 °C for 3 hours. The solution was evaporated and the resulting oil was dissolved in DMF. Slow evaporation of this solution resulted in formation of uncoloured crystals of compound $[Zn((S,S)-L^{CH2Ph})_2 \cdot 2DMF \cdot H_2O]_n$ (4) (0.160 g, 67 % yield). IR (KBr, cm⁻¹): 3460 (m,br), 3139 (m), 3101 (m), 3063 (m), 3030 (m), 2931 (s), 2864 (m), 1699 (m), 1664 (vs), 1642 (vs), 1557 (m), 1497 (m), 1455 (m), 1411 (m), 1385 (s), 1345 (m), 1298 (m), 1182 (w), 1162 (m), 1096 (m), 1031 (w), 962 (w), 944 (w), 926

(w), 841 (w), 773 (w), 750 (m), 724 (m), 700 (f), 662 (m), 571 (w), 502 (w), 458 (w). $[\alpha]_D^{25} = -85.3^{\circ}$ (H₂O). ¹H NMR (D₂O, 300 Hz): δ 2.75 (s, 3H, *CH*₃, DMF), 2.91 (s, 3H, *CH*₃, DMF), 3.07 (dd, 2H, ¹J_{HH} = 14.5, ³J_{HH} = 11 Hz, *CH*H'Ph), 3.41 (dd, 2H, ¹J_{HH} = 14.5, ³J_{HH} = 4 Hz, CH*H*'Ph), 4.95 (dd, 2H, ¹J_{HH} = 11, ³J_{HH} = 4 Hz, *CH*CH₂Ph), 6.90 (m, 4H, *CH*, Ph), 7.18 (m, 6H, *CH*, Ph), 7.20 (s, 2H, NC*H*CHN), 7.82 (s, 1H, *H*CO, DMF), 8.43 (s, 1H, NC*H*N). ¹³C{¹H} NMR (CD₃OD, 75.47 Hz): δ 30.3 (s, *C*H₃, DMF), 35.6 (s, *C*H₃, DMF), 39.6 (s, *C*H₂Ph), 66.3 (s, *C*HCH₂Ph), 121.2 (s, NCHCHN), 126.8, 128.5, 128.6 (s, *C*H, Ph), 136.1 (s, NCHN), 163.5 (s, HCO, DMF), 172.1 (s, COO). Elemental Anal. Calculated for ZnC₄₂H₃₈N₄O₈·2 DMF·H₂O: C, 60.28; H, 5.69; N, 8.79 %. Found: C, 59.49; H, 5.62; N, 9.04 %. No good microanalytical results were obtained for this complex and here is stated the best one.

[Cu((*S*,*S*)-L^{Me})₂]_n (5). Over a solution of HL^{Me} (1.061 g, 5 mmol) in a 3:1 mixture of EtOH/H₂O (3 mL) was added copper(II) acetate dihydrated (0.501 g, 2.5 mmol). The resulting mixture was stirred at 60 °C during 2 h. The solution was evaporated and the resulting oil was dissolved in a 1:3 mixture of H₂O/DMF. Slow evaporation of this solution afforded blue crystals of compound [Cu((*S*,*S*)-L^{Me})₂]_n (5) (0.9985 g, 82 % yield). IR (KBr, cm⁻¹): 3135 (w), 2972 (w), 1651 (vs), 1565 (m), 1549 (m), 1462 (m), 1390 (s), 1361 (s), 1340 (s), 1258 (m), 1232 (m), 1166 (m), 1101 (w), 1040 (w), 970 (w), 892 (w), 839 (w), 781 (m), 681 (m), 638 (w). [α]_D²⁵ = +62.4° (H₂O). μ_{eff} = 1.72 μ_{B} at 26 °C. Elemental Anal. Calc. for CuC₁₈H₂₂N₄O₈: C, 44.49; H, 4.56; N, 11.53. Found: C, 44.96; H, 4.35; N, 11.52 %.

[Cu((*S*,*S*)-L^{iPr})₂]_n (6) and [Cu((*R*,*R*)-L^{iPr})₂]_n (7). Following the same synthetic procedure of complex **5** were prepared these compounds starting from HL^{iPr} (0.5 mmol) in 74 %, **6**, and 50 %, **7**, yield, respectively. [Cu((*S*,*S*)-L^{iPr})₂]_n (**6**): IR (KBr, cm⁻¹): 3165 (vw), 2965 (s), 2875 (s), 1662 (m), 1639 (vs), 1468 (vw) 1366 (s), 1303 (m), 1230 (m), 1185 (w), 1153 (m), 1117 (w), 919 (w), 854 (vw), 817 (vw), 759 (m), 648 (w), 510 (w), 425 (vw). $[\alpha]_D^{25} = +112.2$ (H₂O). $\mu_{eff} = 1.72 \,\mu_B$ at 26 °C. Elemental Anal. Calc. for CuC₂₆H₃₈N₄O₈: C, 52.21; H, 6.40; N, 9.37. Found: C, 52.75; H, 6.27; N, 9.52 %. [Cu((*R*,*R*)-L^{iPr})₂]_n (**7**): IR (KBr, cm⁻¹): 3468 (s, br), 3217 (m), 3134 (m), 2965 (s), 2876 (w), 1662 (m), 1644 (vs), 1530 (m), 1467 (m) 1418 (m), 1384 (vs), 1315 (m), 1303 (m), 1230 (m), 1185 (m), 1153 (s), 1116 (w), 1029 (w), 978 (vw), 919 (w), 853 (m), 817 (m), 759 (s), 648 (w), 546 (w), 509 (w), 476 (w), 424 (w). [α]_D²⁵ = -107.2 (H₂O). $\mu_{eff} = 1.70 \,\mu_B$ at 26 °C. Elemental Anal. Calc. for CuC₂₆H₃₈N₄O₈: C, 52.21; H, 6.40; N, 9.37. Found: C, 51.67; H, 6.38; N, 9.44 %.

 $[Cu((S,S)-L^{CH2Ph})_2]_n$ (8). Over a solution of (S,S)-HL^{CH2Ph} (0.1822 g, 0.5 mmol) in MeOH (5 mL) was added a solution of sodium acetate (0.0410 g, 0.5 mmol). Over this solution was added a solution of copper chloride dihydrated (0.0426 g, 0.25 mmol). The resulting mixture was stirred at 60 °C during 2 h. A blue precipitate was formed and isolated by filtration. The solid was dissolved in hot MeOH (10 mL) and DMF (10 mL) was added. Slow evaporation of this solution generated blue crystals of compound $[Cu((S,S)-L^{CH2Ph})_2]_n$ (8) in 20 % yield. IR (KBr, cm⁻¹): 3469 (m,br), 3150 (m),3130 (m), 3101 (m), 3064 (m), 3028 (m), 2929 (s), 1720 (w), 1645 (vs), 1564 (m), 1550 (m), 1497 (m), 1455 (m), 1419 (m), 1384 (s), 1347 (m), 1282 (m), 1253 (m), 1198 (m), 1184 (m), 1155 (w), 1104 (m), 1084 (m), 1032 (w), 944 (w), 894 (w), 862 (w), 833 (w), 812 (w), 752 (w), 731 (m), 701 (m), 677 (m), 636 (m), 574 (w), 504 (w).

X-ray crystallography

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

A summary of the crystallographic data and structure refinement results for compounds 2, 4-6 and 8 is given in Table S1. Crystals of suitable size for X-ray diffraction analysis were coated with dry perfluoropolyether and mounted on glass fibers and fixed in a cold nitrogen stream (T = 213 K) to the goniometer head. Data collection was performed on a Bruker-Nonius X8Apex-II CCD diffractometer, using monochromatic radiation λ (Mo K_a) = 0.71073 Å, by means of ω and φ scans with a width of 0.50 degree. The data were reduced (SAINT¹⁹) and corrected for absorption effects by the multi-scan method (SADABS).²⁰ The structures were solved by direct methods (SIR-2002²¹) and refined against all F^2 data by full-matrix leastsquares techniques (SHELXL-2016/6²²) minimizing $w[F_0^2 - F_c^2]^2$. All non-hydrogen atoms were refined anisotropically. The hydrogen atoms were included from calculated positions and refined riding on their respective carbon atoms with isotropic displacement parameters. A search for solvent accessible voids in 2 and 8 crystals using PLATON- SQUEEZE,²³ showed potential solvent volumes of 486 $Å^3$ (equating to 9 molecules of water per unit cell) and 2116 $Å^3$ (equating to 42 molecules of water per unit cell), respectively, impossible to model even with the most severe restraints. The corresponding CIF data represent SQUEEZE treated structures with the water molecules handling as a diffuse contribution to the overall scattering, without specific atom position and excluded from the structural model. The SQUEEZE output results were embedded within the CIF by SHELXL-2016/6. CCDC 1407226 (2), 1493049 (4), 1493050 (5), 1407227 (6) and 1493051 (8) contain the supplementary crystallographic data for this paper.

X-ray powder diffraction

X-ray powder diffraction (XRPD) patterns were collected on a Bruker D8 advance A25 diffractometer in a Bragg-Brentano reflection configuration by using the Lynxeye detector. A summary of the crystallographic data and structure refinement results for compounds **3** and **7** is given in Table S2. XRPD patterns were auto-indexed using the DICVOL06 program,²⁴ in an orthorhombic cell for $[Zn((R,R)-L^{iPr})_2 H_2O]_n$ (**3**) and in a triclinic unit cell for $[Cu((R,R)-L^{iPr})_2]_n$ (**7**). Their crystal structures were determined by the Rietveld method,²⁵ using the structures of $[Zn((S,S)-L^{iPr})_2 H_2O]_n$ (**2**) and $[Cu((S,S)-L^{iPr})_2]_n$ (**6**) as starting models, respectively. The Rietveld refinement was carried out by using the GSAS package²⁶ and the graphic interface EXPGUI²⁷ with soft constraints to maintain chemically reasonable geometries for the imidazolium moiety and the alkyl chain: N^{...}C and C^{...}C, 1.50(1), C^{...}C, 2.50(2), C^{...}O, 1.23(1), O^{...}O, 2.21(2), C^{...}O, 2.36(2) Å, for the CH[(CH(CH₃)₂]COO moiety. Hydrogen atoms were not included in the refinements. The final weight factors for the soft constrains were fixed at 10. Only one isotropic atomic displacement parameter was refined.

Computational details

The electronic structure and geometries of the conformers of $[(S,S)-L^R]^-$ anions $(R = Me, {}^iPr,$ CH₂Ph) were computed using density functional theory at the B3LYP level,²⁸ with the 6-311G+(d,p) basis set for all the atoms. This combination of method and basis sets provides a good structural description of the (S,S)-HL^R compounds according to the comparison of the structural parameters of the optimised structures of (S,S)-HL^{Me} and (S,S)-HL^{CH2Ph}, with those of the reported crystal structures (CSD refcodes LUYSAA and LUYRON, respectively; see Table S3).¹² The conformers of the enantiomerically related $[(R,R)-L^{iPr}]^{-1}$ anion were also computed and the resulting structure and energies were compared with those of the [(S,S)- L^{iPr} anion. Since no noteworthy differences were found, other $[(R,R)-L^{R}]^{-1}$ anions were not theoretically analyzed. Molecular geometries of all $[L^{R}]^{-}$ compounds were optimised without symmetry constraints. Frequency calculations were carried out at the same level of theory to identify all of the stationary points as minima (zero imaginary frequencies) and to provide the thermal correction to free energies at 298.15 K and 1 atm. The DFT calculations were performed using the Gaussian 09 suite of programs.²⁹ The cartesian coordinates and energetic parameters of all optimised compounds are reported in the Electronic Supplementary Information (ESI).

General procedure for enantioselective Zn-catalyzed synthesis of diethyl ((4-methoxyphenyl)((4-methoxyphenyl)amino)methyl)phosphonate

The reactor (a 50 mL vial equipped with a Young valve and containing a stirrer flea) was charged with 4-methoxybenzaldehyde (0,5 mmol), 4-methoxyaniline (0,5 mmol), diethyl phosphonate (0,5 mmol), Zn(II) coordination polymer **1** or **2** (0.05 mmol) and the reaction solvent (1 mL), in the aforementioned order. Zn-catalyzed *in-situ* reaction were performed using the same procedure, but introducing zinc acetate (0.05 mmol) and HL^R (0.05 mmol), instead the preformed Zn complex. The reactor was sealed and maintained at the working temperature, with constant stirring (600 rpm) in a thermostated bath for the duration of the reaction. Upon completion the resulting mixture was evaporated to dryness affording a residue that was extracted with dichloromethane (10 mL) and then filtered with 0.45 μ m nylon syringe filter. An aliquot (1 mL) of the resulting solution was evaporated to dryness and then the residue was analyzed by ¹H and ³¹P{¹H} NMR. Finally, an aliquot (1 mL) of the dichloromethane solution was analyzed by HPLC (by diluting with 10 mL of isopropanol).

Results and discussion

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

Syntheses of complexes 1-8

The treatment of solutions of zinc acetate or copper acetate with HL^R compounds results in the formation, after the appropriate work up, of colourless or blue crystals of compounds **1-4** (Zn) or **5-8** (Cu), respectively (Scheme 2). Broad IR bands centred at around 1640 cm⁻¹, due to the antisymmetric COO vibrations, are indicative of the presence of carboxylate groups in these complexes. These bands are slightly shifted to lower wavenumbers with respect those of the HL^R compounds (for instance, 1684 cm⁻¹ for compound HL^{iPr}), in agreement with ligand [L^R]⁻ coordination. A comparable absorption was reported for the analogous zinc derivatives with the [L^H]⁻ and [L^{Me}]⁻ ligands7^{a,c,16} or for copper compounds with the [L^H]⁻ ligand.8 The attribution of the absorptions within the range 1390-1370 cm⁻¹ to the symmetric COO vibrations allowed to determine a $\Delta(vCOO_{asym}-vCOO_{sym})$ value of *ca*. 250 cm⁻¹. This value is compatible with the κ^1 -O coordination of the carboxylate group observed in the structural characterization of these compounds (see below). NMR spectra (¹H and ¹³C{¹H}) for complexes **2-4** are similar to those reported for HL^R compounds¹³ and only small

Dalton Transactions

displacements of the $[L^R]^-$ signals are observed with respect to the parent HL^R compounds. Spectroscopic properties of complex **1**, IR and NMR spectra, are similar to those reported in the bibliography.¹⁶ Compounds **5-8** are paramagnetic and the μ_{eff} values, close to 1.7 μ B, are in agreement with the presence of a square planar copper(II) centre. To our knowledge, there are no structurally characterised examples of *d*-metal compounds with the chiral ligands $[L^{iPr}]^$ and $[L^{CH2Ph}]^-$ and, in order to establish the crystal structures of these complexes, single-crystal X-ray diffraction studies were carried out with complexes **2**, **4**, **6** and **8**. Additionally, complex **5** ($[L^{Me}]^-$ ligand) was also structurally characterised by single-crystal X-ray methods. Characterization by X-ray powder diffraction, XRPD, was performed with complexes **1**, **3** and **7**.



Scheme 2 Synthesis of zinc (1-4) and copper (5-8) complexes

Structural characterization of zinc complexes

Compounds 2 and 4 crystallise in the $P2_12_12_1$ space group, orthorhombic, and in the $P3_2$ space group, trigonal, respectively, and both are two-dimensional coordination polymers in which the zinc ions are interconnected by bridging $[L^R]^-$ (R = ⁱPr, 2; CH₂Ph, 4) anions through the carboxylate groups. Selected structural parameters of 2 and 4 are collected in Table 1. Both carboxylate groups are κ^1 -coordinated and this monodentate bonding mode is consequent with the asymmetry observed between the two C-O distances of the carboxylate

moieties (compare, for instance in 2, the observed ranges of 1.18-1.24 and 1.26-1.29 Å for the C=O and C-O bonds, respectively). This μ - κ^{1} , κ^{1} -coordination mode was also observed in the related zinc derivatives with the non-chiral $[L^{H}]^{-}$ and chiral $[L^{Me}]^{-}$ anions.7^{a,c,16} The coordination geometry of zinc is tetrahedral in 2 and 4, with values for the four-coordinate geometry index³⁰ τ_4 of 0.9 and 0.8, respectively. The Zn-O distances found in the four κ^{1} -carboxylate moieties are within the range 1.93-1.99 Å, in agreement with the experimental lengths observed for other zinc carboxylates (Cambridge Structural Database search³¹). The asymmetric unit of 2 (Fig. 1a) consists of two zinc ions and four [L^{iPr}]⁻ anions, in agreement with the composition Zn:2L^{iPr}, analytically determined. The relative orientation of the 3methylbutanoate arms with respect the imidazolium ring is dissimilar within the asymmetric unit. Only one L^{iPr} ligand shows a roughly *cis* disposition of the carboxylate groups (torsion angle C_{carboxy}-C_{chiral}-C'_{chiral}-C'_{carboxy} of *ca*. 50°). The three remaining [L^{iPr}]⁻ ligands display these torsion angles close to ca. 85°. This feature is responsible of the construction of the twodimensional polymer (Fig. 2b). This 2D distribution found in 2 differs from that observed in the structure of the related complex $[Zn(L^{Me})_2]_n$,¹⁶ in which there are two relative orientation of the dicarboxylate arms within the asymmetric unit with C_{carboxy}-C_{chiral}-C'_{chiral}-C'_{carboxy} torsion angles around 70° and 175° (trans disposition). The asymmetric unit of 4 (Fig. 2a) consists of one zinc ion and two [L^{CH2Ph}]⁻ anions and the relative orientation of the 3phenylpropanoate arms with respect the imidazolium ring is roughly *cis* with torsion angles C_{carboxy}-C_{chiral}-C'_{chiral}-C'_{carboxy} close to 20°. The resulting 2D coordination polymer is slightly different to that of complex 2 (Fig. 2b). In both complexes, 2 and 4, the 3D crystal distribution was produced by the packing of the 2D lamellar sheets due to non-covalent interactions through the alkyl hydrophobic part of the $[L^R]^-$ anions (see, for example, Fig. S1 for 2 in the ESI).

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.



Fig. 1 (a) Asymmetric unit of compound 2, hydrogen atoms were omitted for clarity. (b) Lamellar sheet in 2 viewed along c axis with tetrahedral polyhedron for zinc atoms.

Distances (A)			2			4	
Zn-O	Zn(1)-O(1)	1.945(7)	Zn(2)-O(9)	1.992(7)	Zn(1)-O(5)	1.934(4)	
	Zn(1)-O(3)	1.966(7)	Zn(2)-O(11)	1.962(7)	Zn(1)-O(7)	1.939(4)	
	Zn(1)-O(5)	1.967(7)	Zn(2)-O(13)	1.935(7)	Zn(1)-O(3)	1.958(4)	
	Zn(1)-O(7)	1.987(7)	Zn(2)-O(15)	1.938(7)	Zn(1)-O(1)	1.963(4)	
C-0	O(1)-C(12)	1.284(11)	O(9)-C(24)	1.265(13)	O(1)-C(1)	1.250(6)	
	O(2)-C(12)	1.233(12)	O(10)-C(24)	1.242(12)	O(2)-C(1)	1.227(6)	
	O(3)-C(13)	1.289(14)	O(11)-C(44)	1.272(13)	O(3)-C(21)	1.240(6)	
	O(4)-C(13)	1.200(14)	O(12)-C(44)	1.233(14)	O(4)-C(21)	1.217(7)	
	O(5)-C(14)	1.253(14)	O(13)-C(48)	1.283(13)	O(5)-C(30)	1.272(7)	
	O(6)-C(14)	1.229(15)	O(14)-C(48)	1.232(13)	O(6)-C(30)	1.219(7)	
	O(7)-C(19)	1.288(13)	O(15)-C(32)	1.293(12)	O(7)-C(42)	1.277(7)	
	O(8)-C(19)	1.214(13)	O(16)-C(32)	1.219(13)	O(8)-C(42)	1.219(7)	
Angles (°)							
O-Zn-O	O(1)-Zn(1)-O(3)	119.2(3)	O(15)-Zn(2)-O(13)	119.2(3)	O(5)-Zn(1)-O(7)	108.37(17)	
	O(1)-Zn(1)-O(5)	108.7(3)	O(15)-Zn(2)-O(11)	111.0(3)	O(5)-Zn(1)-O(3)	107.1(2)	
	O(3)-Zn(1)-O(5)	115.7(3)	O(13)-Zn(2)-O(11)	110.8(3)	O(7)-Zn(1)-O(3)	121.9(2)	
	O(1)-Zn(1)-O(7)	96.9(3)	O(15)-Zn(2)-O(9)	97.4(3)	O(5)-Zn(1)-O(1)	121.7(2)	
	O(3)-Zn(1)-O(7)	111.3(3)	O(13)-Zn(2)-O(9)	105.5(3)	O(7)-Zn(1)-O(1)	108.46(19)	
	O(5)-Zn(1)-O(7)	102.1(3)	O(11)-Zn(2)-O(9)	112.0(3)	O(3)-Zn(1)-O(1)	89.23(15)	
C-O-Zn	C(12)-O(1)-Zn(1)	118.7(6)	C(24)-O(9)-Zn(2)	120.7(6)	C(1)-O(1)-Zn(1)	140.8(4)	
	C(13)-O(3)-Zn(1)	114.1(7)	C(44)-O(11)-Zn(2)	122.8(7)	C(21)-O(3)-Zn(1)	143.3(4)	
	C(14)-O(5)-Zn(1)	121.5(7)	C(48)-O(13)-Zn(2)	115.8(7)	C(30)-O(5)-Zn(1)	116.2(4)	
	C(19)-O(7)-Zn(1)	114.4(7)	C(32)-O(15)-Zn(2)	128.8(7)	C(42)-O(7)-Zn(1)	115.4(4)	
0-C-0	O(2)-C(12)-O(1)	126.9(10)	O(10)-C(24)-O(9)	127.8(10)	O(2)-C(1)-O(1)	123.4(4)	
	O(2)-C(12)-C(8)	118.8(8)	O(16)-C(32)-O(15)	126.9(9)	O(4)-C(21)-O(3)	123.2(5)	
	O(4)-C(13)-O(3)	126.9(10)	O(12)-C(44)-O(11)	126.6(11)	O(6)-C(30)-O(5)	126.6(5)	
	O(6)-C(14)-O(5)	127.1(11)	O(14)-C(48)-O(13)	126.7(9)	O(8)-C(42)-O(7)	126.4(6)	

 Table 1 Selected structural parameters for zinc complexes 2 and 4.

Symmetry transformations used to generate equivalent atoms: complex **2**: #1 x-1/2, -y+3/2, -z+1. #2 x-1/2, -y+1/2, -z+1. #3 x, y-1, z. #4 x+1/2, -y+1/2, -z+1. #5 x, y+1, z. #6 x+1/2, -y+3/2, -z+1. Complex **4**: #1 x-1, y, z; #2 x-1, y-1, z; #3 x+1, y, z; #4 x+1, y+1, z.



Fig. 2 (a) Asymmetric unit of compound 4, hydrogen atoms and solvent crystallization molecules were omitted for clarity. (b) 2D sheet in 4 viewed along c axis.

Concerning complexes 1 and 3, they were characterised by XRPD methods. The experimental X-ray diffractogram of 1 shows no significant differences with respect to the simulated powder diffraction pattern calculated from the single-crystal data previously reported (refcode VITHUD¹⁶) and no requires further comments. Rietveld refinement from the XRPD of complex 3 was performed and its final Rietveld plot is given in Fig. S2. From these results, we deduced that the overall structure of complex 3 is completely similar to that of its enantiomer 2 (both showed identical XRPDs, Fig. S3), which was previously solved by single crystal X-ray methods.

Structural characterization of copper complexes

Complexes **5** and **6** crystallise in the P_1 space group, triclinic, while **8** crystallises in the $C222_1$ space group, orthorhombic. All are two-dimensional coordination polymers in which the copper ions are bridged by monodentate carboxylate groups of L^R anions. Selected structural parameters of **5**, **6** and **8** are collected in Table 2. The environment of copper in the three compounds is square planar with values lower than 0.1 for the four-coordinate geometry index τ_4 . In these compounds, the Cu-O bonding distances are within the range 1.92-2.00 Å, in agreement with the experimental lengths observed for other copper carboxylates.³¹ The distances between copper and the second oxygen atom of the carboxylate group cover the range 2.58-2.76 Å. Although these distances are slightly lower than the sum of the van der Waals radii, they were essentially considered as non-bonding distances (mean value for the Cu-O bond length in Cu(κ^1 -OC(O)-R complexes is 2.01(2) Å³¹). The asymmetric unit of complex **5** shows two non-symmetry related copper atoms, Cu1 and Cu2, that originate two different Cu-L^{Me} polymers with C_{carboxy}-C_{chiral}-C' _{carboxy} torsion angles of *ca*. 55 and 41°,

more in agreement with a roughly *cis* disposition of the CHMeCOO groups with respect the imidazolium ring. Fig. 3a shows the common $\mu^2 \cdot \kappa^1 \cdot O^1, \kappa^1 \cdot O^3$ coordination mode found for the bridging $[L^{Me}]^-$ anion in both Cu1 and Cu2 atoms, while the resulting 2D disposition along *c* axis is depicted in the Fig. 3b. This 2D structure is formed by the accommodation of the two different Cu-L^{Me} polymers of Cu1 and Cu2 atoms, which are illustrated in Fig. 3c. These two different Cu-L^{Me} polymers are connected by weak C–H···O non-classical hydrogen bonds between both the C_{chiral}-H and C²_{imidazol}-H bonds of the bridging $[L^{Me}]^-$ fragments, belonging to one polymer, and the oxygen atoms of the carboxylate moiety of the $[L^{Me}]^-$ ligands corresponding to the adjacent polymer.

The relative orientation of the 3-methylbutanoate arms with respect the imidazolium ring is approximately *trans* in complex **6**, with torsion angles $C_{carboxy}-C_{chiral}-C'_{chiral}-C'_{carboxy}$ of *ca.* 97 and 117°. In contrast with the related zinc complex **2**, two different $\mu^2 - \kappa^1 - O^1, \kappa^1 - O^3$ coordination modes were found for bridged $[L^{iPr}]^-$ anions, corresponding to monodentate κ^1 -*syn*, κ^1 -*syn* and κ^1 -*anti*, κ^1 -*anti* coordination (Fig. 4a). This variation produces a slightly different 2D sheet of **6** in comparison to **2** (Fig. 4b). In **6**, the 3D packing occurred by non-covalent interactions between the ⁱPr groups of the sheets (Fig. S4). Additionally, the supramolecular arrangement is reinforced by the presence of weak C–H…O non-classical hydrogen bonds between the C_{chiral}-H fragments belonging to one 2D sheet and carbonyl groups of the carboxylate moiety of the neighbour sheet. Finally, the general $\mu^2 - \kappa^1 - O^1, \kappa^1 - O^3$ coordination mode is also present in the complex **8**. Fig. 5a showed the crystal packing of complex **8** along *c* axis. The C_{carboxy}-C_{chiral}-C'_{chiral}-C'_{carboxy} torsion angles are close to *ca.* 40°, in agreement with a roughly *cis* disposition of the CH(CH₂Ph)COO groups with respect the imidazolium ring, as occurred in the related zinc complex **4**. This disposition generates two-dimensional coordination polymers, which are arranged parallel to the *b* axis (see Fig. 5b).

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.





Fig. 3 (a) Common $\mu^2 - \kappa^1 - O^1, \kappa^1 - O^3$ coordination mode of $[L^{Me}]^-$ ligand in **5** for the two nonsymmetry related copper atoms, Cu1 and Cu2. (b) Lamellar sheet in **5** viewed along *c* axis. (c) Crystal packing of **5** (left) and separation of the two different Cu-L^{Me} polymers for Cu1 and Cu2 atoms (right).

Dalton Transactions Accepted Manuscript



Fig. 4 (a) Two different $\mu^2 - \kappa^1 - O^1, \kappa^1 - O^3$ coordination modes of $[L^{iPr}]^-$ ligand in **6**. (b) Lamellar sheet in **6** viewed along *c* axis.



Fig. 5 (a) Crystal packing of complex 8, viewed along c axis, within the unit cell. (b) 3D arrangement of the 2D lamellar sheets, viewed along b axis.

$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	28	Dalton Transactions							
Distances (Å) 5 6 8 Cu-O Cu(1)-O(1) 1.9525(18) Cu(1)-O(1) 1.949(7) Cu(1)-O(1) 1.941(4) Cu(2)-O(7) Cu(2)-O(7) 1.963(7) Cu(1)-O(1) 1.949(7) Cu(1)-O(1) 1.941(4) Cu(2)-O(7) Cu(2)-O(7) 2.0073(14) Cu(1)-O(7) 1.924(7) Cu(1)-O(1) 1.941(4) Cu(2)-O(7) C-O O(1)-C(5) 1.267(3) O(1)-C(9) 1.294(7) Cu(1)-O(1) 1.935(4) Cu(2)-O(7) O(3)-C(9) 1.283(2) O(3)-C(20) 1.229(12) O(1)-C(18) 1.161(15) O(8)-C(34) O(4)-C(9) 1.227(3) O(4)-C(20) 1.241(12) O(4)-C(13) 1.225(7) O(10)-C(43) O(6)-C(14) 1.228(3) O(5)-C(25) 1.231(12) O(6)-C(22) 1.229(7) C(1A)-O(2A) O(7)-C(16) 1.275(2) O(7)-C(26) 1.279(13) O(7)-C(34) 1.272(7) O(1B)-O(2B) O(3)-C(1)-O(1)#1 84.56(13) O(7)-Cu(1)-O(1) 89.4(3) O(1)-Cu(1)-O(1)#1 90.3(2) O(7)#4-Cu(2)-O	Table 2 Salas	stad atmustural param	ators for con	nor complexes (6 and 9				
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	Distances (Å)	5		6		8			
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	Cu-O	Cu(1)-O(1)	1.9525(18)	Cu(1)-O(1)	1.949(7)	Cu(1)-O(1)	1.941(4)	Cu(1)-O(5)	1.95
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		Cu(1)-O(3)	1.9736(16)	Cu(1)-O(3)	1.948(7)	Cu(1)-O(9)	1.941(4)	Cu(2)-O(7)	1.94
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		Cu(2)-O(5)	1.9602(14)	Cu(1)-O(5)	1.963(7)	Cu(1)-O(11)#1	1.959(4)	Cu(2)-O(3)#3	1.95
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		Cu(2)-O(7)	2.0073(14)	Cu(1)-O(7)	1.924(7)				
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	C-0	O(1)-C(5) 1.	267(3)	O(1)-C(9)	1.296(12)	O(1)-C(1B)	1.161(15)	O(8)-C(34)	1.21
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $		O(2)-C(5) 1.	218(3)	O(2)-C(9)	1.229(12)	O(1)-C(1A)	1.303(13)	O(9)-C(43)	1.23
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $		O(3)-C(9) 1.	283(2)	O(3)-C(20)	1.261(12)	O(3)-C(13)	1.285(7)	O(10)-C(43)	1.24
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $		O(4)-C(9) 1.	227(3)	O(4)-C(20)	1.243(12)	O(4)-C(13)	1.232(7)	O(11)-C(55)	1.2
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		O(5)-C(14) 1.	268(3)	O(5)-C(25)	1.265(12)	O(5)-C(22)	1.280(6)	O(12)-C(55)	1.23
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		O(6)-C(14) 1.	225(3)	O(6)-C(25)	1.231(12)	O(6)-C(22)	1.229(7)	C(1A)-O(2A)	1.22
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		O(7)-C(16) 1.	275(2)	O(7)-C(26)	1.279(13)	O(7)-C(34)	1.272(7)	C(1B)-O(2B)	1.23
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $		O(8)-C(16) 1.	229(2)	O(8)-C(26)	1.239(12)				
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	Angles (°)								
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	O-Cu-O	O(1)-Cu(1)-O(1)#1	84.56(13)	O(7)-Cu(1)-O(1)) 89.4(3)	O(1)-Cu(1)-O(9)	92.45(19)	O(7)#4-Cu(2)-O(7)	89.
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		O(1)-Cu(1)-O(3)	92.67(8)	O(7)-Cu(1)-O(3)	3) 176.3(4)	O(1)-Cu(1)-O(11)#1	90.3(2)	O(7)#4-Cu(2)-O(3)#3	91.2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		O(3)#1-Cu(1)-O(3)	90.61(10)	O(1)-Cu(1)-O(3)	3) 90.3(3)	O(9)-Cu(1)-O(11)#1	177.1(2)	O(7)-Cu(2)-O(3)#3	179
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		O(5)-Cu(2)-O(5)#3	92.79(10)	O(7)-Cu(1)-O(5	5) 90.4(3)	O(1)-Cu(1)-O(5)	176.6(2)	O(7)- $Cu(2)$ - $O(3)$ #2	91.2
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$		O(5)-Cu(2)-O(7)#3	173.99(7)	O(1)-Cu(1)-O(3)	173.9(3)	O(9)-Cu(1)-O(5)	90.9(2)	O(3)#2-Cu(2)-O(3)#3	88.4
$\begin{array}{c c c c c c c c c c c c c c c c c c c $		O(5)-Cu(2)-O(7)	91.79(6)	O(3)-Cu(1)-O(3)	b) 90.3(3)	O(11)#1-Cu(1)-O(5)	86.27(15)		
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0.0.0	O(7)#3-Cu(2)-O(7)	83.92(8)		1017(()	$C(1\mathbf{P}) O(1) C(1)$	102 4(0)	O(24) O(7) O(0)	115
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	C-O-Cu	C(5)-O(1)-Cu(1)	123.03(15)	C(9)-O(1)-Cu(1)	1) 121.7(6)	C(1B)-O(1)-Cu(1)	123.4(8)	C(34)-O(7)-Cu(2)	115
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		C(9)-O(3)-Cu(1) C(14) O(5) Cu(2)	102.37(13) 114.72(12)	C(20)-O(3)-Cu(3)	$\begin{array}{cccc} 1 & 100.7(0) \\ (1) & 108.7(5) \end{array}$	C(1A)-O(1)-Cu(1) C(12) O(2) Cu(2)#5	11/.1(0) 108.2(4)	C(43)-O(9)-Cu(1) C(55) O(11) Cu(1)#2	118
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		C(14)-O(3)-Cu(2) C(16) O(7) Cu(2)	114.73(13) 102.05(13)	C(25)-O(5)-Cull	(1) 100.7(3) (1) 125.1(5)	C(13)-O(3)-Cu(2)#3 C(22) O(5) Cu(1)	100.3(4) 107.3(3)	C(33)-O(11)-Cu(1)#2	105
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0.0.0	O(2) C(5) O(1)	102.93(13) 107.7(2)	O(2) C(0) O(1)	(1) 123.1(3) 124.6(7)	O(2A) O(3)-O(1)	107.3(3) 124.6(10)	O(8) C(24) O(7)	126
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	0-0-0	O(2)-O(3)-O(1) O(4) C(0) O(2)	121.1(2) 124.6(2)	O(2) - O(3) - O(1)	124.0(7)	O(2A)-O(1A)-O(1)	124.0(10) 120.2(12)	O(0) - C(34) - O(7) O(0) - C(42) - O(10)	120
0(0) - 0(17) - 0(0) - 120 - 10(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) - 0(0) -		O(4) - O(3) - O(3) O(6) - O(14) - O(5)	124.0(2) 126.1(2)	O(4) - O(20) - O(3)	5) 123.2(0) 5) 126.0(8)	O(1)-O(1D)-O(2D) O(4)-O(13)-O(3)	120.3(13) 124.4(5)	O(3) - O(43) - O(10) O(12) - C(55) - O(11)	124
(1,1,2,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1		O(8)-C(14)-O(3)	120.1(2) 125 1(2)	O(8)-C(25)-O(2)	(120.0(0)) (123.0(0))	O(4)-O(3)	124.4(3) 124.9(5)	O(12) - O(33) - O(11)	120

 Table 2 Selected structural parameters for copper complexes 5, 6 and 8.

Symmetry transformations used to generate equivalent atoms: complex 5: #1 -x+2,-y,z. #2 x-1/2,-y+1/2,-z. #3 -x+1,-y,z. #4 -x+3/2,y+1/2,-z+1. #5 x+1/2,-y+1/2,-z. #6 -x+3/2,y-1/2,-z+1. Complex 6: #1 x, y+1, z. #2 x, y, z+1. #3 x, y-1, z. #4 x, y, z-1. Complex 8: #1 x+1/2,-y+1/2,-z+1. #2 x-1/2,-y+1/2,-z+1. #3 x-1/2,y-1/2,z. #4 x,-y,-z+1. #5 x+1/2,y+1/2,z.

Complex 7 was characterised by XRPD methods and showed identical X-ray diffractogram that its enantiomer 6 (Fig. S5). Rietveld refinement was performed and its final Rietveld plot is given in Fig. S6. From these results, we deduced that the overall structure of 7 is completely similar to that of its enantiomer 6, solved by single crystal X-ray methods, and no further discussion is required.

TGA study of complexes 2-8

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

With the aim of study the decomposition pathway of these coordination polymers, thermogravimetric analyses were carried out with complexes 2-8 (Fig. 6). TGA analysis of compound **1** was previously reported.¹⁶ As expected, the enantiomerically related zinc compounds 2 and 3, $[Zn(L^{iPr})_2 \cdot H_2O]_n$, showed identical TGA profiles in which, after the crystallization water loss (ca. 5 % weight loss), the resulting $[Zn(L^{iPr})_2]_n$ material was thermally stable up to ca. 280 °C. Then, a simultaneous consecutive decarboxylation and imidazolium decomposition with complete loss of [L^{iPr}]⁻ ligand was observed in the 280-500 °C range (ca. 85 % weight loss). Complex 4, $[Zn((S,S)-L^{CH2Ph})_2 \cdot 2DMF \cdot H_2O]_n$, showed a similar TGA profile. A weight loss of about 18 % was observed in the initial stage, at ca. 150 °C, attributable to the loss of the crystallization DMF and water molecules. Subsequently, the elimination of the [L^{CH2Ph}]⁻ ligand by decarboxylation and imidazolium decomposition was completed at 500 °C. There was no weight loss for 2-4 after 550 °C, and the final product was attributed to the corresponding zinc oxide. The thermogravimetric behavior of compounds 2-4 is closely connected with that previously reported for the parent HL^R compounds. In fact, controlled hydrothermal decarboxylation of these HL^R compounds was proposed as new approach toward the synthesis of new ionic liquids.3 The TGA analysis of copper compounds 5-8 (Fig. 6, bottom) gave analogous results than those discussed for Zn derivatives. For example, the TGA profile of compound 5, $[Cu(L^{Me})_2]_n$, was very alike to that reported for complex $[Zn(L^{Me})_2]_n$, 1.¹⁶ Complexes 6 and 7, $[Cu(L^{iPr})_2]_n$, were stable until 250 °C with a decomposition pathway completely similar to that found for their zinc analogues 2 and 3. After a weight loss of about 85 % (550 °C), the resulting material was the corresponding oxide with no additional weigh loss after 600 °C. In particular, for the copper complex 6 the formation of the copper oxide, CuO, after heating at 600 °C, was confirmed by PXRD methods (Fig. S7, ESI). Finally, for compound $\mathbf{8}$ a stepwise decarboxylation and imidazolium decomposition can be discerned with an inflection point at around 320 °C.



Fig. 6 Thermogravimetric analyses for zinc 2-4 (upper) and copper complexes 5-8 (bottom).

DFT calculations of [L^R]⁻ anions

In order to gain further information about the coordination features of the $[L^R]^-$ anions and their conformational flexibility, DFT calculations³² were carried out for these anions with R = Me, ⁱPr and CH₂Ph. The discussion was centred in the *S*,*S* isomers. Three conformers were located for each $[(S,S)-L^R]^-$ anion and the corresponding structures show different orientations of the 1,3-substituted arms with respect the imidazolium ring. Fig. 7 depicted the optimised structures of the three conformers located for the $[(S,S)-L^{iPr}]^-$ anion and those for other $[(S,S)-L^R]^-$ anions (R = Me and CH₂Ph) were collected at the ESI (Fig. S8). The maximum energy

Dalton Transactions

difference, measured as relative Gibbs free energy, between these conformers was computed to be < 1 kcal·mol⁻¹ and this fact was in agreement with the expected free rotation along the C_{chiral} -N vector and with the miscellaneous orientations of these arms found in the structures of compounds **2**, **4-6**, **8**. These theoretical results confirm the conformational flexibility as ditopic linker of the $[L^R]^-$ ligands. Their coordination capabilities were analyzed and, for instance, the HOMO to HOMO-3 orbitals, for the most stable conformer of $[L^{iPr}]^-$ anion, display principally the *in-phase* and *out-of-phase* combinations of σ O lone pairs (Fig. S9).



Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

Fig. 7 Optimised structures of the three conformers located for the $[(S,S)-L^{iPr}]^{-}$ anion: frontal (up) and lateral views (bottom).

Preliminary studies in asymmetric catalysis: synthesis of chiral α-aminophosphonates

Chiral α -aminophosphonates, considered as phosphorous analogues of α -amino acids, are of special interest due to their broad range of biological activities³³ and their promising use in medicinal chemistry and the pharmaceutical sciences.³⁴ Among other approaches,³⁵ Kabachnik-Fields reaction represent the most efficient methods for the synthesis of α -aminophosphonates, which is a one-pot, three-component procedure using carbonyl compound, amine and dialkyl phosphonate,³⁶ generally promoted by acid catalyst.^{34b,37} Despite the importance of the Kabachnik-Fields reaction, enantioselective versions for the preparation of enantiomerically enriched α -aminophosphonates are scarce.³⁸ Only a few examples has been reported, including both organocatalysis³⁹ and metal catalysis.⁴⁰ It is worth mentioning that in 2011 Ohara *et al.* reported an efficient method that allows the synthesis of enantioenriched α -aminophosphonates using a zinc(II) complex of 1,2-bis(imidazolin-2-ly)pyridine.⁴¹ Recently, we have explored the use of imidazolium-based zwitterionic dicarboxylic ligands, [L^R]⁻, as chiral inductors for the catalytic asymmetric oxidation of

sulfides with hydrogen peroxide using oxoperoxo-Mo(VI) catalysts.⁴² On these bases, we decided to explore the use of Zn(II) homochiral coordination polymers **1** and **2** as catalysts in order to produce enantiomerically enriched α -aminophosphonates. The synthetic efficiency of the Zn-catalyst in the Kabachnik–Fields reaction was evaluated in the one-pot, three-component procedure using 4-methoxybenzaldehyde, 4-methoxyaniline and diethyl phosphonate as model reaction (Scheme 3).



Scheme 3 Model reaction for the synthesis of chiral α -aminophosphonates

Table 3 reports selected results from the reactions carried out in a reactor with a 1:1:1:0.1 ratio of aldehyde:aniline:phosphonate:catalyst. Several solvents were tested at different reaction temperature in order to evaluate its effects on the enantioselectivity of the process. The best conversion and selectivity to the α -aminophosphonate was obtained in toluene at 80 °C, however, unfortunately, no enantioselectivity was observed in these reactions. Better enantioselectivities were obtained in MeOH, though the ees were very low, *ca.* 1-5%, and with poor selectitivities to the α -aminophosphonate. Particularly, in MeOH at 60 °C and employing catalyst **1** (entry 5, Table 3), afforded 14% of the α -aminophosphonate with a 4.5% of enantiomeric excess. The fast on/off exchange of the ligand from the coordination sphere of the Zn(II) is perhaps the key reason why only poor enantioselectivities are obtained using these complexes as catalysts for Kabachnik-Fields reaction.

Entry	Catalyst	Solvent	T (°C)	t (h)	Conversion ^b	Selectivity ^b	%ee ^c
1	1	Toluene	80	18	86	93	-0.2
2	2	Toluene	80	18	90	59	-0.2
3	1	CHCl ₃	50	18	95	43	+0.3
4	2	CHCl ₃	50	18	80	14	+1
$5^{\rm e}$	1 ^d	MeOH	60	18	99	53	+2.6
6 ^e	1	MeOH	30	42	98	14	+4.5
7 ^e	1^{d}	MeOH	30	42	92	19	+3.3

Table 3 Synthesis of α -aminophosphonates using zinc(II) coordination polymers 1 and 2 as catalysts.^a

^a Reaction conditions: catalyst 0.05 mmol, 4-methoxybenzaldehyde 0.5 mmol, 4-methoxyaniline 0.5 mmol, diethyl phosphonate 0.5 mmol, solvent 1.0 mL. ^b Determined by ¹H NMR analysis. ^c Determined by HPLC. ^d Catalyst prepared *in-situ* by mixing Zn(AcO)₂ (0.10 mmol) and (*S,S*)-HL^{Me} (0.20 mmol). ^e Solvent 2.0 mL.

Conclusions

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

Two-dimensional homochiral coordination polymers of general formulation $[M(L^R)_2]_n$ were obtained by reaction between zinc acetate or copper acetate with enantiomerically pure HL^R compounds and they were characterised and structurally identified by X-ray diffraction methods (single crystal and powder). In all the zinc, **1**-**4**, and the copper, **5**-**8**, complexes the metals were bridged by the chiral imidazolium-based dicarboxylate ligands, $[L^R]^-$, through the two carboxylate groups. The observed general $\mu^2 - \kappa^1 - O^1, \kappa^1 - O^3$ coordination mode afforded tetrahedral metal coordination environment for zinc and square planar one for copper complexes. DFT calculations on the $[L^R]^-$ anions confirmed the conformational flexibility of the carboxylate arms linked to the imidazolium moiety and this fact makes these $[L^R]^-$ ligands good ditopic linkers for the formation of different coordination polymers with four-coordinated metal centres. Although the catalytic performance of Zn complexes **1** and **2** in the synthesis of chiral α -aminophosphonates is very poor, further studies are in progress.

Acknowledgements

Financial support from the Junta de Andalucía (Proyecto de Excelencia FQM-7079) is gratefully acknowledged. R. P. C. acknowledges to the Junta de Andalucía the postdoctoral contract associated to FQM-7079 project. We thank to the Centro de Servicios de Informática y Redes de Comunicaciones (CSIRC), Universidad de Granada, for providing the computing time.

References

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:12.

- 1 Z. Fei, D. Zhao, T. J. Geldbach, R. Scopelliti and P. J. Dyson, *Chem.–Eur. J.*, 2004, **10**, 4886-4893.
- 2 (a) H. Debus, Justus Liebigs Ann. Chem. 1858, 107, 199-208. (b) B. Radzisewski, Ber. Dtsch. Chem. Ges. 1882, 15, 2706-2708.
- 3 S. Kirchhecker, M. Antonietti and D. Esposito, Green Chem., 2014, 16, 3705-3709.
- 4 J. Sun, L. Han, W. Cheng, J. Wang, X. Zhang and S. Zhang, *ChemSusChem*, 2011, 4, 502-507.
- 5 S. J. Yang, R. Rothe, S. Kirchhecker, D. Esposito, M. Antonietti and N. Fechler, *Carbon* 2015, **94**, 641-645.
- 6 Selected examples are: (a) V. Gandin, M. Pellei, M. Marinelli, C. Marzano, A. Dolmella, M. Giorgetti and C. Santini, *J. Inorg. Biochem.*, 2013, 129, 135-144. (b) W. A. Herrmann, L. J. Goossen and M. Spiegler, *J. Organomet. Chem.*, 1997, 547, 357-366. (c) W.-H. Yang, C.-S. Lee, S. Pal, Y.-N. Chen, W.-S. Hwang, I. J. B. Lin and J.-C. Wang, *J. Organomet. Chem.*, 2008, 693, 3729-3740. (d) M. Pellei, V. Gandin, M. Marinelli, C. Marzano, M. Yousufuddin, H. V. R. Dias and C. Santini, *Inorg. Chem.*, 2012, 51, 9873-9882.
- Selected examples are: (a) Z. Fei, W. H. Ang, T. J. Geldbach, R. Scopelliti and P. J. Dyson, *Chem. Eur. J.*, 2006, 12, 4014-4020. (b) Z. Fei, T. J. Geldbach, D. Zhao, R. Scopelliti and P. J. Dyson, *Inorg. Chem.*, 2005, 44, 5200-5202. (c) Z. Fei, D. Zhao, T. J. Geldbach, R. Scopelliti, P. J. Dyson, S. Antonijevic and G. Bodenhausen, *Angew. Chem. Int. Ed.*, 2005, 44, 5720-5725.
- See for example: (a) P. Suresh, C. N. Babu, N. Sampath and G. Prabusankar, *Dalton Trans.*, 2015, 44, 7338-7346. (b) P. Suresh, C. N. Babu and G. Prabusankar, *Polyhedron*, 2015, 89, 322-329. (c) P. Suresh, S. Radhakrishnan, C. N. Babu, A. Sathyanarayana, N. Sampath and G. Prabusankar, *Dalton Trans.*, 2013, 42, 10838-10846. (d) B. F. Abrahams, H. E. Maynard-Casely, R. Robson and K. F. White, *CrystEngComm*, 2013, 15, 9729-9737.
- 9 Z. Fei, T. J. Geldbach, R. Scopelliti and P. J. Dyson, Inorg. Chem., 2006, 45, 6331-6337.
- Selected examples are: (a) L. Han, S. Zhang, Y. Wang, X. Yan and X. Lu, *Inorg. Chem.*, 2009, 48, 786-788. (b) X. W. Wang, L. Han, T. J. Cai, Y. Q. Zheng, J. Z. Chen and Q. Deng, *Cryst. Growth Des.*, 2007, 7, 1027-1030. (c) X.-C. Chai, Y.-Q. Sun, R. Lei, Y.-P. Chen, S. Zhang, Y.-N. Cao and H.-H. Zhang, *Cryst. Growth Des.*, 2010, 10, 658-668.

12.

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:

- 11 D. Esposito, S. Kirchhecker and M. Antonietti, Chem. Eur. J., 2013, 19, 15097-15100.
- 12 O. Kühl and G. Palm, Tetrahedron: Asymmetry, 2010, 21, 393-397.
- 13 O. Kühl, S. Millinghaus and P. Wehage, Cent. Eur. J. Chem., 2010, 8, 1223-1226.
- 14 (a) A. Ferry, K. Schaepe, P. Tegeder, C. Richter, K. M. Chepiga, B. J. Ravoo and F. Glorius, *ACS Catal.*, 2015, 5, 5414-5420. (b) M. A. Reynoso-Esparza, I. I. Rangel-Salas, A. A. Peregrina-Lucano, J. G. Alvarado-Rodríguez, F. A. López-Dellamary-Toral, R. Manríquez-González, M. L. Espinosa-Macías and S. A. Cortes-Llamas, *Polyhedron*, 2014, 81, 564-571.
- 15 X. Wang, X.-B. Li, R.-H. Yan, Y.-Q. Wang and E.-Q. Gao, *Dalton Trans.*, 2013, **42**, 10000-10010.
- 16 C. N. Babu, A. Sathyanarayana, S. M. Mobin and G. Prabusankar, *Inorg. Chem. Commun.*, 2013, 37, 222-224.
- 17 L. Ma, C. Abney and W. Lin, Chem. Soc. Rev., 2009, 38, 1248-1256.
- 18 M. Yoon, R. Srirambalaji and K. Kim, Chem. Rev., 2012, 112, 1196-1231.
- 19 Bruker. APEX2. Bruker AXS Inc., Madison, Wisconsin, USA, 2007.
- 20 Bruker Advanced X-ray Solutions, Bruker AXS Inc., Madison, Wisconsin, USA, 2001.
- C. M. Burla, M. Camalli, B. Carrozzini, G. L. Cascarano, C. Giacovazzo, G. Polipori and R. Spagna, SIR2002: the program, *J. Appl. Cryst.*, 2003, 36, 1103.
- 22 G. M. Sheldrick, Acta Cryst., 2015, C71, 3-8.
- 23 (a) A. L. Spek, J. Appl. Crystallogr., 2003, 36, 7-13. (b) A. L. Spek, Acta Cryst. 2009, D65, 148-155. (c) A. L. Spek, Acta Cryst. 2015, C71, 9-18.
- 24 A. Boultif and D. Louer, J. Appl. Cryst., 2004, 37, 724-731.
- 25 H. M. Rietveld, J. Appl. Cryst., 1969, 2, 65–71.
- 26 A. C. Larson and R. B. von Dreele, General Structure Analysis System (GSAS), Los Alamos National Laboratory Report LAUR 86-748, 2004.
- 27 B. H. Toby, J. Appl. Cryst., 2001, 34, 210-213.
- 28 (a) A. D. Becke, J. Chem. Phys., 1993, 98, 5648-5652. (b) C. Lee, W. Yang and R. G. Parr, Phys. Rev. B, 1988, 37, 785-789.
- 29 Gaussian 09, Revision E.01, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta,

F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R.
Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J.
Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken,
C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R.
Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G.
A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B.
Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2009.

30 L. Yang, D. R. Powell and R. P. Houser, Dalton Trans., 2007, 955-964.

12

Published on 01 December 2016. Downloaded by University of Regina on 07/12/2016 12:05:

- Cambridge Structural Database System, Cambridge Crystallographic data Centre, 12
 Union Road, Cambridge, CB2 1EZ, UK. C. R. Groom, I. J. Bruno, M. P. Lightfoot and S.
 C. Ward, *Acta Crystallogr.*, 2016, **B72**, 171–179.
- 32 R. G. Parr and W. Yang, *Density Functional Theory of Atoms and Molecules*, Oxford University Press, New York, 1989.
- 33 (a) P. P. Giannousis and P. A. Bartlett, *J. Med. Chem.*, 1987, **30**, 1603–1609. (b) J.
 Hiratake and J. Oda, *Biosci. Biotech. Biochem.*, 1997, **61**, 211–218. (c) E. D. Naydenova,
 P. T. Todorov and K. D. Troev, *Amino Acids*, 2010, **38**, 23–30.
- 34 (a) A. K. Bhattacharya, D. S. Raut, K. C. Rana, I. K. Polanki, M. S. Khan and S. Iram, *Eur. J. Med. Chem.*, 2013, 66, 146–152. (b) S. A. R. Mulla, M. Y. Pathan, S. S. Chavan, S. P. Gample and D. Sarkar, *RSC Adv.* 2014, 4, 7666-7672.
- 35 (a) Y. Q. Yu, *Synth.*, 2013, 45, 2545–2550. (b) Y. Zhao, X. Chen, T. Chen, Y. Zhou, S. Yin and L. Han, *J. Org. Chem.*, 2015, 80, 62–69. (c) B. Das, G. Satyalakshmi, K. Suneel and K. Damodar, *J. Org. Chem.*, 2009, 74, 8400–8402.
- 36 (a) N. S. Zefirov and E. D. Matveeva, *Arkivoc*, 2008, 2008, 1–17. (b) G. Keglevich and E. Bálint, *Molecules*, 2012, 17, 12821–12835.
- 37 T. Akiyama, M. Sanada and K. Fuchibe, Synlett, 2003, 2003, 1463–1464.
- 38 J.-A. Ma, Chem. Soc. Rev., 2006, 35, 630-636.
- (a) L. Wang, S. M. Cui, W. Meng, G. W. Zhang, J. Nie and J. A. Ma, *Chinese Science Bull.*, 2010, 1729–1731. (b) W. Xu, S. Zhang, S. Yang, L. H. Jin, P. S. Bhadury, D. Y. Hu and Y. Zhang, *Molecules*, 2010, **15**, 5782–5796. (c) P. S. Reddy, M. V. K. Reddy and P. V. G. Reddy, *Chinese Chem. Lett.*, 2016, **27**, 943-947. (d) X. Cheng, R. Goddard, G. Buth and B. List, *Angew. Chemie Int. Ed.*, 2008, **47**, 5079–5081. (e) P. B. Thorat, S. V. Goswami, R. L. Magar, B. R. Patil and S. R. Bhusare, *Eur. J. Org. Chem.*, 2013, **2**, 5509–5516.

Dalton Transactions

- 40 X. Zhou, D. Shang, Q. Zhang, L. Lin, X. Liu and X. Feng, *Org. Lett.*, 2009, **11**, 1401–1404.
- 41 M. Ohara, S. Nakamura and N. Shibata, Adv. Synth. Catal., 2011, 353, 3285-3289.
- 42 C. J. Carrasco, F. Montilla and A. Galindo, Catal. Commun., 2016, 84, 134-136.

Dalton Transactions Accepted Manuscript

FOR the Table of Contents

Synthesis and structural characterization of homochiral 2D coordination polymers of zinc and copper with conformationally flexible ditopic imidazolium-based dicarboxylate ligands

Antonio I. Nicasio, Francisco Montilla,* Eleuterio Álvarez, Rosario P. Colodrero and Agustín Galindo*

Chiral imidazolium-based dicarboxylate ligands behave as bridging ditopic linkers capable of forming 2D coordination polymers with zinc and copper metals.

