

# An efficient and new protocol for phosphine-free Suzuki coupling reaction using palladium-encapsulated and air-stable MIDA boronates in an aqueous medium

Cite this: *RSC Adv.*, 2014, 4, 28148

Joaquim Fernando Mendes da Silva,<sup>\*a</sup> Andres Felipe Yepes Perez<sup>ab</sup> and Natália Pinto de Almeida<sup>a</sup>

Received 20th April 2014  
Accepted 13th June 2014

DOI: 10.1039/c4ra03586k

www.rsc.org/advances

A simple methodology that uses a system based on polyurea microencapsulated palladium (PdEnCat 30™) and aryl or (2-pyridyl) MIDA boronates for Suzuki–Miyaura cross-coupling reactions of (hetero)aryl halides in water–alcohol under phosphine-free conditions was developed.

## Introduction

The palladium catalyzed Suzuki–Miyaura cross-coupling reaction is one of the most powerful and convenient approaches for the synthesis of biaryl compounds.<sup>1</sup> Palladium-catalyzed cross-coupling processes for carbon–carbon bond formation of (het) aryl halides have become an indispensable tool for synthetic and medicinal chemists in their search for new derivatives with wide-ranging therapeutic potential.<sup>2</sup> During the last two decades, there has been an increasing interest in the use of water as solvent for many homogeneously catalyzed reactions.<sup>3</sup> Cost, environmental benefits and safety are among the reasons most often argued to justify the replacement of organic solvents by water in organic transformations. The use of water in Pd-catalyzed cross-coupling reactions goes back to the early development of Suzuki–Miyaura cross-coupling reactions.<sup>4</sup>

On another hand, although boronic acids can serve as excellent building blocks for the synthesis of a wide range of natural products, pharmaceuticals, and materials,<sup>5</sup> some of the potentially most useful boronic acids, including 2-heterocyclic<sup>6–8</sup> derivatives, are inherently unstable, which can significantly limit their benchtop storage and/or efficient cross-coupling. *N*-Methyliminodiacetic acid (MIDA) boronates<sup>9,10</sup> represent the first general solution to this problem by virtue of their stability and remarkable capacity for *in situ slow release* of unstable boronic acids. We herein report the use of (aryl) or (2-pyridyl) *N*-methyliminodiacetic acid (MIDA) boronate for

Suzuki–Miyaura cross-coupling reactions of highly challenging aryl and heteroarylchlorides, bromides or iodides.

Also, as part of our ongoing interest in palladium-catalyzed reactions, we describe the Suzuki–Miyaura cross-coupling reaction of aryl(heteroaryl)halides with the corresponding MIDA boronates catalyzed by an air-stable and available palladium-encapsulated catalyst (Pd EnCat30™). PdEnCat catalyst significantly facilitates these reactions, benefiting from a much simplified workup procedure: the heterogeneous catalyst is readily removed by filtration.

Therefore, these transformations are often mediated by Pd species in the presence of phosphine ligands, and this may result in difficulties associated with the removal of homogeneous material from reaction mixtures. We have developed an effective methodology that use phosphine-free supported Pd catalyst entrapped in a polyurea matrix, with potential for both economic and environmental advantages. PdEnCat 30™ have proven to be effective as recyclable catalysts for use in Suzuki type cross-coupling reactions and do not require expensive and toxic phosphines as co-ligands.<sup>11–13</sup>

On the other hand, due to their significant and varied biological activities, the design and development of novel methods for the construction of 2-thienyl, 2-pyridil, 3-pyridil, 4-pyridil and 2-phenyl functionalized pyridines have drawn remarkable interest from the synthetic, organic as well as medicinal chemists.<sup>14</sup> 2-(2'-Thienyl)pyridine derivatives have been used frequently to obtain complexes with several transition metals with a wide range of applications in solar energy conversion devices as photocatalysts, photosensitizers and electroluminescent emitters.<sup>15</sup> Also, derivatives of 2-thienylpyridine skeleton exhibited considerable topoisomerase I and II inhibitory activity<sup>16</sup> and are being used in the design of various anticancer agents.<sup>17</sup> Secondly, phenyl pyridines have emerged as the integral backbone of several potent azapeptide HIV protease inhibitors with anti HIV activity such as Atazanavir (BMS-232632), an orally

<sup>a</sup>Instituto de Química, Universidade Federal do Rio de Janeiro, Rua Hélio de Almeida 40, Pólo de Xistoquímica Prof. Claudio Costa Neto, PXQmed A 107, Cidade Universitária, Rio de Janeiro, CEP 21941-614, Brazil. E-mail: joaquim@iq.ufrj.br; andre\_yepes@hotmail.com; Fax: +55 21 2560 2069

<sup>b</sup>Facultad de Ciencias Físicas Exactas y Naturales Universidad de Santander (UDES). Centro Investigación en Ciencias Químicas y Tecnologías Sostenibles-QUIMISOST Cidade Universitária, Campus Lagos del Cacique Bucaramanga, Colombia. E-mail: andre\_yepes@hotmail.com; Fax: +57 7 6516492




Fig. 1 Structure of 2-thienylpyridines,  $\alpha$ -terpyridine and 4,6-diaryl-2,4'-bipyridine derivatives.

available azapeptide.<sup>18</sup> Bipyridine compounds as  $\alpha$ -terpyridine or 4,6-diaryl-2,4'-bipyridine molecule can act as tridentate ligands and form stable complexes by chelating a broad variety of transition metal ions. The numerous reports on DNA binding property and antitumor activity of terpyridine complexes have attracted multiple researchers.<sup>19</sup> (Fig. 1)

One of the main objectives of eco-friendly methods in organic chemistry is to develop cost-effective and environmentally benign synthetic protocols which have become one of the main themes of contemporary synthetic chemistry. Thus, we developed a mild, phosphine-free single-step procedure for the Suzuki coupling of readily available hetaryl(aryl)halides in water medium to the corresponding biaryl, bipyridyl and thienylpyridine as a part of our ongoing research in the development of new ecofriendly protocols.

## Results and discussion

In order to optimize the reaction conditions for the Suzuki–Miyaura synthesis we studied the coupling reaction of 4-bromoacetophenone with 2-formylphenylboronic acid MIDA or 6-methoxyphenylboronic acid MIDA ester (1.5 equiv.) in the presence of  $K_2CO_3$  (3.0 equiv.) as a model.

Two parameters were investigated: (1) optimal ratio of aqueous solvent used and (2) palladium catalyst concentration (Table 1).

Initially, the reaction was carried out using only ethanol as solvent, Pd EnCat30 (10 mol%) as the catalyst and  $K_2CO_3$  (3 equiv.) as base, heating under reflux for 3–9 h, when we isolated the Suzuki product in very low yields, the major products being the homocoupling and the dehalogenation products, alongside with the formation of black palladium. However, yields increased substantially when significant volumes of water were added to the reaction. Then, we tested several volumetric ratios of ethanol :  $H_2O$  (1 : 1, 2 : 1, 3 : 1, 4 : 1, 6 : 1, 8 : 1) and we found that in the reaction conditions used, the highest conversion to **1** was observed when the volumetric ratio of ethanol :  $H_2O$  was 4 : 1, together with the lowest

occurrence of the debrominated or homocoupling side products, while the formation of black palladium was not observed. In the other ratios tested, the coupling reactions were accompanied by different extents of hydrodehalogenation or homocoupling, which were isolated always as major products in these cases where formation of black palladium was observed.

Heterocyclic compounds are of particular interest to the pharmaceutical industry,<sup>20</sup> but the application of heterocyclic compounds in cross-coupling reactions remains a challenge.<sup>21</sup> To the best of our knowledge, only a few successful examples of aqueous-phase Suzuki–Miyaura cross-coupling reactions of heteroaryl chlorides are known.<sup>22</sup>

Thus, there is a need for the development of efficient procedures for the cross-coupling reactions of heteroaryl chlorides with arylboronic acids in water. To carry out an economical large-scale synthesis of pharmaceutical intermediates or other bulk fine chemicals, a low loading catalyst is needed. So then, we finally investigated the minimum quantity of catalyst required to achieve a fast and complete conversion. We

Table 1 Suzuki–Miyaura reaction of 4-bromoacetophenone with 2-formylphenylboronic acid MIDA ester as a model to optimize the conditions

Entry <sup>a</sup>	Volumetric ratios EtOH : $H_2O$	Catalyst loading (mol%)	Suzuki-coupling product ( <b>1a</b> ) yield <sup>b</sup> (%)
1	1 : 1	10% mol	(51%)
2	2 : 1	10% mol	(56%)
3	3 : 1	10% mol	(60%)
4	4 : 1	10% mol	(76%)
5	4 : 1	8% mol	(77%)
6	4 : 1	6% mol	(80%)
7	4 : 1	2.5% mol	(70%)
8	4 : 1	1% mol	(67%)
9	6 : 1	10% mol	(61%)
10	8 : 1	10% mol	(63%)

<sup>a</sup> 3.0 equiv. of  $K_2CO$  were used. <sup>b</sup> Isolated yield.

Table 2 Reuse of Pd-catalyst in the Suzuki–Miyaura reaction of 4-bromoacetophenone with 2-formylphenylboronic acid MIDA ester<sup>a</sup>

% Yield <b>1a</b>	1st recycle	2nd recycle	3rd recycle
	78	77	74

<sup>a</sup> Conditions:  $K_2CO_3$  (3.0 mmol), EtOH :  $H_2O$  (4 : 1), reflux, 8 h.

**Table 3** Suzuki–Miyaura reaction of aryl (hetaryl) halides with 2-formylphenylboronic acid MIDA ester and two 2-pyridyl boronic acid MIDA ester<sup>a</sup>

Entry <sup>c</sup>	R-X	R <sup>1</sup>	Reaction time (h)	Suzuki-coupling product (1) yield <sup>b</sup> (%)
1			(8)	<b>1a</b> (80%)
2			(5)	<b>1b</b> (90%)
3			(3)	<b>1c</b> (98%)
4			(9)	<b>1d</b> (78%)
5			(8)	<b>1e</b> (80%)
6 <sup>d</sup>			(8)	<b>1f</b> (91%)
7			(9)	<b>1g</b> (97%)
8			(8)	<b>1h</b> (85%)
9			(5)	<b>1i</b> (84%)
10			(9)	<b>1j</b> (81%)
11			(3)	<b>1k</b> (82%)
12			(8)	<b>1l</b> (84%)
13			(8)	<b>1m</b> (96%)
14			(9)	<b>1n</b> (91%)
15			(9)	<b>1o</b> (72%)
16			(3)	<b>1p</b> (76%)
17			(8)	<b>1q</b> (85%)
18			(8)	<b>1r</b> (76%)
19			(3)	<b>1s</b> (80%)

**Table 3** (Contd.)

Entry <sup>c</sup>	R-X	R <sup>1</sup>	Reaction time (h)	Suzuki-coupling product (1) yield <sup>b</sup> (%)
20			(8)	<b>1t</b> (75%)

<sup>a</sup> Reaction conditions: 1 equiv. of aryl halide (1 mmol), 1.5 equiv. of MIDA boronate, 6 mol% Pd EnCat30, 3 equiv. of K<sub>2</sub>CO<sub>3</sub>, 0.1 M in 4 : 1 ethanol–H<sub>2</sub>O. <sup>b</sup> Isolated yield. <sup>c</sup> Cross-couplings were run at reflux. <sup>d</sup> 4-Bromopyridine hydrochloride was used.

compared three conditions with decreasing amounts of Pd EnCat30 (without phosphine) and using K<sub>2</sub>CO<sub>3</sub> as base and ethanol : H<sub>2</sub>O (4 : 1) as solvent. Then, carrying out the coupling reaction at palladium loadings of 10, 8, 6, 2.5 or 1 mol% always led to the desired coupling product in similar yields after 8 h at reflux. The best results were obtained in the presence of a 6 mol% catalyst concentration.

Additionally, we performed a study of the catalytic activity of Pd EnCat30 for our reaction under the best conditions found using the reaction of 4-bromoacetophenone with 2-formylphenylboronic acid MIDA ester as model. The results showed that when the catalyst was recycled 3 times at the reaction conditions established no significant loss of activity or yield occurred. During these reactions, the formation of black palladium was not observed (Table 2).

Under the optimized reaction conditions, we investigated the reaction of some aryl halides with 2-formylphenylboronic acid and 6-methoxy-2-pyridylboronic acids MIDA esters. This investigation showed that the reaction successfully occurred without any phosphine ligand to give the corresponding Suzuki-coupling products **1a–t** in good to excellent yields (Table 3).

Then, a wide array of aryl halides reacted efficiently with 2-formylphenylboronic acid MIDA ester and 6-methoxy-2-pyridylboronic acid MIDA ester respectively to provide the corresponding products **1a–t**. For example, 4-bromobenzaldehyde and 4-iodonitrobenzene reacted with 2-formylphenylboronic acid MIDA ester to yield in 90% and 98% respectively, the coupling products **1b** and **1c**.

The coupling reactions of representative heteroaromatic halides were also effective with this protocol. Under the same conditions, 3-bromopyridine reacted efficiently with 6-methoxy-2-pyridylboronic acid MIDA ester within 8 h, providing 85% yield of the desired product **1q**, while, the quinoline **1s** was prepared by the reaction of 2-bromoquinoline with 6-methoxy-2-pyridylboronic acid MIDA ester to give the desired product in good yield (80%). The 2,2'-bipyridine unit is a key structural feature found in some important natural products that show antibiotic and cytotoxic activities.<sup>23</sup>

The feasibility of our strategy was demonstrated in the synthesis of thio-derivatives, when 2-bromothiophene was treated with 2-formylphenylboronic acid MIDA ester or 6-methoxyphenylboronic acid MIDA ester (1.5 equiv.) under the conditions already established. The reaction proceeded well to give 2-(thiophen-2-yl)benzaldehyde (**1h**) and 2-methoxy-6-(thiophen-2-yl)pyridine (**1t**) respectively, in good yields *via* C–C bond formation. Since most of the strategies for the synthesis of these phenyl-pyridines or thienylpyridines by condensation and substitution reactions frequently afford the desired products in low yields and with low regioselectivity towards the 2-position, our synthetic strategy allows the effective construction of large libraries of 2-thienyl and 2-phenylpyridinederivatives in an efficient and eco-friendly method.

## Conclusions

In conclusion, a convenient methodology was developed for the palladium catalyzed Suzuki–Miyaura reaction in water. The reaction is catalyzed by a system based on Pd- micro-encapsulated in a polyurea matrix (Pd EnCat30), through a simple and efficient protocol for the phosphine-free Suzuki coupling of aryl (hetaryl) halides with arylboronic acids MIDA esters. The advantages of this procedure are simple operation, good to excellent yields, short reaction time, absence of phosphine-ligands and mild and environmentally friendly conditions. A great diversity of substrates was tolerated for this method.

## Experimental procedure

### General information

All reagents were purchased from Sigma Aldrich and used without further purification. Ethanol were purchased from local suppliers. The analyses by Gas Chromatography-Mass Spectrometry (GC-MS) were performed on a Shimadzu GCMS-QP2010S with a RTX-5MS capillary column (30 m × 0.25 mm × 0.25 μm, with the stationary phase 5% diphenyl and 95% dimethylpolysiloxane), mass detector in mode electron ionization (EI) standard (70 eV), with helium as a carrier gas. The <sup>1</sup>H- and <sup>13</sup>C NMR spectra were recorded on a Bruker AMX-200 spectrometer operating at 200 MHz (<sup>1</sup>H) and 50 MHz (<sup>13</sup>C) in CDCl<sub>3</sub> with TMS as an internal standard. The <sup>1</sup>H- and <sup>13</sup>C NMR spectra were processed using MESTREC/NMR.

### General method for carrying out the Suzuki-coupling (Table 3)

Under ambient atmosphere, to a 15 mL vial equipped with a stir bar was added the halide (1.0 mmol), 2-formylphenyl or 2-pyridyl MIDA boronate (1.5 mmol), K<sub>2</sub>CO<sub>3</sub> (3.0 mmol), Pd enCat30 (6 mol%) and ethanol : H<sub>2</sub>O (4 : 1 v/v). The reaction mixture was stirred at reflux for 3–9 h. The mixture was cooled to room temperature and then was transferred to a 60 mL separatory funnel and was added brine (10 mL). The mixture was extracted with AcOEt (3 × 10 mL). The combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo*. The organic residue was adsorbed onto silica gel, and then purified by column flash

chromatography (hexane/ethyl acetate as eluent) to afford the desired product **1a–t**. All the compounds prepared are known compounds.<sup>24–42</sup> The products were confirmed by comparing the <sup>1</sup>H NMR and mass spectral data with authentic samples reported in the literature.

**4'-Acetyl-[1,1'-biphenyl]-2-carbaldehyde (1a).**<sup>24</sup> Following the general procedure for Table 3, using 4-bromoacetophenone (70 mg, 0.351 mmol), 2-formylphenylboronic acid MIDA ester (110 mg, 0.422 mmol), potassium carbonate (145 mg, 1.054 mmol), Pd EnCat30 (53 mg, 0.021 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1a** as a colorless oil (63 mg, 80% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.67 (s, 3H), 7.43–7.61 (m, 3H), 7.63–7.72 (m, 2H), 8.02–8.09 (m, 3H), 9.97 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 26.9, 128.3, 128.6, 128.7, 129.7, 130.0, 130.3, 130.5, 130.8, 133.9, 136.8, 142.9, 144.7, 191.9, 197.7.

**4'-Nitro-[1,1'-biphenyl]-2-carbaldehyde (1b).**<sup>25</sup> Following the general procedure for Table 3, using 4-iodonitrobenzene (70 mg, 0.281 mmol), 2-formylphenylboronic acid MIDA ester (88 mg, 0.337 mmol), potassium carbonate (117 mg, 0.843 mmol), Pd EnCat30 (43 mg, 0.017 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 5 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1b** as a light yellow solid (54 mg, 90% yield), mp 118–120 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.43 (d, *J* = 7.6 Hz, 1H), 7.57–7.67 (m, 4H), 7.72 (t, *J* = 7.5 Hz, 1H), 8.04 (d, *J* = 7.6 Hz, 1H), 8.34 (d, *J* = 8.3 Hz, 1H), 9.97 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 123.8, 129.1, 129.3, 130.8, 131.0, 133.8, 134.1, 143.2, 145.0, 147.9, 191.2.

**[1,1'-Biphenyl]-2,4'-dicarbaldehyde (1c).**<sup>26</sup> Following the general procedure for Table 3, using 4-bromobenzaldehyde (80 mg, 0.432 mmol), 2-formylphenylboronic acid MIDA ester (170 mg, 0.648 mmol), potassium carbonate (180 mg, 1.30 mmol), Pd EnCat30 (64.8 mg, 0.026 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 3 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 7 : 1 hexane : ethyl acetate) to provide **1c** as a white solid (87 mg, 98% yield), mp 94–96 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.27–7.74 (m, 5H), 7.98–8.08 (m, 3H), 9.98 (s, 1H), 10.11 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 128.4, 128.8, 129.8, 130.7, 130.8, 133.8, 133.9, 136.0, 144.3, 144.4, 191.6, 191.8.

**4'-Methoxy-[1,1'-biphenyl]-2-carbaldehyde (1d).**<sup>27</sup> Following the general procedure for Table 3, using 4-iodoanisole (85 mg, 0.363 mmol), 2-formylphenylboronic acid MIDA ester (142 mg, 0.540 mmol), potassium carbonate (151 mg, 1.09 mmol), Pd EnCat30 (54 mg, 0.022 mmol), 4 : 1 aq. ethanol (11 mL) were heated under reflux for 9 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 15 : 1 hexane : ethyl acetate) to provide **1d** as a colorless oil (60 mg, 78% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.88 (s, 3H), 7.01 (d, *J* = 8.8 Hz, 2H), 7.31 (d, *J* = 8.8 Hz, 2H), 7.42–7.46 (m, 2H), 7.62 (t, *J* = 7.6 Hz, 1H), 8.01 (d, *J* = 7.6 Hz, 1H), 10.0 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 55.6, 114.2, 127.6, 127.8, 130.3, 131.0, 131.5, 133.7, 134.0, 145.9, 159.9, 192.9.

**4'-Methyl-[1,1'-biphenyl]-2-carbaldehyde (1e).**<sup>28</sup> Following the general procedure for Table 3, using 4-bromotoluene

(75 mg, 0.44 mmol), 2-formylphenylboronic acid MIDA ester (171 mg, 0.657 mmol), potassium carbonate (181 mg, 1.31 mmol), Pd EnCat30 (66 mg, 0.026 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 20 : 1 hexane : ethyl acetate) to provide **1d** as a light yellow oil (68 mg, 80% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.40 (s, 3H), 7.22–7.30 (m, 4H), 7.39 (d, *J* = 8.2 Hz, 1H), 7.48 (d, *J* = 7.5 Hz, 1H), 7.60 (td, *J* = 7.5, 1.2 Hz, 1H), 8.00 (dd, *J* = 8.2, 1.2 Hz, 1H), 9.97 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 55.5, 114.4, 127.3, 127.6, 130.6, 130.9, 131.5, 133.9, 134.1, 146.0, 159.8, 192.8.

**2-(Pyridin-4-yl)benzaldehyde (1f).**<sup>29</sup> Following the general procedure for Table 3, using 4-bromopyridine hydrochloride (95 mg, 0.49 mmol), 2-formylphenylboronic acid MIDA ester (192 mg, 0.734 mmol), potassium carbonate (203 mg, 1.47 mmol), Pd EnCat30 (74 mg, 0.029 mmol), 4 : 1 aq. ethanol (12 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 3 : 2 hexane : ethyl acetate) to provide **1f** as a light orange solid (80 mg, 91% yield); mp 64–66 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.32–7.35 (m, 2H), 7.43 (d, *J* = 7.5 Hz, 1H), 7.60 (t, *J* = 7.5 Hz, 1H), 7.73 (dt, *J* = 7.5, 1.5 Hz, 1H), 8.07 (d, *J* = 7.5 Hz, 1H), 8.72 (dd, *J* = 6.2, 1.1 Hz, 2H), 9.98 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 124.9, 128.6, 129.2, 130.5, 133.6, 134.0, 142.9, 146.1, 150.0, 191.3.

**2-(Pyridin-3-yl)benzaldehyde (1g).**<sup>30</sup> Following the general procedure for Table 3, using 3-bromopyridine (70 mg, 0.44 mmol), 2-formylphenylboronic acid MIDA ester (139 mg, 0.531 mmol), potassium carbonate (184 mg, 1.33 mmol), Pd EnCat30 (66 mg, 0.026 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 9 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 5 : 1 hexane : ethyl acetate) to provide **1g** as a colorless oil (80 mg, 91% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.40–7.46 (m, 2H), 7.58 (d, *J* = 7.5 Hz, 1H), 7.60 (t, *J* = 7.5 Hz, 1H), 7.73 (d, *J* = 7.5 Hz, 1H), 8.06 (d, *J* = 7.5 Hz, 1H), 8.70 (dd, *J* = 6.4, 1.5 Hz, 2H), 9.99 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 123.3, 128.7, 128.9, 131.2, 133.8, 133.9, 134.1, 137.4, 142.0, 149.5, 150.3, 191.5.

**2-(Thiophen-2-yl)benzaldehyde (1h).**<sup>30</sup> Following the general procedure for Table 3, using 2-bromothiophene (100 mg, 0.61 mmol), 2-formylphenylboronic acid MIDA ester (240 mg, 0.92 mmol), potassium carbonate (253 mg, 1.83 mmol), Pd EnCat30 (92 mg, 0.036 mmol), 4 : 1 aq. ethanol (13 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 30 : 1 hexane : ethyl acetate) to provide **1h** as a light yellow oil (98 mg, 85% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.07–7.14 (m, 1H), 7.15–7.18 (m, 1H), 7.46–7.49 (m, 3H), 7.60 (d, *J* = 8.2 Hz, 1H), 8.01 (d, *J* = 7.5 Hz, 1H), 10.20 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 127.5, 127.9, 128.0, 128.4, 129.7, 131.5, 133.7, 134.4, 138.2, 138.9, 192.2.

**[1,1'-Biphenyl]-2,2'-dicarbaldehyde (1i).**<sup>31</sup> Following the general procedure for Table 3, using 2-bromobenzaldehyde (60 mg, 0.324 mmol), 2-formylphenylboronic acid MIDA ester (127 mg, 0.49 mmol), potassium carbonate (134 mg, 0.972 mmol), Pd EnCat30 (48 mg, 0.020 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 5 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1i** as a light yellow oil (56 mg,

84% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.34 (dd, *J* = 1.0, 7.8 Hz, 2H), 7.58 (t, *J* = 7.6 Hz, 2H), 7.66 (dt, *J* = 1.2, 7.6 Hz, 2H), 8.06 (dd, *J* = 1.0, 7.8 Hz, 2H), 9.83 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 191.1, 141.3, 134.7, 133.5, 131.8, 128.9, 128.7.

**4'-Hydroxy-2'-methyl-[1,1'-biphenyl]-2-carbaldehyde (1j).**<sup>32</sup> Following the general procedure for Table 3, using 4-chloro-3-methylphenol (70 mg, 0.491 mmol), 2-formylphenylboronic acid MIDA ester (192 mg, 0.73 mmol), potassium carbonate (203 mg, 1.47 mmol), Pd EnCat30 (73 mg, 0.0294 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 1 : 1 hexane : ethyl acetate) to provide **1j** as a white solid (81 mg, 81% yield), mp 148–150 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.51 (s, 3H), 7.27–7.17 (m, 2H), 7.35 (d, *J* = 7.4 Hz, 1H), 7.60–7.55 (m, 2H), 7.87–7.90 (m, 2H), 9.86 (s, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 55.5, 114.1, 121.4, 128.6, 128.7, 132.3, 133.3, 133.6, 133.7, 134.8, 135.4, 139.7, 155.6, 195.9.

**2-Methoxy-6-(4-nitrophenyl)pyridine (1k).**<sup>33</sup> Following the general procedure for Table 3, using 4-iodonitrobenzene (85 mg, 0.34 mmol), 6-methoxy-pyridylboronic acid MIDA ester (135 mg, 0.51 mmol), potassium carbonate (142 mg, 1.02 mmol), Pd EnCat30 (51 mg, 0.020 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 3 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1k** as a light yellow solid (64 mg, 82% yield); mp 104–106 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.05 (s, 3H), 6.80 (d, *J* = 8.0 Hz, 1H), 7.42 (d, *J* = 8.0 Hz, 1H), 7.69 (t, *J* = 8.0 Hz, 1H), 8.18–8.32 (m, 4H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.6, 111.5, 114.0, 124.1, 127.5, 139.6, 145.2, 148.2, 152.2, 164.2.

**4-(6-Methoxypyridin-2-yl)acetophenone (1l).**<sup>34</sup> Following the general procedure for Table 3, using 4-bromoacetophenone (50 mg, 0.25 mmol), 6-methoxy-pyridylboronic acid MIDA ester (99 mg, 0.38 mmol), potassium carbonate (104 mg, 0.75 mmol), Pd EnCat30 (38 mg, 0.015 mmol), 4 : 1 aq. ethanol (8 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1l** as a white solid (48 mg, 84% yield); mp 73–75 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.64 (s, 3H), 4.05 (s, 3H), 6.75 (d, *J* = 8.0 Hz, 1H), 7.38 (d, *J* = 8.0 Hz, 1H), 7.66 (t, *J* = 8.0 Hz, 1H), 8.01–8.16 (m, 4H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 26.9, 53.5, 110.6, 113.7, 126.9, 128.9, 137.2, 139.5, 143.5, 153.4, 164.1, 198.0.

**4-(6-Methoxypyridin-2-yl)benzaldehyde (1m).**<sup>35</sup> Following the general procedure for Table 3, using 4-bromobenzaldehyde (60 mg, 0.32 mmol), 6-methoxy-pyridylboronic acid MIDA ester (128 mg, 0.48 mmol), potassium carbonate (134 mg, 0.96 mmol), Pd EnCat30 (49 mg, 0.019 mmol), 4 : 1 aq. ethanol (8 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1m** as a white solid (66 mg, 96% yield); mp 74–76 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.05 (s, 3H), 6.77 (d, *J* = 7.6 Hz, 1H), 7.42 (d, *J* = 7.6 Hz, 1H), 7.67 (t, *J* = 7.6 Hz, 1H), 7.96 (d, *J* = 7.6 Hz, 2H), 8.22 (d, *J* = 7.6 Hz, 2H), 10.07 (s, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.5, 110.9, 113.9, 127.3, 130.2, 136.5, 139.5, 144.8, 153.2, 164.1, 192.1.

**2-Methoxy-6-(4-methoxyphenyl)pyridine (1n).**<sup>36</sup> Following the general procedure for Table 3, using 4-iodoanisole (50 mg,

0.21 mmol), 6-methoxyppyridylboronic acid MIDA ester (84 mg, 0.32 mmol), potassium carbonate (89 mg, 0.63 mmol), Pd EnCat30 (32 mg, 0.013 mmol), 4 : 1 aq. ethanol (8 mL) were heated under reflux for 9 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1n** as a colorless oil (42 mg, 91% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.85 (s, 3H), 4.02 (s, 3H), 6.63 (d, *J* = 7.8 Hz, 1H), 6.97 (d, *J* = 8.0 Hz, 1H), 7.26 (d, *J* = 8.0 Hz, 1H), 7.55 (t, *J* = 8.0 Hz, 1H), 7.98–8.02 (m, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.3, 55.5, 108.5, 112.1, 114.2, 114.4, 127.9, 128.2, 132.0, 139.3, 154.6, 160.6, 163.9.

**6-Methoxy-2,2'-bipyridine (1o).**<sup>37</sup> Following the general procedure for Table 3, using 2-bromopyridine (60 mg, 0.38 mmol), 6-methoxyppyridylboronic acid MIDA ester (150 mg, 0.57 mmol), potassium carbonate (157 mg, 1.14 mmol), Pd EnCat30 (57 mg, 0.023 mmol), 4 : 1 aq. ethanol (8 mL) were heated under reflux for 9 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 15 : 1 hexane : ethyl acetate) to provide **1o** as a colorless oil (51 mg, 72% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.05 (s, 3H), 6.76 (d, *J* = 8.0 Hz, 1H), 7.28–7.33 (m, 1H), 7.74 (t, *J* = 8.0 Hz, 1H), 7.80 (td, *J* = 7.4, 1.0 Hz, 1H), 8.00 (d, *J* = 7.4 Hz, 1H), 8.41 (d, *J* = 7.5 Hz, 1H), 8.65 (dd, *J* = 7.4, 1.0 Hz, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.4, 111.2, 113.9, 121.2, 123.7, 136.9, 139.6, 149.3, 153.7, 156.3, 163.7.

**6'-Methoxy-5-nitro-2,2'-bipyridine (1p).**<sup>38</sup> Following the general procedure for Table 3, using 2-chloro-4-nitropyridine (65 mg, 0.41 mmol), 6-methoxyppyridylboronic acid MIDA ester (163 mg, 0.62 mmol), potassium carbonate (170 mg, 1.23 mmol), Pd EnCat30 (62 mg, 0.025 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 3 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 10 : 1 hexane : ethyl acetate) to provide **1p** as a white solid (72 mg, 76% yield); mp 114–116 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.06 (s, 3H), 6.88 (d, *J* = 7.8 Hz, 1H), 7.76 (t, *J* = 7.6 Hz, 1H), 8.13 (d, *J* = 7.4 Hz, 1H), 8.57–8.64 (m, 2H), 9.45 (s, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.6, 113.4, 115.7, 121.0, 132.0, 139.8, 144.0, 144.9, 151.3, 161.1, 164.0.

**6-Methoxy-2,3'-bipyridine (1q).**<sup>39</sup> Following the general procedure for Table 3, using 3-bromopyridine (60 mg, 0.38 mmol), 6-methoxyppyridylboronic acid MIDA ester (150 mg, 0.57 mmol), potassium carbonate (157 mg, 1.14 mmol), Pd EnCat30 (57 mg, 0.023 mmol), 4 : 1 aq. ethanol (8 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 5 : 1 hexane : ethyl acetate) to provide **1q** as a colorless oil (60 mg, 85% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.03 (s, 3H), 6.74 (d, *J* = 8.0 Hz, 1H), 7.35–7.41 (m, 2H), 7.66 (t, *J* = 8.0 Hz, 1H), 7.80 (dt, *J* = 8.0, 1.0 Hz, 1H), 8.62 (dd, *J* = 8.0, 1.0 Hz, 1H), 9.25 (s, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.5, 110.4, 114.0, 123.6, 134.1, 134.7, 139.5, 148.4, 149.8, 152.1, 164.2.

**6-Methoxy-2,4'-bipyridine (1r).**<sup>40</sup> Following the general procedure for Table 3, using 4-bromopyridine hydrochloride (80 mg, 0.41 mmol), 6-methoxyppyridylboronic acid MIDA ester (163 mg, 0.61 mmol), potassium carbonate (170 mg, 1.23 mmol), Pd EnCat30 (62 mg, 0.025 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>,

2 : 1 hexane : ethyl acetate) to provide **1r** as a colorless oil (58 mg, 76% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.97 (s, 3H), 6.72 (dd, *J* = 8.3, 0.6 Hz, 1H), 7.34 (dd, *J* = 7.4, 0.6 Hz, 1H), 7.60 (t, *J* = 7.4 Hz, 1H), 7.81–7.87 (m, 2H), 8.59–8.66 (m, 2H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.5, 111.7, 113.6, 121.0, 139.5, 146.3, 140.4, 151.9, 164.2.

**2-(6-Methoxyppyridin-2-yl)quinoline (1s).**<sup>41</sup> Following the general procedure for Table 3, using 2-bromoquinoline (70 mg, 0.33 mmol), 6-methoxyppyridylboronic acid MIDA ester (133 mg, 0.50 mmol), potassium carbonate (139 mg, 1.00 mmol), Pd EnCat30 (50 mg, 0.020 mmol), 4 : 1 aq. ethanol (10 mL) were heated under reflux for 3 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 20 : 1 hexane : ethyl acetate) to provide **1s** as a white solid (63 mg, 80% yield); mp 79–81 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.02 (s, 3H), 6.83 (d, *J* = 8.1 Hz, 1H), 7.54 (t, *J* = 7.4 Hz, 1H), 7.68–7.80 (m, 2H), 7.84 (d, *J* = 8.1 Hz, 1H), 8.17 (d, *J* = 8.5 Hz, 1H), 8.17–7.30 (m, 2H), 8.57 (d, *J* = 8.5 Hz, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.5, 111.5, 114.7, 119.2, 126.8, 127.7, 128.4, 129.6, 130.0, 136.7, 139.6, 148.1, 153.9, 156.3, 163.8.

**2-Methoxy-6-(thiophen-2-yl)pyridine (1t).**<sup>42</sup> Following the general procedure for Table 3, using 2-bromothiophene (100 mg, 0.61 mmol), 6-methoxyppyridylboronic acid MIDA ester (240 mg, 0.92 mmol), potassium carbonate (253 mg, 1.83 mmol), Pd EnCat30 (91 mg, 0.036 mmol), 4 : 1 aq. ethanol (13 mL) were heated under reflux for 8 h. After work-up the residue was purified by flash column chromatography (SiO<sub>2</sub>, 20 : 1 hexane : ethyl acetate) to provide **1t** as a light yellow oil (88 mg, 75% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.05 (s, 3H), 7.23 (dd, *J* = 3.6, 0.9 Hz, 1H), 7.69 (dd, *J* = 5.1, 0.9 Hz, 1H), 7.92 (dd, *J* = 5.1, 0.6 Hz, 1H), 8.02 (dd, *J* = 8.0, 1.2 Hz, 1H), 8.04 (t, *J* = 8.0, 1.2 Hz, 1H), 8.49 (dd, *J* = 7.8, 1.2 Hz, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 53.4, 108.9, 111.3, 124.3, 127.1, 127.9, 139.1, 144.9, 150.0, 163.5.

## Acknowledgements

The authors acknowledge CAPES and FAPERJ for financial support for this work.

## Notes and references

- (a) A. Suzuki, *Angew. Chem., Int. Ed.*, 2011, **50**, 6722; (b) A. Suzuki and Y. Yamamoto, *Chem. Lett.*, 2011, **40**, 894; (c) A. Fihri, M. Bouhrara, B. Nekoueishahraki, J.-M. Basset and V. Polshettiwar, *Chem. Soc. Rev.*, 2011, **40**, 5181; (d) A. Balanta, C. Godard and C. Claver, *Chem. Soc. Rev.*, 2011, **40**, 4973; (e) N. Miyaura and A. Suzuki, *Chem. Rev.*, 1995, **95**, 2457; (f) F. A. Littke and G. C. Fu, *Angew. Chem., Int. Ed.*, 2002, **41**, 4176; (g) J. Dupont, C. S. Consorti and J. Spencer, *Chem. Rev.*, 2005, **105**, 2527; (h) A. V. Gaikwad, A. Holuigue, M. B. Thathagar, J. E. Elshof and G. Rothenberg, *Chem.-Eur. J.*, 2007, **13**, 6908; (i) N. Miyaura and A. Suzuki, *Chem. Rev.*, 1995, **95**, 2457.
- (a) X.-F. Wu, P. Anbarasan, H. Neumann and M. Beller, *Angew. Chem., Int. Ed.*, 2010, **49**, 9047; (b) M. Seki, *Synthesis*, 2006, 2975; (c) *Cross-Coupling Reactions. A*

- practical Guide Topics Curr. Chem.*, ed. N. Miyaura, Springer Verlag, Berlin, 2002, vol. 219.
- 3 (a) M. O. Simon and C. J. Li, *Chem. Soc. Rev.*, 2012, **41**, 1415; (b) M. Lamblin, L. Nassar-Hardy, J. C. Hierso, E. Fouquet and F. X. Felpin, *Adv. Synth. Catal.*, 2010, **352**, 33; (c) R. N. Butler and A. G. Coyne, *Chem. Rev.*, 2010, **110**, 6302; (d) K. H. Shaughnessy, *Chem. Rev.*, 2009, **109**, 643; (e) U. M. Lindström, *Chem. Rev.*, 2002, **102**, 2751.
- 4 N. Miyaura, K. Yamada, H. Suginome and A. Suzuki, *J. Am. Chem. Soc.*, 1985, **107**, 972.
- 5 D. G. Hall, *Boronic Acids*, Wiley-VCH, Weinheim, Germany, 2005.
- 6 S. L. Buchwald, *J. Am. Chem. Soc.*, 2007, **129**, 3358.
- 7 For some examples of coupling of 2-heterocyclic boronic acids with activated aryl and acid chlorides, see: (a) Ref. 2; (b) C. A. Fleckenstein and H. Plenio, *J. Org. Chem.*, 2008, **73**, 3236; (c) M. J. Burns, I. J. S. Fairlamb, A. R. Kapdi, P. Sehnal and R. J. K. Taylor, *Org. Lett.*, 2007, **9**, 5397; (d) B. Xin, Y. Zhang and K. Cheng, *J. Org. Chem.*, 2006, **71**, 5725; (e) N. Pagano, J. Maksimoska, H. Bregman, D. S. Williams, R. D. Webster, F. Xue and E. Meggers, *Org. Biomol. Chem.*, 2007, **5**, 1218, For some representative examples of coupling of 2-heterocyclic boronic acids with aryl iodides and bromides, see: (f) K. Takimiya, Y. Kunugi, Y. Toyoshima and T. Otsubo, *J. Am. Chem. Soc.*, 2005, **127**, 3605; (g) S. W. Thomas, K. Venkatesan, P. Mueller and T. M. Swager, *J. Am. Chem. Soc.*, 2006, **128**, 16641; (h) G. E. Collis, A. K. Burrell, E. J. Blandford and D. L. Officer, *Tetrahedron*, 2007, **63**, 11141; (i) P. Qin, H. Zhu, T. Edvinsson, G. Boschloo, A. Hagfeldt and L. Sun, *J. Am. Chem. Soc.*, 2008, **130**, 8570; (j) H. Maeda, Y. Haketa and T. Nakanishi, *J. Am. Chem. Soc.*, 2007, **129**, 13661.
- 8 E. Tyrrell and P. Brookes, *Synthesis*, 2003, 469.
- 9 T. Mancilla, R. Contreras and B. Wrackmeyer, *J. Organomet. Chem.*, 1986, **307**, 1.
- 10 (a) E. P. Gillis and M. D. Burke, *J. Am. Chem. Soc.*, 2007, **129**, 6716; (b) S. J. Lee, K. C. Gray, J. S. Paek and M. D. Burke, *J. Am. Chem. Soc.*, 2008, **130**, 466; (c) E. P. Gillis and M. D. Burke, *J. Am. Chem. Soc.*, 2008, **130**, 14084; (d) B. E. Uno, E. P. Gillis and M. D. Burke, *Tetrahedron*, 2009, **65**, 3130; (e) S. G. Ballmer, E. P. Gillis and M. D. Burke, *Org. Synth.*, 2009, **86**, 344; (f) E. P. Gillis and M. D. Burke, *Aldrichimica Acta*, 2009, **42**, 17.
- 11 C. K. Lee, A. B. Holmes, S. V. Ley, I. F. McConvey, B. Al-Duri, G. A. Leeke, R. C. Santos and J. P. K. Seville, *Chem. Commun.*, 2005, 2175.
- 12 A. Molnar, *Chem. Rev.*, 2011, **111**, 2251.
- 13 I. R. Baxendale, C. M. Griffiths-Jones, S. V. Ley and G. K. Tranmer, *Chem.-Eur. J.*, 2006, 4407.
- 14 G. R. Dick, E. M. Woerly and M. D. Burke, *Angew. Chem., Int. Ed.*, 2012, **51**, 2667.
- 15 (a) Y. Fuchita, H. Ieda, S. Wada, S. Kameda and M. Mikuriya, *J. Chem. Soc., Dalton Trans.*, 1999, 4431; (b) J. Schmidt, J. Strasser and H. Yersin, *Inorg. Chem.*, 1997, **36**, 3957; (c) W. S. Tang, X. X. Lu, K. M. C. Wong and V. W. W. Yam, *J. Mater. Chem.*, 2005, **15**, 2714; (d) H. Yersin, S. Schützenmeir, H. Wiedenhofer and A. Von Zelewsky, *J. Phys. Chem.*, 1993, **97**, 13496; (e) E. Chung, D. Bizzotto and M. O. Wolf, *Chem. Commun.*, 2002, 3026; (f) P. I. Kvam, M. V. Puzyk, V. S. Cotlyr, K. P. Balashev and J. Songstad, *Acta Chem. Scand.*, 1995, **49**, 645; (g) R. Argazzi, C. A. Bignozzi, T. A. Heimer and G. J. Meyer, *Inorg. Chem.*, 1997, **36**, 2; (h) H. Hori, F. P. A. Johnson, K. Koike, O. Ishitani and T. J. Ibuski, *J. Photochem. Photobiol., A*, 1996, **96**, 171.
- 16 J. A. Holden, *Curr. Med. Chem.: Anti-Cancer Agents*, 2001, **1**, 1.
- 17 (a) L. X. Zhao, Y. S. Moon, A. Basnet, E. K. Kim, Y. Jahng, J. G. Park, T. C. Jeong, W. J. Cho, S. U. Choi, C. O. Lee, S. Y. Lee, C. S. Lee and E.-S. Lee, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 1333; (b) A. Basnet, P. Thapa, R. Karki, H. Choi, J. H. Choi, M. Yun, B.-S. Jeong, Y. Jahng, Y. Na, W.-J. Won-Jea Cho, Y. Kwon, C.-S. Lee and E.-S. Lee, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 42; (c) P. Thapa, R. Karki, U. Thapa, Y. Jahng, M. J. Jung, J. M. Nam, Y. Na, Y. Kwon and E.-S. Lee, *Bioorg. Med. Chem.*, 2010, **18**, 377.
- 18 M. F. Siddiqui and A. I. Levey, *Drugs Future*, 1999, **24**, 375.
- 19 (a) U. S. Schubert and C. Eschbaumer, *Angew. Chem., Int. Ed.*, 2002, **41**, 2892; (b) G. Lowe, A. S. Droz, T. Vilaivan, G. W. Weaver, J. J. Park, J. M. Pratt, L. Tweedale and L. R. Kelland, *J. Med. Chem.*, 1999, **42**, 3167; (c) P. J. Carter, C. C. Cheng, H. H. Thorp and J. Am, *Chem. Soc.*, 1998, **120**, 632; (d) A. McCoubrey, H. C. Latham, P. R. Cook, A. Rodger and G. Lowe, *FEBS Lett.*, 1996, **380**, 73.
- 20 (a) D. Zhao, J. You and C. Hu, *Chem.-Eur. J.*, 2011, **17**, 5466; (b) V. F. Slagt, A. H. M. de Vries, J. G. de Vries and R. M. Kellogg, *Org. Process Res. Dev.*, 2010, **14**, 30; (c) R. A. Hughes and C. J. Moody, *Angew. Chem., Int. Ed.*, 2007, **46**, 7930.
- 21 (a) K. L. Billingsley and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2008, **47**, 4695; (b) R. Ghosh, N. N. Adarsh and A. Sarkar, *J. Org. Chem.*, 2010, **75**, 5320; (c) K. Billingsley and S. L. Buchwald, *J. Am. Chem. Soc.*, 2007, **129**, 3358; (d) A. Thakur, K. Zhang and J. Louie, *Chem. Commun.*, 2012, **48**, 203; (e) K. L. Billingsley, K. W. Anderson and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2006, **45**, 3484; (f) T. Noël and A. J. Musacchio, *Org. Lett.*, 2011, **13**, 5180; (g) N. Kudo, M. Perseghini and G. C. Fu, *Angew. Chem., Int. Ed.*, 2006, **45**, 1282.
- 22 (a) S. Lou and G. C. Fu, *Adv. Synth. Catal.*, 2010, **352**, 2081; (b) D. H. Lee, M. Choi, B. W. Yu, R. Ryoo, A. Taher, S. Hossain and M. J. Jin, *Adv. Synth. Catal.*, 2009, **351**, 2912; (c) K. W. Anderson and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2005, **44**, 6173; (d) X.-X. Zhou and L.-X. Shao, *Synthesis*, 2011, 3138; (e) D.-H. Lee, J.-Y. Jung and M.-J. Jin, *Green Chem.*, 2010, **12**, 2024.
- 23 (a) F. Trécourt, B. Gervais, M. Mallet and G. J. Quéguiner, *J. Org. Chem.*, 1996, **61**, 1673; (b) F. Trécourt, B. Gervais, G. J. Quéguiner, O. Mongin, C. Le Gal and F. Mongin, *J. Org. Chem.*, 1998, **63**, 2892.
- 24 (a) O. Dogan, N. Gurbuz, I. Ozdemir and B. Cetinkaya, *Heteroat. Chem.*, 2008, **19**, 569; (b) V. Farina, B. Krishnan, D. R. Marshall and G. P. Roth, *J. Org. Chem.*, 1993, **58**, 5434; (c) O. Dogan, N. Gurbuz, I. Ozdemir and B. Cetinkaya, *J. Heterocycl. Chem.*, 2009, **46**, 186.

- 25 A. L. Korich and T. S. Hughes, *Org. Lett.*, 2008, **10**, 5405.
- 26 S. Debasree, C. Kalicharan and R. Brindaban, *Tetrahedron Lett.*, 2009, **50**, 1003.
- 27 A. K. Steib, O. M. Kuzmina, S. Fernandez, D. Flubacher and P. Knochel, *J. Am. Chem. Soc.*, 2013, **135**, 15346.
- 28 S. Laval, W. Dayoub, L. Pehlivan, E. Metay, D. Delbrayelle, G. Mignani and M. Lemaire, *Tetrahedron Lett.*, 2014, **55**, 23.
- 29 Y. Fei, S. Yi, Z. Lei, X. Qing, Z. Yan and W. Jianbo, *Org. Lett.*, 2011, **13**, 5020.
- 30 Y. Junfeng, L. Sijia, Z. Jian-Feng and Z. Jianrong, *Eur. J. Org. Chem.*, 2012, **2012**, 6248.
- 31 M. Gary and S. Deidre, *Encyclopedia of Reagents for Organic Synthesis*, 2007.
- 32 G. Solladie, P. Hugel and R. Bartsch, *J. Org. Chem.*, 1998, **63**, 3895.
- 33 L. D. Smirnov, M. R. Avezov, V. P. Lezina, R. O. Alieva and K. M. Dyumaev, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1971, 626.
- 34 L. D. Cantin, S. Choi, R. Clark, B. Hentemann, F. Martin, M. Xin, R. Joachim, L. Sidney, X. Ma, A. Christiana, L. Rico and C. Libing, *et al. PCT Int. Appl.*, WO 2004058174 A2, 2004.
- 35 T. Tetsuo, S. Tsuyoshi, S. Fumiyasu, K. Kenichi, S. Yuki, H. Makoto, H. Akira, T. Masato, U. Fumihito and O. Takahiro, *PCT Int. Appl.*, WO 2014021281 A1 20140206, 2014.
- 36 N. A. Isley, F. Gallou and B. H. Lipshutz, *J. Am. Chem. Soc.*, 2013, **135**, 17707.
- 37 L. Lian-Yan, K. Xing-Rui and D. Xin-Fang, *J. Org. Chem.*, 2014, **79**, 777.
- 38 S. Sakashita, M. Takizawa, J. Sugai, H. Ito and Y. Yamamoto, *Org. Lett.*, 2013, **15**, 4308.
- 39 D. X. Yang, S. L. Colletti, K. Wu, M. Song, G. Y. Li and H. C. Shen, *Org. Lett.*, 2009, **11**, 381.
- 40 S. M. Jen and T. K. Yih, *Heterocycles*, 1990, **31**, 637.
- 41 P. Gros and Y. Fort, *J. Chem. Soc., Perkin Trans. 1*, 1998, **21**, 3515.
- 42 D. Meenakshi, M. Debasis and O. Michael, *Chem.-Eur. J.*, 2010, **16**, 4279.