


Arylboration of 1-Arylalkenes by Cooperative Nickel/Copper Catalysis

Kazuhiko Semba,* Yasuhiro Ohtagaki, and Yoshiaki Nakao*

Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Katsura, Nishikyō-ku, Kyoto 615-8510, Japan

Supporting Information




ABSTRACT: A method for the arylboration of 1-arylalkenes with bis(pinacolato)diboron and aryl chlorides or tosylates by cooperative Ni/Cu catalysis has been developed, which affords 2-boryl-1,1-diaryllalkanes in high regio- and stereoselectivity. Under the applied conditions, this method is tolerant toward various functional groups, including silyl ether, alkoxycarbonyl, and aminocarbonyl moieties.

Alkylboranes are useful synthetic intermediates in organic synthesis, as the C–B bonds can be readily converted into C–C, C–O, or C–N bonds.¹ Conventionally, such C–B bonds are prepared via the hydroboration of alkenes² or the nucleophilic substitution of B(OR)₃ with organolithium or organomagnesium reagents.³ Transition-metal-catalyzed carboborations⁴ of alkenes represent another powerful tool for the preparation of alkylboranes, mostly on account of their high functional group tolerance and the readily available variety of alkenes.⁵ To date, both the intra- and intermolecular carboborations of alkenes have been accomplished by catalytic methods based on a single metal. Recently, our group and that of Brown independently reported the intermolecular arylboration of alkenes with bis(pinacolato)diboron (B₂(pin)₂) and aryl halides by cooperative Pd/Cu catalysis.⁶ However, the use of Pd catalysts is not cost-effective, and phenol-derived electrophiles, which are generally more readily available compared to aryl halides, are restricted to aryl triflates. Conversely, aryl tosylates, which are easier to access and handle than aryl triflates, cannot be used as substrates in such reactions. Herein, we report the arylboration of alkenes with aryl chlorides or tosylates and B₂(pin)₂ by cooperative Ni/Cu catalysis.^{7,8}


To test the viability of a cooperative Ni/Cu catalysis, we initially examined the cross-coupling of alkylcopper **1a**,⁹ which was prepared by a borylcupration of styrene (**2a**), with *p*-chlorotoluene (**3a**) (eq 1). Even though **4a** was gratifyingly obtained in 41% yield, alkenylboronate **5a** and toluene were also formed in 40% and 41% yield, respectively. In the absence of Ni(cod)₂, the formation of **4a** was not observed. These preliminary results encouraged us to investigate and develop cross-coupling reactions based further on cooperative Ni/Cu catalysis.

For that purpose, we carried out the arylboration of **2a** with **3a** and B₂(pin)₂ in the presence of Ni(cod)₂ (2.0 mol %), tricyclopentylphosphine (PCyp₃, 4.0 mol %), (IPr)CuCl (1.0 mol %), and LiOtBu (1.5 mmol). The reaction proceeded




catalytically and furnished **4a** in 53% yield. Subsequently, we screened the reaction conditions with respect to various nickel and copper sources, phosphine ligands, and bases. As a result, we found that the arylboration of **2a** (1.0 mmol) with **3a** (1.0 mmol) and B₂(pin)₂ (1.0 mmol) afforded **4a** in 72% yield in the presence of Ni(acac)₂ (2.0 mol %), CuCl (1.0 mol %), PCyp₃ (6.0 mol %), and LiOtBu (1.5 mmol), under concomitant formation of small amounts of **5a**, **6a**, and **7a** (entry 1, Table 1). The effects of varying the monodentate phosphine ligand are summarized in entries 2–6 (Table 1).¹⁰ When PPh₃, PnPr₃, and PtBu₃ were employed, the formation of **4a** was not observed (entries 2, 3, and 6). The use of P*t*Pr₃ and PCy₃ resulted in the formation of **4a**, albeit in lower yield compared to using PCyp₃ (entries 4 and 5). Bidentate ligands such as 1,2-bis(diphenylphosphino)ethane (dppe) and 2,2'-bipyridine (bpy) did not promote the formation of **4a** (entries 7 and 8). Moreover, the presence of LiOtBu was found to be crucial in order to obtain **4a** in good yield. Other bases such as LiOMe, NaOtBu, or KOtBu afforded **4a** in very low yield (entries 9–11). The arylboration of **2a** did not proceed in the absence of Ni(acac)₂, and without CuCl, the yield of **4a**

Received: June 9, 2016



Scheme 1. A Plausible Mechanism for the Arylboration of 1-Arylalkenes by Cooperative Ni/Cu Catalysis



In conclusion, we have developed a method for the arylboration of alkenes with aryl chlorides or tosylates and $\text{B}_2(\text{pin})_2$ by cooperative Ni/Cu catalysis. The reaction proceeds regio- and stereoselectively to afford 2-boryl-1,1-diarylalkanes, which represent potent intermediates for the generation of 1,1-diarylalkanes. This novel Ni/Cu catalysis should provide valuable insights for the development of advanced cooperative base metal catalysis.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b01675.

Detailed experimental procedures including spectroscopic and analytical data (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: semba.kazuhiko.Sn@kyoto-u.ac.jp.

*E-mail: nakao.yoshiaki.8n@kyoto-u.ac.jp.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by a Grant-in-Aid for Young Scientists (B) (26810058) from MEXT and the CREST program "Establishment of Molecular Technology towards the Creation of New Functions" Area from JST.

■ REFERENCES

- (a) Jana, R.; Pathak, T. P.; Sigman, M. S. *Chem. Rev.* **2011**, *111*, 1417. (b) Ohishi, T.; Zhang, L.; Nishiura, M.; Hou, Z. *Angew. Chem., Int. Ed.* **2011**, *50*, 8114. (c) Mlynarski, S. N.; Karns, A. S.; Morken, J. P. *J. Am. Chem. Soc.* **2012**, *134*, 16449. (d) Scott, H. K.; Aggarwal, V. K. *Chem. - Eur. J.* **2011**, *17*, 13124. (e) Thomas, S. P.; French, R. M.; Jheengut, V.; Aggarwal, V. K. *Chem. Rev.* **2009**, *9*, 24.
- (2) For reviews on stoichiometric and catalytic hydroborations, see: (a) Brown, H. C. *Pure. Appl. Chem.* **1976**, *7*, 49. (b) Beletskaya, I.; Pelter, A. *Tetrahedron* **1997**, *53*, 4957. (c) Crudden, C. M.; Edwards, D. *Eur. J. Org. Chem.* **2003**, *2003*, 4695.
- (3) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457.
- (4) For selected examples on the carboboration of alkynes and dienes, see: (a) Yang, F.-Y.; Wu, M.-Y.; Cheng, C.-H. *J. Am. Chem. Soc.* **2000**, *122*, 7122. (b) Sugimoto, M.; Yamamoto, A.; Murakami, M. *J. Am. Chem. Soc.* **2003**, *125*, 6358. (c) Sugimoto, M.; Yamamoto, A.; Murakami, M. *Angew. Chem., Int. Ed.* **2005**, *44*, 2380. (d) Sugimoto, M.; Shirakura, M.; Yamamoto, A. *J. Am. Chem. Soc.* **2006**, *128*, 14438. (e) Yamamoto, A.; Sugimoto, M. *J. Am. Chem. Soc.* **2005**, *127*, 15706. (f) Daini, M.; Yamamoto, A.; Sugimoto, M. *J. Am. Chem. Soc.* **2008**, *130*, 2918. (g) Daini, D.; Sugimoto, M. *Chem. Commun.* **2008**, 5224. (h) Zhou, Y.; You, W.; Smith, K. B.; Brown, M. K. *Angew. Chem., Int. Ed.* **2014**, *53*, 3475. (i) Zhang, L.; Cheng, J.; Carry, B.; Hou, Z. *J. Am. Chem. Soc.* **2012**, *134*, 14314. (j) Alfaro, R.; Parra, A.; Alemán, J.; Ruano, J. L. G.; Tortosa, M. *J. Am. Chem. Soc.* **2012**, *134*, 15165. (k) Okuno, Y.; Yamashita, M.; Nozaki, K. *Angew. Chem., Int. Ed.* **2011**, *50*, 920. (l) Okuno, Y.; Yamashita, M.; Nozaki, K. *Eur. J. Org. Chem.* **2011**, *2011*, 3951. (m) Chen, C.; Kehr, G.; Fröhlich, R.; Erker, G. *J. Am. Chem. Soc.* **2010**, *132*, 13594. (n) Lesieur, M.; Bidal, D. Y.; Lazreg, F.; Nahra, F.; Cazin, C. S. *J. ChemCatChem* **2015**, *7*, 2108. For reviews, see: (o) Sugimoto, M. *Chem. Rev.* **2010**, *10*, 348. (p) Kehr, G.; Erker, G. *Chem. Commun.* **2012**, 48, 1839.
- (5) (a) Kubota, K.; Yamamoto, E.; Ito, H. *J. Am. Chem. Soc.* **2013**, *135*, 2635. (b) Zhong, C.; Kunii, S.; Kosaka, Y.; Sawamura, M.; Ito, H. *J. Am. Chem. Soc.* **2010**, *132*, 11440. (c) Ito, H.; Toyoda, T.; Sawamura, M. *J. Am. Chem. Soc.* **2010**, *132*, 5990. (d) Daini, M.; Sugimoto, M. *J. Am. Chem. Soc.* **2011**, *133*, 4758. (e) Kuznetsov, N. Y.; Starikova, Z. A.; Averkiev, B.; Bubnov, Y. N. *Russ. Chem. Bull.* **2005**, *54*, 678. (f) Yoshida, H.; Kageyuki, I.; Takaki, H. *Org. Lett.* **2013**, *15*, 952. (g) Bubnov, Y. N.; Nesmeyanova, O. A.; Rudashevskaya, T. Y.; Mikhailov, B. M.; Kazansky, B. A. *Tetrahedron Lett.* **1971**, *12*, 2153. (h) Kageyuki, I.; Yoshida, H.; Takaki, H. *Synthesis* **2014**, 46, 1924. (i) Su, W.; Gong, T.-J.; Lu, X.; Xu, M.-Y.; Yu, C.-G.; Xu, Z.-Y.; Yu, H.-Z.; Xiao, B.; Fu, Y. *Angew. Chem., Int. Ed.* **2015**, *54*, 12957. (j) Jia, T.; Cao, P.; Wang, B.; Lou, Y.; Yin, X.; Wang, M.; Liao, J. *J. Am. Chem. Soc.* **2015**, *137*, 13760. (k) Yang, K.; Song, Q. *J. Org. Chem.* **2016**, *81*, 1000.
- (6) (a) Semba, K.; Nakao, Y. *J. Am. Chem. Soc.* **2014**, *136*, 7567. (b) Smith, K. B.; Logan, K. M.; You, W.; Brown, M. K. *Chem. - Eur. J.* **2014**, *20*, 12032. (c) Logan, K. M.; Smith, K. B.; Brown, M. K. *Angew. Chem., Int. Ed.* **2015**, *54*, 5228.
- (7) For leading references on cooperative Ni/Cu catalysis, see: (a) Beletskaya, I. P.; Latyshev, G. V.; Tsvetkov, A. V.; Lukashev, N. V.

Tetrahedron Lett. **2003**, *44*, 5011. (b) Wang, L.; Li, P.; Zhang, Y. *Chem. Commun.* **2004**, 514. (c) Vechorkin, O.; Barmaz, D.; Proust, V.; Hu, X. *J. Am. Chem. Soc.* **2009**, *131*, 12078. (d) Vechorkin, O.; Proust, V.; Hu, X. *Angew. Chem., Int. Ed.* **2010**, *49*, 3061. (e) Yi, J.; Lu, X.; Sun, Y.-Y.; Xiao, B.; Liu, L. *Angew. Chem., Int. Ed.* **2013**, *52*, 12409. (f) Gallego, G.; Brück, A.; Irran, E.; Meier, F.; Kaupp, M.; Driess, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2013**, *135*, 15617. (g) Zarate, C.; Martin, R. *J. Am. Chem. Soc.* **2014**, *136*, 2236. (h) Niwa, T.; Ochiai, H.; Watanabe, Y.; Hosoya, T. *J. Am. Chem. Soc.* **2015**, *137*, 14313. (i) Wang, Y.; Wu, S.-B.; Shi, W.-J.; Shi, Z.-J. *Org. Lett.* **2016**, *18*, 2548.

(8) For selected examples on cooperative nonprecious metal catalysis, see: (a) Karunananda, M. K.; Parmelee, S. R.; Waldhart, G. W.; Mankad, N. P. *Organometallics* **2015**, *34*, 3857. (b) Mazzacano, T. J.; Mankad, N. P. *J. Am. Chem. Soc.* **2013**, *135*, 17258. (c) Jayarathne, U.; Mazzacano, T. J.; Bagherzadeh, S.; Mankad, N. P. *Organometallics* **2013**, *32*, 3986. (d) Shirakawa, E.; Yamagami, T.; Kimura, T.; Yamaguchi, S.; Hayashi, T. *J. Am. Chem. Soc.* **2005**, *127*, 17164. (e) Shirakawa, E.; Ikeda, D.; Masui, S.; Yoshida, M.; Hayashi, T. *J. Am. Chem. Soc.* **2012**, *134*, 272. (f) Zhao, Y.; Weix, D. J. *J. Am. Chem. Soc.* **2014**, *136*, 48. (g) Ackerman, L. K. G.; Anka-Lufford, L. L.; Naodovic, M.; Weix, D. J. *Chem. Sci.* **2015**, *6*, 1115. (h) Zhao, Y.; Weix, D. J. *J. Am. Chem. Soc.* **2015**, *137*, 3237.

(9) For a reference on the preparation of alkylcopper **1a**, see: Laitar, D. S.; Tsui, E. Y.; Sadighi, J. P. *Organometallics* **2006**, *25*, 2405.

(10) When a mixture of pregenerated complexes $\text{Ni}(\text{cod})_2/2\text{PPh}_3$ and $\text{CuCl}/2\text{PCy}_3$ or $\text{Ni}(\text{cod})_2/2\text{PCy}_3$ and $\text{CuCl}/2\text{PPh}_3$ were used in the aryloboration of **2a** with **3h** and $\text{B}_2(\text{pin})_2$, the crossover of PPh_3 and PCy_3 on nickel and copper was observed. See the [Supporting Information](#) for details.

(11) The borylcupration of 1-hexene with $(\text{IPr})\text{CuB}(\text{pin})$ proceeds sluggishly; see: Reference 9.

(12) For stereoselective aryloborations, see: Reference 6c.

(13) For a reference of configurational isomerization of an alkylcopper, see: Morozova, V.; Moriya, K.; Mayer, P.; Knochel, P. *Chem. - Eur. J.* **2016**, *22*, 9962.