

# Multicomponent Coupling Reactions of Two N-Tosyl Hydrazones and **Elemental Sulfur: Selective Denitrogenation Pathway toward** Unsymmetric 2,5-Disubstituted 1,3,4-Thiadiazoles

Zhen Zhou, Yang Liu, Jiangfei Chen, En Yao, and Jiang Cheng\*

School of Petrochemical Engineering, Jiangsu Key Laboratory of Advanced Catalytic Materials & Technology, Jiangsu Province Key Laboratory of Fine Petrochemical Engineering, Changzhou University, Changzhou 213164, P. R. China

**S** Supporting Information



ABSTRACT: A copper-mediated, three-component reaction between two different N-Ts hydrazones and elemental sulfur was developed, leading to a series of unsymmetric 2,5-disubstituted 1,3,4-thiadiazoles in moderate yields with good functional group compatibility. This procedure features the employment of elemental sulfur and the selective denitrogenation between aryl and alkyl aldehyde N-tosyl hydrazones, allowing rapid access to unsymmetric 2,5-disubstituted 1,3,4-thiadiazoles frameworks with chemical diversity and complexity.

1,3,4-Thiadiazoles possess a broad spectrum of biological activities.<sup>1</sup> For example, they could serve as antidepressant,<sup>2</sup> anxiolytic,<sup>3</sup> antimicrobial,<sup>4</sup> antituberculosis,<sup>5</sup> anti-inflammatory,<sup>6</sup> antidiabetic,<sup>7</sup> anticancer,<sup>8</sup> antihypertensive,<sup>9</sup> and antifungal drugs.<sup>10</sup> Therefore, numerous versatile and flexible approaches have been established to access 1,3,4-thiadiazoles, including the sulfuration of the corresponding 1,4-dicarbonyl and acyl precursors (in situ formed or not) by  $P_2S_5$  or Lawesson's reagent,<sup>11</sup> cyclization of thiohydrazine,<sup>12</sup> and transformation from other heterocycles.<sup>13</sup> However, the sulfuration still suffers from harsh reaction conditions. P2S5 and Lawesson's reagent are sensitive to moisture have a strong odor, despite some modifications.<sup>14</sup> Moreover, the synthesis of S-substituted substrate was tedious and, in most cases, ineffective to access the 2,5-disubstituted analogues with carbon-centered groups. Therefore, the development of new methods to access 1,3,4-thiadiazoles via novel pathways and reaction partners remains an urgent goal for organic chemists.

The multicomponent coupling reactions (MCRs) between elemental sulfur and two N-tosyl hydrazones as readily available starting materials show that denitrogenation<sup>15</sup> is an ideal pathway toward such frameworks with chemical diversity and complexity.<sup>16</sup> However, in the case of two N-tosyl hydrazones, a great challenge arises from the selective denitrogen step between two N-tosyl hydrazone molecules, which may result in various distributions of two homoproducts and one crossproduct.<sup>17</sup> Herein, we describe a three-component reaction of elemental sulfur toward unsymmetric 1,3,4-thiadiazoles with 2,5-disubstituted carbon-centered groups via a selective

denitrogen pathway based on the reactivity discrimination between aryl and alkyl aldehyde N-tosyl hydrazones.

Initially, we tested the combination of benzaldehyde N-tosyl hydrazone (1.0 equiv), n-butanal N-tosyl hydrazone (1.2 equiv), and sublimed sulfur in the presence of CuI (20 mol %) and  $K_2CO_3$  (2 equiv) in dimethylacetamide (DMAC, 2 mL) as the model reaction. To our delight, the cross-annulation product 2-phenyl-5-propyl-1,3,4-thiadiazole 3aa via selective denitrogenation was isolated in 50% yield, and no homoannulated isomers were detected at 120 °C after 24 h under air (Table 1, entry 1). Subsequently, other bases such as  $Na_2CO_3$ , <sup>t</sup>BuONa, <sup>t</sup>BuOK, NaOH, Ag<sub>2</sub>CO<sub>3</sub>, and NaH<sub>2</sub>PO<sub>4</sub>, were examined. However, they were all inferior to K<sub>2</sub>CO<sub>3</sub>, generating **3aa** in lower yields (Table 1, entries 2-7). Pleasingly, Cs<sub>2</sub>CO<sub>3</sub> worked well with an acceptable 62% yield (Table 1, entry 8). In the absence of copper, **3aa** was isolated in 44% yield (Table 1, entry 9). Other tested copper, such as CuCl (39%), and CuBr (36%), decreased the reaction efficiency (Table 1, entries 10 and 11). The tested solvents, such as DMF, CH<sub>3</sub>CN, and dioxane, were inferior to DMAC (24-47%, Table 1, entries 12-14). The reaction efficiency decreased under elevated or lowered temperature (Table 1, entry 15), as did the procedure conducted under either N2 or O2, respectively (Table 1, entry 15). Thus, the optimized conditions were established as follows: benzaldehyde N-tosyl hydrazone (0.1 mmol), nbutanal N-tosyl hydrazone (0.12 mmol), and sublimed sulfur

Received: August 29, 2016

Table 1. Screening of the Optimized Reaction Conditions<sup>a</sup>

	NHTs +	NNHTs IL + 🋐 -		→ <sup>n</sup> Pr S Ph
тег <b>1</b> а	a Pn	⊓ 2a		3aa
entry	copper	base	solvent	yield <sup>b</sup> (%)
1	CuI	K <sub>2</sub> CO <sub>3</sub>	DMAC	50
2	CuI	Na <sub>2</sub> CO <sub>3</sub>	DMAC	28
3	CuI	<sup>t</sup> BuONa	DMAC	39
4	CuI	<sup>t</sup> BuOK	DMAC	29
5	CuI	NaOH	DMAC	31
6	CuI	Ag <sub>2</sub> CO <sub>3</sub>	DMAC	<5
7	CuI	NaH <sub>2</sub> PO <sub>4</sub>	DMAC	24
8	CuI	Cs <sub>2</sub> CO <sub>3</sub>	DMAC	62
9		$Cs_2CO_3$	DMAC	44
10	CuCl	Cs <sub>2</sub> CO <sub>3</sub>	DMAC	39
11	CuBr	Cs <sub>2</sub> CO <sub>3</sub>	DMAC	36
12	CuI	$Cs_2CO_3$	DMF	47
13	CuI	$Cs_2CO_3$	CH <sub>3</sub> CN	34
14	CuI	Cs <sub>2</sub> CO <sub>3</sub>	dioxane	24
15	CuI	Cs <sub>2</sub> CO <sub>3</sub>	DMAC	44, <sup>c</sup> 28, <sup>d</sup> 52, <sup>e</sup> 25 <sup>f</sup>

<sup>*a*</sup>Reaction conditions: **1a** (0.12 mmol), **2a** (0.1 mmol),  $S_8$  (0.0625 mmol), CuX (20 mol %) in solvent (2 mL) under air at 100 °C. <sup>*b*</sup>Isolated yield. <sup>*c*</sup>90 °C. <sup>*d*</sup>110 °C. <sup>*c*</sup>Under N<sub>2</sub>. <sup>*f*</sup>Under O<sub>2</sub>.

(0.0625 mmol) in the presence of CuI (20 mol %) and  $Cs_2CO_3$  (0.2 mmol) in dimethylacetamide (DMAC, 2 mL).

After the establishment of the optimized reaction conditions, the substrates scope and limitation of alkyl aldehyde *N*-tosyl hydrazones in the reaction with benzaldehyde *N*-tosyl hydrazone and sublimed sulfur were tested, as shown in Scheme 1. As expected, besides *n*-propyl, this procedure allowed rapid construction of 2-phenyl-1,3,4-thiadiazoles substituted with 5-primary carbon groups, such as *n*-butyl (**3ga**, 54%), *n*-pentyl (**3ha**, 41%), and ethyl (**3ia**, 57%) in moderate yields (**3ga**-ia). Importantly, the secondary carbon groups, such as isobutyl (**3ba**, 51%), isopropyl (**3da**, 60%), 2-





<sup>a</sup>Reaction conditions: **1a** (0.12 mmol), **2a** (0.1 mmol),  $S_8$  (0.0625 mmol),  $Cs_2CO_3$  (0.2 mmol), CuI (20 mol %), DMAC (2 mL), air, 100 °C, 12 h.

ethylethyl (3ca, 52%), cyclohexanyl (3ea, 52%), and (2-methyl)ethyl (3fa, 47%) were facilely introduced to the 5-position of 2-phenyl-1,3,4-thiadiazoles.

Next, the scope of aromatic aldehyde *N*-tosyl hydrazones in the reaction with *n*-butanal *N*-tosyl hydrazone and sublimed sulfur was studied (Scheme 2). This procedure was compatible





<sup>a</sup>Reaction conditions: **1a** (0.12 mmol), **2a** (0.1 mmol),  $S_8$  (0.0625 mmol),  $Cs_2CO_3$  (0.2 mmol), CuI (20 mol %), DMAC (2 mL), air, 100 °C, 12 h.

with some functional groups on the phenyl ring, such as methyl (**3ab**, 51%), methoxy (**3ac**, 53%), trifluoromethyl (**3ad**, 45%), fluoro (**3ae**, 45%), chloro (**3af**, 48%), and cyano (**3ag**, 43%), which provided a facile handle for potentially further functionalization. 2-Naphthaldehyde *N*-tosyl hydrazine was also a good reaction partner, giving the corresponding product **3ai** in 52% yield. Notably, 2-thienaldehyde *N*-tosyl hydrazone served as a reaction partner, and **3aj** was isolated in 55% yield.

Notably, this procedure was also applicable in the synthesis of symmetric 1,3,4-thiadiazoles (Scheme 3). For example, 2,5-diphenyl-1,3,4-thiadiazoles 4 and 2,5-dicyclohexyl-1,3,4-thiadiazoles 5 were isolated in 67% and 44% yields under the standard procedure, respectively.

To gain insight into the mechanism, control experiments were conducted. No <sup>15</sup>N was detected in the 2-phenyl-5isopropyl-1,3,4-thiadiazoles by the reaction of isobutanal *N*-tosyl hydrazone, <sup>15</sup>N-labeled benzaldehyde *N*-tosyl hydrazone, and sublimed sulfur under the standard procedure (Scheme 4, eq 1). However, <sup>15</sup>N was totally incorporated in 2-phenyl-5-



"Reaction conditions: N-Ts hydrazone (0.1 mmol), S $_8$  (0.1 mmol), Cs $_2$ CO $_3$  (0.2 mmol), CuI (20 mol %), DMAC (2 mL), air, 100 °C, 12 h.

### Scheme 4. Preliminary Mechanism Study



isopropyl-1,3,4-thiadiazoles by the reaction of <sup>15</sup>N-labeled isobutanal *N*-tosyl hydrazone, benzaldehyde *N*-tosyl hydrazone, and sublimed sulfur (Scheme 4, eq 2). This result confirmed the selective denitrogenation took place in benzaldehyde *N*tosyl hydrazone rather than isobutanal *N*-tosyl hydrazone. Moreover, 1,4-diphenylformalazine 6 was subjected to the standard procedure and no 1,3,4-thiadiazoles was detected, ruling out the possibility of benzaldehyde 2-butylidenehydrazone as the intermediate (Scheme 4, eq 3).

On the basis of these experimental results, a tentative pathway of this reaction is outlined in Scheme 5. In the case of

#### Scheme 5. Tentative Mechanism



the blank experiment, in step 1, under basic conditions, the sequential elimination of proton and  $Ts^-$  in benzaldehyde *N*-tosyl hydrazone took place leading to a diazo species, which transformed to carbene species by the extrusion of  $N_2$ .<sup>17–19</sup> Then, the reaction of carbene species with elemental sulfur produced benzothialdehyde 7.<sup>20</sup> Meanwhile, in the presence of base, the anion species 8 was formed. In step 2, the nucleophilic attack of 8 to 7 provided intermediate 9. Then, the intramolecular nucleophilic attack of 9 produced intermediate 10. Finally, the elimination of  $Ts^-$  in intermediate 10 delivered intermediate 11, which transformed to the final product by aromatization. In the presence of CuI, the copper carbene may be involved in the procedure.

In conclusion, we have developed a copper-promoted selective denitrogenation MCR between aryl aldehyde *N*-tosyl hydrazone, alkyl aldehyde *N*-tosyl hydrazone, and elemental sulfur toward 2-aryl-5-alkyl-1,3,4-thiadiazoles with chemical diversity and complexity in moderate yields. In this procedure, the selective denitrogenation took place in alkyl *N*-tosyl hydrazone rather than in aryl aldehyde *N*-tosyl hydrazine. Notably, this procedure was also applicable in the synthesis of symmetric 2,5-disubstituted-1,3,4-thiadiazoles.

# ASSOCIATED CONTENT

## **Supporting Information**

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.or-glett.6b02583.

Experimental procedures and NMR spectra (PDF)

## AUTHOR INFORMATION

#### Corresponding Author

\*E-mail: jiangcheng@cczu.edu.cn.

#### Notes

The authors declare no competing financial interest.

#### ACKNOWLEDGMENTS

We thank the National Natural Science Foundation of China (Nos. 21272028 and 21572025), "Innovation & Entrepreneurship Talents" Introduction Plan of Jiangsu Province, the Key University Science Research Project of Jiangsu Province (15KJA150001), Jiangsu Key Laboratory of Advanced Catalytic Materials & Technology (BM2012110), and Advanced Catalysis and Green Manufacturing Collaborative Innovation Center for financial support.

## REFERENCES

(1) For reviews, see: (a) Vadivelu, A.; Gopal, V.; Reddy, C. U. M.; Evanjelene, V. K. J. Chem. Pharm. Res. 2014, 6, 855. (b) Haider, S.; Alam, M. S.; Hamid, H. Eur. J. Med. Chem. 2015, 92, 156. (c) Hu, Y.; Li, C.-Y.; Wang, X.-M.; Yang, Y.-H.; Zhu, H.-L. Chem. Rev. 2014, 114, 5572. (d) Jain, A. K.; Sharma, S.; Vaidya, A.; Ravichandran, V.; Agrawal, R. K. Chem. Biol. Drug Des. 2013, 81, 557. (e) Christoforou, I. C.; Koutentis, P. A.; Michaelidou, S. S. ARKIVOC 2006, 7, 207. (f) Jain, A. K.; Sharma, S.; Vaidya, A.; Ravichandran, V.; Agrawal, R. K. Chem. Biol. Drug Des. 2013, 81, 557. (g) Khazi, I. A. M.; Gadad, A. K.; Lamani, R. S.; Bhongade, B. A. Tetrahedron 2011, 67, 3289.

(2) (a) Jatav, V.; Mishra, P.; Kashaw, S.; Stables, J. P. *Eur. J. Med. Chem.* **2008**, 43, 1945. (b) Sharma, R.; Prasad, Y.; Mishra, G. P.; Chaturvedi, S. C. *Med. Chem. Res.* **2014**, 23, 252.

(3) (a) Clerici, F.; Pocar, D.; Guido, M.; Loche, A.; Perlini, V.; Brufani, M. J. Med. Chem. 2001, 44, 931. (b) Jubie, S.; Dhanabal, P.; Azam, M. A.; Kumar, N. S.; Ambhore, N.; Kalirajan, R. Med. Chem. Res. 2015, 24, 1605.

(4) (a) Manjunatha, K.; Poojary, B.; Kumar, V.; Lobo, P. L.; Fernandes, J.; Chandrashekhar, C. *Pharm. Chem.* **2015**, *7*, 207. (b) Farghaly, T. A.; Abdallah, M. A.; Masaret, G. S.; Muhammad, Z. A. *Eur. J. Med. Chem.* **2015**, *97*, 320.

(5) (a) Kolavi, G.; Hegde, V.; Khazi, I.; Gadad, P. *Bioorg. Med. Chem.* 2006, 14, 3069. (b) Oruc, E. E.; Rollas, S.; Kandemirli, F.; Shvets, N.; Dimoglo, A. S. J. Med. Chem. 2004, 47, 6760.

(6) (a) Salgin-Goksen, U.; Gokhan-Kelekci, N.; Goktas, O.; Koysal, Y.; Kilic, E.; Isik, S.; Aktay, G.; Ozalp, M. *Bioorg. Med. Chem.* **2007**, *15*, 5738. (b) Chidananda, N.; Poojary, B.; Sumangala, V.; Kumari, N. S.; Shetty, P.; Arulmoli, T. *Eur. J. Med. Chem.* **2012**, *51*, 124.

(7) (a) Lee, J.; Lee, S.-H.; Seo, H. J.; Son, E.-J.; Lee, S. H.; Jung, M. E.; Lee, M. W.; Han, H. K.; Kim, J.; Kang, J.; Lee, J. *Bioorg. Med. Chem.* **2010**, 18, 2178. (b) Pattan, S. R.; Kekare, P.; Dighe, N. S.; Nirmal, S. A.; Musmade, D. S.; Parjane, S. K.; Daithankar, A. V. J. Chem. Pharm. Res. 2009, 1, 191.

(8) (a) Kumar, D.; Maruthi Kumar, N.; Chang, K. H.; Shah, K. *Eur. J. Med. Chem.* **2010**, *45*, 4664. (b) Megally Abdo, N. Y.; Kamel, M. M. *Chem. Pharm. Bull.* **2015**, *63*, 369.

(9) (a) Hasui, T.; Matsunaga, N.; Ora, T.; Ohyabu, N.; Nishigaki, N.; Imura, Y.; Igata, Y.; Matsui, H.; Motoyaji, T.; Tanaka, T.; Habuka, N.; Sogabe, S.; Ono, M.; Siedem, C. S.; Tang, T. P.; Gauthier, C.; De Meese, L. A.; Boyd, S. A.; Fukumoto, S. J. Med. Chem. 2011, 54, 8616. (b) Samel, A. B.; Pai, N. R. J. Chin. Chem. Soc. 2010, 57, 1327.

(10) (a) Zoumpoulakis, P.; Camoutsis, C.; Pairas, G.; Sokovic, M.; Glamoclija, J.; Potamitis, C.; Pitsas, A. *Bioorg. Med. Chem.* **2012**, *20*, 1569. (b) Rezki, N.; Al-Yahyawi, A. M.; Bardaweel, S. K.; Al-Blewi, F. F.; Aouad, M. R. *Molecules* **2015**, *20*, 16048.

(11) (a) Mazzone, G.; Puglisi, G.; Bonina, F.; Corsaro, A. J. Heterocycl. Chem. 1983, 20, 1399. (b) Stolle, R.; Kind, W. J. Prakt. Chem. 1904, 70, 423. (c) Augustine, J. K.; Vairaperumal, V.; Narasimhan, S.; Alagarsamy, P.; Radhakrishnan, A. Tetrahedron 2009, 65, 9989.

(12) (a) Ferrari, S.; Morandi, F.; Motiejunas, D.; Nerini, E.; Henrich, S.; Luciani, R.; Venturelli, A.; Lazzari, S.; Calo, S.; Gupta, S.; Hannaert, V.; Michels, P. A. M.; Wade, R. C.; Costi, M. P. J. Med. Chem. 2011, 54, 211. (b) Sayed, A. R. Tetrahedron Lett. 2010, 51, 4490.

(13) (a) Padmavathi, V.; Reddy, S. N.; Reddy, G. D.; Padmaja, A. *Eur. J. Med. Chem.* **2010**, *45*, 4246.

(14) Kaleta, Z.; Makowski, B. T.; Soos, T.; Dembinski, R. Org. Lett. 2006, 8, 1625.

(15) (a) Xiao, L.; Li, Y.; Liao, L.; Liu, L. New J. Chem. 2013, 37, 1874.
(b) Cao, Z.; Hou, B.; Chen, W.; Zhao, Q. Pet. Sci. Technol. 2007, 25, 705.

(16) Reaction of 2 equiv of N-tosyl hydrazone with  $S_2Cl_2$  leaing to symmetric 2,5-disubstituted 1,3,4-thiadiazoles in relative low yields: Okuma, K.; Nagakura, K.; Nakajima, Y.; Kubo, K.; Shioji, K. Synthesis **2004**, 1929.

(17) For reviews on the nitro group retaining reaction of diazo compounds, see: (a) Qiu, D.; Qiu, M.; Ma, R.; Zhang, Y.; Wang, J. *Huaxue Xuebao* **2016**, *74*, 472. For recent examples of the selective nitro group retaining reaction of diazo compounds, see: (b) Xu, G.; Zhu, C.; Gu, W.; Li, J.; Sun, J. *Angew. Chem., Int. Ed.* **2015**, *54*, 883. (c) Guo, H.; Zhang, D.; Zhu, C.; Li, J.; Xu, G.; Sun, J. Org. Lett. **2014**, *16*, 3110.

(18) For reviews on N-Ts hydrazones, see: (a) Zhang, Y.; Wang, J. Top. Curr. Chem. 2012, 327, 239. (b) Jadhav, A. P.; Ray, D.; Rao, V. U. B.; Singh, R. P. Eur. J. Org. Chem. 2016, 2016, 2369. (c) Liu, Z.; Wang, J. J. Org. Chem. 2013, 78, 10024. (d) Liu, Z.; Zhang, Y.; Wang, J. Youji Huaxue 2013, 33, 687. (e) Xiao, Q.; Zhang, Y.; Wang, J. Acc. Chem. Res. 2013, 46, 236. (f) Shao, Z.; Zhang, H. Chem. Soc. Rev. 2012, 41, 560.

(19) For selected recent examples, see: (a) Zhang, Z.; Liu, Y.; Ling, L.; Li, Y.; Dong, Y.; Gong, M.; Zhao, X.; Zhang, Y.; Wang, J. J. Am. Chem. Soc. 2011, 133, 4330. (b) Zhou, L.; Ye, F.; Zhang, Y.; Wang, J. J. Am. Chem. Soc. 2010, 132, 13590. (c) Zhang, Z.; Liu, Y.; Gong, M.; Zhao, X.; Zhang, Y.; Wang, J. Angew. Chem., Int. Ed. 2010, 49, 1139. (d) Zhou, P.-X.; Ye, Y.-Y.; Liang, Y.-M. Org. Lett. 2013, 15, 5080. (e) Liu, X.; Ma, X.; Huang, Y.; Gu, Z. Org. Lett. 2013, 15, 4814. (f) Khanna, A.; Maung, C.; Johnson, K. R.; Luong, T. T.; Van Vranken, D. L. Org. Lett. 2012, 14, 3233. (g) Tsai, A. S.; Curto, J. M.; Rocke, B. N.; Dechert-Schmitt, A.-M. R.; Ingle, G. K.; Mascitti, V. Org. Lett. 2016, 18, 508. (h) Zhou, P.-X.; Ye, Y.-Y.; Ma, J.-W.; Zheng, L.; Tang, Q.; Qiu, Y.-F.; Song, B.; Qiu, Z.-H.; Xu, P.-F.; Liang, Y.-M. J. Org. Chem. 2014, 79, 6627. (i) Wu, X.-X.; Zhou, P.-X.; Wang, L.-J.; Xu, P.-F.; Liang, Y.-M. Chem. Commun. 2014, 50, 3882. (j) Chen, Z.; Yan, Q.; Liu, Z.; Xu, Y.; Zhang, Y. Angew. Chem., Int. Ed. 2013, 52, 13324. (20) Latif, N.; Fathy, I. J. Org. Chem. 1962, 27, 1633.