


Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. Rast, B. Modéc, M. Stephan and B. Mohar, *Org. Biomol. Chem.*, 2016, DOI: 10.1039/C5OB02352A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Organic & Biomolecular Chemistry

PAPER

 γ -Sultam-cored N,N-ligands in ruthenium(II)-catalyzed asymmetric transfer hydrogenation of aryl ketones†Slavko Rast,^a Barbara Modec,^b Michel Stephan^{a,c} and Barbara Mohar^{*a}Received 00th January 20xx,
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

www.rsc.org/

The synthesis of new enantiopure *syn*- and *anti*-3-(α -aminobenzyl)-benzo- γ -sultam ligands **6** and their application in the ruthenium(II)-catalyzed asymmetric transfer hydrogenation (ATH) of ketones using formic acid/triethylamine is described. In particular, benzo-fused cyclic ketones afforded excellent enantioselectivities in reasonable time employing low loading of the *syn* ligand-containing catalyst. A never-before-seen dynamic kinetic resolution (DKR) during reduction of a γ -keto carboxylic ester (**S7**) derivative of 1-indanone is realized leading as well to excellent induction.

Introduction

The efficient asymmetric transfer hydrogenation (ATH) of ketones employing the HCO₂H/Et₃N binary mixture can be currently achieved under mild conditions by three generations of RSO₂-DPEN-based chiral Ru(II) complexes (available in both enantiomeric forms, DPEN = *trans*-1,2-diphenylethylenediamine) (Fig. 1).^{1–5} Noyori and co-workers' chiral [RuCl(TsDPEN)(η^6 -arene)]-type complexes (1st generation) were the starting point of such catalyzed asymmetric transformation both on the applied and fundamental levels.^{2,3} Following, intra covalent tethering of the diamine and η^6 -arene ligand units (2nd and 3rd generations) led to an increased longevity of the catalytic species improving thus the turnover number.^{4,5}

Fig. 1 "SO₂DPEN"-Embedded ATH-efficient chiral Ru(II) complexes.

Exploring the origin of the stereocontrol by the structural stereoarray of the 1st generation ligands and aiming to enhance the enantioselectivity and catalyst activity, empirical modifications of the chiral elements were undertaken (Fig. 2). In particular, TsDPEN skeletal alteration at the level of its ethylene-bridge substituents on position 1 or 2 revealed the critical importance of their aromatic nature (and inherent steric bulk) as well as the advantage of their *anti* disposition.^{2a,6}

^a National Institute of Chemistry, Hajdrihova 19, 1001 Ljubljana, Slovenia.^b Faculty of Chemistry and Chemical Technology, University of Ljubljana, Večna pot 113, 1000 Ljubljana, Slovenia.^c Present address: PhosPhoenix SARL, 115, rue de l'Abbé Groult, 75015 Paris, France.† Electronic Supplementary Information (ESI) available: X-ray crystallographic data for racemic *trans*-**5**, [*syn*-(3*S*,1'*R*)-**6**](*S*)-CSA, [*anti*-(3*R*,1'*R*)-**6**](*S*)-CSA, and **S7** ATH major reduction product, HPLC and GC chromatograms of ATH products, ¹H, ¹³C NMR and HMBC spectra. See DOI: 10.1039/x0xx00000x




Fig. 2 Representative results of [RuCl(TsDPEN or "altered TsDPEN")](*p*-cymene)-catalyzed ATH of acetophenone using HCO₂H/Et₃N 5:2.

In our ongoing research interest in this area, we present hereafter the synthesis of 5-membered cyclic minimalist TsDPEN analogs (Fig. 3) and their investigation in the ATH of various classes of ketones. This new design possesses a partial degree of stereochemical rigidity, maintaining however TsDPEN structural elements of the *vic*-diaryls disposition and the mono *N*-sulfonyl group. Such structure gives rise to two possible pairs of *syn* and *anti* diastereomers which preparation was targeted.





Fig. 3 Possible heterocyclic regioisomers of "structurally-simplified TsDPEN".

Results and discussion

Synthesis of the New 5-Membered Cyclic *N,C*-SO₂-DPEN Ligands. In this study, we opted for a non-asymmetric synthetic strategy emphasizing on a convenient access to these ligands with a late-stage resolution. Thus, the NaOMe-mediated Wittig reaction of sodium *ortho*-formylbenzenesulfonate with benzyltriphenylphosphonium chloride afforded pure sodium (*E*)- β -styrylbenzenesulfonate [(*E*)-1] in 59% yield after recrystallization (Scheme 1). Conversion into the (*E*)-sulfonamide (*E*)-2 followed by epoxidation with *m*-CPBA and reaction with LiOMe in MeOH led to the regioselective formation of the 6-*endo-tet*-cyclized *trans*- δ -sultam *trans*-4 in high overall yield (82%). Such regioselectivity is supported by ¹H, ¹³C-HMBC analysis from a correlation between the proximal aromatic hydrogen of the 1,1-dioxo-benzo-1,2-thiazinane core and the carbon atom bearing the hydroxyl group (see the Supporting Information).⁷ Tandem *in situ* *O*-mesylation–intra-*N*-alkylation furnished in 76% yield the *trans*-configured aziridine-cored product *trans*-5 (for its X-ray structure showing the aromatic rings in *trans* and a chiral angular N atom, see the Supporting Information). Alternatively to this circuitous approach, a straightforward single-step conversion of (*E*)-2 into *trans*-5 via Rh₂(OAc)₄-catalyzed aziridination using PhI(OAc)₂⁸ was achieved in 57% yield. Further on, consecutive aziridine highly regioselective ring-opening with sodium azide in MeCN/H₂O (4:1) and Pd/C-catalyzed hydrogenation gave the *syn*-3-(α -aminobenzyl)-benzo- γ -sultam *syn*-6. The (3*S*,1'*R*)- and (3*R*,1'*S*)-configured enantiomeric ligands **6** were separated by preparative chiral HPLC in >99% ee and 78% combined yields. The absolute configuration of the 1st eluting enantiomer was determined by X-ray of its (*S*)-CSA salt (Fig. 4).⁹

Organic & Biomolecular Chemistry

PAPER



Scheme 1 Preparation of the *syn*-3-(α -aminobenzyl)-benzo- γ -sultam ligand **6** enantiomers.





Fig. 4. ORTEP drawing of the $(1S)$ -camphor-10-sulfonic acid ((S) -CSA) salt of the HPLC 1st eluting *syn*-**6** enantiomer at the 50% probability level ((S) -CSA was omitted for clarity; for full details, see the Supporting Information).

Next, the complementary diastereomer *anti*-3-(α -aminobenzyl)-benzo- γ -sultam *anti*-**6** was prepared analogously, however by resorting to NaHMDS as the base¹⁰ in the Wittig step (Scheme 2). The resulting $\sim 1:1$ (E/Z)-isomeric mixture **1** was directly engaged in the further transformation (via the sulfonamide **2**) into the aziridine **5** upon $\text{Rh}_2(\text{OAc})_4$ -catalyzed aziridination. The *cis* and *trans* diastereomers **5** were separated by silica gel chromatography at this stage of the sequence (28% yield for *cis*-**5**) and the former was converted as above into the corresponding racemic *anti*-3-(α -aminobenzyl)-benzo- γ -sultam *anti*-**6**. Its $(3R,1'R)$ - and $(3S,1'S)$ -configured enantiomers were separated by preparative chiral Supercritical Fluid Chromatography (SFC) affording the ligands in >99% ee and 44% combined yields. The absolute configuration of the 1st eluting enantiomer was equally determined by X-ray of its (S) -CSA salt (Fig. 5).¹¹

Organic & Biomolecular Chemistry

PAPER



Scheme 2 Preparation of the *anti*-3-(α -aminobenzyl)-benzo- γ -sultam ligand **6** enantiomers.




Fig. 5 ORTEP drawing of the (1*S*)-camphor-10-sulfonic acid ((*S*)-CSA) salt of the SFC 1st eluting *anti*-**6** enantiomer at the 50% probability level ((*S*)-CSA was omitted for clarity; for full details, see the Supporting Information).

Evaluation of the New Sulfonamido–Amine Ligands **6 in ATH.** The assessment of the efficiency in ATH of the Ru(II) complexes incorporating the new 3-(α -aminobenzyl)-benzo- γ -sultam ligands was first conducted on the representative benchmark substrates acetophenone (**S1**), ethyl benzoylacetate (**S2**), 1-indanone (**S3**), and α -tetralone (**S4**) (Table 1). The Ru complexes were prepared from [RuCl₂(η^6 -arene)]₂ precursor and the enantiopure sulfonamido–amine ligand **6** (*syn* or *anti*; 1.1 equiv to Ru atom) at 40 °C (1 h) in 1,2-dichloroethane. The catalysts' screening with an S/C = 200 was performed at 40 °C using HCO₂H/Et₃N 5:2.

The outcome of this exploratory profiling clearly revealed the faster reduction rate using the new sulfonamido–amine ligands **6** versus the more flexible TsDPEN and “altered TsDPEN” ligands, or Wills’ conformationally locked indane-

cored sulfonamido–amine ligands of Fig. 2. Also, it was noticed for TsDPEN that supplemental amount of HCO₂H/Et₃N 5:2 was required in order to revitalize the reduction and drive it to completion. However, acetophenone (**S1**) and its α -ethoxycarbonyl-substituted derivative **S2** afforded lower enantioselectivities (up to 85% ee and 92% ee, respectively) to the ones with the TsDPEN reference. Noteworthy, as (*R*)-1-phenylethanol was the major resulting enantiomer, the sense of induction on acetophenone (**S1**) of the *syn*-(3*R*,1'*S*)-**6** and *anti*-(3*R*,1'*R*)-**6** revealed to be in line with the one expected with the *syn*-(1*R*,2*S*)-TsDPEN (as its *syn*-(1*S*,2*R*)-enantiomer led to (*S*)-1-phenylethanol) or observed with *anti*-(*R,R*)-TsDPEN (Fig. 2).

Noticeably, the performance of *syn*-(3*R*,1'*S*)-**6** was particularly good against the benzo-fused ketones, 1-indanone (**S3**) and α -tetralone (**S4**), as up to 99% ee with full conversion was obtained in reasonable times leading as well to (*R*)-configured alcohols.^{2a} Also, (*R*)-configured products (95 to >99% ee) were formed employing the *anti*-(3*R*,1'*R*)-**6** demonstrating hence that the chirality on the C(3) atom (bearing the sulfonamido group) predominantly determines the enantiofacial discrimination.

Table 1 Ru(II)-catalyzed ATH of benchmark ketones with the enantiopure ligands *syn*-(3*R*,1'*S*)-**6** (HPLC 2nd eluting) and *anti*-(3*R*,1'*R*)-**6** (SFC 1st eluting)^a

Ketone	Ligand	η^6 -arene	t (h)	Conv. (%)	Ee (%)
	<i>syn</i> - 6	<i>p</i> -cymene	4 3 ^b	>99 100	78 85
	<i>syn</i> - 6	mesitylene	7	80	72
	<i>anti</i> - 6	<i>p</i> -cymene	4	>99	58
	<i>anti</i> - 6	mesitylene	7	90	72
	TsDPEN	<i>p</i> -cymene	4 4 ^b 20 ^b	25 25 99	97 97 97
	<i>syn</i> - 6	<i>p</i> -cymene	3	100	92
	<i>syn</i> - 6	mesitylene	3	100	86
	<i>anti</i> - 6	<i>p</i> -cymene	3	100	54
	<i>anti</i> - 6	mesitylene	3	100	82
	TsDPEN	<i>p</i> -cymene	7 7 ^b	85 >99	98 98
	<i>syn</i> - 6	<i>p</i> -cymene	4	100	99
	<i>anti</i> - 6	<i>p</i> -cymene	5	>99	95
	<i>anti</i> - 6	mesitylene	7	90	95
	TsDPEN	<i>p</i> -cymene	4 ^b 20 ^b	50 >99	- 99
	<i>syn</i> - 6	<i>p</i> -cymene	3	100	99
	<i>syn</i> - 6	mesitylene	7	80	98
	<i>anti</i> - 6	<i>p</i> -cymene	7	95	>99
	<i>anti</i> - 6	mesitylene	7	85	>99
	TsDPEN	<i>p</i> -cymene	3 3 ^b 20 ^b	25 40 98	99 99 98

^a Reaction conditions: S/C = 200, ketone (1.0 mmol), 1,2-dichloroethane (1 mL), HCO₂H/Et₃N 5:2 (250 μ L), 40 °C. Conversion was determined by ¹H NMR of the extracted crude. Isolated yields were 96–98%. Ees were determined by chiral GC or HPLC. (*R*)-Configured alcohols were obtained [(*R,R*)-TsDPEN used]. For further details, see the Experimental Section. ^b Reaction in neat HCO₂H/Et₃N 5:2 (500 μ L).

Therefore, the [RuCl₂(*p*-cymene)]₂/(3*R*,1'*S*)-**6** complex was further screened on a series of methoxycarbonyl-substituted 1-indanone (**S6–S7**) and methoxycarbonyl-substituted α -tetralone (**S8–S10**) (Table 2).

Table 2 Ru(II)-catalyzed ATH of benzo-fused cyclic ketones with the *syn*-(3*R*,1'*S*)-**6** (HPLC 2nd eluting) ligand^a

Ketone	S/C	t (h)	Conv. (%)	Product	
				<i>cis/trans</i>	Ee (%)
	1000	16	95	-	99
	1000	16	>99	-	99
	1000	12	100	-	99
	200	4	100	97:3	>99 (<i>cis</i>)
	200	6	100	95:5	99 (<i>cis</i>) >99 (<i>trans</i>)
	200	6	>99	98:2	>99 (<i>cis</i>)
	100	6	>99	50:50	99 (<i>cis</i>) ^b
	200	3	100	50:50	>99 (<i>cis</i>) >99 (<i>trans</i>)

^a Reaction conditions: [RuCl₂(*p*-cymene)]₂/(3*R*,1'*S*)-**6**, ketone (1.0 mmol), 1,2-dichloroethane (1 mL), HCO₂H/Et₃N 5:2 (250 μ L), 40 °C. Conversion and *cis/trans* ratio (dr) were determined by ¹H NMR of the extracted crude. Isolated yields were 94–99%. Ees were determined by chiral GC or HPLC. (*R*)-Configured alcohols were obtained in all cases. For further details, see the Experimental Section. ^b Determined by ¹⁹F NMR of the (*R*)-Mosher ester; *trans*-stereoisomers were not separated.

Employing an S/C = 1000, a 99% ee coupled with a high conversion was obtained within 16 h for 1-indanone (**S3**), α -tetralone (**S4**), and 4-chromanone (**S5**).

Shifting to functionalized variants, ATH of racemic 2- or 3-methoxycarbonyl-1-indanone (**S6** and **S7**) using an S/C = 200 resulted in high diastereoselectivities (*cis/trans* 97:3 and 95:5, respectively) and 99% enantioselectivity indicating a dynamic kinetic resolution (DKR)^{12,13} occurring during their reduction. Regioisomeric methoxycarbonyl- α -tetralones **S8–S10** displayed as well an excellent ATH outcome. A high enantioselectivity (>99% ee) was attained in all cases, with **S8** undergoing DKR¹² in 98:2 *cis/trans* ratio. Noteworthy, **S9** exhibited some kinetic resolution (KR) during ATH as, at ~75% conversion, ¹H NMR indicated a 60:40 *cis/trans* ratio.¹⁴


Most interestingly, in a related fashion to the reported DKR during ATH of readily enolizable β -keto esters (such as

ARTICLE

Journal Name

S6 and **S8**),¹² a first-time DKR during reduction of a γ -keto ester as *rac*-3-methoxycarbonyl-1-indanone¹⁵ (**S7**) was encountered (Scheme 3). We hypothesize that this DKR occurs by keto-enol tautomerization of this 5-membered cyclic γ -keto ester via the racemization-prone dual benzylic and allylic carbon. This stereolability could be accentuated in the transition state (TS) influenced by the electrostatic interaction as shown in Scheme 3. We assume that the configuration of the GC-detected *trans*-product (5.3%) is as depicted supported by the 99% ee obtained with 1-indanone (**S3**). By contrast, the 6-membered δ -keto ester higher homolog **S10** is unable to undergo racemization (under the test conditions) of its methoxycarbonyl-borne benzylic carbon precluding thus DKR.

The [Ru(TsDPEN)(η^6 -arene)]-catalyzed ATH mechanism was established by Noyori, Ikariya and co-workers who considered multiple CH... π attractive electrostatic interactions in the Ru(II)-hydride-ketone TS.^[16] In the case of racemic ketone **S7**, a more facile reduction of the (*S*)-enantiomer (vs. the (*R*)-enantiomer) is attributable to the CO₂Me group outwards orientation minimizing the steric interference in the TS.



Scheme 3 DKR of *rac*-3-methoxycarbonyl-1-indanone (**S7**) during ATH (Table 2).

Conclusions

We have successfully prepared diastereo- and enantiopure *syn*- and *anti*-(α -aminobenzyl)-benzo- γ -sultam ligands **6** and explored them in Ru(II)-catalyzed ATH of conventional ketones in the presence of formic acid/triethylamine 5:2. The ATH rate and enantioselectivity using the *syn* isomer ligand were better than with the *anti* isomer. High enantioselectivities (99 to >99% ee) were obtained for the benzo-fused cyclic ketones **S3–S10** based on 1-indanone or

α -tetralone. Namely, a first-time ever DKR of a γ -keto carboxylic ester occurred during reduction. In fact, ketone **S7** derived from 1-indanone afforded the corresponding γ -hydroxy ester in high *cis/trans* ratio (95:5) and excellent ee (99% ee for *cis*, >99% ee for *trans*).

Finally, this study is yet another example of how skeletal changes in a chiral ligand design can translate into unpredictable catalyst properties.

Experimental section

Materials and methods. The following non-commercial ketones were prepared according to literature procedures: 2-methoxycarbonyl-1-indanone¹⁷ and 2-methoxycarbonyl-1-tetralone¹⁸ from 1-indanone and α -tetralone, respectively, using dimethyl carbonate; 3-methoxycarbonyl-1-indanone¹⁹ by esterification of 3-oxo-1-indanecarboxylic acid; 3-methoxycarbonyl-1-tetralone²⁰ starting by a Stobbe condensation of benzaldehyde with dimethyl succinate; 4-methoxycarbonyl-1-tetralone¹⁹ from 2-phenyl-glutaric anhydride.

All reactions were conducted under an inert atmosphere (nitrogen or argon) using anhydrous solvents. HCO₂H/Et₃N 5:2 (azeotrope) was prepared by adding Et₃N (280 mL, 2 mol) to HCO₂H (190 mL, 5 mol) at 0 °C under nitrogen atmosphere and used as such. Analytical thin-layer chromatography (TLC) was performed using Silica Gel 60 F254 pre-coated plates (0.25 mm thickness); *R_f* values are reported and visualization was accomplished by irradiation with an UV lamp (254 nm) and/or staining with KMnO₄ solution. Silica Gel 60 (40–63 μ m) was used for flash column chromatography. ¹H (299.9 MHz; internal Me₄Si) and ¹³C (75.4 MHz; internal CDCl₃, δ 77.00) spectra were recorded for solutions in CDCl₃ if not stated otherwise. HRMS measurements were obtained on a Q-TOF instrument equipped with an orthogonal Z-spray ESI interface.

Sodium (*E*)-2- β -styrylbenzenesulfonate (= sodium *trans*-*o*-stilbenesulfonate) [(*E*)-1**].** To a freshly prepared solution of NaOMe in MeOH (80 mL, 54 mmol) was added 2-formylbenzenesulfonic acid sodium salt (10.20 g, 49.0 mmol) and the mixture cooled on an ice-bath. Following, benzyltriphenylphosphonium chloride (19.05 g, 49.0 mmol) in MeOH (50 mL) was added dropwise. The resulting pale yellow mixture was allowed to reach rt during 1 h and left to stir for 16 h. The mixture was concentrated and the residue suspended in cold H₂O (50 mL). The solid was filtered, washed with cold H₂O (20 mL) and CH₂Cl₂ until TLC showed no triphenylphosphine oxide present. After drying, a white powder (12.5 g) was obtained as an (*E/Z*)-isomeric mixture in 85:15 ratio [by ¹H NMR in DMSO-*d*₆; δ 6.54 (d, *J* = 12.3 Hz) is the characteristic signal for the (*Z*)-isomer and δ 8.25 (d, *J* = 16.6 Hz) for the (*E*)-isomer]. Recrystallization from MeCN led to the title product as a white powder (8.14 g, 59% yield). ¹H NMR (300 MHz, DMSO-*d*₆): δ = 7.13 (d, 16.6 Hz, 1 H), 7.18–7.41 (m, 5 H), 7.50–7.53 (m, 2 H), 7.75–7.81 (m, 2 H), 8.25 (d, *J* = 16.6 Hz, 1 H). ¹³C NMR (76 MHz, DMSO-*d*₆): δ = 125.3, 126.52, 126.58, 127.1, 127.5, 128.1, 128.3, 128.7,

128.9, 134.3, 137.8, 145.7. HRMS–ESI (m/z): $[M]^+$ calcd for $C_{14}H_{11}^{23}NaO_3S$, 282.0327; found, 282.0326.

Sodium (*E/Z*)-2- β -styrylbenzenesulfonate [(*E/Z*)-1]. To a cold (0 °C) suspension of benzyltriphenylphosphonium chloride (11.67 g, 30 mmol) in THF (250 mL) was added dropwise under stirring NaHMDs (2 M in THF, 15 mL, 30 mmol). After stirring for 3 h at rt, the reaction mixture was cooled to –78 °C and a solution of 2-formylbenzenesulfonic acid sodium salt (6.25 g, 30 mmol) in MeOH (70 mL) was added dropwise. After 2 h, the mixture was allowed to reach rt and then concentrated. The residue was suspended in EtOAc/2-PrOH (200 mL), filtered, and purified by column chromatography eluting with hexane/EtOAc/MeOH (7:2:1) then EtOAc/MeOH (4:1). Fractions containing the product/triphenylphosphine oxide were concentrated, and the solid suspended in H₂O (400 mL) and stirred at rt for 3 h. The insoluble matter was filtered off and the filtrate concentrated and dried under high vacuum at 60 °C affording the title product with a ~1:1 *E/Z* ratio [by ¹H NMR in DMSO-*d*₆; δ 6.54 (d, J = 12.3 Hz) is the characteristic signal for the (*Z*)-isomer and δ 8.25 (d, J = 16.6 Hz) for the (*E*)-isomer] (7.70 g, 91% yield) as a pale yellow powder.

(*E*)-2- β -Styrylbenzenesulfonamide [(*E*)-2]. To a cold (0 °C) suspension of (*E*)-1 (6.40 g, 22.68 mmol) in CH₂Cl₂ (160 mL) was added under stirring thionyl chloride (6.5 mL, 90 mmol) followed by DMF (50 μ L). After heating at 50 °C for 5 h, the resulting solution was cooled to 0 °C and added dropwise to a cold (0 °C) solution of NH₄OH (25%, 100 mL)/acetone (50 mL) keeping the internal temperature below 5 °C. Following, the mixture was stirred at 0 °C for 2 h, and CH₂Cl₂ (100 mL) and brine (50 mL) were added. The aqueous layer was re-extracted with CH₂Cl₂ (2 \times 100 mL), and the combined organic layers were dried (Na₂SO₄) and concentrated. The crude was filtered through a bed of silica gel eluting with CH₂Cl₂/EtOAc (4:1). After concentration and trituration with *i*Pr₂O, the title product (5.17 g, 88%) was obtained as colorless crystals; mp 145–146 °C. ¹H NMR (300 MHz, CDCl₃): δ = 4.72 (s, 2 H), 7.07 (d, J = 16 Hz, 1 H), 7.28–7.51 (m, 4 H), 7.50–7.65 (m, 3 H), 7.73 (d, J = 8 Hz, 1 H), 7.97 (d, J = 16 Hz, 1 H), 8.07 (dd, J = 8, 1 Hz, 1 H). ¹³C NMR (76 MHz, CDCl₃): δ = 124.4, 127.1, 127.5, 128.0, 128.2, 128.7, 128.9, 133.0, 134.6, 136.3, 136.5, 138.9. HRMS–ESI (m/z): $[M + H]^+$ calcd for $C_{14}H_{14}NO_2S$, 260.0745; found, 260.0748.

(*E/Z*)-2- β -Styrylbenzenesulfonamide [(*E/Z*)-2]. Prepared from (*E/Z*)-1 (~1:1) (4.00 g, 14.18 mmol) following a similar procedure as for (*E*)-2. The title product (3.17 g, 86% yield; *E/Z* ~1:1) was obtained as an amber-colored oil.

***trans*- α -Sulfamoyl-stilbene oxide (*trans*-3).** To a solution of (*E*)-2 (4.50 g, 17.36 mmol) in CH₂Cl₂ (50 mL) was added at rt a solution of *m*-CPBA (5.00 g, 77%; dried over MgSO₄) in CH₂Cl₂ (50 mL). After stirring for 16 h, the mixture was washed with sat. aq NaHCO₃ (2 \times 100 mL). The organic phase was filtered through a bed of silica gel/Na₂SO₄ and concentrated affording the title product (4.59 g, 96% yield) as a slightly yellowish syrup which crystallized upon standing; mp 86–89 °C. ¹H NMR (300 MHz, CDCl₃): δ = 3.88 (d, J = 2 Hz, 1 H), 4.62 (d, J = 2 Hz, 1 H), 4.87 (br s, 2 H), 7.34–7.51 (m, 6

H), 7.57–7.71 (m, 2 H), 8.04 (d, J = 8 Hz, 1 H). ¹³C NMR (76 MHz, CDCl₃): δ = 59.9, 62.3, 125.7, 126.8, 128.0, 128.2, 128.8, 133.1, 135.5, 136.0, 140.0. HRMS–ESI (m/z): $[M + H]^+$ calcd for $C_{14}H_{14}NO_3S$, 276.0694; found, 276.0689.

***trans*-4-Hydroxy-3-phenyl-1,1-dioxo-benzo-1,2-thiazinane (*trans*-4).** To a cold (0 °C) solution of *trans*-3 (2.00 g, 7.26 mmol) in MeOH (30 mL) was added a freshly prepared solution of LiOMe in MeOH (0.5 M, 14.8 mL, 7.4 mmol) and the resulting solution allowed to reach rt. After heating at 55 °C for 2 h, it was brought to rt and the concentrated residue was partitioned between CH₂Cl₂ (100 mL) and 0.5 M HCl (40 mL). The aq layer was re-extracted with CH₂Cl₂ (20 mL), and the combined organic layers were filtered through a bed of silica gel/Na₂SO₄ and concentrated affording the title product (1.94 g, 97% yield) as a white crispy foam. The structure is supported by ¹H, ¹³C-HMBC analysis from a correlation between the proximal aromatic hydrogen of the 1,1-dioxo-benzo-1,2-thiazinane core and the carbon atom bearing the hydroxyl group (see the Supporting Information). ¹H NMR (300 MHz, CDCl₃): δ = 2.30 (d, J = 6 Hz, 1 H), 4.56–4.68 (m, 1 H), 4.92 (d, J = 9 Hz, 1 H), 5.12 (dd, J = 10 and 5 Hz, 1 H), 7.41–7.55 (m, 6 H), 7.60–7.70 (m, 1 H), 7.77–7.83 (m, 1 H), 7.86 (dd, J = 8 and 1 Hz, 1 H). ¹³C NMR (76 MHz, CDCl₃): δ = 63.7, 69.8, 123.0, 126.8, 127.7, 128.4, 129.1, 129.2, 132.6, 137.1, 137.3, 137.4. HRMS–ESI (m/z): $[M + H]^+$ calcd for $C_{14}H_{14}NO_3S$, 276.0694; found, 276.0690.

***trans*-2,3-Benzylidene-1,1-dioxo-benzo-1,2-thiazolidine (*trans*-5).** To a cold (0 °C) solution of *trans*-4 (1.50 g, 5.45 mmol) in THF (30 mL) was added Et₃N (1.7 mL, 12 mmol) followed by dropwise addition of methanesulfonyl chloride (0.69 g, 6.0 mmol). The resulting white suspension was left to stir at 0 °C for 0.5 h then at rt for 1 h. The mixture was quenched at 0 °C with H₂O (20 mL), EtOAc (20 mL) was added and layers were separated. The aq layer was re-extracted with EtOAc (2 \times 20 mL) and the combined organic layers were washed with brine (20 mL), dried (Na₂SO₄) and concentrated. The crude was purified on silica gel eluting with petroleum ether (40–60)/EtOAc (9:1, 8:2, 7:3 then 6:4) affording the title product (1.07 g, 76% yield) as colorless crystals. Its structure was determined by X-ray analysis (see the Supporting Information). ¹H NMR (300 MHz, CDCl₃): δ = 3.48 (d, J = 3 Hz, 1 H), 4.18 (d, J = 3 Hz, 1 H), 7.39 (m, 5 H), 7.54–7.70 (m, 3 H), 7.74–7.81 (m, 1 H). ¹³C NMR (76 MHz, CDCl₃): δ = 51.2, 57.5, 123.4, 125.5, 126.4, 128.7, 129.0, 130.3, 133.5, 133.6, 134.0, 136.0. HRMS–ESI (m/z): $[M + H]^+$ calcd for $C_{14}H_{12}NO_2S$, 258.0589; found, 258.0586.

***cis*-2,3-Benzylidene-1,1-dioxo-benzo-1,2-thiazolidine (*cis*-5).** To a degassed solution of (*E/Z*)-2 (~1:1) (4.00 g, 15.4 mmol) in CH₂Cl₂ (160 mL) was successively added at rt Al₂O₃ (3.9 g, 38.6 mmol), molecular sieves 4 Å (4 g), Rh₂(OAc)₄ (341 mg, 0.77 mmol) and PhI(OAc)₂ (7.46 g, 23.17 mmol). The resulting mixture was stirred at 40 °C for 4 h, brought to rt and filtered through a plug of silica gel eluting with CH₂Cl₂. The crude was purified on silica gel eluting with toluene/EtOAc (24:1) affording *trans*-5 (0.97 g, 25% yield) and *cis*-5 (1.1 g, 28% yield) as an oil. *cis*-5: ¹H NMR (300 MHz, CDCl₃): δ = 4.31 (d, J = 5 Hz, 1 H), 4.62 (d, J = 5 Hz, 1 H),

6.92–7.02 (m, 2 H), 7.03–7.21 (m, 3 H), 7.35–7.52 (m, 2 H), 7.61–7.70 (m, 1 H), 7.71–7.82 (m, 1 H). ^{13}C NMR (76 MHz, CDCl_3): δ = 47.3, 55.8, 122.5, 126.4, 127.2, 127.8, 128.5, 130.3, 131.0, 133.2, 134.1, 137.1. HRMS–ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{14}\text{H}_{12}\text{NO}_2\text{S}$, 258.0589; found, 258.0588.

syn-3-(α -Aminobenzyl)-1,1-dioxo-benzo-1,2-thiazolidine (syn-6). A solution of *trans*-5 (910 mg, 3.53 mmol) and NaN_3 (0.56 g, 8.56 mmol) in $\text{MeCN}/\text{H}_2\text{O}$ (4:1, 30 mL) was heated at 55 °C for 1 h, then brought to rt and concentrated. The residue was partitioned between H_2O (pH 4.5) and CH_2Cl_2 , and the combined organic layers were dried (Na_2SO_4) and concentrated. To the resulting oil (1.3 g) in EtOAc (30 mL), 5% Pd/C (130 mg) was added and the mixture was hydrogenated using Parr apparatus at 20 psig H_2 for 1 h. After filtration through Celite and concentration, the residue was purified on silica gel eluting successively with CH_2Cl_2 , $\text{CH}_2\text{Cl}_2/\text{EtOAc}$ (4:1 then 1:2) and EtOAc/EtOH (4:1), affording *syn*-6 (910 mg, 94% yield) as a yellowish syrup. ^1H NMR (300 MHz, CDCl_3): δ = 2.11 (br s, 3 H), 4.36 (d, J = 6 Hz, 1 H), 4.90 (d, J = 6 Hz, 1 H), 7.02–7.19 (m, 1 H), 7.28–7.62 (m, 7 H), 7.66–7.86 (m, 1 H). ^{13}C NMR (76 MHz, CDCl_3): δ = 59.0, 63.1, 121.4, 125.8, 127.1, 128.2, 128.9, 129.6, 132.5, 136.6, 137.0, 140.6. HRMS–ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{14}\text{H}_{15}\text{N}_2\text{O}_2\text{S}$, 275.0854; found, 275.0849. The *syn*-6 enantiomers were separated by preparative HPLC on ChiralPak IC (5 μm , 250 \times 30 mm), mobile phase $\text{CH}_2\text{Cl}_2/\text{EtOH}$ (98:2), flow rate 42.5 mL/min, 25 °C, UV detection at 260 nm. Analytical HPLC conditions: ChiralPak IC (5 μm , 250 \times 4.6 mm), mobile phase: $\text{CH}_2\text{Cl}_2/\text{EtOH}$ (98:2), flow rate: 1 mL/min, 25 °C, UV detection at 230 nm.²¹

1st eluting *syn* enantiomer (3S,1'R)-6: beige-colored powder, 374 mg (40%); t_R 15.9 min; >99% ee; $[\alpha]_D^{32}$ –11.5 (c 1.0 in CHCl_3); chemical purity (area at 230 nm) = 96.9%. The indicated absolute configuration was determined by X-ray analysis of the (1S)-camphor-10-sulfonic acid salt (colorless plates from MeOH).

2nd eluting *syn* enantiomer (3R,1'S)-6: beige-colored powder, 373 mg (38%); t_R 26.9 min; >99% ee; $[\alpha]_D^{25}$ +11.6 (c 1.0 in CHCl_3); chemical purity (area at 230 nm) = 93.3%.

anti-3-(α -Aminobenzyl)-1,1-dioxo-benzo-1,2-thiazolidine (anti-6). Prepared from *cis*-5 (1.10 g, 4.27 mol) following a similar procedure as for *syn*-6. The title compound (1.06 g, 90% yield) was obtained as a yellowish crispy foam. ^1H NMR (300 MHz, CDCl_3): δ = 4.25 (d, J = 6 Hz, 1 H), 4.79 (d, J = 6 Hz, 1 H), 5.51 (br s, 1 H), 6.58–6.95 (m, 1 H), 7.29–7.63 (m, 7 H), 7.68–7.99 (m, 1 H). ^{13}C NMR (76 MHz, CDCl_3): δ = 59.3, 63.7, 121.4, 124.9, 127.2, 128.4, 129.0, 129.6, 132.7, 136.5, 137.5, 141.2. HRMS–ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{14}\text{H}_{15}\text{N}_2\text{O}_2\text{S}$, 275.0854; found, 275.0858. The *anti*-6 enantiomers were separated by preparative SFC on ChiralPak AY (20 μm , 300 \times 30 mm), mobile phase 'A' for CO_2 and 'B' for 2-PrOH (0.1% NH_3 in H_2O), gradient 'B' 40%; flow rate 70 mL/min, back pressure 100 bar, 38 °C, UV detection at 220 nm. Analytical SFC conditions: ChiralPak AY (150 \times 4.6 mm), mobile phase 'A' for CO_2 and 'B' for EtOH (0.05% Et_2NH), gradient 'B' 40%, flow rate 2.4 mL/min, UV detection at 210 nm.²²

1st eluting *anti* enantiomer (3R,1'R)-6: slightly greenish powder, 177 mg (20%); t_R 1.82 min; >99% ee; $[\alpha]_D^{23}$ –7.9 (c 1.0 in CHCl_3); chemical purity (area at 210 nm) = 100%. The indicated absolute configuration was determined by X-ray analysis of the (1S)-camphor-10-sulfonic acid salt (colorless needles from EtOAc/MeOH).

2nd eluting *anti* enantiomer (3S,1'S)-6: slightly greenish powder, 215 mg (24%); t_R 2.87 min; >99% ee; $[\alpha]_D^{23}$ +7.7 (c 1.0 in CHCl_3); chemical purity (area at 210 nm) = 100%.

General ATH procedure for ketones listed in Tables 1 and 2.

Catalyst preparation. A mixture of the enantiopure ligand **6** (1.1 equiv to Ru atom) and $[\text{RuCl}_2(\eta^6\text{-arene})]_2$ in 1,2-dichloroethane (1 mL/1.0 mmol of ketone) was degassed with N_2 back-filling and stirred at 40 °C for 1 h. The resulting orange-colored solution of the catalyst was used directly in ATH or was concentrated to dryness when the ATH was performed in neat $\text{HCO}_2\text{H}/\text{Et}_3\text{N}$ 5:2.

ATH of ketones. To the above preformed solution of $[\text{RuCl}(\text{ligand } \mathbf{6})(\eta^6\text{-arene})]$ complex (1 mL; with S/C = 100: 0.01 mmol Ru atom; with S/C = 200: 0.005 mmol Ru atom; with S/C = 1000: 0.001 mmol Ru atom) was added $\text{HCO}_2\text{H}/\text{Et}_3\text{N}$ 5:2 (250 μL ; 500 μL when no cosolvent was used) followed by the ketone (1.0 mmol). This mixture was stirred at 40 °C with continuous mild N_2 sweeping. The reaction progress was monitored by ^1H NMR. Workup: the reaction mixture was partitioned between Et_2O (5 mL) and H_2O (5 mL). The aq layer was re-extracted with Et_2O (5 mL) and the combined organic layers were filtered through a bed of silica gel/ Na_2SO_4 and concentrated.

Preparation of racemic alcohol standards. Samples of the racemic alcohols were prepared by reduction of the corresponding ketones with NaBH_4 (3 equiv) in MeOH (EtOH in the case of ethyl benzoylacetate) at 0 °C to rt. Workup: H_2O was added, the mixture neutralized with aq HCl, and the product extracted with Et_2O . The Et_2O layer was successively washed with H_2O and brine, filtered through a bed of silica gel/ Na_2SO_4 , and concentrated.

Characterization and ee determination of the ATH prepared chiral alcohols.

(R)-1-Phenylethanol. Colorless oil (117 mg, 96% yield); 78% ee. ^1H NMR (300 MHz, CDCl_3): δ = 1.47 (d, J = 6.4 Hz, 3 H), 2.04 (br s, 1 H), 4.86 (q, J = 6.4 Hz, 1 H), 7.35–7.33 (m, 5 H). Ee was determined by chiral GC analysis^{23a} on Chiralsil-DEX CB column (25 $\text{m} \times 0.25$ mm), 120 °C (isothermal), t_R 4.6 min (R), 4.8 min (S).

Ethyl (R)-3-hydroxy-3-phenylpropionate. Colorless oil (190 mg, 98% yield); 92% ee. ^1H NMR (300 MHz, CDCl_3): δ = 1.27 (t, J = 7.2 Hz, 3 H), 2.65–2.83 (m, 2 H), 3.00 (q, J = 7.2 Hz, 1 H), 4.19 (q, J = 7.1 Hz, 2 H), 5.14 (dd, J = 8.3 and 4.5 Hz, 1 H), 7.28–7.42 (m, 5 H). Ee was determined by chiral GC analysis^{23b} on Chiralsil-DEX CB (25 $\text{m} \times 0.25$ mm), 140 °C (isothermal); t_R 14.6 min (S), 14.9 min (R).

(R)-1-Indanol. Colorless oil which crystallized upon standing (130 mg, 97% yield); 99% ee. ^1H NMR (300 MHz, CDCl_3): δ = 1.75 (br s, 1 H), 1.86–2.02 (m, 1 H), 2.41–2.58 (m,

1 H), 2.74–2.91 (m, 1 H), 3.06 (m, J = 15.9, 8.5, 4.8 Hz, 1 H), 5.25 (t, J = 6.0 Hz, 1 H), 7.17–7.26 (m, 3 H), 7.33–7.47 (m, 1 H). Ee was determined by chiral HPLC analysis^{23a} on Chiralcel OD column (25 cm). Eluent hexane/2-PrOH (98:2), flow rate 1.0 mL/min, UV detection at 254 nm; racemate: t_R 22.6 min (S), 26.2 min (R).

(R)-1-Tetralol. Colorless oil (194 mg, 96% yield); 99% ee. ¹H NMR (300 MHz, CDCl₃): δ = 1.70 (br s, 1 H), 1.74–2.06 (m, 4 H), 2.66–2.89 (m, 2 H), 4.66–4.90 (m, 1 H), 7.05–7.15 (m, 1 H), 7.15–7.24 (m, 2 H), 7.43 (dd, J = 5.1, 3.9 Hz, 1 H). Ee was determined by chiral GC analysis^{23a} on Chiralsil-DEX CB column (25 m \times 0.25 mm), 130 °C (isothermal); racemate: t_R 13.3 min (S), 13.8 min (R).

(R)-4-Chromanol. White crystals (145 mg, 97% yield); 99% ee. ¹H NMR (300 MHz, CDCl₃): δ = 1.72 (br s, 1 H), 1.95–2.22 (m, 2 H), 4.19–4.40 (m, 2 H), 4.80 (t, J = 4.0 Hz, 1 H), 6.77–7.02 (m, 2 H), 7.17–7.24 (m, 1 H), 7.31 (dd, J = 7.6, 1.4 Hz, 1 H). Ee was determined by chiral HPLC analysis^{23a} on Chiralcel OD-H column (25 cm). Eluent hexane/2-PrOH (95:5), flow rate 1.0 mL/min, UV detection at 220 nm; racemate: t_R 10.0 min (S), 11.7 min (R).

(1R,2R)-2-Methoxycarbonyl-1-indanol. Yellowish oil which crystallized upon standing (186 mg, 97% yield); *cis/trans* 97:3 by ¹H NMR; >99% ee (*cis*). ¹H NMR (300 MHz, CDCl₃): δ = 2.77 (d, J = 6.1 Hz, 1 H), 2.97–3.21 (m, 1 H), 3.34–3.53 (m, 2 H), 3.79 (s, 3 H), 5.35 (t, J = 5.9 Hz, 1 H), 7.23–7.32 (m, 3H), 7.39–7.48 (m, 1H). Ee was determined by chiral GC analysis^{12,23a} on Chiralsil-DEX CB column (25 m \times 0.25 mm), 130 °C (isothermal); racemates (*cis/trans* = 55:45): t_R 33.7 min [*cis*-(R,R)], 34.8 min [*cis*-(S,S)], 45.0 min (*trans*-1), 50.0 min (*trans*-2).

(1R,3S)-3-Methoxycarbonyl-1-indanol. Slightly yellowish powder (184 mg, 96% yield); *cis/trans* 95:5 by ¹H NMR; 98.8% ee (*cis*), >99% ee (*trans*); mp 105–108 °C; [α]_D²⁰ –37.5 (c 1.15 in CHCl₃). ¹H NMR (CDCl₃): δ = 2.20 (m, 0.05 H, *trans*), 2.33 (app t, J = 14.5, 8.0 and 6.8 Hz, 1 H), 2.59 (ddd, J = 14.5, 8.0 and 6.8 Hz, 1 H), 2.84 (m, 0.05 H, *trans*) 3.14 (br s, 1 H), 3.72 (s, 0.15 H, *trans*), 3.76 (s, 3 H), 4.02 (dd, J = 8.0 and 2.8 Hz, 1 H), 4.24 (dd, 0.05 H, *trans*), 5.16 (m, 1 H), 5.46 (t, 0.05 H, *trans*), 7.30–7.39 (m, 3 H), 7.50–7.53 (m, 1 H). ¹³C NMR (CDCl₃): δ = 38.5, 48.4, 52.6, 75.2, 124.8, 125.4, 128.6, 128.9, 140.3, 145.4, 175.8. HRMS–ESI (m/z): [M + Na]⁺ calcd for C₁₁H₁₂O₃²³Na, 215.0684; found, 215.0686. Ees were determined by chiral GC analysis on Chiralsil-DEX CB column (25 m \times 0.25 mm), 130 °C (isothermal); racemates (*cis/trans* = 97:3): t_R 16.8 min [*cis*-(1S,3R)], 18.6 min [*cis*-(1R,3S)], 24.1 min (*trans*-1), 25.2 min (*trans*-2). Recrystallization from hexane/CH₂Cl₂ afforded colorless needles (162 mg, 84% yield, >99.9% de, >99.9% ee) which were used for absolute configuration determination by X-ray analysis (see the Supporting Information).

(1R,2R)-2-Methoxycarbonyl-1-tetralol. Colorless oil (202 mg, 98% yield); *cis/trans* = 97.6:2.4 by ¹H NMR; 99.4% ee (*cis*). ¹H NMR (300 MHz, CDCl₃): δ = 2.07–2.31 (m, 2 H), 2.75–2.99 (m, 4 H), 3.70 (s, 3 H), 4.80 (d, J = 9.0 Hz; CH, *trans* diastereomer), 5.05 (m, 1 H), 7.12–7.14 (m, 1 H), 7.19–7.25 (m, 2 H), 7.37–7.42 (m, 1 H). Ee was determined by chiral

HPLC analysis on Chiralpak IB-3 column (25 cm). Eluent hexane/2-PrOH (97:3), flow rate 1.0 mL/min, UV detection at 220 nm; racemates (mixture prepared by weighing both corresponding ATH products derived from enantiomeric catalysts): t_R 15.4 min (*trans*-1), 17.4 min [*cis*-(R,R)], 19.7 min (*trans*-2), 32.7 min [*cis*-(S,S)].¹²

(1R,3RS)-3-Methoxycarbonyl-1-tetralol. Light violet-colored syrup (204 mg, 99% yield); *cis/trans* = 50:50 by ¹H NMR; 99% ee (*cis*). ¹H NMR (300 MHz, CDCl₃): δ = 1.70–1.71 (m, 1 H), 1.92–2.04 (m, 2 H), 2.22 (d, J = 8.1 Hz, 1 H), 2.35–2.42 (m, 1 H), 2.44–2.52 (m, 1 H), 2.85–3.19 (m, 6 H), 3.73 and 3.75 (2s, 2 \times 3 H), 4.82–4.91 (m, 2 H), 7.10–7.28 (m, 6 H), 7.32–7.37 (m, 1 H), 7.54–7.56 (m, 1 H). HRMS–ESI (m/z): [M + Na]⁺ calcd for C₁₂H₁₄O₃²³Na, 229.0841; found, 229.0846. Racemates (*cis/trans* = 96:4). Ee was determined by ¹⁹F NMR (CDCl₃) of the (R)-Mosher ester (prepared from (S)-Mosher's acid chloride): δ 71.95 [*cis*-(1R,3R)], 72.00 [*cis*-(1S,3S)], 72.24 (*trans*; stereoisomers not separated). Absolute configuration was assigned based on the general observed trend of enantioselectivity.

(1R,4RS)-4-Methoxycarbonyl-1-tetralol. Colorless syrup (202 mg, 98% yield); *cis/trans* = 50:50 by ¹H NMR; >99% ee (*cis*), >99% ee (*trans*). ¹H NMR (300 MHz, CDCl₃): δ = 1.72–1.80 (m, 1 H), 2.31–1.88 (m, 7 H), 2.62 (br s, 2 H), 3.67 and 3.69 (2s, 2 \times 3 H), 3.75–3.83 (m, 2 H), 4.65–4.73 (m, 2 H), 7.23–7.27 (m, 6 H), 7.39–7.47 (m, 2 H). HRMS–ESI (m/z): [M + Na]⁺ calcd for C₁₂H₁₄O₃²³Na, 229.0841; found, 229.0845. Ees were determined by chiral GC analysis on Chiralsil-DEX CB column (25 m \times 0.25 mm), 150 °C (isothermal); racemates (*cis/trans* = 72:28): t_R 21.1 min [*cis*-(1R,4S)], 21.9 min [*trans*-(1R,4R)], 22.7 min [*cis*-(1S,4R)], 24.4 min [*trans*-(1S,4S)]. Absolute configuration was assigned based on the general observed trend of enantioselectivity.

Acknowledgements

This work was supported by the Ministry of Higher Education, Science, and Technology of the Republic of Slovenia (grant number P1-242).

Notes and references

- For reviews, see: (a) S. Gladiali and E. Alberico, *Chem. Soc. Rev.*, 2006, **35**, 226–236; (b) T. Ikariya, K. Murata and R. Noyori, *Org. Biomol. Chem.*, 2006, **4**, 393–406; (c) T. Ikariya and A. J. Blacker, *Acc. Chem. Res.*, 2007, **40**, 1300–1308; (d) C. Wang, X. Wu and J. Xiao, *Chem. Asian J.*, 2008, **3**, 1750–1770; (e) D. Wang and D. Astruc, *Chem. Rev.*, 2015, **115**, 6621–6686; (f) F. Foubelo, C. Najera and M. Yus, *Tetrahedron: Asymmetry*, 2015, **15–16**, 769–790.
- For various ATH ArSO₂-DPEN ligands, see: (a) A. Fujii, S. Hashiguchi, N. Uematsu, T. Ikariya and R. Noyori, *J. Am. Chem. Soc.*, 1996, **118**, 2521–2522; therein, {Ru[(S,S)-TsDPEN](mesitylene)} (S/C = 200, 28 °C, 48 h) led with 1-indanone (**53**) or α -tetralone (**54**) to 99% ee (S) (>99% conv.) but testing **54** at 60 °C led to 98% ee (>99% conv.) within 6 h. (b) Y. Ma, H. Liu, L. Chen, X. Cui, J. Zhu and J. Deng, *Org. Lett.*, 2003, **5**, 2103–2106; (c) D. Šterk, M. Stephan and B. Mohar, *Tetrahedron Lett.*, 2004, **45**, 535–537; (d) P. N. Liu, P. M. Gu, F. Wang and Y.

- Q. Tu, *Org. Lett.*, 2004, **6**, 169–172; (e) Z. Zhou, Y. Sun and A. Zhang, *Cent. Eur. J. Chem.*, 2011, **9**, 175–179.
- 3 (a) B. Mohar, A. Valleix, J.-R. Desmurs, M. Felemez, A. Wagner and C. Mioskowski, *Chem. Commun.*, 2001, **24**, 2572–2573; (b) D. Šterk, M. Stephan and B. Mohar, *Tetrahedron: Asymmetry*, 2002, **13**, 2605–2608; (c) Z. Luo, F. Qin, S. Yan and X. Li, *Tetrahedron: Asymmetry*, 2012, **23**, 333–338; (d) W.-W. Wang, Z.-M. Li, L. Su, Q.-R. Wang and Y.-L. Wu, *J. Mol. Cat. A: Chem.*, 2014, **387**, 92–102; (e) X. Liu, C. Chen, Y. Xiu, A. Chen, L. Guo, R. Zhang, J. Chen and Z. Hou, *Catal. Commun.*, 2015, **67**, 90–94.
- 4 For 2nd generation Ru(II) complexes, see: (a) J. Hannedouche, G. J. Clarkson and M. Wills, *J. Am. Chem. Soc.*, 2004, **126**, 986–987; (b) A. Kišić, M. Stephan and B. Mohar, *Org. Lett.*, 2013, **15**, 1614–1617; (c) A. Kišić, M. Stephan and B. Mohar, *Adv. Synth. Catal.*, 2014, **356**, 3193–3198.
- 5 For 3rd generation Ru(II) complexes, see: (a) A. M. Hayes, D. J. Morris, G. J. Clarkson and M. Wills, *J. Am. Chem. Soc.*, 2005, **127**, 7318–7319; (b) T. Touge, T. Hakamata, H. Nara, T. Kobayashi, N. Sayo, T. Saito, Y. Kayaki and T. Ikariya, *J. Am. Chem. Soc.*, 2011, **133**, 14960–14963; (c) R. Soni, K. E. Jolley, G. J. Clarkson and M. Wills, *Org. Lett.*, 2013, **15**, 5110–5113; (d) R. Soni, T. H. Hall, B. P. Mitchell, M. R. Owen and M. Wills, *J. Org. Chem.*, 2015, **80**, 6784–6793; (e) A. Kišić, M. Stephan and B. Mohar, *Adv. Synth. Catal.*, 2015, **357**, 2540–2546.
- 6 (a) A. Hayes, G. Clarkson and M. Wills, *Tetrahedron: Asymmetry*, 2004, **15**, 2079–2084; (b) B. Zhang, H. Wang, G.-Q. Lin and M.-H. Xu, *Eur. J. Org. Chem.*, 2011, 4205–4211; (c) M. J. Palmer, J. A. Kenny, T. Walsgrove, A. M. Kawamoto and M. Wills, *J. Chem. Soc., Perkin Trans. 1*, 2002, 416–427; (d) P. Roszkowski, J. K. Maurin and Z. Czarnocki, *Tetrahedron: Asymmetry*, 2013, **24**, 643–650.
- 7 The preparation of (*R*)-3-hydroxymethyl-1,1-dioxo-benzo-1,2-thiazolidine was described following a related synthetic strategy. Therein, in addition to a 5-*exo*-type cyclization, the formation of *trans*-(3*R*,4*S*)-4-hydroxy-3-hydroxymethyl-1,1-dioxo-benzo-1,2-thiazinane derived from 6-*endo*-type cyclization occurred as well. For this, see: K. H. Ahn, H. H. Baek, S. J. Lee and C.-W. Cho, *J. Org. Chem.*, 2000, **65**, 7690–7696.
- 8 J.-L. Liang, S.-X. Yuan, P. W. H. Chan and C.-M. Che, *Org. Lett.*, 2002, **4**, 4507–4510.
- 9 C₂₄H₃₀N₂O₆S₂, *F_w* 506.62, monoclinic, space group *P* 2₁ (No. 4), *a* = 12.5236(3), *b* = 7.53800(17), *c* = 12.9539(3) Å, *β* = 92.676(2)°, *V* = 1221.55(5) Å³, *Z* = 2, *T* = 150(2) K, *d*_{calcd} = 1.377 g cm⁻³, *μ* = 0.261 mm⁻¹, 11729 measured reflections, 5599 unique reflections (*R*_{int} = 0.0280), 324 refined parameters, *R*₁ [*I* > 2σ(*I*)] = 0.0482, *wR*₂ [all data] = 0.1230.
- 10 P. A. Byrne and D. G. Gilheany, *J. Am. Chem. Soc.*, 2012, **134**, 9225–9239.
- 11 C₂₄H₃₀N₂O₆S₂, *F_w* 506.62, monoclinic, space group *P* 2₁ (No. 4), *a* = 13.1356(2), *b* = 7.03520(10), *c* = 14.6189(2) Å, *β* = 112.8645(8)°, *V* = 1244.81(3) Å³, *Z* = 2, *T* = 293(2) K, *d*_{calcd} = 1.352 g cm⁻³, *μ* = 0.256 mm⁻¹, 20767 measured reflections, 5599 unique reflections (*R*_{int} = 0.0280), 323 refined parameters, *R*₁ [*I* > 2σ(*I*)] = 0.0474, *wR*₂ [all data] = 0.1300.
- 12 For DKR during Ru-TsDPEN-catalyzed (*S*/*C* = 200) ATH of 2-ethoxycarbonyl-1-indanone (**S6** analog) (6 days, 81% yield, *cis/trans* >99:1, 99% ee for *cis*) and 2-ethoxycarbonyl-1-tetralone (**S8** analog) (6 days, 90% yield, *cis/trans* >99:1, 99% ee for *cis*) using HCO₂H/Et₃N 5:2 at rt, see: A. Ros, A. Magriz, H. Dietrich, J. M. Lassaletta and R. Fernández, *Tetrahedron*, 2007, **63**, 7532–7537.
- 13 For a review on DKR in ATH, see: (a) H. Pellissier, *Tetrahedron*, 2011, **67**, 3769–3802. For selected recent DKR examples in ATH, see: (b) B. Seashore-Ludlow, F. Saint-Dizier and P. Somfai, *Org. Lett.*, 2012, **14**, 6334–6337; (c) K. M. Steward, E. C. Gentry and J. S. Johnson, *J. Am. Chem. Soc.*, 2012, **134**, 7329–7332; (d) S.-M. Son and H.-K. Lee, *J. Org. Chem.*, 2014, **79**, 2666–2681; (e) F. Xu, M. J. Zacuto, Y. Kohmura, J. Rosen, A. Gibb, M. Alam, J. Scott and D. Tschaen, *Org. Lett.*, 2014, **16**, 5422–5425; (f) P.-G. Echeverria, J. Cornil, C. Féraud, A. Guérinot, J. Cossy, P. Phansavath and V. Ratovelomanana-Vidal, *RSC Advances*, 2015, **5**, 56815–56819; (g) T. Cheng, Q. Ye, Q. Zhao and G. Liu, *Org. Lett.*, 2015, **17**, 4972–4975.
- 14 For a kinetic resolution of racemic *cis*-3-hydroxymethyl-indanol or -tetralol by oxidation in acetone using [Ru(TsDPEN)(*p*-cymene)], see: M. Caro, M. Torrado, C. F. Masaguer and E. Raviña, *Tetrahedron: Asymmetry*, 2003, **14**, 3689–3696.
- 15 A preparation of enantiomeric 3-ethoxycarbonyl-1-indanone (96% ee) by asymmetric hydroacylation of *o*-formyl-atropic acid ethyl ester using [Rh[(*R*)-BINAP]ClO₄] has been achieved. For this, see: K. Kundu, J. V. McCullagh and A. T. Morehead, Jr., *J. Am. Chem. Soc.*, 2005, **127**, 16042–16043.
- 16 (a) P. A. Dub and T. Ikariya, *J. Am. Chem. Soc.*, 2013, **135**, 2604–2619; (b) A. Matsuoka, C. A. Sandoval, M. Uchiyama, R. Noyori and H. Naka, *Chem. Asian J.*, 2015, **10**, 112–115.
- 17 H. O. House and C. B. Hudson, *J. Org. Chem.*, 1970, **35**, 647–651.
- 18 L.-Q. Cui, Z.-L. Dong, K. Liu and C. Zhang, *Org. Lett.*, 2011, **13**, 6488–6491.
- 19 C. J. O'Donnell, R. A. Singer, J. D. Brubaker and J. D. McKinley, *J. Org. Chem.*, 2004, **69**, 5756–5759.
- 20 H. Malik, F. P. J. T. Rutjes and B. Zwanenburg, *Tetrahedron*, 2010, **66**, 7198–7203.
- 21 Separation of *rac syn*-**6** enantiomers by preparative HPLC was outsourced to Chiral Technologies Europe (France).
- 22 Separation of *rac anti*-**6** enantiomers by preparative SFC was outsourced to WuXi AppTec Co., Ltd (China).
- 23 For chiral GC analysis, see: (a) ref. 5e. (b) ref. 3b.
- ‡ CCDC-1436151 [for *syn*-(3*S*,1'*R*)-**6**-(*S*)-CSA], CCDC-1436152 [for *anti*-(3*R*,1'*R*)-**6**-(*S*)-CSA], CCDC-1436153 [for *trans*-**5**], and CCDC-1436154 [for **S7**-reduced] contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.cam.ac.uk_request.cif.