Highly fluorescent and electroactive molecular squares containing perylene bisimide ligands

Frank Würthner* and Armin Sautter

Abteilung Organische Chemie II, Universität Ulm, Albert-Einstein-Allee 11, 89081 Ulm, Germany. E-mail: frank.wuerthner@chemie.uni-ulm.de

Received (in Cambridge, UK) 16th December 1999, Accepted 7th February 2000, Published on the Web, 29th February 2000

Large functional molecular squares have been assembled with ditopic perylene bisimide bridging ligands and $Pt(\pi)$ and $Pd(\pi)$ phosphine corner units and their optical, electrochemical and spectroelectrochemical properties have been studied.

Interest in functional supramolecular assemblies1 has been triggered by examples from nature like the light harvesting system (LHS) in purple bacteria.² The supramolecular functionality of this natural assembly is realized by chlorophyll molecules organized in cyclic arrangements which control the electronic coupling of the dye molecules and allow accommodation of the reaction center of photosynthesis within the largest ring of LHS.² A straightforward access to cyclic structures in high yields was introduced by Fujita and Ogura³ and Stang and Olenyuk⁴ via the concept of molecular squares using ditopic bridging ligands and *cis*-coordinating square planar Pt(II) or Pd(II) corners. Some very interesting examples have been reported based on pyridyl-substituted porphyrin ligands,⁵ where functionality is introduced by the ligand but porphyrin fluorescence is quenched upon metal complexation. Here, we report on a new ligand based on perylene tetracarboxylic acid bisimides^{6,7} which exhibits fluorescence quantum yields $\Phi_{\rm F}$ near 100%⁶ as well as redox activity⁸ and which conserves these properties in the metal-assembled state: a giant photo- and redox-active molecular square (metal-metal diagonal ca. 3.4 nm).

The perylene ligand **1** was synthesized in 66% yield by condensation of the corresponding perylene bisanhydride⁷ with 4-aminopyridine in quinoline with catalytic amounts of zinc(II) acetate. The ligand exhibits an intense absorption band at λ_{max} = 585 nm and red fluorescence at λ_{max} = 618 nm with $\Phi_{\rm F}$ = 0.94 for **1** in chloroform.⁹

Equimolar mixing of ditopic perylene ligand 1 with $dpppM(OTf)_2$ [M = Pt(II) or Pd(II), dppp = 1,3-bis(diphenylphosphino)propane; OTf = $CF_3SO_3^{-1}$ in CH_2Cl_2 leads exclusively to the formation of molecular squares 2a,b in 90 and 95% isolated yield, respectively (Scheme 1). The squares were characterized by elemental analysis, ¹H and ³¹P{¹H} NMR spectroscopy and electrospray mass spectrometry (ESI-MS) $(\mathbf{\hat{2}a})$. † In the ³¹P{¹H} NMR spectra of **2a**,**b** only one singlet is observed owing to the high symmetry of the squares, additionally there are Pt-satellites resulting from ¹⁹⁵Pt-³¹P spin-spin coupling in the spectrum of 2a. A typical change in chemical shifts of the phosphorus nuclei of ca. 10 ppm results upon complexation with 1.10 The 1H NMR spectra show only a single set of signals for the ligand 1 and the dppp moiety with significant changes in the chemical shifts of the α - and β pyridyl protons (ca. 0.3 ppm) owing to metal complexation.¹⁰ In the ESI-MS of 2a (acetone solution) we observed signals at m/z2575.0 which we assign to the M-3OTf species with +3 charge state (separation of peaks by 1/3 mass unit) which corresponds to a molecular mass of 7725 for the square minus three triflate anions in excellent agreement with the calculated mass of 7724.89 $(C_{409}H_{360}N_{16}O_{47}F_{15}P_8S_5Pt_4)$. Furthermore, we observed signals at m/z 1894, 1485 and 1213 which we assign to (M-4OTf)⁴⁺, (M-5OTf)⁵⁺ and (M-6OTf)⁶⁺ species (THFethyl acetate solution). The less stable corresponding Pd-square



Scheme 1 Reagents and conditions: i, dpppM(OTf)_2, CH_2Cl_2, 24 h, 20 $^\circ C.$

2b has not, as yet, been characterized by the ESI–MS technique.

The optical and electrochemical properties of the perylene ligand **1** remain almost unchanged upon formation of the complexes **2**. We ascribe this to the fact that in perylene tetracarboxylic acid bisimides, the imide substituents have only a small electronic coupling to the HOMO and LUMO of the perylene chromophore.^{8,11} Therefore, the chromophoric perylene unit is largely decoupled of the binding site and only small bathochromic shifts of $\Delta \lambda = 5-7$ nm in the absorption and emission spectra are observed upon complexation. The absorption maximum of **2a** is located at $\lambda_{max} = 591$ nm, the emission occurs at 625 nm. An even more important consequence is negligible quenching of the perylene fluorescence in these complexes with $\Phi_{\rm F}$ (**2a**, CHCl₃) = 0.88.⁹

The ligand 1 and platinum complex 2a exhibit reversible electrochemistry which was investigated by cyclic voltammetry in CH₂Cl₂. 1 displays two, reversible reductions at potentials of -1.08 and -1.23 V (vs. Fc/Fc⁺), but oxidation is irreversible causing adsorption of 1 on the platinum electrode. The molecular square 2a shows two reversible waves in the reductive cycle at similar potentials (-1.01, -1.14 V vs. Fc/ Fc⁺) but the oxidative cycle, in contrast to 1 shows a reversible wave at +0.93 V, probably as a consequence of the pyridyl nitrogen lone pairs which are now blocked by platinumcomplexation (Fig. 1). The electrochemical studies of the palladium complex 2b revealed a more complicated behavior displaying an irreversible reduction at E < -1.15 V which we attribute to electron transfer processes involving the Pd(n) corners.

A closer insight into the electrochemical processes was gained by spectroelectrochemical investigations of 2a and 1. By increasing the potential in a stepwise manner to the first reduction wave of 2a (-1.01 V) the absorption bands of the perylene bisimide chromophores at 591 nm decrease while three bands in the UV–VIS–NIR appear with a maximum absorption at 791 nm [Fig. 2(a)]. Based on identical spectroscopic changes observed for the free perylene bisimide radical anionic species. After a further increase of the potential to the second reduction



Fig. 1 Cyclic voltammogram of perylene–platinum square **2a** in CH₂Cl₂; scan rate 100 mV s⁻¹. Working electrode: Pt disc, auxiliary electrode: Pt wire, reference electrode: Ag/AgCl; concentration **2a** = 2.5×10^{-4} M; supporting electrolyte NBu₄PF₆ (0.1 M).



Fig. 2 Spectroelectrograms of perylene–platinum square **2a** in CH₂Cl₂ with NBu₄PF₆ (0.1 M) as supporting electrolyte. Working electrode: Pt disc; auxiliary electrode: Pt wire; reference electrode: Ag/AgCl; concentration **2a** = 1.4×10^{-4} M. Stepwise increase of the applied potential to radical anionic perylene species (a) and to dianionic perylene species (b).

wave of 2a (-1.14 V) the perylene radical anion bands disappear completely whereas a broad and intense band appears, with a maximum absorption at 679 nm [Fig. 2(b)], corresponding to dianionic perylene bisimides. Oxidation to the radical cationic state results in less pronounced spectral changes in the visible but new broad bands appear in the NIR at 809, 908 and 1225 nm. The distinct isosbestic points in the absorption spectra for all redox couples confirm the reversibility of all oxidation and reduction processes. Therefore, we conclude that (i) the redox processes take place exclusively at the perylene ligands with the Pt(II) corner units being inert in the applied potential range and acting solely as structural building blocks, (ii) the perylene ligands assembled in the square do not interact with one another even in a charged state, and (iii) each pervlene ligand in the square is reduced and oxidized at the same time at the given potentials.12

In summary, we report on a new functional ditopic perylene ligand 1 which has been used to construct nanosized molecular squares 2 with Pt(π) and Pd(π) phosphine corner units. The

optical and electrical properties of the ligands are conserved in the metal-assembled state 2a which shows a fluorescence quantum yield of almost unity and multiple, fully reversible redox couples. The giant inner volume of nanosquares 2 seems ideal to accommodate large functional guest molecules for studies on redox and photocatalytic processes.

We thank Dr Peter Weber (Scripps, La Jolla) for ESI mass spectrometry and helpful discussions. We gratefully acknowledge the Fonds der Chemischen Industrie, the BMBF, the DFG and the Ulmer Universitätsgesellschaft for financial support, the BASF AG and Degussa-Hüls AG for the donation of chemicals.

Notes and references

† Selected data for **2a**: mp > 330 °C. ¹H NMR (CDCl₃, 500 MHz), δ 9.13 (br s, 16H), 8.12 (s, 16H), 7.69 (br s, 32H), 7.40 (m, 32H), 7.32 (m, 16H), 7.20 (d, 32H, *J* 8.4 Hz), 7.09 (d, 16H, *J* 5.9 Hz), 6.77 (d, 32H, *J* 8.4 Hz), 3.29 (br s, 16H), 2.20 br s, 8H), 1.24 (s, 144H). ³¹P{¹H} NMR (CDCl₃, 85% H₃PO₄, 202 MHz), δ –15.11 (s). UV–VIS(CH₂Cl₂), λ (ε) 591 (217000), 550 (135000), 459 nm (72000 dm³ mol⁻¹ cm⁻¹). $\lambda_{\rm em, max}$ (CH₂Cl₂) 625 nm. MS (ESI, acetone) Calc. (M–3OTf)³⁺ *m*/z 2575.7. Found 2575.0. Found: C, 59.29; H, 4.59; N, 2.54; S, 3.08. C₄₁₂H₃₆₀N₁₆O₅₆F₂₄P₈Pt₄S₈·8H₂O requires C, 59.50; H, 4.56; N, 2.69; S, 3.08%.

- J. M. Lehn, Supramolecular Chemistry, VCH, Weinheim, 1995; D. Philp and J. F. Stoddart, Angew. Chem., Int. Ed. Engl., 1996, 35, 1154; Comprehensive Supramolecular Chemistry, ed. J. M. Lehn, J. L. Atwood, J. E. D. Davies, D. D. MacNicol and F. Vögtle, Pergamon, Oxford, 1996, vol. 1–11.
- 2 G. McDermott, S. M. Prince, A. A. Freer, A. M. Hawthornthwaite-Lawless, M. Z. Papiz, R. J. Cogdell and N. W. Isaacs, *Nature*, 1995, **374**, 517; X. Hu and K. Schulten, *Phys. Today*, 1997, 28.
- 3 M. Fujita and K. Ogura, Bull. Chem. Soc. Jpn., 1996, 69, 1471.
- 4 P. J. Stang and B. Olenyuk, Acc. Chem. Res., 1997, 30, 502.
- 5 J. Fan, J. A. Whiteford, B. Olenyuk, M. D. Levin, P. J. Stang and E. B. Fleischer, J. Am. Chem. Soc., 1999, **121**, 2741; C. M. Drain and J. M. Lehn, J. Chem. Soc., Chem. Commun., 1994, 2313; R. V. Slone and J. T. Hupp, Inorg. Chem., 1997, **36**, 5422; C. M. Drain, F. Nifiatis, A. Vasenko and J. D. Batteas, Angew. Chem., Int. Ed., 1998, **37**, 2344.
- 6 G. Seybold and G. Wagenblast, *Dyes Pigments*, 1989, **11**, 303; R. Gvishi and B. Reisfeld, *Chem. Phys. Lett.*, 1993, **213**, 338.
- 7 D. Dotcheva, M. Klapper and K. Müllen, *Macromol. Chem. Phys.*, 1994, **195**, 1905; F. Würthner, C. Thalacker and A. Sautter, *Adv. Mater.*, 1999, **11**, 754.
- 8 S. K. Lee, Y. Zu, A. Herrmann, Y. Geerts, K. Müllen and A. J. Bard, J. Am. Chem. Soc., 1999, **121**, 3513.
- 9 Fluorescence quantum yields were determined relative to 1,6,7,12-tetraphenoxy-*N*,*N*'-bis(2,6-diisopropylphenyl)perylene-3,4,9,10-tetracarboxylic acid bisimide ($\Phi_{\rm R} = 0.96$, CHCl₃)⁶ by the optically dilute method ($A \le 0.035$) in chloroform; J. N. Demas and G. A. Crosby, *J. Phys. Chem.*, 1971, **75**, 991.
- 10 P. J. Stang, D. H. Cao, S. Saito and A. M. Arif, J. Am. Chem. Soc., 1995, 117, 6273.
- 11 H. Langhals, S. Demmig and H. Huber, Spectrochim. Acta, Part A, 1988, 44, 1189.
- 12 This conclusion is drawn from spectroelectrochemistry which shows complete disappearance of the bands of the neutral and the radicalanionic perylene species at the respective redox potentials.

Communication a909892e