

## Synthesis of a carborane-containing cholesterol derivative and evaluation as a potential dual agent for MRI/BNCT applications†

Cite this: *Org. Biomol. Chem.*, 2014, **12**, 2457

Diego Alberti,<sup>‡a</sup> Antonio Toppino,<sup>‡b</sup> Simonetta Geninatti Crich,<sup>\*a</sup> Chiara Meraldi,<sup>b</sup> Cristina Prandi,<sup>b</sup> Nicoletta Protti,<sup>c,d</sup> Silva Bortolussi,<sup>c,d</sup> Saverio Altieri,<sup>c,d</sup> Silvio Aime<sup>a</sup> and Annamaria Deagostino<sup>b</sup>

In this study the synthesis and characterization of a new dual, imaging and therapeutic, agent is proposed with the aim of improving the efficacy of Boron Neutron Capture Therapy (BNCT) in cancer treatment. The agent (Gd-B-AC01) consists of a carborane unit (ten boron atoms) bearing a cholesterol unit on one side (to pursue the incorporation into the liposome bi-layer) and a Gd(III)/1,4,7,10-tetraazacyclododecane monoamide complex on the other side (as a MRI reporter to attain the quantification of the B/Gd concentration). In order to endow the BNCT agent with specific delivery properties, the liposome embedded with the MRI/BNCT dual probes has been functionalized with a pegylated phospholipid containing a folic acid residue at the end of the PEG chain. The vector allows the binding of the liposome to folate receptors that are overexpressed in many tumor types, and in particular, in human ovarian cancer cells (IGROV-1). An *in vitro* test on IGROV-1 cells demonstrated that Gd-B-AC01 loaded liposomes are efficient carriers for the delivery of the MRI/BNCT probes to the tumor cells. Finally, the BNCT treatment of IGROV-1 cells showed that the number of surviving cells was markedly smaller when the cells were irradiated after internalization of the folate-targeted GdB10-AC01/liposomes.

Received 3rd December 2013,  
Accepted 11th February 2014

DOI: 10.1039/c3ob42414f

www.rsc.org/obc

## Introduction

Boron Neutron Capture Therapy (BNCT) is a binary treatment that combines low energy neutron irradiation with the presence of boron-containing compounds at the target cells. This methodology has been proposed for several pathologies, although most of the efforts have been focused on cancer.<sup>1–3</sup> BNCT has been under investigation for many years, but it has not entered the routine clinical practice yet. Since this treatment is effective only on cells internalizing large amounts of boron (at least 20–30 ppm), the principal cause of its limited success mainly depends on the insufficient delivery of NCT compounds to the target cells. A key issue deals with the measurement of the local boron concentration that is crucial to determine the optimal neutron irradiation time and to

calculate the radiation dose. One of the problems confronting the successful clinical implementation of BNCT is the difficulty of quantitatively mapping the distribution of the boron containing molecules in patients before and during the treatment. To date there has been no non-invasive way to evaluate the boron concentration in diseased and healthy tissues of the subject undergoing the irradiation. Thus, dose calculations are based on boron content values in blood, tumour, and normal tissue obtained from biodistribution studies performed beforehand.<sup>4,5</sup> Blood samples can be taken just before and even during irradiation to measure the tissue boron concentration, assuming the tumour/blood ratios assessed in previous biodistribution studies. However the boron distribution varies among patients and therefore large uncertainties exist in the tumour-to-blood boron concentration ratio. The recent development of more sensitive imaging techniques and molecular medicine protocols, based on the use of site specific agents able to carry large amounts of both therapeutic and diagnostic agents, can contribute to enhance the BNCT efficacy and to become more competitive with the established tumour treatment protocols.<sup>6</sup> Among the compounds developed for BNCT, much attention has been devoted to carboranes, icosahedral cages containing several boron atoms, which could increase the payload of boron in the tumour

<sup>a</sup>Department of Molecular Biotechnology and Health Sciences, University of Torino, via Nizza 52, 10126 Torino, Italy. E-mail: simonetta.geninatti@unito.it; Fax: +39 011 6705687

<sup>b</sup>Department of Chemistry, University of Torino, via P. Giuria 7, 10125 Torino, Italy

<sup>c</sup>Department of Nuclear and Theoretical Physics, University of Pavia, Pavia, Italy

<sup>d</sup>Nuclear Physics National Institute (INFN), Section of Pavia, Pavia, Italy

†Electronic supplementary information (ESI) available: <sup>1</sup>H NMR, <sup>13</sup>C NMR and DEPT spectra of products 1–5 and 8. See DOI: 10.1039/c3ob42414f

‡These authors contributed equally.

cells.<sup>7–11</sup> Boron containing compounds can be further functionalized with lipophilic moieties in order to permit their specific and efficient delivery through the incorporation into targeted nanoparticles. The synthesis of new dual agents for applications in MRI/BNCT where a carborane unit is linked to a lipophilic chain and to a Gd-DOTA complex through amidic bonds has been developed in our laboratory (AT101).<sup>12</sup> According to the synthetic design, the lipophilic probe AT101 binds to LDLs (low density lipoproteins) and accumulates in tumor cells characterized by an up-regulation of LDL transporters. The exploitation of LDLs as biological vectors was indeed the key to obtain a high concentration of B atoms per g in the tumor cells together with a good selectivity between tumor and healthy tissues. *In vivo* MR image acquisition showed that the amount of boron taken up in the tumor region was above the threshold required for a successful NCT treatment. More recently, with the purpose of simplifying the synthetic procedure but maintaining, and possibly improving, the favorable properties of AT101, we designed a new structure containing a triazole linker instead of the amidic function (MEA/01).<sup>13</sup> Its adducts were tested “*in vitro*” on B16 melanoma cells and the obtained intracellular Gd/B concentration appears sufficient for further “*in vivo*” BNCT studies.


An alternative promising approach to deliver efficiently boron containing agents to tumors consists of the use of liposomes. In fact, there are many examples in the literature of formulations containing boron compounds encapsulated into the aqueous cavity or, after conjugation with lipids, incorporated into the liposomal bilayer.<sup>14–16</sup> A specific class of liposomes, named “long time circulating” polyethylene glycol (PEG)-coated liposomes, is able to accumulate in solid tumors as a consequence of microvascular permeability and defective lymphatic drainage.<sup>17–19</sup> The extent of passive extravasation is directly dependent on the prolonged residence time of liposomes in the bloodstream. As far as the internalization step is concerned, one should note that in the absence of a receptor mediated mechanism, the internalization of the intact liposome into tumor cells does not take place. Therefore, in the absence of liposome active targeting the boron containing molecule has first to be released in the tumor extracellular matrix in order to allow its diffusion into cells. Alternatively, the active targeting of liposomes to tumor cells is obviously a promising strategy to improve drug delivery to tumors. To this purpose, liposomes have to be functionalized with specific ligands able to bind and to internalize through receptors expressed on tumor vasculature or on tumor cells. Recently, Nakamura and co-workers developed liposomes containing mercaptoundecahydrododecaborate (BSH) in the inner aqueous cavity and 10% distearoyl boron lipid (DSBL) incorporated into the phospholipid bilayer.<sup>20,21</sup> The proposed system appears particularly efficient because the liposome shell itself contains boron in addition to the boron encapsulated in the aqueous cavity. Furthermore, they performed the *in vivo* real-time tracking of 10% DSBL liposomes by MRI using liposomes encapsulating Gd-DOTA, observing a high signal intensity enhancement in the tumor region. The

disadvantage of the use of therapeutic and imaging agents as independent molecules is that they maintain the same biodistribution only if the liposome remains intact. After liposome degradation their biodistribution will be different and dependent on their size, hydrophilic character, residual charges, *etc.* In order to overcome this problem, in this study a dual agent, in which the imaging (Gd-DOTA) and the BNCT (carborane) components are part of the same molecule, is proposed. The incorporation of the molecule into the liposome bilayer has been pursued by a functionalization with a cholesterol residue. Cholesterol is a standard component of the liposome bi-layer usually introduced into the phospholipid formulation in order to increase the particle stability. In contrast, complexes bearing only a palmitic chain, as the parent AT101 previously investigated by our group, cannot be steadily incorporated into the liposome bi-layer and the complex may be released randomly. Synthetic mimics of cholesterol represent an important molecular tool in the fields of bioorganic/medicinal chemistry and chemical biology. Furthermore, carborane cholesterol derivatives were incorporated into unilamellar liposomes showing selective localization in tumours<sup>22,23</sup> and they might be useful vehicles for transporting boron to the tumor site. In this study, the liposomes embedded with MRI/BNCT dual probes have been functionalized with a pegylated phospholipid, containing a folic acid residue at the end of the PEG chain, as a specific carrier to transport large amounts of B/Gd dual agents to tumor cells, in particular human ovarian cancer cells overexpressing folate receptors. Finally, in order to improve the efficacy of the dual agent, the DOTA-monoamide ligand has been complexed with gadolinium enriched in the <sup>157</sup>Gd isotope (95%) that has the largest (255 000 b) thermal neutron capture cross-section among all the stable isotopes, and it is about 66 times as large as <sup>10</sup>boron (3838 b).<sup>24</sup>

## Results and discussion

### Synthesis of the B/Gd dual agent

Since the hydroxyl group of cholesterol is sterically hindered and then unreactive, in the first synthetic step it has been functionalized, by a nucleophilic substitution, with a hydroxyl propyl group in order to make easier the linkage to the carborane cage. As shown in Scheme 1, cholesterol was transformed into the corresponding mesylate (**1**) in quantitative yields (as assessed by the detection of the signal at 2.97 ppm in the <sup>1</sup>H spectrum pertinent to the mesylate CH<sub>3</sub>). Intermediate **1** was reacted with propan-1,3-diol in anhydrous dioxane at 120 °C overnight. Despite the harsh applied conditions, the product was obtained in 27% yield after chromatographic purification because of the steric hindrance of the alcoholic group. Alcohol **2** was then oxidized to the corresponding carboxylic acid, following the classical Jones reaction conditions with an overall yield of 78%. The formation of 3-(cholesten-5-en-3 $\beta$ -yloxy)-propionic acid (**3**, Scheme 1) was confirmed by the detection of a resonance at 170 ppm in the <sup>13</sup>C spectrum pertinent to the carbonyl group. The COOH moiety was then coupled to C-(2-



**Scheme 1** C-(2-Benzyloxyethyl)-C'-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-o-carborane (**4**). Reaction conditions and yields: (a) MsCl, Et<sub>3</sub>N, Et<sub>2</sub>O, 0 °C  $\rightarrow$  r.t. (>99%); (b) HO(CH<sub>2</sub>)<sub>3</sub>OH, dioxane, 120 °C (27%); (c) 3 M CrO<sub>3</sub> in H<sub>2</sub>SO<sub>4</sub>, acetone, r.t. (80%); (d) dicyclohexylcarbodiimide, CH<sub>2</sub>Cl<sub>2</sub>, rt (79%) or *N*-hydroxysuccinimide, dicyclohexylcarbodiimide, CH<sub>2</sub>Cl<sub>2</sub>, rt (64%).


benzyloxy)-ethyl-C'-aminomethyl-o-carborane **7** (prepared as previously reported).<sup>12</sup> Firstly, the coupling was attempted according to the procedure proposed by Kaminski,<sup>25</sup> but the desired product was not obtained. A two-step protocol, where the acid was activated with *N*-hydroxysuccinimide (NHS) in the presence of DCC, was then applied with the formation of product **4** (64% yield after chromatographic purification). <sup>1</sup>H NMR analysis confirmed the proposed structure. In particular, the resonance of the  $\alpha$ -methylene group shifted from 3.39 ppm in the amino derivative **7** to 4.06 ppm in the coupling product.

At this point the functionalization of the other arm of the o-carborane has been tackled. As reported in Scheme 2, the benzylic protecting group was selectively removed by Pd/C catalyzed hydrogenation.<sup>26</sup> Different catalyst concentrations and solvents were tested in order to avoid the concomitant hydrogenation of the cholesterol double bond. It was found that 5% of Pd/C in EtOH-CH<sub>2</sub>Cl<sub>2</sub> 2/1 provide the best conditions, affording after 24 h the desired product **5** in 92% overall yield.


Alcohol **5** was readily oxidized to the corresponding carboxylic acid **6** by CrO<sub>3</sub> in acetone-sulfuric acid solution (Scheme 2).

The structure of derivative **6** was supported by the <sup>1</sup>H and <sup>13</sup>C NMR data. The <sup>1</sup>H NMR spectrum showed the disappearance of the signal at 3.75 ppm assigned to the HOCH<sub>2</sub>-group, with the concomitant growth in the <sup>13</sup>C NMR spectrum of the signal at 172 ppm corresponding to the carbonyl group.

In order to obtain the molecule containing both the BNCT and MRI moieties, derivative **6** was reacted with *N*-tert-butyl-DOTAMA-C<sub>6</sub>-NH<sub>2</sub> using the two-step procedure previously described for the preparation of C-(2-benzyloxyethyl)-C'-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-o-carborane (**4**) (Scheme 2). Carborane **6** was then activated *via* NHS in the presence of DCC and after filtration of DCU, reacted with the suitable amine affording C-[*N*-tert-butyl(DOTAMA-C<sub>6</sub>)-carbamoylmethyl]-C'-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-o-carborane (**8**) in 67% overall yield. The structure was completely characterized and confirmed by ESI mass spectrometry with the detection of the characteristic ion at 1327.87 (M + H<sup>+</sup>). After removal of *tert*-butyl ester groups (CF<sub>3</sub>CO<sub>2</sub>H/



**Scheme 2** Synthesis of C-[(DOTAMA-C<sub>6</sub>)carbamoylmethyl]-C'-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-o-carborane (**9**). Reaction conditions and yields: (a) H<sub>2</sub>, Pd/C, EtOH, CH<sub>2</sub>Cl<sub>2</sub>, rt (92%); (b) 3 M CrO<sub>3</sub> in H<sub>2</sub>SO<sub>4</sub>, acetone, r.t. (80%); (c) *N*-hydroxysuccinimide, dicyclohexylcarbodiimide, *i*-PrEt<sub>2</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, rt (42%); (d) CF<sub>3</sub>COOH, CH<sub>2</sub>Cl<sub>2</sub>, r.t., (95%).



**Scheme 3** Synthesis of Gd(III)-C-[N-(DOTAMA-C<sub>6</sub>)carbamoylmethyl]-C'-2-[1,3-bis(hexadecyloxy)propan-2-yloxy]acetoamidomethyl-o-caborane complex (**10**, Gd-B-AC01). (a) GdCl<sub>3</sub>, H<sub>2</sub>O/MeOH.


CH<sub>2</sub>Cl<sub>2</sub>), the Gd(III) complex **10** was finally obtained by adding stoichiometric amounts of GdCl<sub>3</sub> in an aqueous solution of the ligand at pH 6.5 (Scheme 3).

### Liposome preparation and characterization

Liposomes were prepared by means of the standard thin lipid film hydration method<sup>27</sup> using the following components: POPC, CHOL, Gd-B-AC01, DSPE-PEG2000, and DSPE-PEG2000-Folate. Three types of liposomes containing the same amount of POPC (71%) and different amounts of cholesterol and Gd-B-AC01, namely 15.5 : 7.5%, 13 : 10%, and 8 : 15%, respectively, were prepared. In order to increase the stability and limit their non-specific phagocytic uptake, all the liposomes contain 6% of pegylated phospholipid (DSPE-PEG2000). The millimolar relaxivity (21.5 MHz, 25 °C) of Gd-B-AC01 incorporated into the liposome formulation was  $15.3 \pm 0.7$  (mmol L<sup>-1</sup>)<sup>-1</sup> s<sup>-1</sup>. This value is relatively high as a consequence of both the decrease of the molecular mobility upon the insertion into the liposome membrane and the relatively fast water exchange rate through the phospholipid bilayer due to the presence in the formulation of a high % of unsaturated phospholipids (POPC). The average liposome hydrodynamic diameter was  $125 \pm 10$  nm for all investigated formulations as detected by dynamic light scattering measurements. In order to accumulate the Gd-B-AC01 loaded liposome selectively in the tumor cells, 1% of DSPE phospholipid conjugated to a folate residue has been added to the phospholipid formulation. Overexpression of folate receptors (FRs) is well established in many cancer cells (ovary, brain, kidney, breast and lung)<sup>28,29</sup> thus identifying this receptor as a potential target for many types of ligand directed cancer therapeutics.<sup>30</sup> FR may be further qualified as a tumor-specific target, since it generally becomes accessible to intravenous administered drugs only after malignant transformation. Folate receptors (FRs) are normally expressed on cancer cells up to 10<sup>7</sup> FRs per cell but the rate of receptor recycling is relatively low.<sup>31,32</sup> Folate conjugates bind to folate receptors with high affinity ( $K_d \approx 10^{-9}$  M) and they are internalized through receptor-mediated endocytosis. The physicochemical properties of the liposome with 1% folate-phospholipid are very similar to the untargeted control liposome (without folic acid).

### NMRD profiles

$1/T_1$  nuclear magnetic resonance dispersion (NMRD) profiles, over a range of Larmor frequencies  $0.01 < \text{MHz} < 70$ , have been



**Fig. 1**  $1/T_1$  <sup>1</sup>H NMRD profile (0.01–70 MHz, pH 7, and 25 °C) of Gd-B-AC01 (○) and of Gd-B-AC01/liposomes (■). The profile was acquired using a 0.32 mM Gd complex solution and normalized to the relaxivity (mm<sup>-1</sup> s<sup>-1</sup>).

recorded for an aqueous solution of Gd-B-AC01 (0.32 mM) and compared with the profile obtained for Gd-B-AC01 embedded in the liposome membrane (Fig. 1). Both profiles show a relaxivity hump at about 30 MHz, typical of slowly moving systems. The relaxivity enhancement observed with respect to the free complex is limited by the relatively long exchange time  $\tau_M$  of water directly coordinated to the metal ion (inner sphere water) as previously shown for several Gd complexes bearing a DOTA-monoamide ligand structure.<sup>33,34</sup> The presence of a relaxivity peak between 10 and 60 MHz of the Gd-B-AC01 aqueous solution confirms that the complex forms stable micelles. A titration ( $T_1$  vs. [Gd-B-AC01]) indicated a critical micelle concentration (cmc) lower than 15  $\mu\text{M}$ . The linear dependence of relaxation rates between 300 and 15  $\mu\text{M}$  indicates that over this concentration range the complex remains in the micellar form. Lower concentrations are under the detection limit of the relaxometric method.

### Uptake experiments on IGROV-1 cells *in vitro* and MRI analysis

Liposomes were tested *in vitro* on Human Ovarian Carcinoma IGROV-1 cells to evaluate the amount of Gd taken up by cells after 24 hours incubation by measuring the metal concentration by ICP-MS analysis. These cells express a significant level of FR $\alpha$  ( $0.5\text{--}1 \times 10^6$  receptors per cell).<sup>35</sup> Liposome stability was demonstrated by the absence of non-specific release of Gd-B-AC01 to tumor cells. For this purpose, cells were incubated, for 24 h at 37 °C, in the presence of non-targeted Gd-B-AC01/liposomes containing increased % of Gd-B-AC01 from 7.5 to 15%. The results reported in Fig. 2A show that the amount of Gd associated non-specifically with IGROV-1 cells is directly proportional to the % loading of Gd-B-AC01 in the liposome. This is due to the high affinity of cholesterol for cell membranes that causes the translocation of a small but significant % of the complex from the liposome to the cell membrane. By reducing the % of Gd-B-AC01 with respect to the






Fig. 2 (A) Gd moles released to IGROV-1 cell membranes by non-targeted Gd-B-AC01/liposomes prepared with an increased amount (%) of GdAC01 after 24 h incubation Gd-B-AC01 in the presence of cells. The amount of cell associated Gd has been determined by ICP-MS. (B) Internalization of Gd chelates in IGROV-1 cells incubated in the presence of an increasing amount of non-targeted (O) and folate-targeted (■) liposomes for 24 hours at 37 °C.

other phospholipids, it is possible to reduce the number of released complexes bound specifically to the cell surface that can cause the decrease of the ratio of the internalized boron by tumors with respect to the surrounding healthy cells. For these reasons the formulation containing 7.5% of Gd-B-AC01 has been selected to perform the FR targeting study on FR expressing IGROV-1 cells. The Gd moles internalized by IGROV-1 cells, measured by ICP-MS after incubation in the presence of the increased concentration of folate-targeted liposomes, were compared with those obtained using non-targeted ones. Fig. 2B shows that by increasing the concentration of targeted and non-targeted liposomes incubated with IGROV-1 cells, the targeted liposomes quickly reach complete saturation of their folate uptake. This behavior demonstrates a high affinity of folate targeted liposomes for the FR receptor. Moreover, it indicates that the non-specific Gd-B-AC01 release from non-targeted liposomes does not invalidate the comparison if one works with relatively low liposome concentrations.

In order to assess whether the amount of Gd internalized in target cells is enough to permit MRI visualization,  $T_1$  weighted images of glass capillaries containing cellular pellets obtained by incubating *ca.*  $1 \times 10^6$  IGROV cells with increasing amounts of targeted and non-targeted Gd-B-AC01/liposomes were acquired at 1 and 7 T (Fig. 3). At 1 T, there is a dominant effect of the long reorientational motion of the paramagnetic macromolecule that causes a dramatic relaxation rate increase with a consequent signal intensity (SI) enhancement increase as shown in the NMRD profiles of Fig. 1. Clearly, the SI enhancements of labeled cells measured at 1 T (SI = 80% after 5  $\mu\text{M}$  Gd incubation) were markedly higher than those measured at 7 T (SI = 45% after 5  $\mu\text{M}$  Gd incubation). The mean SI enhancement (%) of the target tissues was calculated according to eqn (1):

$$\text{SI \% enhancement} = \frac{(\text{mean SI}_{\text{POST}} - \text{mean SI}_{\text{PRE}})}{\text{mean SI}_{\text{PRE}}} \times 100 \quad (1)$$

where  $\text{SI}_{\text{PRE}}$  and  $\text{SI}_{\text{POST}}$  correspond to the SI measured in untreated or control cells ( $\text{SI}_{\text{PRE}}$ ) and in liposome treated cells ( $\text{SI}_{\text{POST}}$ ), respectively.




Fig. 3 Weighted spin echo images, recorded at 1 (A) and 7 (B) T, of an agar phantom containing IGROV-1 cells labeled with folate-targeted and non-targeted Gd-B-AC01/liposomes: (a) non-incubated control cells; (b) and (c) cells incubated for 24 h with non-targeted liposomes of 5 (b) and 10 (c)  $\mu\text{M}$ ; cells incubated with folate targeted liposomes of 5 (d) and 10 (e)  $\mu\text{M}$ . Cells are pelleted at the bottom of glass capillaries.

The recorded SI of the targeted liposome in both cell lines is higher with respect to non-targeted ones. This demonstrates the specific accumulation of folate targeted liposomes in tumor cells with respect to control liposomes. The amount of internalized boron measured by ICP-MS was 38 and 9 ppm for folate-targeted and non-targeted liposomes, respectively. The former amount is significantly higher than the minimum boron concentration necessary to perform BNCT. We can conclude that Gd-B-AC01 loaded liposomes are efficient carriers for the delivery of MRI/BNCT probes to tumor cells.

#### BNCT treatment of IGROV-1 cells

Fig. 4 shows the percent of cells that survived the neutron irradiation. Two cell controls have been used in this experiment. The first did not receive any neutron irradiation or boron and the second control was irradiated without boron. The number of surviving cells was significantly lower when cells had been irradiated after folate-targeted GdB10-AC01/liposome internalization whereas in the absence of boron the effect of neutron irradiation is almost negligible. Furthermore,




Fig. 4 (A) Number of IGROV-1 cells that survived the BNCT treatment; (B) proliferation curve of IGROV-1 cells re-plated one day after the BNCT treatment.

the proliferation rate of the boron treated cells that survived the irradiation is significantly lower than both control cells.

Although the cells have not been completely killed, the purpose of the experiments has been satisfied because these measurements were meant to verify that the new formulations had potentiality as vectors for BNCT. Further studies with dose escalation are planned.

## Conclusions

In summary, the dual MRI/BNCT probe described in this paper can be exploited to accumulate B and Gd in target tumor cells. The new formulation has shown to have relevant characteristics: (1) the ability to selectively concentrate high amounts of B in the tumor cells; (2) the possibility to quantify the boron concentration by MRI; (3) the cholesterol versatility to be incorporated into different particles can be exploited to perform a multi-step treatment based on the administration of different drugs targeted to different overexpressed receptors; (4) the amount of internalized B is enough to perform an efficient BNCT treatment and the selective targeting of B using folate-targeted liposomes is reported to reduce significantly the uptake by healthy cells in the surrounding regions; (5) the use of liposomes as nanoplateforms for the delivery of Gd and B agents will permit the simultaneous delivery of standard

anti-tumor drugs (such as doxorubicin) in order to improve the efficacy of the treatment.

## Experimental procedures

### General methods

Flasks and all equipment used for the generation and reaction of moisture-sensitive compounds were dried using an electric heater under Ar. THF was distilled from benzophenone ketyl, anhydrous Et<sub>2</sub>O was distilled from LiAlH<sub>4</sub> and anhydrous CH<sub>2</sub>Cl<sub>2</sub> from CaH<sub>2</sub> prior to use. BuLi (1.6 M in hexanes) was obtained from Aldrich. (Bmim)<sup>+</sup>Cl<sup>−</sup> was purchased from Solvent Innovation GmbH. Decaborane was bought from KATCHEM spol. s r.o. All commercially obtained reagents and solvents were used as received. Products were purified by preparative column chromatography on Macherey Nagel silica gel for flash chromatography, 0.04–0.063 mm/230–400 mesh.

Reactions were monitored by TLC using silica gel on TLC-PET foils Fluka, 2–25 mm, layer thickness 0.2 mm, medium pore diameter 60 Å. Carboranes and their derivatives were visualized on TLC plates using a 5% PdCl<sub>2</sub> aqueous solution in HCl. <sup>1</sup>H NMR spectra were recorded at 200 MHz, and <sup>13</sup>C NMR spectra at 50.2 MHz. Data were reported as follows: chemical shifts in ppm with tetramethylsilane as an internal standard, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, dd = double-doublet, m = multiplet, br = broad), coupling constants (Hz), and assignment. <sup>13</sup>C NMR spectra were measured with complete proton decoupling. Chemical shifts were reported in ppm with the residual solvent as an internal standard. GC-MS spectra were obtained on a mass selective detector HP 5970 B instrument operating at an ionizing voltage of 70 eV connected to a HP 5890 GC with a cross-linked methyl silicone capillary column (25 m × 0.2 mm × 0.33 mm film thickness). ESI MS spectra were obtained on a Waters micromass ZQ spectrometer equipped with an ESI ion source. IR spectra were recorded on a Perkin Elmer BX FT-IR. *C*-(2-Benzyloxyethyl)-*C'*-aminomethyl-*o*-carborane (7) and *N*-*tert*-butylDOTAMA-C<sub>6</sub>-NH<sub>2</sub> were synthesized as described in the literature and their spectroscopic data correspond to those reported.<sup>12</sup>

**(3β)-Cholest-5-en-3-methanesulfonate (1).** Cholesterol (12.9 mmol, 5 g) was dissolved in 100 mL of anhydrous Et<sub>2</sub>O and cooled to 0 °C in a 250 mL three-necked round bottom flask, and then Et<sub>3</sub>N (1.5 equiv., 19.4 mmol, 2.7 mL) followed by MsCl (19.4 mmol, 1.5 mL) were added. The reaction mixture was stirred for 1 h, quenched with H<sub>2</sub>O and then extracted with Et<sub>2</sub>O (3 × 30 mL). The combined extracts were washed with H<sub>2</sub>O (2 × 30 mL), dried and evaporated under reduced pressure leaving 5.9 g (98%) of a pale yellow oil which was at once used for the following reaction. Found C, 72.37; H, 10.40; S, 6.92. Calc. for C<sub>28</sub>H<sub>48</sub>O<sub>3</sub>S: C, 72.36; H, 10.41; S, 6.90.  $\nu_{\max}$  (neat)/cm<sup>−1</sup> 2940, 1467, 1353, 1172;  $\delta_{\text{H}}$  (200 MHz; CDCl<sub>3</sub>, Me<sub>4</sub>Si): 0.65 (3H, s, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>), 0.70–2.05 (40H, m, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>, (CH<sub>3</sub>)<sub>2</sub>CH, CH<sub>3</sub> ring, CH<sub>2</sub> ring, CH ring), 2.50 (2H, m, CH<sub>2</sub>CH=C), 3.03 (1H, m, CH<sub>3</sub>SO<sub>2</sub>), 4.50 (1H, m, CHO), 5.41

(1H, bd,  $\text{CH}_2\text{CH}=\text{C}$ ,  $J = 5.5$  Hz);  $\delta_{\text{C}}$  (50.2 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 11.6 ( $\text{CH}_3$ ), 18.5 ( $\text{CH}$ ), 18.9 ( $\text{CH}$ ), 20.8 ( $\text{CH}_2$ ), 22.4 ( $\text{CH}$ ), 22.6 ( $\text{CH}$ ), 23.6 ( $\text{CH}_2$ ), 24.0 ( $\text{CH}_2$ ), 27.7 ( $\text{CH}_3$ ), 28.0 ( $\text{CH}_2$ ), 28.7 ( $\text{CH}_2$ ), 31.5 ( $\text{CH}$ ), 31.6 ( $\text{CH}_2$ ), 35.5 ( $\text{CH}$ ), 35.9 ( $\text{CH}_2$ ), 36.1 ( $\text{CH}_2$ ), 36.6 ( $\text{CH}$ ), 38.4 ( $\text{CH}_2$ ), 38.9 ( $\text{CH}_2$ ), 39.3 ( $\text{CH}_2$ ), 39.4 ( $\text{CH}_2$ ), 42.1 (Cq), 49.7 ( $\text{CH}_3$ ), 55.9 ( $\text{CH}$ ), 56.4 ( $\text{CH}$ ), 81.7 ( $\text{CH}$ ), 123.4 (Cq), 138.5 ( $\text{CH}$ ).  $m/z$  (ESI+) 465.33  $[\text{M} + \text{H}]^+$ .

**3-((3 $\beta$ )-Cholest-5-en-3-yloxy)-propanol (2).** According to the literature procedure,<sup>27</sup> (3 $\beta$ )-cholest-5-en-3-methanesulfonate (1) (12.8 mmol, 5.93 g) was dissolved in 100 mL of anhydrous dioxane in a 250 mL three-necked round bottom flask. 1,3-Propanediol (319 mmol, 24.3 g) was added and the resulting mixture was stirred overnight at 120 °C. Then the solution was cooled down at room temperature and the solvent was removed under reduced pressure. The residue was dissolved in 50 mL of  $\text{CH}_2\text{Cl}_2$  and washed with  $\text{H}_2\text{O}$ . The organic layer was separated, washed with  $\text{NaHCO}_3$  (1  $\times$  50 mL), water (1  $\times$  50 mL), and brine (1  $\times$  50 mL), dried over anhydrous  $\text{K}_2\text{CO}_3$ , and evaporated under reduced pressure. The crude was purified on silica gel (petroleum ether–diethyl ether 50/50) giving 1.54 g (27%) of a white solid. Found C, 81.05; H, 11.80. Calc. for  $\text{C}_{30}\text{H}_{52}\text{O}_2$ : C, 81.02; H, 11.79.  $\nu_{\text{max}}$  (neat)/ $\text{cm}^{-1}$  3434, 2934, 2098, 1642;  $\delta_{\text{H}}$  (200 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 0.67 (3H, s,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ), 0.70–2.49 (42H, m,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ,  $(\text{CH}_3)_2\text{CH}$ ,  $\text{CH}_3$  ring,  $\text{CH}_2$  ring,  $\text{CH}$  ring), 2.60 (1H, bs, OH), 3.19 (1H, tt,  $J = 8.5$ , 5.2, CHO), 3.67 (2H, t,  $J = 5.8$  Hz,  $\text{HOCH}_2\text{CH}_2\text{O}$ ), 3.77 (2H, t,  $J = 5.4$  Hz,  $\text{HOCH}_2\text{CH}_2\text{O}$ ), 5.33 (1H, bd,  $J = 5.2$  Hz,  $\text{CH}_2\text{CH}=\text{C}$ );  $\delta_{\text{C}}$  (50.2 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 11.6 ( $\text{CH}_3$ ), 18.5 ( $\text{CH}_3$ ), 19.0 ( $\text{CH}_3$ ), 20.8 ( $\text{CH}_2$ ), 22.3 ( $\text{CH}$ ), 22.5 ( $\text{CH}$ ), 23.8 ( $\text{CH}_2$ ), 24.0 ( $\text{CH}_2$ ), 27.6 ( $\text{CH}$ ), 28.0 ( $\text{CH}$ ), 31.6 ( $\text{CH}_2$ ), 32.3 ( $\text{CH}_2$ ), 35.6 ( $\text{CH}$ ), 36.0 ( $\text{CH}_2$ ), 36.4 (Cq), 37.0 ( $\text{CH}_2$ ), 38.8 ( $\text{CH}_2$ ), 39.2 ( $\text{CH}_2$ ), 39.5 ( $\text{CH}_2$ ), 42.0 (Cq), 49.9 ( $\text{CH}$ ), 56.0 ( $\text{CH}$ ), 56.4 ( $\text{CH}$ ), 60.2 ( $\text{CH}_2$ ), 65.8 ( $\text{CH}_2$ ), 76.4 ( $\text{CH}$ ), 121.3 ( $\text{CH}$ ), 140.3 (Cq);  $m/z$  (ESI+) 467.70  $[\text{M} + \text{Na}]^+$ .

**3-(Cholest-5-en-3 $\beta$ -yloxy)-propionic acid (3).** 3-((3 $\beta$ )-Cholest-5-en-3-yloxy)-propanol (2) (0.63 g, 1.41 mmol) was dissolved in 21 mL of acetone, and then a 3 M solution of  $\text{CrO}_3$  (4 equiv., 5.6 mmol, 0.56 g) in  $\text{H}_2\text{SO}_4$  (7 mL) was added dropwise at 0 °C. The reaction mixture was stirred overnight at rt, and then quenched with  $\text{H}_2\text{O}$ . The solvent was evaporated under reduced pressure and the mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (5  $\times$  10 mL); the organic layers were washed once with brine (1  $\times$  10 mL), dried and evaporated giving 0.49 g (76%) of a white solid. Found C, 78.57; H, 11.00. Calc. for  $\text{C}_{30}\text{H}_{50}\text{O}_3$ : C, 78.55; H, 10.99.  $\nu_{\text{max}}$  (neat)/ $\text{cm}^{-1}$  3400, 2938, 1715;  $\delta_{\text{H}}$  (200 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 0.68 (3H, s,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ), 0.70–2.05 (38H, m,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ,  $(\text{CH}_3)_2\text{CH}$ ,  $\text{CH}_3$  ring,  $\text{CH}_2$  ring,  $\text{CH}$  ring), 2.05–2.50 (2H, m,  $\text{CH}_2\text{CH}_2\text{COOH}$ ), 2.62 (2H, t,  $J = 6.4$  Hz,  $\text{CH}_2\text{COOH}$ ), 3.21 (1H, m, CHO), 3.76 (2H, t,  $J = 6.4$  Hz,  $\text{CH}_2\text{OH}$ ), 5.35 (1H, bd,  $J = 5.2$  Hz,  $\text{CH}_2\text{CH}=\text{C}$ );  $\delta_{\text{C}}$  (50.2 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 11.6 ( $\text{CH}_3$ ), 18.6 ( $\text{CH}_3$ ), 19.1 ( $\text{CH}_3$ ), 20.8 ( $\text{CH}_2$ ), 22.3 ( $\text{CH}_3$ ), 22.6 ( $\text{CH}_3$ ), 23.6 ( $\text{CH}_2$ ), 24.0 ( $\text{CH}_2$ ), 27.8 ( $\text{CH}_2$ ), 28.0 ( $\text{CH}$ ), 31.6 ( $\text{CH}_2$ ), 31.8 ( $\text{CH}$ ), 35.0 ( $\text{CH}_2$ ), 35.6 ( $\text{CH}$ ), 36.0 ( $\text{CH}_2$ ), 36.6 ( $\text{CH}_2$ ), 36.9 ( $\text{CH}_2$ ), 38.7 ( $\text{CH}_2$ ), 39.4 ( $\text{CH}_2$ ), 39.5 ( $\text{CH}_2$ ), 42.1 (Cq), 49.9 ( $\text{CH}$ ), 55.9 ( $\text{CH}$ ), 56.6 ( $\text{CH}$ ), 62.8 ( $\text{CH}_2$ ), 79.3 ( $\text{CH}$ ), 121.5 ( $\text{CH}$ ), 140.4 (Cq), 176.9 (Cq);  $m/z$  (ESI+) 459.72  $[\text{M} + \text{H}]^+$ .

**C-(2-Benzyloxyethyl)-C'-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-o-carborane (4).** In a 100 mL three-necked round bottom flask, 3-(cholesten-5-en-3 $\beta$ -yloxy)-propionic acid (3) (1.27 mmol, 0.58 g) was dissolved in 40 mL of dry  $\text{CH}_2\text{Cl}_2$ , and then C-(benzyloxy)-ethyl-C'-aminomethyl-o-carborane (1.26 mmol, 0.38 g) and *N,N'*-dicyclohexylcarbodiimide (1.51 mmol, 0.31 g) were added at room temperature. The resulting mixture was stirred at room temperature overnight. The reaction mixture was filtered and was washed with 3%  $\text{NaHCO}_3$  (1  $\times$  30 mL) and brine (1  $\times$  30 mL). The organic layer was dried, and the solvent evaporated under reduced pressure. The crude was purified on silica gel (petroleum ether–diethyl ether 80/20) giving 0.7 g (73%) of a white solid.

M.p. = 76–79 °C. Found C, 67.45; H, 9.84; S. Calc. for  $\text{C}_{42}\text{H}_{73}\text{B}_{10}\text{NO}_3$ : C, 67.43; H, 9.83.  $\nu_{\text{max}}$  (neat)/ $\text{cm}^{-1}$  3332, 2581, 1660;  $\delta_{\text{H}}$  (200 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 0.68 (3H, s,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ), 0.70–2.49 (40H + 10H, m,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ,  $(\text{CH}_3)_2\text{CH}$ ,  $\text{CH}_3$  ring,  $\text{CH}_2$  ring,  $\text{CH}$  ring, BH), 2.40 (2H, t,  $J = 5.2$  Hz,  $\text{CH}_2\text{CH}_2\text{CONH}$ ), 2.69 (2H, t,  $J = 6.0$  Hz,  $\text{PhCH}_2\text{OCH}_2\text{CH}_2$ ), 3.23 (1H, m, CHO), 3.68 (4H, m,  $\text{PhCH}_2\text{OCH}_2\text{CH}_2$ ,  $\text{OCH}_2(\text{CH}_2)_2\text{CONH}$ ), 4.07 (2H, d,  $J = 6.4$  Hz,  $\text{CONHCH}_2$ ), 4.51 (2H, s,  $\text{PhCH}_2\text{O}$ ), 5.38 (1H, bd,  $J = 4.8$  Hz,  $\text{CH}_2\text{CH}=\text{C}$ ), 7.31 (1H, t,  $J = 6.0$  Hz,  $\text{CONH}$ ), 7.39 (5H, bs, Ph);  $\delta_{\text{C}}$  (50.2 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 11.6 ( $\text{CH}_3$ ), 18.6 ( $\text{CH}_3$ ), 19.2 ( $\text{CH}_3$ ), 20.9 ( $\text{CH}_2$ ), 22.4 ( $\text{CH}_3$ ), 22.7 ( $\text{CH}_3$ ), 23.7 ( $\text{CH}_2$ ), 24.1 ( $\text{CH}_2$ ), 27.8 ( $\text{CH}$ ), 28.1 ( $\text{CH}_2$ ), 31.7 ( $\text{CH}$ ), 31.8 ( $\text{CH}_2$ ), 35.1 ( $\text{CH}_2$ ), 35.6 ( $\text{CH}_2$ ), 36.0 ( $\text{CH}$ ), 36.7 ( $\text{CH}_2$ ), 36.8 (Cq), 36.9 ( $\text{CH}_2$ ), 38.8 ( $\text{CH}_2$ ), 39.3 ( $\text{CH}_2$ ), 39.6 ( $\text{CH}_2$ ), 41.2 ( $\text{CH}_2$ ), 42.1 (Cq), 50.0 ( $\text{CH}$ ), 56.0 ( $\text{CH}$ ), 56.6 ( $\text{CH}$ ), 63.2 ( $\text{CH}_2$ ), 68.2 ( $\text{CH}_2$ ), 72.9 ( $\text{CH}_2$ ), 78.4 (Cq), 79.1 (Cq), 79.4 ( $\text{CH}$ ), 121.9 ( $\text{CH}$ ), 127.6 ( $\text{CH}$ ), 127.7 ( $\text{CH}$ ), 128.3 ( $\text{CH}$ ), 137.3 (Cq), 140.1 (Cq), 171.2 (Cq);  $m/z$  (ESI+) 749.15  $[\text{M} + \text{H}]^+$ .

**C-(Hydroxyethyl)-C'-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-o-carborane (5).** In a 50 mL two-necked round bottom flask, C-(2-benzyloxyethyl)-C'-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-o-carborane (4) (0.18 g, 0.24 mmol) was dissolved in 20 mL of a mixture of  $\text{EtOH}$ – $\text{CH}_2\text{Cl}_2$  (50–50), and then Pd/C (10%) was wetted with a few drops of water and added. The reaction mixture was stirred overnight at rt in a  $\text{H}_2$  saturated atmosphere, filtered and then the solvent was evaporated under reduced pressure giving 0.14 g (88%) of a white solid. M.p. = 103–106 °C. Found C, 63.90; H, 10.25; S. Calc. for  $\text{C}_{35}\text{H}_{67}\text{B}_{10}\text{NO}_3$ : C, 63.88; H, 10.26;  $\nu_{\text{max}}$  (neat)/ $\text{cm}^{-1}$  3433, 2583, 1655, 1587;  $\delta_{\text{H}}$  (200 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 0.70 (3H, s,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ), 0.80–2.40 (41H + 10H, m,  $\text{CH}_3\text{CH}(\text{CH}_2)_3$ ,  $(\text{CH}_3)_2\text{CH}$ ,  $\text{CH}_3$  ring,  $\text{CH}_2$  ring,  $\text{CH}$  ring, OH, BH), 2.50 (2H, t,  $J = 5.0$  Hz,  $\text{HOCH}_2\text{CH}_2$ ), 2.63 (2H, t,  $J = 6.2$  Hz,  $\text{CH}_2\text{CH}_2\text{CONH}$ ), 3.24 (1H, m, CHO), 3.73 (2H, t,  $J = 6.4$  Hz,  $\text{OCH}_2(\text{CH}_2)_2\text{CONH}$ ), 3.80 (2H, t,  $J = 5.2$  Hz,  $\text{HOCH}_2\text{CH}_2$ ), 4.11 (2H, d,  $J = 6.6$  Hz,  $\text{CONHCH}_2$ ), 5.39 (1H, bs,  $\text{CH}_2\text{CH}=\text{C}$ ), 7.36 (1H, bt,  $\text{CONH}$ );  $\delta_{\text{C}}$  (50.2 MHz;  $\text{CDCl}_3$ ,  $\text{Me}_4\text{Si}$ ): 11.6 ( $\text{CH}_3$ ), 18.5 ( $\text{CH}_3$ ), 19.2 ( $\text{CH}_3$ ), 20.9 ( $\text{CH}_2$ ), 22.4 ( $\text{CH}_3$ ), 22.6 ( $\text{CH}_3$ ), 23.6 ( $\text{CH}_2$ ), 24.1 ( $\text{CH}_2$ ), 27.8 ( $\text{CH}$ ), 28.0 ( $\text{CH}_2$ ), 31.7 ( $\text{CH}$ ), 31.8 ( $\text{CH}_2$ ), 33.7 ( $\text{CH}$ ), 35.6 ( $\text{CH}_2$ ), 36.0 ( $\text{CH}$ ), 36.6 ( $\text{CH}_2$ ), 36.8 (Cq), 37.4 ( $\text{CH}_2$ ), 38.7 ( $\text{CH}_2$ ), 39.3 ( $\text{CH}_2$ ), 39.5 ( $\text{CH}_2$ ), 41.4 ( $\text{CH}_2$ ), 42.1 (Cq), 49.9 ( $\text{CH}$ ), 55.9 ( $\text{CH}$ ), 60.8 ( $\text{CH}_2$ ), 63.1 ( $\text{CH}_2$ ), 78.1 (Cq), 79.0



(Cq), 79.6 (CH), 122.0 (CH), 140.0 (Cq), 171.7 (Cq);  $m/z$  (ESI<sup>+</sup>) 659 [M + H]<sup>+</sup>.

[Cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)-*o*-carboranyl]-acetic acid (**6**). *C*-(Hydroxyethyl)-*C'*-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-*o*-carborane (**5**) (0.48 g, 0.73 mmol) was dissolved in 21 mL of acetone, and then a 3 M solution of CrO<sub>3</sub> (4 equiv., 2.9 mmol, 0.29 g) in H<sub>2</sub>SO<sub>4</sub> (7 mL) was added carefully at 0 °C. The reaction mixture was left overnight at rt, and then quenched with H<sub>2</sub>O. The solvent was evaporated under reduced pressure and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (5  $\times$  10 mL), the organic layers were washed once with brine (1  $\times$  10 mL), dried and evaporated giving 0.42 g (85%) of pale yellow oil. Found C, 62.53; H, 9.76; N, 2.09. Calc. for C<sub>35</sub>H<sub>65</sub>B<sub>10</sub>N<sub>4</sub>O<sub>4</sub>: C, 62.56; H, 9.75; N, 2.08;  $\nu_{\max}$  (neat)/cm<sup>-1</sup> 3316, 2932, 2613, 2577, 1739, 1586, 1552;  $\delta_{\text{H}}$  (200 MHz; CDCl<sub>3</sub>, Me<sub>4</sub>Si): 0.70 (3H, s, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>), 0.80–2.45 (41H + 10H, m, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>, (CH<sub>3</sub>)<sub>2</sub>CH, CH<sub>3</sub> ring, CH<sub>2</sub> ring, CH ring, BH), 2.63 (2H, t,  $J$  = 5.6 Hz, CH<sub>2</sub>CH<sub>2</sub>CONH), 3.24 (1H, m, CHO), 3.79 (2H, t,  $J$  = 5.6 Hz, OCH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CONH), 4.11 (2H, d,  $J$  = 4.4 Hz, CONHCH<sub>2</sub>), 5.39 (1H, bs, CH<sub>2</sub>CH=C  $\delta_{\text{c}}$  (50.2 MHz; d<sub>6</sub>DMSO, Me<sub>4</sub>Si): 12.3 (CH<sub>3</sub>), 19.0 (CH<sub>3</sub>), 19.6 (CH<sub>3</sub>), 22.9 (CH<sub>2</sub>), 23.1 (CH<sub>3</sub>), 23.2 (CH<sub>3</sub>), 24.5 (CH<sub>2</sub>), 25.9 (CH<sub>2</sub>), 28.0 (CH), 29.7 (CH<sub>2</sub>), 32.0 (CH), 33.9 (CH<sub>2</sub>), 35.8 (CH), 36.2 (CH<sub>2</sub>), 36.3 (CH), 36.9 (CH<sub>2</sub>), 37.0 (Cq), 39.0 (CH<sub>2</sub>), 40.6 (CH<sub>2</sub>), 41.6 (CH<sub>2</sub>), 42.4 (CH<sub>2</sub>), 42.7 (CH<sub>2</sub>), 43.1 (Cq), 48.1 (CH), 50.2 (CH), 56.2 (CH<sub>2</sub>), 56.8 (CH<sub>2</sub>), 76.0 (Cq), 78.7 (Cq), 81.6 (CH), 121.5 (CH), 141.0 (Cq), 168.9 (Cq);  $m/z$  (ESI<sup>+</sup>) 696 [M + Na + H]<sup>+</sup>.

*C*-[*N*-*tert*-Butyl(DOTAMA-C<sub>6</sub>)-carbamoylmethyl]-*C'*-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-*o*-carborane (**8**). According to the previous procedure described for compound **3**, [cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)-*o*-carboranyl]-acetic acid (**6**) (0.2 g, 0.29 mmol) was reacted with *N*-*tert*-butylDOTAMA-C<sub>6</sub>-NH<sub>2</sub> (0.19 g, 0.29 mmol). The crude was purified on silica gel (eluant: CH<sub>2</sub>Cl<sub>2</sub>–MeOH 96–4, then CH<sub>2</sub>Cl<sub>2</sub>–MeOH 80–20) affording 0.15 g (42%) of pale yellow oil. Found C, 62.57; H, 9.82; N, 7.41. Calc. for C<sub>69</sub>H<sub>129</sub>B<sub>10</sub>N<sub>7</sub>O<sub>10</sub>: C, 62.55; H, 9.81; N, 7.40;  $\nu_{\max}$  (neat)/cm<sup>-1</sup> 3294, 2933, 2575, 1730, 1665, 1552;  $\delta_{\text{H}}$  (200 MHz; CDCl<sub>3</sub>, Me<sub>4</sub>Si): 0.70 (3H, s, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>), 0.80–2.50 (78H + 10H, m, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>, (CH<sub>3</sub>)<sub>2</sub>CH, CH<sub>3</sub> ring, CH<sub>2</sub> ring, CH ring, NHCH<sub>2</sub>CH<sub>2</sub>NH, NHCOCH<sub>2</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>2</sub>NHCO, BH), 1.50 (27H, s, COOtBu), 3.10–3.80 (13H, m, CH<sub>2</sub>COOtBu, NHCOCH<sub>2</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>2</sub>NHCO, CHO), 3.79 (2H, bt, OCH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CONH), 4.11 (2H, d,  $J$  = 6.2 Hz, CONHCH<sub>2</sub>), 5.39 (1H, bd, CH<sub>2</sub>CH=C), 8.20 (1H, bs, NHCO), 8.70 (2H, m, 2  $\times$  NHCO);  $\delta_{\text{c}}$  (50.2 MHz; CDCl<sub>3</sub>, Me<sub>4</sub>Si): 11.6 (CH<sub>3</sub>), 18.5 (CH<sub>3</sub>), 19.3 (CH<sub>3</sub>), 20.9 (CH<sub>2</sub>), 22.4 (CH<sub>3</sub>), 22.7 (CH<sub>3</sub>), 23.7 (CH<sub>2</sub>), 24.1 (CH<sub>2</sub>), 25.2 (CH<sub>2</sub>), 27.8 (9  $\times$  CH<sub>3</sub>), 28.4 (CH<sub>2</sub>), 28.4 (CH<sub>2</sub>), 28.5 (CH<sub>2</sub>), 29.5 (CH<sub>2</sub>), 31.6 (CH), 31.7 (CH<sub>2</sub>), 32.9 (CH<sub>2</sub>), 35.6 (CH), 36.0 (CH<sub>2</sub>), 36.9 (Cq), 37.1 (CH<sub>2</sub>), 38.1 (CH<sub>2</sub>), 38.5 (CH<sub>2</sub>), 38.9 (CH<sub>2</sub>), 39.3 (CH<sub>2</sub>), 39.6 (CH<sub>2</sub>), 41.4 (CH<sub>2</sub>), 41.6 (CH<sub>2</sub>), 42.0 (2  $\times$  CH<sub>2</sub>), 46.0–54.0 (8  $\times$  CH<sub>2</sub>), 50.0 (CH), 55.4 (2  $\times$  CH<sub>2</sub>), 56.0 (CH), 56.6 (CH), 57.0 (Cq), 63.8 (2  $\times$  CH<sub>2</sub>), 65.8 (Cq), 75.5 (Cq), 79.0 (2  $\times$  CH), 80.0 (Cq), 81.7 (Cq), 81.8 (Cq), 121.7 (CH), 140.7 (Cq), 166.3 (Cq), 171.1 (Cq), 171.3 (Cq), 171.4 (Cq), 171.9 (Cq), 172.2, (Cq);  $m/z$  (ESI<sup>+</sup>) 1350 [M + Na + H]<sup>+</sup>.

*C*-[(DOTAMA-C<sub>6</sub>)-carbamoylmethyl]-*C'*-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-*o*-carborane (**9**). In a 25 mL round bottom flask, 247 mg (0.186 mmol) of *C*-[*N*-*tert*-butyl (DOTAMA-C<sub>6</sub>)-carbamoylmethyl]-*C'*-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-*o*-carborane (**8**) were cooled to 0 °C, dissolved in 5 mL of a mixture of CF<sub>3</sub>COOH–CH<sub>2</sub>Cl<sub>2</sub> (50–50) and stirred for 4 h at rt. After evaporation of CF<sub>3</sub>COOH–CH<sub>2</sub>Cl<sub>2</sub>, 204 mg (1.99 mmol) of viscous colorless oil were obtained (95%). Found C, 59.21; H, 9.17; N, 8.49. Calc. for C<sub>57</sub>H<sub>105</sub>B<sub>10</sub>N<sub>7</sub>O<sub>2</sub>: C, 59.19; H, 9.15; N, 8.48;  $\nu_{\max}$  (neat)/cm<sup>-1</sup> 3390, 2931, 2581, 1741, 1676, 1586;  $\delta_{\text{H}}$  (400 MHz; CD<sub>3</sub>SOCD<sub>3</sub>, Me<sub>4</sub>Si): 0.65 (3H, s, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>), 0.80–2.40 (51H + 10H, m, CH<sub>3</sub>CH(CH<sub>2</sub>)<sub>3</sub>, (CH<sub>3</sub>)<sub>2</sub>CH, CH<sub>3</sub> ring, CH<sub>2</sub> ring, CH ring, NHCH<sub>2</sub>CH<sub>2</sub>NH, NHCOCH<sub>2</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>2</sub>NHCO, BH), 3.10–3.80 (13H, m, CH<sub>2</sub>COOtBu, NHCOCH<sub>2</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>2</sub>NHCO, CHO), 3.79 (2H, bt, OCH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CONH), 4.00 (2H, d,  $J$  = 6.2 Hz, CONHCH<sub>2</sub>), 5.30 (1H, bs, CH<sub>2</sub>CH=C), 7.30 (1H, m, NHCO), 8.60 (2H, m, 2  $\times$  NHCO);  $\delta_{\text{c}}$  (100.4 MHz; CD<sub>3</sub>SOCD<sub>3</sub>, Me<sub>4</sub>Si): 11.6 (CH<sub>3</sub>), 18.5 (CH<sub>3</sub>), 19.3 (CH<sub>3</sub>), 20.8 (CH<sub>2</sub>), 22.4 (CH<sub>3</sub>), 22.7 (CH<sub>3</sub>), 23.1 (CH<sub>2</sub>), 23.8 (CH<sub>2</sub>), 24.7 (CH<sub>2</sub>), 28.4 (CH<sub>2</sub>), 28.4 (CH<sub>2</sub>), 28.5 (CH<sub>2</sub>), 29.5 (CH<sub>2</sub>), 31.6 (CH), 31.7 (CH<sub>2</sub>), 32.9 (CH<sub>2</sub>), 35.6 (CH), 36.0 (CH<sub>2</sub>), 36.9 (Cq), 37.1 (CH<sub>2</sub>), 38.1 (CH<sub>2</sub>), 38.5 (CH<sub>2</sub>), 38.9 (CH<sub>2</sub>), 39.3 (CH<sub>2</sub>), 39.6 (CH<sub>2</sub>), 41.4 (CH<sub>2</sub>), 41.6 (CH<sub>2</sub>), 42.0 (2  $\times$  CH<sub>2</sub>), 46.0–54.0 (8  $\times$  CH<sub>2</sub>), 50.0 (CH), 55.4 (2  $\times$  CH<sub>2</sub>), 56.0 (CH), 56.6 (CH), 57.0 (Cq), 63.8 (2  $\times$  CH<sub>2</sub>), 65.8 (Cq), 75.5 (Cq), 79.0 (2  $\times$  CH), 80.0 (Cq), 121.7 (CH), 140.7 (Cq), 166.3 (Cq), 171.1 (Cq), 171.3 (Cq), 171.4 (Cq), 171.9 (Cq), 172.2, (Cq);  $m/z$  (ESI<sup>+</sup>) 1350 [M + Na + H]<sup>+</sup>.

Gd(III)-*C*-[*N*-(DOTAMA-C<sub>6</sub>)-carbamoylmethyl]-*C'*-2-[1,3-bis(hexadecyloxy)propan-2-yloxy]acetoamidomethyl-*o*-carborane complex (**10**). In a 100 mL round bottom flask, 70 mg of *C*-[(DOTAMA-C<sub>6</sub>)-carbamoylmethyl]-*C'*-[cholesten-5-en-3 $\beta$ -yloxy-(3-propionacetoamidomethyl)]-*o*-carborane (**9**) (0.055 mmol) and 1 equivalent of GdCl<sub>3</sub> were dissolved in water at rt. The pH solution was checked and maintained at 6.5 with 1 M NaOH aqueous solution. The solution was stirred for 24 h, and then the pH was adjusted to 8.5 with 1 M NaOH aqueous solution and the mixture was stirred for 2 h, and then filtered with a 0.2 mm syringe filter, the pH was adjusted to 7 with a 1 M HCl aqueous solution and the solvent evaporated. Inorganic salts were removed by gel filtration on a Sephadex® G10 column. The solvent was removed affording 63 mg of the complex (0.043 mmol, 40%). Found C, 46.68; H, 7.88; S. Calc. for C<sub>12</sub>H<sub>24</sub>B<sub>10</sub>O<sub>2</sub>: C, 46.73; H, 7.84;  $\nu_{\max}$  (neat)/cm<sup>-1</sup> 3329, 2923, 2540, 1681, 1620, 1409, 1180;  $m/z$  (ESI<sup>+</sup>) 1451.90 [M + H]<sup>+</sup>.

### Liposome preparation and characterization

All the phospholipids used in the liposome preparation (POPC: 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine, DSPE-PEG 2000 methoxy: 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[methoxy(polyethylene glycol)-2000] ammonium salt, DSPE-PEG 2000-Folate {1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[folate(polyethylene glycol)-2000] ammonium salt} were purchased from Avanti Polar Lipids (Alabaster, AL). Cholesterol (Chol) was purchased from Sigma-Aldrich (St. Louis, MO). Gd-B-AC01 was synthesized



according to the above procedure. Liposomes were prepared by following the thin lipid film hydration method.<sup>36</sup> The following liposome formulations were prepared using POPC : Chol : Gd-B-AC01 : DSPE-PEG2000 methoxy : DSPE-PEG2000-folate; Lipo-Ctrl 1: 71:8:15:6:0 molar ratio. Lipo-Ctrl 2: 71:13:10:6:0 molar ratio. Lipo-Ctrl 3: 71:15.5:7.5:6:0 molar ratio. Liposomes with folate (Lipo-Folate) were prepared by adding 1% molar ratio of DSPE-PEG 2000 Folate to all previous described formulations; moreover DSPE-PEG 2000 was added in 5% molar ratio instead of 6%. Briefly, all lipids were dissolved in chloroform and the organic solution was slowly evaporated for removing the solvent until a thin film was formed. The film was then hydrated at 55 °C with 1.5 mL of a saline buffer containing 5 mM Hepes and 150 mM NaCl, pH = 7.4 (HBS). The suspension of multilamellar vesicles (MLV) was extruded (Lipex extruder, Northern Lipids Inc., Canada) four times on a polycarbonate filter with a pore diameter of 400 nm, four times on a 200 nm filter and three times on a 100 nm filter. The final liposome suspension was dialyzed two times at 4 °C against a 100-fold volume of HBS. The amount of Gd-B-AC01 incorporated into the liposome was determined by <sup>1</sup>H NMR *T*<sub>1</sub> measurements at 21.5 MHz, 25 °C (Stelar Spinmaster, Mede, Italy) of the mineralized complex solution (in HCl 6 M at 120 °C for 16 h). The hydrated mean diameter of liposomes was determined using a Malvern dynamic light-scattering spectrophotometer (Malvern Instruments, Malvern, UK). All samples were analyzed at 25 °C in filtered (cutoff = 200 nm) HBS buffer (pH = 7). The polydispersity index (PDI) for all the liposomes prepared in this work was smaller than 0.2. The liposomes were used within two weeks from the preparation and stored at 4 °C under argon.

**<sup>1</sup>H relaxation rate measurements.** The longitudinal water proton relaxation rates were measured on the Stelar Spinmaster spectrometer (Stelar, Mede, Italy) operating at 21.5 MHz by means of the standard inversion recovery technique (16 experiments, 2 scans). A typical 90° pulse width was 3.5 ms and the reproducibility of the *T*<sub>1</sub> data was ±0.5%. The 1/*T*<sub>1</sub> nuclear magnetic relaxation dispersion (NMRD) profiles of water protons were measured over a continuum of magnetic field strength from 0.00024 to 0.5 T (corresponding to 0.01–20 MHz proton Larmor frequency) on the fast field-cycling relaxometer (Stelar Spinmaster FFC 2000; Stelar) equipped with a silver magnet. The relaxometer operates under complete computer control with an absolute uncertainty in the 1/*T*<sub>1</sub> values of ±1%. The typical field sequences were used: the non-polarized sequence between 40 and 8 MHz and the pre-polarized sequence between 8 and 0.01 MHz. The observation field was set at 13 MHz. Sixteen experiments of two scans each were used for the *T*<sub>1</sub> determination for each field.

**Cell culture and uptake experiments.** Human ovarian carcinoma cell line (IGROV-1) was kindly provided by Dr Claudia Cabella (Bracco Imaging, Colliere Giososa TO). IGROV-1 cells were cultured in RPMI (Lonza) supplemented with 10% (v/v) FBS, 2 mM glutamine, 100 U mL<sup>-1</sup> penicillin, and 100 U mL<sup>-1</sup> streptomycin. Cells were incubated at 37 °C in a humidified atmosphere of 5% CO<sub>2</sub>. At 80% confluence, cells were

detached with 0.02% EDTA (Lonza). For the *in vitro* uptake experiments, about 4.5 × 10<sup>5</sup> of IGROV were seeded in 6 cm diameter culture dishes. After 6 h, the medium was removed and it was replaced with RPMI w/o folate. After 24 h, cells were incubated for 24 h with Lipo-Folate or Lipo-Ctrl. At the end of incubation, cells were washed three times with 5 mL ice-cold PBS, detached with 0.05% trypsin and 0.02% EDTA in PBS and transferred into glass capillaries for MRI analysis (see below). The Gd content of IGROV-1 cells was determined by inductively coupled plasma mass spectrometry (ICP-MS) (Element-2; Thermo-Finnigan, Rodano, Milan, Italy). Sample digestion was performed with 2 mL of concentrated HNO<sub>3</sub> (70%) under microwave heating (Milestone MicroSYNTH Microwave labstation equipped with an optical fiber temperature control and HPR-1000/6 M six position high pressure reactor, Bergamo, Italy). After digestion, sample volumes were brought to 3 mL with ultrapure water and samples were analyzed by ICP-MS. The protein concentration was determined from cell lysates by the Bradford assay, using bovine serum albumin as a standard.

**MRI.** MR images were acquired at 1 and 7 T field. MR images at 1 T were acquired with an Aspect M2 High Performance MRI System (Aspect Magnet Technologies Ltd, Netanya, Israel) consisting of a NdFeB magnet, equipped with a 35 mm solenoid Tx/Tr coil of inner diameter 35 mm. This system is equipped with fast gradient coils (gradient strength, 450 mT m<sup>-1</sup> at 60 A; ramp time, 250 μs at 160 V) with a field homogeneity of 0.2–0.5 gauss. MR images at 7 T were acquired on a Bruker Avance300 spectrometer equipped with a Micro 2.5 microimaging probe (Bruker BioSpin, Ettlingen, Germany) using two birdcage resonators with 30 and 10 mm inner diameters.

For recording MR images *in vitro*, cells were pelleted at the bottom of glass capillaries placed in a phantom embedded of high gelling agarose gel (1% in PBS). MR images were acquired using a standard *T*<sub>1</sub> weighted multislice spin echo sequence, using the following parameters: TR/TE/NEX 250/4/6, resolution 78 μm, slice thickness 1 mm at 7 T and TR/TE/NEX 250/7/16, resolution 78 μm, slice thickness 1 mm at 1 T, respectively. At both field strengths a standard saturation recovery protocol was used to measure *T*<sub>1</sub> relaxation times.

### Cell irradiation

Seven flasks, four with IGROV-1 cells previously incubated for 24 h in the presence of folate targeted GdAC01/liposomes, at 10 μM Gd concentration, and three with non-treated control cells were irradiated in the thermal column of the TRIGA Mark II reactor at the University of Pavia, Italy. The cells cultured in the presence of liposomes were washed with cold PBS before irradiation. At the end of the irradiation, the medium was removed, it was replaced with RPMI and flasks were placed at 37 °C in a humidified atmosphere of 5% CO<sub>2</sub>.

The irradiation position had been previously characterized from the point of view of neutron flux distribution by means of thin activation foils.<sup>37</sup> At a reactor power of 250 kW the thermal neutron flux in air at that position is (1.17 ± 0.03) × 10<sup>10</sup> cm<sup>-2</sup> s<sup>-1</sup>, while the epithermal and fast components are at least two orders of magnitude lower. The flux is roughly

constant (less than 1%) along the vertical direction, thus the flasks were superposed and irradiated at the same time. The irradiation time has been fixed at 15 minutes at a reactor power of 30 kW, corresponding to a thermal neutron fluence of  $1.26 \times 10^{12} \text{ cm}^{-2}$ . The radiation dose absorbed by cells treated with 10  $\mu\text{M}$  Gd was 6.2 Gy and the dose absorbed by control cells was 0.6 Gy.

**Proliferation assay.** The day after irradiation cells were detached with 0.02% EDTA and the trypan blue exclusion test of cell viability was performed. Then, around  $6 \times 10^5$  IGROV-1 cells from each differently treated flask were seeded in 10 cm diameter culture dishes. After 1, 2, 3 and 4 days, cells were washed with PBS, detached with 0.05% trypsin and 0.02% EDTA in PBS and transferred into falcon tubes. Then, cells were sonicated for 30" at 30% power in ice and the protein concentration from cell lysates was determined by the Bradford method, using bovine serum albumin as a standard.

## Conflict of interest

The authors declare no competing financial interest.

## Acknowledgements

This research was performed in the framework of the EU COST Action TD1004, and supported by MIUR (PRIN 2009235JB7), by the University of Torino (code D15E11001710003 project: Innovative Nanosized Theranostic Agents), by the University of Genova (Progetto San Paolo; Title: Validazione di molecole per il rilascio tumore specifico di farmaci e la valutazione contestuale della risposta mediante imaging funzionale), and by Consorzio Interuniversitario di Ricerca in Chimica dei Metalli dei Sistemi Biologici (CIRCMSB). We thank Franco Fedeli for the synthesis of *N*-tert-butylDOTAMA- $\text{C}_6\text{-NH}_2$ .

## References

- 1 R. F. Barth, *J. Neuroonc.*, 2003, **62**, 1–5.
- 2 R. F. Barth, J. A. Coderre, M. G. H. Vicente and T. E. Blue, *Clin. Cancer Res.*, 2005, **11**, 3987–4002.
- 3 M. F. Hawthorne and M. W. Lee, *J. Neuroonc.*, 2003, **62**, 33–45.
- 4 J. Laakso, M. Kulvik, I. Ruokonen, J. Vahatalo, R. Zilliacus, M. Farkkila and M. Kallio, *Clin. Chem.*, 2001, **47**, 1796–1803.
- 5 M. Kortensniemi, T. Seppala, I. Auterinen and S. Savolainen, *Appl. Radiat. Isot.*, 2004, **61**, 823–827.
- 6 S. Geninatti-Crich, A. Deagostino, A. Toppino, D. Alberti, P. Venturello and S. Aime, *Anti-Cancer Agents Med. Chem.*, 2012, **12**, 543–553.
- 7 A. H. Soloway, W. Tjarks, B. A. Barnum, F. G. Rong, R. F. Barth, I. M. Codogni and J. G. Wilson, *Chem. Rev.*, 1998, **98**, 1515–1562.
- 8 J. F. Valliant, K. J. Guenther, A. S. King, P. Morel, P. Schaffer, O. O. Sogbein and K. A. Stephenson, *Coord. Chem. Rev.*, 2002, **232**, 173–230.
- 9 G. Wu, R. F. Barth, W. Yang, R. J. Lee, W. Tjarks, M. V. Backer and J. M. Backer, *Anti-Cancer Agents Med. Chem.*, 2006, **6**, 167–184.
- 10 E. L. Crossley, E. J. Ziolkowski, J. A. Coderre and L. M. Rendina, *Mini Rev. Med. Chem.*, 2007, **7**, 303–313.
- 11 I. Viñas and C. Teixidor, *Future Med. Chem.*, 2013, **5**, 617–619.
- 12 S. Aime, A. Barge, A. Crivello, A. Deagostino, R. Gobetto, C. Nervi, C. Prandi, A. Toppino and P. Venturello, *Org. Biomol. Chem.*, 2008, **6**, 4460–4466.
- 13 A. Toppino, M. E. Bova, S. G. Crich, D. Alberti, E. Diana, A. Barge, S. Aime, P. Venturello and A. Deagostino, *Chemistry*, 2013, **19**, 720–727.
- 14 P. J. Kueffer, C. A. Maitz, A. A. Khan, S. A. Schuster, N. I. Shlyakhtina, S. S. Jalisatgi, J. D. Brockman, D. W. Nigg and M. F. Hawthorne, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 6512–6517.
- 15 H. Nakamura, in *Methods Enzym.*, ed. N. Duzgunes, 2009, pp. 179–208.
- 16 S. Altieri, M. Balzi, S. Bortolussi, P. Bruschi, L. Ciani, A. M. Clerici, P. Faraoni, C. Ferrari, M. A. Gadan, L. Panza, D. Pietrangeli, G. Ricciardi and S. Ristori, *J. Med. Chem.*, 2009, **52**, 7829–7835.
- 17 V. P. Torchilin, *Adv. Drug Delivery Rev.*, 2008, **60**, 548–558.
- 18 A. P. Pathak, D. Artemov, B. D. Ward, D. G. Jackson, M. Meeman and Z. M. Bhujwalla, *Cancer Res.*, 2005, **65**, 1425–1432.
- 19 M. R. Dreher, W. G. Liu, C. R. Michelich, M. W. Dewhirst, F. Yuan and A. Chilkoti, *J. Natl. Cancer Inst.*, 2006, **98**, 335–344.
- 20 H. Koganei, M. Ueno, S. Tachikawa, L. Tasaki, H. S. Ban, M. Suzuki, K. Shiraishi, K. Kawano, M. Yokoyama, Y. Maitani, K. Ono and H. Nakamura, *Bioconjugate Chem.*, 2013, **24**, 124–132.
- 21 H. Nakamura, *Future Med. Chem.*, 2013, **5**, 715–730.
- 22 M. Bialek-Pietras, A. B. Olejniczak, S. Tachikawa, H. Nakamura and Z. J. Lesnikowski, *Bioorg. Med. Chem.*, 2013, **21**, 1136–1142.
- 23 G. L. Pan, S. Oie and D. R. Lu, *Pharm. Res.*, 2004, **21**, 1257–1262.
- 24 N. Dewi, *et al.*, *Biomed. Pharmacother.*, 2013, **67**, 451–457.
- 25 Z. J. Kaminski, *Synthesis*, 1987, 917–920.
- 26 J. Schimmel, M. I. P. Eleuterio, G. Ritter and R. R. Schmidt, *Eur. J. Org. Chem.*, 2006, 1701–1721.
- 27 A. Bajaj, P. Kondaiah and S. Bhattacharya, *Bioconjugate Chem.*, 2008, **19**, 1640–1651.
- 28 E. I. Sega and P. S. Low, *Cancer Metast. Rev.*, 2008, **27**, 655–664.
- 29 W. Xia and P. S. Low, *J. Med. Chem.*, 2010, **53**, 6811–6824.
- 30 Y. J. Lu and P. S. Low, *Adv. Drug Delivery Rev.*, 2002, **64**, 342–352.
- 31 B. A. Kamen and A. K. Smith, *Adv. Drug Delivery Rev.*, 2004, **56**, 1085–1097.

- 32 C. M. Paulos, J. A. Reddy, C. P. Leamon, M. J. Turk and P. S. Low, *Mol. Pharm.*, 2004, **66**, 1406–1414.
- 33 L. Tei, A. Barge, S. G. Crich, R. Pagliarin, V. Negri, D. Ramella, G. Cravotto and S. Aime, *Chemistry*, 2010, **16**, 8080–8087.
- 34 C. Grange, S. Geninatti-Crich, G. Esposito, D. Alberti, L. Tei, B. Bussolati, S. Aime and G. Camussi, *Cancer Res.*, 2010, **70**, 2180–2190.
- 35 N. Kamaly, T. Kalber, M. Thanou, J. D. Bell and A. D. Miller, *Bioconjugate Chem.*, 2009, **20**, 648–655.
- 36 F. Olson, C. A. Hunt, F. C. Szoka, W. J. Vail and D. Papahadjopoulos, *Biochim. Biophys. Acta*, 1979, **557**, 9–23.
- 37 N. Protti, S. Bortolussi, M. I. M. Prata, P. Bruschi, S. Altieri and D. W. Nigg, *Trans. Am. Nucl. Soc.*, 2012, **107**, 1269–1272.