


This article was downloaded by: [University of Connecticut]

On: 08 October 2014, At: 17:49

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

π -Face-Selective 1,3-Dipolar Cycloadditions of 3,4-Di-tert-butylthiophene 1-Oxide with 1,3-Dipoles

Juzo Nakayama^a, Tomohiro Furuya^a, Yasuko Suzuki^a, Suguru Hiraiwa^a & Yoshiaki Sugihara^a

^a Department of Chemistry, School of Science and Engineering, Saitama University, Saitama, Japan
Published online: 28 Apr 2009.

To cite this article: Juzo Nakayama, Tomohiro Furuya, Yasuko Suzuki, Suguru Hiraiwa & Yoshiaki Sugihara (2009) π -Face-Selective 1,3-Dipolar Cycloadditions of 3,4-Di-tert-butylthiophene 1-Oxide with 1,3-Dipoles, Phosphorus, Sulfur, and Silicon and the Related Elements, 184:5, 1175-1183, DOI: [10.1080/10426500902856297](https://doi.org/10.1080/10426500902856297)

To link to this article: <http://dx.doi.org/10.1080/10426500902856297>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or

indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

π -Face-Selective 1,3-Dipolar Cycloadditions of 3,4-Di-*tert*-butylthiophene 1-Oxide with 1,3-Dipoles

Juzo Nakayama, Tomohiro Furuya, Yasuko Suzuki, Suguru Hiraiwa, and Yoshiaki Sugihara

Department of Chemistry, School of Science and Engineering, Saitama University, Saitama, Japan

3,4-Di-tert-butylthiophene 1-oxide underwent 1,3-dipolar cycloadditions with 1,3-dipoles such as nitrile oxide, diazomethane, nitrile imide, nitron, and azomethine ylide at its syn- π -face with respect to the S=O bond.

Keywords 1,3-Dipolar cycloaddition; π -face selectivity; heterocycles; stereochemistry; thiophene 1-oxide

INTRODUCTION

π -Facial selectivity (diastereofacial selectivity) in Diels–Alder reactions has been attracting considerable attention. It has been investigated most extensively by using 5-substituted cyclopentadienes as the diene.¹ Thiophene 1-oxides, which have the general structure shown in Figure 1, are no longer aromatic and hence highly reactive. They serve as a type of cyclic diene that possesses two π -faces, i.e., *syn*- and *anti*-faces, for Diels–Alder reactions, with respect to the S=O bond. We have reported that 3,4-di-*tert*-butylthiophene 1-oxide (**1**) undergoes *syn*- π -face-selective Diels–Alder reactions with a wide range of dienophiles.^{2–4} Recently, we also reported that the S₂O, formed by retro-Diels–Alder reaction of bicyclic compound **2**, disproportionates to S₃ (triatomic sulfur) and SO₂, and the resulting S₃ undergoes a *syn*- π -face-selective 1,3-dipolar cycloaddition to **1**, the counterpart of the S₂O formation, to give the final product **3**; the stereochemistry of **3** was determined by X-ray diffraction analysis (Scheme 1).⁵ We have now examined 1,3-dipolar

Received 27 December 2007; accepted 29 February 2008.

Dedicated to Professor Marian Mikołajczyk, CBMiM PAN in Łódź, Poland, on the occasion of his 70th birthday.

Address correspondence to Juzo Nakayama, Department of Chemistry, School of Science and Engineering, Saitama University, Sakura-ku, Saitama, Saitama 338-8570, Japan. E-mail: nakayama@mail.saitama-u.ac.jp





FIGURE 1 General structure and two π -faces of thiophene 1-oxides.


cycloadditions of **1** with a series of 1,3-dipoles to know more about the stereochemical course of the reaction. To our knowledge, π -facial selectivity in 1,3-dipolar cycloadditions has been far less investigated.^{6–9}



SCHEME 1 *Syn*- π -face-selective 1,3-dipolar cycloaddition of S_3 to thiophene 1-oxide **1**.

Initially, 1,3-dipolar cycloadditions of **1** with propargyl-allenyl type 1,3-dipoles were examined. The reaction of **1** with acetonitrile oxide, generated in situ by reaction of nitroethane with phenyl isocyanate in the presence of Et_3N in CH_2Cl_2 at room temperature,¹⁰ furnished the single 1,3-dipolar cycloadduct **4a**¹¹ in 80% yield (Scheme 2). On the other hand, the reaction of **1** with benzonitrile oxide in CH_2Cl_2 , generated by treatment of α -chlorobenzaldehyde oxime ($PhCCl=NOH$) with Et_3N ,¹² afforded a 10:1 mixture of the two diastereomers **4b** and **4b'**,¹¹ which were isolated in 80% and 7% yields, respectively. The ratio **4b**:**4b'** showed solvent dependency; it was 23:1 and 24:1 in benzene and toluene, respectively.


The regiochemistry of the addition in the formation of **4a** was determined by observation of the coupling (1.1 Hz) between H_a and the



SCHEME 2 1,3-Dipolar cycloaddition of **1** with nitrile oxides.


oxazole methyl protons. The configuration of the sulfinyl group was determined on the basis of aromatic solvent-induced shift (ASIS) and $\text{Eu}(\text{thd})_3$ -induced shift [$\text{Eu}(\text{thd})_3$; europium tris(2,2,6,6-tetramethyl-3,5-heptanedionate)]. It is well documented that benzene coordinates to the sulfur atom of the sulfinyl group,^{13–17} whereas the europium atom of $\text{Eu}(\text{thd})_3$ coordinates to the oxygen atom. Thus, for the adduct **4a**, the ^1H NMR spectrum in C_6D_6 would result in the high field shift of H_a and H_b , while the oxazole methyl signal remains virtually unchanged. On the other hand, $\text{Eu}(\text{thd})_3$ would bring about the low field shift of H_a , H_b , and the methyl protons. Indeed, the spectrum in C_6D_6 resulted in the high field shift of H_a and H_b by 0.60 and 0.43 ppm, respectively, keeping the methyl signal virtually unchanged. Meanwhile, $\text{Eu}(\text{thd})_3$ (0.2 molar amount) brought about the low field shift of H_a , H_b , and methyl signals by 0.52, 0.45, and 0.56 ppm, respectively. The ASIS and $\text{Eu}(\text{thd})_3$ -induced shift for **4b** were similar to those described above. Meanwhile, H_a of **4b'** remote from the coordinated benzene showed a negligibly small ASIS (0.03 ppm low field shift), whereas H_b moved to a high field by 0.59 ppm. The above methods were also used for the determination of the stereochemistry of the other 1,3-dipolar cycloadducts described below.

The reaction of **1** with an equimolar amount of diazomethane at 0°C produced the single adduct **4c**¹¹ in 93% yield (Scheme 3). In the ^1H NMR of **4c**, H_a appeared as a d/d ($J = 9.5/8.0$ Hz) due to the coupling with the methylene protons in agreement with the assigned regiochemistry. Incidentally, diphenyldiazomethane, trimethylsilyl azide, and phenyl azide failed to react with **1**.



SCHEME 3 1,3-Dipolar cycloaddition of **1** with diazomethane.

Benzonitrile *N*-phenylimide (**6**), generated from **5**,¹⁸ added to **1** to furnish **4d**¹¹ in about 50% yield (Scheme 4).¹⁹ The structure of **4d** was determined by X-ray diffraction analysis (Figure 2).²⁰ Interestingly, methyl protons and methyl carbons of the *tert*-butyl group attached to the sp^3 carbon appeared as a broad singlet both in the ^1H and ^{13}C NMR spectra, probably because of the restricted rotation.





SCHEME 4 1,3-Dipolar cycloaddition of **1** with nitrile imine **6**.

Next, reactions with allyl type 1,3-dipoles were investigated. Heating equimolar amounts of **1** and nitron **7** in refluxing toluene provided the sole adduct **4e**¹¹ in 61% yield (Scheme 5). Similarly, the reaction with nitron **8** gave the adduct **4f**¹¹ in 73% yield.


The reaction of **1** with azomethine ylide **9**,²¹ generated from three molar amounts of amine **10**, gave a 10:1 diastereomeric mixture of **4g** and **4g'**¹¹ in good combined yield; only **4g** was isolated in pure form in 73% yield (Scheme 6).

For the attempted reaction with carbonyl ylide **11**, its precursor tetracyanoethylene oxide **12** reduced **1** to give thiophene **13** and carbonyl cyanide **14**, as was reported by us.²² The reaction with ozone proceeded smoothly, but gave a complex mixture after treatment with Zn in acetic acid, from which a crystalline product, the structure of which could not be determined unambiguously, was isolated in low yield.





SCHEME 5 1,3-Dipolar cycloaddition of **1** with nitrones **7** and **8**.



SCHEME 6 1,3-Dipolar cycloaddition of **1** with azomethine imide **9**.

A Mulliken population analysis (B3LYP/6-31G* level²³) of **1** predicted that the positive charges are located at the 3- and 4-positions, while the negative charges are located at the 2- and 5-positions (Figure 3). Thus, the observed regioselectivity is in agreement with the electronic demand, where the positive end of 1,3-dipoles adds to the 2-position, and the negative end adds to the 3-position. The observed regioselectivity is also in harmony with the steric demand that is brought about by bulky *tert*-butyl group. Only in the case of diazomethane, the steric demand governs the regiochemistry, where the less bulky negative end (diazo nitrogen) added to the congested 3-position, while the bulkier negative end (CH₂) added to the less congested 2-position.

As described above, the 1,3-dipolar cycloadditions of **1** took place preferentially at the *syn*- π -face to the S=O bond. This would best be explained as follows. The 1-oxide **1** has a bent structure at C₁ and C₄ with a tilt angle of 9.3° (Figure 4). Thus, for the *syn*- π -face addition, the transition state can be easily reached with a smaller conformational




FIGURE 2 ORTEP plot of molecular structure of **4d** in the crystal. Ellipsoids are drawn at 50% probability level.




FIGURE 3 Mulliken population analysis of **1** (B3LYP/6-31G* level).




FIGURE 4 Conformational changes required at the transition states.

change of **1**, whereas, for the *anti*-face addition, a larger conformational change is required to reach the transition state, where the inversion at C₁ and C₄ is required.²⁻⁴ Accordingly, the activation energy of the reaction would be smaller for the *syn*- π -face addition than for the *anti*- π -face addition, thus making the former process more favorable.

REFERENCES

- [1] W. Oppolzer, In: *Comprehensive Organic Synthesis*, L. A. Paquette (Ed.) (Pergamon Press, Oxford, 1991), Vol. 5, Chapter 4.1.
- [2] T. Otani, J. Takayama, Y. Sugihara, A. Ishii, and J. Nakayama, *J. Am. Chem. Soc.*, **125**, 8255 (2003).
- [3] J. Takayama, S. Fukuda, Y. Sugihara, A. Ishii, and J. Nakayama, *Tetrahedron Lett.*, **44**, 5159 (2003).
- [4] J. Takayama, Y. Sugihara, T. Takayanagi, and J. Nakayama, *Tetrahedron Lett.*, **46**, 4165 (2005).
- [5] J. Nakayama, S. Aoki, J. Takayama, A. Sakamoto, Y. Sugihara, and A. Ishii, *J. Am. Chem. Soc.*, **126**, 9085 (2004).
- [6] M. Burdisso, R. Gandolfi, P. Pevarello, A. L. Poppi, and A. Rastelli, *Tetrahedron Lett.*, **26**, 4653 (1985).
- [7] H. Landen, B. Margraf, and H.-D. Martin, *Tetrahedron Lett.*, **29**, 6593 (1988).
- [8] H. Landen, B. Margraf, H.-D. Martin, and A. Steigel, *Tetrahedron Lett.*, **29**, 6597 (1988).
- [9] H. Hake, H. Landen, H.-D. Martin, B. Mayer, A. Steigel, G. Distefano, and A. Modelli, *Tetrahedron Lett.*, **29**, 6601 (1988).
- [10] T. Mukaiyama and T. Hoshino, *J. Am. Chem. Soc.*, **82**, 5339 (1960).
- [11] Supporting information of cycloadducts **4a-4g** (satisfactory elemental analyses were obtained except **4d** whose analysis sample could not be supplied¹⁹). **4a**: colorless crystals; mp 193–194 °C; IR (KBr) ν 1048 cm⁻¹ (S=O); ¹H NMR (CDCl₃) δ 1.05 (s, 9H), 1.35 (s, 9H), 2.18 (d, 3H, *J* = 1.1 Hz), 4.86 (d, 1H, *J* = 1.1 Hz, H_a), 6.61 (s, 1H, H_b); ¹H NMR (C₆D₆) δ 1.74 (s, 9H), 1.11 (s, 9H), 2.14 (d, 3H, *J* = 1.5 Hz), 4.26 (d, 1H, *J* = 1.5 Hz, H_a), 6.18 (s, 1H, H_b); ¹H NMR (CDCl₃/0.2 mol eq. Eu(thd)₃) δ 1.19 (s, 9H), 1.47 (s, 9H), 2.73 (broad s, 3H), 5.38 (broad s, 1H, H_a), 7.06 (broad s, 1H, H_b); ¹³C NMR (CDCl₃) δ 13.9, 27.9, 32.6, 36.1, 36.6, 74.9, 108.4, 134.4, 153.2, 163.1. **4b**: colorless crystals; mp 167–168 °C; IR (KBr) ν 1048 cm⁻¹ (S=O); ¹H NMR (CDCl₃) δ 1.14 (s, 9H), 1.41 (s, 9H), 5.54 (s, 1H, H_a), 6.61 (s, 1H, H_b), 7.35–7.40 (m, 3H), 7.68–7.75 (m, 2H); ¹H NMR (C₆D₆) δ 0.80 (s, 9H), 1.12 (s, 9H), 5.14 (s, 1H, H_a), 6.14 (s, 1H, H_b), 7.02–7.12 (m, 3H), 7.80 (d, 2H, *J* = 1.1); ¹H NMR (CDCl₃/0.2 mol eq. Eu(thd)₃) δ 1.29 (s, 9H), 1.53 (s, 9H), 6.04 (broad s, 1H, H_a), 7.10 (broad s, 1H, H_b), 7.32–7.35 (m, 3H), 8.32 (br, 2H); ¹³C NMR (CDCl₃) δ 27.6, 32.5, 36.2, 37.6, 73.4, 111.0, 127.1, 128.3, 128.7, 130.0, 134.4, 153.0, 163.5. **4b'**: colorless crystals; mp 169–170 °C; IR (KBr) ν 1039 cm⁻¹ (S=O); ¹H NMR (CDCl₃) δ 1.25 (s, 9H), 1.41 (s, 9H), 4.80 (s, 1H, H_a), 6.63 (s, 1H, H_b), 7.44–7.50 (m, 3H), 7.79–7.86 (m, 2H); ¹H NMR (C₆D₆) δ 1.06 (s, 9H), 1.08 (s, 9H), 4.83 (s, 1H, H_a), 6.04 (s, 1H, H_b), 7.01–7.04 (m, 3H), 7.90–7.93 (m, 2H); ¹H NMR (CDCl₃/0.2 mol eq. Eu(thd)₃) δ 1.39 (s, 9H), 1.46 (s, 9H), 5.25 (s, 1H, H_a), 6.81 (s, 1H, H_b), 7.43–7.47 (m, 3H), 7.90–7.95 (m, 2H); ¹³C NMR (CDCl₃) δ 27.2, 32.5, 36.7, 37.3, 79.8, 111.0, 126.9, 127.8, 129.1, 130.3, 130.7, 151.8, 166.8. **4c**: colorless crystals; mp 153–154 °C; IR (KBr) ν 1031 cm⁻¹ (S=O); ¹H NMR (CDCl₃) δ 1.12 (s, 9H), 1.34 (s, 9H), 3.83 (dd, 1H, *J* = 8.0, 9.5 Hz,

H_a), 4.78 (dd, 1H, $J = 8.0$, 18.7 Hz), 4.81 (dd, 1H, $J = 9.5$, 18.7 Hz), 6.35 (s, 1H, H_b); ¹H NMR (C₆D₆) δ 0.74 (s, 9H), 1.03 (s, 9H), 2.99 (dd, 1H, $J = 8.3$, 9.8 Hz, H_a), 4.43 (dd, 1H, $J = 9.8$, 19.0 Hz), 4.93 (dd, 1H, $J = 8.1$, 18.8 Hz), 5.87 (s, 1H, H_b); ¹H NMR (CDCl₃/0.2 mol eq. Eu(thd)₃) δ 1.16 (s, 9H), 1.49 (s, 9H), 4.35 (t, 1H, $J = 9.0$ Hz, H_a), 5.30 (dd, 1H, $J = 9.5$, 18.8 Hz), 5.92 (br, 1H), 6.98 (broad s, 1H, H_b); ¹³C NMR (CDCl₃) δ 28.4, 33.4, 36.4, 37.6, 60.1, 74.1, 119.4, 134.7, 161.7. **4d**: colorless crystals; ¹H NMR (CDCl₃) δ 0.84 (s, 9H), 1.25 (broad s, 9H), 4.87 (s, 1H, H_a), 6.84 (s, 1H, H_b), 7.28–7.31 (m, 4H), 7.33–7.38 (m, 2H), 7.59–7.61 (m, 2H, $J = 7.6$ Hz), 7.80–7.82 (m, 2H, $J = 7.8$ Hz); ¹³C NMR (CDCl₃) δ 28.1 (broad s), 33.7, 37.2, 40.1, 74.6, 97.0, 125.7, 128.2, 128.66, 128.72, 128.9, 131.2, 131.8, 135.4, 143.5, 146.9, 168.6. **4e**: colorless crystals; mp 127–128 °C; IR (KBr) ν 1038 cm⁻¹ (S=O); ¹H NMR (CDCl₃): δ 1.09 (s, 9H), 1.40 (s, 9H), 2.61 (s, 3H), 4.33 (d, 1H, $J = 7.6$ Hz, H_c), 4.61 (d, 1H, $J = 7.6$ Hz, H_a), 6.41 (s, 1H, H_b), 7.20–7.37 (m, 3H), 7.42–7.48 (m, 2H); ¹H NMR (C₆D₆) δ 0.87 (s, 9H), 1.20 (s, 9H), 2.46 (s, 3H), 4.35 (d, 1H, $J = 8.3$ Hz, H_a), 4.57 (d, 1H, $J = 8.8$ Hz, H_c), 6.04 (s, 1H, H_b), 7.07–7.22 (m, 3H), 7.67 (d, 2H, $J = 6.8$ Hz); ¹H NMR (CDCl₃, 0.2 mol eq. Eu(fod)₃. fod = 6,6,7,7,8,8,8-heptafluoro-2,2-dimethyl-3,5-octanedionate) δ 1.27 (s, 9H), 1.55 (s, 9H), 2.91 (s, 3H), 4.34 (broad s, 1H, H_c), 5.37 (m, 1H, H_a), 7.02 (broad s, 1H, H_b), 7.32–7.46 (m, 3H), 8.27 (broad s, 2H); ¹³C NMR (CDCl₃) δ 28.0, 33.1, 35.9, 37.7, 43.1, 70.8, 78.0, 104.0, 128.0, 128.5, 128.7, 132.8, 137.1, 165.1. **4f**: colorless crystals; mp 193.5–194.5 °C; IR (KBr) ν 1040 cm⁻¹ (S=O); ¹H NMR (CDCl₃) δ 1.02 (s, 9H), 1.34 (s, 9H), 1.68–1.83 (m, 2H), 1.95–2.14 (m, 2H), 2.77–2.89 (m, 1H, $J = 13.6$ Hz), 3.38–3.48 (m, 1H, $J = 13.6$ Hz), 4.04 (d, 1H, $J = 3.3$ Hz, H_a), 4.18–4.26 (m, 1H, $J = 4.2$ Hz, H_c), 6.39 (s, 1H, H_b); ¹H NMR (C₆D₆) δ 0.79 (s, 9H), 1.26 (s, 9H), 1.52–1.84 (m, 4H), 2.39–2.54 (m, 1H, $J = 13.6$ Hz), 3.15–3.25 (m, 1H, $J = 13.6$ Hz), 3.60 (d, 1H, $J = 3.4$ Hz, H_a), 4.45–4.54 (m, 1H, $J = 4.2$ Hz, H_c), 6.14 (s, 1H, H_b); ¹H NMR (CDCl₃/0.2 mol eq. Eu(thd)₃) δ 1.08 (s, 9H), 1.39 (s, 9H), 1.69–1.84 (m, 2H), 2.00–2.19 (m, 2H), 2.84–2.95 (m, 1H), 3.44–3.53 (m, 1H), 4.24 (broad s, 1H, H_a), 4.58–4.69 (m, 1H, H_c), 6.61 (s, 1H, H_b); ¹³C NMR (CDCl₃) δ 25.0, 28.1, 32.3, 32.4, 35.90, 35.91, 57.0, 62.4, 78.4, 104.0, 133.5, 166.0. **4g**: colorless crystals; mp 128–129 °C; IR (KBr) ν 1044 cm⁻¹ (S=O); ¹H NMR (CDCl₃) δ 1.00 (s, 9H), 1.27 (s, 9H), 2.66 (d, 1H, $J = 9.9$ Hz), 2.84–3.03 (m, 3H), 3.61 (s, 2H), 4.12 (t, 1H, $J = 8.3$ Hz, H_a), 6.35 (s, 1H, H_b), 7.23–7.31 (m, 5H); ¹H NMR (C₆D₆) δ 0.71 (s, 9H), 0.96 (s, 9H), 2.50 (d, 1H, $J = 9.6$ Hz), 2.65–2.72 (m, 2H), 3.24–3.41 (m, 3H), 3.66 (t, 1H, $J = 8.1$ Hz, H_a), 6.07 (s, 1H, H_b), 7.23–7.31 (m, 5H); ¹H NMR (CDCl₃/0.2 mol eq. Eu(thd)₃) δ 1.09 (s, 9H), 1.36 (s, 9H), 2.84 (m, 1H), 3.07–3.20 (m, 3H), 3.71 (s, 2H), 4.43 (broad s, 1H, H_a), 6.70 (broad s, 1H, H_b), 7.23–7.35 (m, 5H); ¹³C NMR (CDCl₃) δ 29.1, 33.5, 36.0, 36.5, 51.7, 60.1, 61.0, 66.4, 76.0, 127.0, 128.2, 128.5, 133.1, 138.4, 166.1. **4g'**: ¹H NMR (CDCl₃) δ 1.15 (s, 9H), 1.29 (s, 9H), 2.42 (dd, 1H, $J = 7.4$, 9.9 Hz), 2.53 (d, 1H, $J = 9.9$ Hz), 2.79 (d, 1H, $J = 9.9$ Hz), 3.11 (t, 1H, $J = 9.2$ Hz, H_a), 3.47–3.54 (m, 3H), 6.50 (s, 1H, H_b), 7.23–7.31 (m, 5H).

- [12] G. W. Perold, A. P. Steyn, and F. V. K. von Reiche, *J. Am. Chem. Soc.*, **79**, 462 (1957).
 [13] C. H. Green and D. G. Hellier, *J. Chem. Soc., Perkin Trans. 2*, 458 (1972).
 [14] C. H. Green and D. G. Hellier, *J. Chem. Soc., Perkin Trans. 2*, 243 (1973).
 [15] C. H. Green and D. G. Hellier, *J. Chem. Soc., Perkin Trans. 2*, 190 (1975).
 [16] G. W. Buchanan and D. G. Hellier, *Can. J. Chem.*, **54**, 1428 (1976).
 [17] J. G. Pritchard and P. C. Lauterbur, *J. Am. Chem. Soc.*, **83**, 2105 (1961).
 [18] R. Huisgen, M. Seidel, G. Wallbillich, and H. Knupfer, *Tetrahedron*, **17**, 3 (1962).
 [19] *Caution!* The precursor compound **5** caused a severe skin eruption. Therefore, the yield of **4d** was not optimized.
 [20] Crystal data for **4d**: C₂₅H₃₀N₂OS, $M = 406.57$, orthorhombic, space group *Pbca*; $a = 16.6749(19)$, $b = 10.7803(6)$, $c = 24.0477(14)$ Å; $Z = 8$; $V = 4322.8(4)$ Å³; $D_c = 1.249$

g/cm^3 , $\mu = 0.168 \text{ mm}^{-1}$; measured reflections 30101, independent reflections 5199 [$R(\text{int}) = 0.0863$], $R = 0.0788$, $R_w = 0.1523$, $\text{GOF} = 1.091$. Crystallographic data for the structural analysis have been deposited at the Cambridge Crystallographic Data Center, CCDC No. 671816 for **4d**. Copies of this information can be obtained from The Director, DDCD, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: +44-1223-336033; e-mail: deposit@ccdc.cam.ac.uk or <http://www.ccdc.cam.ac.uk>).

- [21] Y. Terao, H. Kotaki, N. Imai, and K. Achiwa, *Chem. Pharm. Bull.*, **33**, 2762 (1985).
- [22] J. Nakayama, A. Tai, S. Iwasa, T. Furuya, and Y. Sugihara, *Tetrahedron Lett.*, **46**, 1395 (2005).
- [23] Although a crystalline product was isolated in low yield, its structure could not be determined unambiguously.
- [24] M. J. Frisch et al., Gaussian, Inc., Pittsburgh, PA, 1998.