This article was downloaded by: [University of Leeds] On: 21 August 2013, At: 09:22 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



### Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lcar20</u>

## Synthesis of $\alpha, \omega$ -Diazidoalditol Derivatives via Both bis- or tris-Cyclic Sulfites and Peracetylated $\alpha, \omega$ -Dibromoalditols as Bielectrophilic Intermediates

Virginie Glaçon<sup>a</sup>, Mohammed Benazza<sup>a</sup>, Aniss El Anzi<sup>a</sup>, Daniel Beaupère<sup>a</sup> & Gilles Demailly<sup>a</sup>

<sup>a</sup> Laboratoire des Glucides, CNRS FRE 2779, Université de Picardie Jules Verne, 33 rue Saint-Leu, F-80039, Amiens, France Published online: 18 Aug 2006.

To cite this article: Virginie Glaon , Mohammed Benazza , Aniss El Anzi , Daniel Beaupre & Gilles Demailly (2004) Synthesis of  $\alpha, \omega$ -Diazidoalditol Derivatives via Both bis- or tris-Cyclic Sulfites and Peracetylated  $\alpha, \omega$ -Dibromoalditols as Bielectrophilic Intermediates, Journal of Carbohydrate Chemistry, 23:2-3, 95-110, DOI: <u>10.1081/CAR-120030470</u>

To link to this article: http://dx.doi.org/10.1081/CAR-120030470

### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

### Synthesis of α,ω-Diazidoalditol Derivatives via Both bis- or tris-Cyclic Sulfites and Peracetylated α,ω-Dibromoalditols as Bielectrophilic Intermediates

Virginie Glaçon, Mohammed Benazza,\* Aniss El Anzi, Daniel Beaupère, and Gilles Demailly

Laboratoire des Glucides, CNRS FRE 2779, Université de Picardie Jules Verne, Amiens, France

#### ABSTRACT

The  $\alpha,\omega$ -diazidoalditol derivatives with *erythro*, *threo*, *xylo*, *ribo*, D-*arabino*, D-*manno*, and D-*gluco* configuration were efficiently synthesized, respectively, from *bis*- or *tris*-cyclic sulfite or peracetylated  $\alpha,\omega$ -dibromoalditol intermediates. The cyclic sulfite intermediates has the advantage to lead directly to the free  $\alpha,\omega$ -diazido- $\alpha,\omega$ -dideoxyalditols.

*Key Words:* Diazidoalditol; Anhydroalditol; Dibromoalditol; Cyclic sulfite; Bielectrophilic intermediate; Azidation.

#### **INTRODUCTION**

The  $\alpha,\omega$ -diazidoalditol derivatives are potential precursors of corresponding diamino derivatives used as copolymers in polyamide syntheses,<sup>[1]</sup> chelating reagents,<sup>[2]</sup> and inhibitors of HIV-1 retrovirus proteases.<sup>[3]</sup>

95

DOI: 10.1081/CAR-120030470 Copyright © 2004 by Marcel Dekker, Inc. 0732-8303 (Print); 1532-2327 (Online) www.dekker.com

<sup>\*</sup>Correspondence: Mohammed Benazza, Laboratoire des Glucides, CNRS FRE 2779, Université de Picardie Jules Verne, 33 rue Saint-Leu F-80039, Amiens, France; Fax: +33(3)22827561; E-mail: mohammed.benazza@sc.u-picardie.fr.

	REPRINTS
--	----------

Azidation reactions are usually carried out by nucleophilic azide substitution for some leaving groups such as sulfonates,<sup>[3,4]</sup> cyclic sulfites,<sup>[5]</sup> and cyclic sulfates,<sup>[6]</sup> by opening of epoxides,<sup>[7]</sup> more directly using PPh<sub>3</sub>/*N*-chlorosuccinimide and azide salts.<sup>[8]</sup> The most commonly used syntheses of  $\alpha, \omega$ -diazidoalditols involve *bis*-epoxides,<sup>[7]</sup> *bis*-sulfonates,<sup>[9]</sup> or *bis*- halogenated<sup>[10]</sup> intermediates obtained from partially protected alditols.

In this article, we report two efficient syntheses of  $\alpha, \omega$ -diazidoalditols (Sch. 1). The first one used the *bis*-cyclic sulfite alditol derivatives (Path 1), and the second one used the peracetylated  $\alpha, \omega$ -dibromoalditol derivatives as key intermediates (Path 2).<sup>[11]</sup>

The alditol cyclic sulfite synthesis was realized with the N,N'-di-imidazolylthione (Im<sub>2</sub>SO) and alditols in THF as solvent. The *bis*-cyclic sulfite of erythritol **2**, D,L-threitol **8**, xylitol **12**, ribitol **20**, and D-arabinitol **28** derivatives (Sch. 2) were obtained, respectively, from the corresponding tetritols (**1** and **7**) and pentitols (**11**, **19**, and **27**) in excellent yields (>95%). With the hexitols [D-mannitol (**33**) and D-glucitol (**39**)], the *tris*-cyclic sulfite derivatives **34** and **40** were isolated in almost quantitative yield (Sch. 3). In each case, a mixture of *bis*- or *tris*-cyclic sulfite derivatives was obtained. The chiral nature of the sulfite, which is responsible for the complex mixture of isomers, could be overcome by oxidizing the products to sulfates. This was especially carried out with tetritol derivatives **2** and **8** which led, respectively, to only one *bis*-cyclic sulfate derivative by oxidation.<sup>[12]</sup> With pentitols and hexitols sulfite, any interesting results were accomplished until now.

Treatment of the *bis*-cyclic sulfite derivative of erythritol **2**, by NaN<sub>3</sub> (6 equiv.) in DMF under optimal conditions (Table 1, entry 1), led to the 1,4-diazido-1,4-dideoxyerythritol (**3**) in excellent yield (80%). A regioselective monoazidation leading to 1-azido-D,L-erythritol **5** (59%) was observed when the reaction was carried out at room temperature during 36 hr (entry 2). With D,L-*threo bis*-cyclic sulfite **8**, the diazido derivative **9** was obtained in excellent yield (89%) (entry 3). No trace of monoazido derivative was observed even when the reaction occurred at room temperature.

With the pentitols, the azidation reaction has appeared slower because 3 hr were needed for a complete disappearance of the substrate. The diazidopentitol derivatives **13** (*xylo*), **21** (*ribo*), and **29** (D-*arabino*) (Sch. 2) were obtained in reasonable to good yields (72%, 56%, and 55%, respectively) (Table 1, entries 4–6). As by-products, the azidoanhydro derivatives **17** (D,L-*xylo*), **25** (D,L-*ribo*), and **31** (D-*arabino*) were observed (entries 4–6). In the case of the compound **32** (acetylated derivative of **31**), the coupling constant  $J_{23} = 0.4$  Hz is in favor of a trans configuration resulting from a regioselective



Copyright @ Marcel Dekker, Inc. All rights reserved



Downloaded by [University of Leeds] at 09:22 21 August 2013



Scheme 2. (i) Im<sub>2</sub>SO, THF; (ii) NaN<sub>3</sub>, DMF; (iii) Ac<sub>2</sub>O, pyridine.

1,4-*O*-heterocyclization. This suggests an initial regioselective nucleophilic attack on the C-5 carbon atom of **28** by an azide ion to give **28a** (Path 1) or **28b** (Path 2) as intermediates (Sch. 4). The subsequent *O*-heterocyclization involving the C-1 as electrophilic site and the O-4 as intramolecular nucleophilic atom led to **31** identified as their peracetylated derivative **32**.

The 1,5-diazido derivative **30** (D-*arabino*) (acetylated derivative of **29**) has a *syn*methine ( $J_{23} = 2.5 \text{ Hz}$ ) and *anti*-methine ( $J_{34} = 8.5 \text{ Hz}$ ) coupling sequence in agreement with the planar and zigzag structure.<sup>[13,14]</sup>

With D-mannitol (**33**) and D-glucitol (**39**) (Sch. 3), the azidation was performed with the corresponding *tris*-cyclic sulfite derivatives **34** and **40**, respectively. With the D-*manno* derivative **34**, the azidation at 130°C gave mainly 1,6-diazido-1,6-dideoxy-D-mannitol (**35**) in 47% yield and as by-product the 1,6-diazido-3,4-O-sulfinyl **37** isolated in 39% yield (Table 2, entry 1). The subsequent acetylation of **37** gave quantitatively **38**.

ORDER		REPRINTS
-------	--	----------





Scheme 3. (i) Im<sub>2</sub>SO, THF; (ii) NaN<sub>3</sub>, DMF; (iii) Ac<sub>2</sub>O, pyridine; (iv) MeONa, MeOH.

At lower temperature (90°C) and with a prolonged time (2 hr) (entry 2), the yield of the 3,4-O-sulfinyl derivative **37** reached 58% while the yield of **35** was decreased to 15%. These results suggest the initial formation of **37** which undergoes subsequently a partial hydrolysis to lead to **35** in 49% yield. The yield of **35** was increased to 67% by treatment of the crude product after azidation by MeONa/MeOH during 24 hr at room temperature (Table 2, entry 3).

The *tris*-cyclic sulfite formation was avoided by a 3,4-di-*O*-benzyl protection leading to **43**<sup>[15]</sup> (Sch. 5). Surprisingly, the azidation of the 3,4-di-*O*-benzyl-1,2 : 5,6-di-*O*-sulfinyl-D-mannitol (**44**) led to the unexpected 2,5-anhydro-6-azido-6-deoxy-D-glucitol (**47**) in excellent yield (89%) through a 2,5-*O*-heterocylization. This compound was identified in its acetylated form **48** which in <sup>1</sup>H NMR showed a coupling constant  $J_{3,4} = 1$  Hz in agreement with a *trans*-configuration of H-3 and H-4.

This 2,5-*O*-heterocyclization could be explained by the mechanism shown in the Sch. 6. Thus, the 5-*exo-tet O*-heterocyclization occurred regioselectively on the C-2 ( $\equiv$ C-5) with an inversion of the configuration from the possible intermediates **44a** (Path 1) or **44b** (Path 2). This type of intramolecular cyclization has already been reported in the literature with

Table 1. Azidation reaction of compounds 2, 8, 12, 20, and 28.

					Isolated yield (%)			
Entry	Substrate	NaN <sub>3</sub> (equiv.)	<i>t</i> (°C)	Time (hr)	Diazido	Monoazido	Anhydro	
1	<b>2</b> (erythro)	6	130	1	<b>3</b> (80)	_	_	
2	2 (erythro)	6	Rt	36	3 (15)	<b>5</b> (59)	_	
3	8 (D,L-threo)	6	130	1	9 (89)		_	
4	<b>12</b> ( <i>xylo</i> )	6	130	3	13 (72)	<b>15</b> (10)	17 (10)	
5	<b>20</b> ( <i>ribo</i> )	6	130	3	21 (56)	<b>23</b> (9)	25 (9)	
6	<b>28</b> (D-arabino)	6	130	3	<b>29</b> (55)	_	<b>31</b> (10)	





Scheme 4.

1,2:5,6-*bis*-epoxyhexane<sup>[16]</sup> and 1,2:5,6-*bis*-aziridine-D-mannitol.<sup>[17]</sup> In all cases the 6-*endo-tet* process are disfavored. This is in agreement with Baldwin's rules.<sup>[18]</sup>

Unlike the *tris*-cyclic sulfite derivative **34** with a *manno* configuration, the azidation of the *tris*-cyclic sulfite of D-glucitol **40** led to a complex mixture of 1,6-diazido-*O*-sulfinyl regio- and stereoisomers (entry 4). When the azidation reaction was followed by methanolysis (MeONa/MeOH), 1,6-diazido-1,6-dideoxy-D-glucitol (**41**) was isolated in 52% yield.

To complement our work on the  $\alpha,\omega$ -regioselective azidation of alditols, another method for diazidoalditol synthesis was considered. This involved the peracetylated  $\alpha,\omega$ -dibromoalditol derivatives **49**, **50**, **51**, **52**, **53**, **54**, and **55** as bielectrophilic intermediates (Table 3). The latter were readily afforded by direct bromination of unprotected alditols using AcBr in 1,4-dioxane.<sup>[11]</sup> The bromination of the alditols **1**, **7**, **19**, **27**, **33**, and **39**, by this reagent, led in a one-pot reaction to complex mixtures of partially acetylated regioisomeric  $\alpha,\omega$ -dibromo- $\alpha,\omega$ -dideoxyalditol derivatives. A subsequent acetylation, followed by a treatment of the resulting crude product by NaN<sub>3</sub> in DMF at 80°C overnight, led to the peracetylated  $\alpha,\omega$ -diazido- $\alpha,\omega$ -dideoxyalditol derivatives **4** (*erythro*, 75%), **10** (D,L-*threo*,

						Isolated yield (%	6)
Entry	Sulfite substrate	NaN <sub>3</sub> (equiv.)	t (°C)	Time	Diazido	Diazido-3,4- <i>O</i> -sulfinyl	Anhydro
1	<b>34</b> (D-manno)	4	130	15 min	<b>35</b> (47)	<b>37</b> (39)	_
2	<b>34</b> (D-manno)	4	90	2 hr	35 (15)	37 (58)	
3	<b>34</b> (D-manno)	4	130	15 min	<b>35</b> $(67)^{a}$		
4	<b>40</b> (D-gluco)	4	130	15 min	Compl	ex mixture	_
5	<b>40</b> (D-gluco)	4	130	15 min	<b>41</b> (52) <sup>a</sup>	—	—

Table 2. Azidation reaction of compounds 34 and 40.

<sup>a</sup>Isolated after treatment with MeONa, MeOH, rt, 24 hr.





Scheme 5. (i) Im<sub>2</sub>SO, THF; (ii) NaN<sub>3</sub>, DMF; (iii) Ac<sub>2</sub>O, pyridine; (iv) MeONa, MeOH.

76%), **14** (*xylo*, 64%), **22** (*ribo*, 50%), **30** (D-*arabino*, 70%), **36** (D-*manno*, 51%), and **42** (D-*gluco*, 42%) previously obtained from cyclic sulfite intermediates (Schs. 2 and 3).

In conclusion, we have reported two short and efficient syntheses of  $\alpha$ , $\omega$ -diazidoalditols from both *bis*- and *tris*-cyclic sulfite derivatives and  $\alpha$ , $\omega$ -dibromoalditols as intermediates. The synthesis via *bis*-cyclic sulfite intermediates was the preferred route because of the high yields of free diazido derivatives obtained in two steps from free alditols. The obtention in excellent yield of the unexpected and interesting 2,5-anhydro-6-azido-6-deoxy derivative **47** (89%) with D-*gluco* configuration from the tandem azidation-2,5-*O*-heterocyclization of 3,4-di-*O*-benzyl-*bis*-*O*-sulfinyl-D-mannitol **44** should also be noted.



Scheme 6.



**Table 3.** Structure of peracetylated  $\alpha, \omega$ -dibromoalditol derivatives.

Br OAc OAc Br	AcO-Br OAc Br	AcOOAc	Br OAc OAc OAc Br	AcO-Br OAc OAc Br	Br AcO AcO OAc	AcO
<b>49</b>	<b>50</b>	51	<b>52</b>	53	D-manno)	55
(Erythro)	(Threo)	(Xylo)	( <i>Ribo</i> )	(D-arabino)		(D-gluco)

#### **EXPERIMENTAL**

#### **General Methods**

Melting points were determined with a Büchi 535 apparatus and are uncorrected. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker 300 WB spectrometer; chemical shifts are reported in  $\delta$  (ppm) relative to Me<sub>4</sub>Si. All <sup>13</sup>C NMR signals were assigned through C,H-correlated spectra with hsqcgrad.routine experiment. TLC was performed on Silica Gel 60 F<sub>254</sub> 230 mesh (E. Merck) with hexane–EtOAc or CH<sub>2</sub>Cl<sub>2</sub>–acetone as eluent, and detection by the vanillin–H<sub>2</sub>SO<sub>4</sub> reagent. The silica gel used in column chromatography was 35–70  $\mu$  (Amicon). Elemental analyses were performed by the "Service de Microanalyse du CNRS" (Laboratoire de Chimie Bioorganique, Université de Reims Champagne Ardenne).

# General Procedure I for the Preparation of *bis*-Cyclic Sulfite Derivatives of Alditols 2, 8, 12, 20, 28, 34, 40, and 44 (GPI)

Freshly distilled thionyl chloride (3 equiv. for tetritols **1** and **7**; 4 equiv. for pentitols and hexitols **11**, **19**, **27**, **33**, **39**, and **44**) was added dropwise to a solution of imidazole (12-16 equiv.) in THF (40 mL) under argon atmosphere at 0°C. The filtrate containing the diimidazolethionyl was directly added dropwise to a solution of alditol (1 g) in THF (10 mL) at  $-10^{\circ}$ C. The mixture was stirred during 30 min. The residue obtained after the removal of the solvent in vacuum was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and washed twice by cold water, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated under reduced pressure at 25°C to obtain the mixture of stereoisomeric *O*-sulfinyl derivatives.

**Di-O-sulfinylerythritol (2).** Reaction of **1** according to GPI gave **2** (1.74 g, 99%); colorless; m.p. 50–53°C;  $R_{\rm f}$  0.45 and 0.55 in 3 : 2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  81.2, 79.6, 79.2, 77.7 (C-2,3), 71.1, 70.3, 69.0, 68.6 (C-1,4); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.14–5.08 (m), 4.85–4.58 (m).

**Di-O-sulfinyl-D,L-threitol (8)**. Reaction of **7** according to GPI gave **8** (1.72 g, 98%); syrup;  $R_{\rm f}$  0.25, 0.31, and 0.49 in 7:3 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  81.1, 80.1, 78.0, 76.9 (C-2,3), 68.6, 68.5, 67.7, 67.2, (C-1,4); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.18–4.90 (m), 4.80–4.39 (m).

**Di-O-sulfinylxylitol Derivatives (12).** Reaction of **11** according to GPI gave **12** (1.54 g, 96%), white powder;  $R_{\rm f}$  0.31–0.54 in 9:1 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  84.7–76.6 (C-2,3,4), 70.5–68.8 (CH<sub>2</sub>OSO–), 63.0–61.2 (C-1,5).

ORDER		REPRINTS
-------	--	----------

**Di-O-sulfinylribitol (20)**. Reaction of **19** according to GPI gave **20** (1.57 g, 98%); white powder;  $R_{\rm f}$  0.44–0.90 in 9 : 1 CH<sub>2</sub>Cl<sub>2</sub>–acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  84.0–75.0 (C-2,3,4), 70.7–66.4 (C-1,5), 61.9–58.0 (CH<sub>2</sub>OH).

**Di-O-sulfinyl-D-arabinitol (28)**. Reaction of **27** according to GPI gave **28** (1.56 g, 97%); white powder;  $R_{\rm f}$  0.45–0.89 in 9 : 1 CH<sub>2</sub>Cl<sub>2</sub>–acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  86.5–76.3 (C-2,3,4), 71.2–67.0 (CH<sub>2</sub>OSO), 62.8–60.0 (C-1,5).

**Tri-O-sulfinyl-D-mannitol (34)**. Reaction of **33** according to GPI gave **34** (1.74 g, 99%); syrup;  $R_{\rm f}$  0.75 in 96:04 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  82.8–77.9 (C-2,3,4,5), 71.0–67.6 (C-1,6).

**Tri-O-sulfinyl-D-glucitol (40)**. Reaction of **39** according to GPI gave **40** (1.74 g, 99%); syrup;  $R_{\rm f}$  0.51–0.76 in 96:04 CH<sub>2</sub>Cl<sub>2</sub>–acetone; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  85.2–78.2 (C-2,3,4,5), 71.5–67.4 (C-1,6).

**3,4-Di-O-benzyl-1,2 : 5,6-di-O-sulfinyl-D-mannitol** (44). Reaction of 43 according to GPI gave 44. A part of the mixture of the stereoisomers crystallized in hexane as solvent. Thus, two different mixtures of stereoisomers were obtained. One resulting from the crystalization (a) and the second resulting from the recovered filtrate after concentration (b). (a): Yield 70 mg (60%); colorless crystal;  $R_{\rm f}$  0.69 and 0.74 in CH<sub>2</sub>Cl<sub>2</sub>; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  137.3–128.2 (Ph), 82.4-77.6 (C-2,3,4,5), 75.5, 75.3, 75.2 (CH<sub>2</sub>Ph), 68.5, 68.2 (C-1,6). (b): 0.45 g; yield 39%;  $R_{\rm f}$  0.54 and 0.63 in CH<sub>2</sub>Cl<sub>2</sub>; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  137.3–128.7 (Ph), 82.7–77.6 (C-2,3,4,5), 75.5–74.2 (CH<sub>2</sub>Ph), 69.2–67.8 (C-1,6).

#### General Procedure II for the Azidation of Alditol Cyclic-Sulfite Derivatives (GPII)

Sodium azide (NaN<sub>3</sub>, 6 equiv., 39.5 mmol) was added to a solution of *bis*-cyclic sulfite derivatives (1 g) in DMF (50 mL) during the desired time. The crude product obtained after concentration was then dissolved in EtOH, filtered, and the filtrate was evaporated and chromatographed on silica gel.

**1,4-Diazido-1,4-dideoxyerythritol (3).** According to GPII, the reaction was carried out at 130°C during 1 hr. The crude product was chromatographed on silica gel (1:4 hexane–EtOAc). Compound **3** (1.13 g, 80%); colorless crystals; m.p. 85–86°C;  $R_{\rm f}$  0.50 in 7:3 hexane–EtOAc; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  73.7 (C-2,3), 56.1 (C-1,4); <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  3.64 (dd, 2H,  $J_{1,2}$ ,  $J_{3,4}$  5.9 Hz,  $J_{1',2}$ ,  $J_{3,4'}$  1.6 Hz, H-2,3), 3.49 (dd, 2H,  $J_{1,1'}$ ,  $J_{4,4'}$  12.8 Hz, H-1',4').

Anal. Calcd for  $C_4H_8N_6O_2$ : C, 27.91; H, 4.68; N, 48.82; O, 18.59. Found: C, 28.01; H, 4.76; N, 48.75.

**1-Azido-1-deoxy-D,L-erythritol (5).** According to GPII, the reaction was carried out at room temperature during 36 hr. The crude product was chromatographed on silica gel (2 : 8 hexane–EtOAc). Compound **5** (0.71 g, 59%); syrup;  $R_{\rm f}$  0.20 in EtOAc; <sup>13</sup>C NMR (C<sub>5</sub>D<sub>5</sub>N):  $\delta$  74.8 (C-3), 74.0 (C-2), 65.2 (C-4), 56.2 (C-1); <sup>1</sup>H NMR (C<sub>5</sub>D<sub>5</sub>N):  $\delta$  4.45 (m, 1H,  $J_{2,3}$  4.9 Hz, H-2), 4.36 (m, 1H,  $J_{3,4}$  5.7 Hz,  $J_{3,4'}$  6.1 Hz, H-3), 4.15 (dd, 1H,  $J_{4,4'}$  12.4 Hz, H-4'), 4.10 (dd, 1H, H-4), 4.01 (dd, 1H,  $J_{1',2}$  1.7 Hz,  $J_{1,1'}$  12.5 Hz, H-1'), 3.93 (dd, 1H,  $J_{1,2}$  5.8 Hz, H-1).

Anal. Calcd for C<sub>4</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub>: C, 32.65; H, 6.17; N, 28.56; O, 32.62. Found: C, 32.49; H, 6.21; N, 28.61.

Downloaded by [University of Leeds] at 09:22 21 August 2013



#### Synthesis of α,ω-Diazidoalditol Derivatives

**1,4-Diazido-1,4-dideoxy-D,L-threitol (9).** According to GPII, the reaction was carried out at 130°C during 1 hr. The crude product was chromatographed on silica gel (2:8 hexane–AcOEt). Compound **9** (1.27 g, 89%); syrup;  $R_{\rm f}$  0.42 in 7:3 hexane–EtOAc; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  70.7 (C-2,3), 53.5(C-1,4); <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  3.69 (dd, 2H,  $J_{1,2}$ ,  $J_{3,4}$  4.8 Hz, H-2,3), 3.45 (dd, 2H,  $J_{1',2}$ ,  $J_{3,4'}$  3.6 Hz, H-1',4'), 3.38 (dd, 2H,  $J_{1,1'}$ ,  $J_{4,4'}$  12.4 Hz, H-1,4).

Anal. Calcd for C<sub>4</sub>H<sub>8</sub>N<sub>6</sub>O<sub>2</sub>: C, 27.91; H, 4.68; N, 48.82; O, 18.59. Found: C, 27.82; H, 4.72; N, 48.94.

**1,5-Diazido-1,5-dideoxyxylitol (13)**. According to GPII, the reaction was carried out at 130°C during 3 hr. The crude product was chromatographed on silica gel (85:15 CH<sub>2</sub>Cl<sub>2</sub>–acetone). Compound **13** (0.96 g, 72%); syrup;  $R_{\rm f}$  0.38 in 7:3 CH<sub>2</sub>Cl<sub>2</sub>–acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  73.8 (C-3), 73.2 (C-2,4), 54.5 (C-1,5); <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  3.83 (dd, 2H,  $J_{1,2}$ ,  $J_{4,5}$  6.1 Hz, H-2,4), 3.51 (t, 1H,  $J_{2,3}$ ,  $J_{3,4}$  4.0 Hz, H-3), 3.41 (dd, 2H,  $J_{1',2}$ ,  $J_{4,5'}$  2.4 Hz, H-1',5'), 3.38 (dd, 2H,  $J_{1,1'}$ ,  $J_{5,5'}$  11.2 Hz, H-1,5).

Anal. Calcd for  $C_5H_{10}N_6O_3$ : C, 29.70; H, 4.99; N, 41.57; O, 23.74. Found: C, 29.81; H, 4.92; N, 41.63.

**1-Azido-1-deoxy-D,L-xylitol (15).** Yield 0.12 g (10%); syrup;  $R_f$  0.33 in 7:3 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  71.2, 70.8, 65.8 (C-2,3,4), 63.3 (C-5), 55.0 (C-1).

Anal. Calcd for C<sub>5</sub>H<sub>11</sub>N<sub>3</sub>O<sub>4</sub>: C, 33.90; H, 6.26; N, 23.72; O, 36.12. Found: C, 33.79; H, 6.43; N, 23.80.

**1,4-Anhydro-5-azido-5-deoxy-D,L-xylitol (17)**. Yield 0.10 g (10%); syrup;  $R_{\rm f}$  0.33 in 7 : 3 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  80.7, 78.5, 77.9 (C-4,3,2), 74.4 (C-1), 51.4 (C-5).

Anal. Calcd for  $C_5H_9N_3O_3$ : C, 37.74; H, 5.70; N, 26.40; O, 30.16. Found: C, 37.63; H, 5.75; N, 26.32.

**1,5-Diazido-1,5-dideoxy-ribitol (21)**. According to GPII, the reaction was carried at 130°C during 3 hr. The crude product was chromatographed on silica gel (85:15 CH<sub>2</sub>Cl<sub>2</sub>-acetone). Compound **21** (0.74 g, 56%); syrup;  $R_{\rm f}$  0.44 in 7:3 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  73.9 (C-3), 73.3 (C-2,4), 54.9 (C-1,5); <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  4.72 (s, 3H, OH), 3.75 (ddd, 2H,  $J_{1,2}$ ,  $J_{4,5}$  6.8 Hz, H-2,4), 3.48 (t, 1H,  $J_{2,3}$ ,  $J_{3,4}$  6.5 Hz, H-3), 3.41 (dd, 2H,  $J_{1',2}$ ,  $J_{4,5'}$  3.2 Hz, H-1',5'), 3.38 (dd, 2H,  $J_{1,1'}$ ,  $J_{5,5'}$  12.7 Hz, H-1,5).

Anal. Calcd for  $C_5H_{10}N_6O_3$ : C, 29.70; H, 4.99; N, 41.57; O, 23.74. Found: C, 29.73; H, 5.07; N, 41.47.

**1-Azido-1-deoxy-D,L-ribitol (23).** Yield 0.10 g (9%); syrup;  $R_{\rm f}$  0.40 in 7:3 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  73.1, 72.9, 65.8 (C-2,3,4), 64.7 (C-5), 56.1 (C-1).

Anal. Calcd for C<sub>5</sub>H<sub>11</sub>N<sub>3</sub>O<sub>4</sub>: C, 33.90; H, 6.26; N, 23.72; O, 36.12. Found: C, 33.74; H, 6.39; N, 23.65.

**1,4-Anhydro-5-azido-5-deoxy-D,L-ribitol (25)**. Yield 0.09 g (9%); syrup;  $R_f$  0.40 in 7:3 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  83.0, 75.3, 75.1 (C-4,3,2), 75.0 (C-1), 54.4 (C-5).

Anal. Calcd for  $C_5H_9N_3O_3$ : C, 37.74; H, 5.70; N, 26.40; O, 30.16. Found: C, 37.70; H, 5.65; N, 26.53.

**1,5-Diazido-1,5-dideoxy-D-arabinitol (29).** According to GPII, the reaction was carried at  $130^{\circ}$ C during 3 hr. The crude product was chromatographed on silica gel (85:15 CH<sub>2</sub>Cl<sub>2</sub>-acetone). Compound **29** (0.73 g, 55%); colorless crystals; m.p.

ORDER		REPRINTS
-------	--	----------

85–86°C,  $[\alpha]_D^{22}$  – 29.5° (*c* 1.10 in MeOH); *R*<sub>f</sub> 0.50 in 7:3 CH<sub>2</sub>Cl<sub>2</sub>–acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD): δ 72.8 (C-3), 71.7 (C-4), 70.4 (C-2), 55.6 (C-5), 55.2 (C-1); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 4.78 (s, 3H, OH), 3.99 (m, 1H, *J*<sub>1,2</sub> 5.1 Hz, *J*<sub>2,3</sub> 2.7 Hz, H-2), 3.80 (m, 1H, *J*<sub>3,4</sub> 8.2 Hz, *J*<sub>4,5</sub>' 2.9 Hz, H-4), 3.54 (dd, 1H, *J*<sub>5,5</sub>' 13.0 Hz, H-5'), 3.42 (m, 3H, H-1',3,5), 3.30 (dd, 1H, *J*<sub>1,1</sub>' 12.5 Hz, H-1).

Anal. Calcd for C<sub>5</sub>H<sub>10</sub>N<sub>6</sub>O<sub>3</sub>: C, 29.70; H, 4.99; N, 41.57; O, 23.74. Found: C, 29.75; H, 4.96; N, 41.70.

**1,4-Anhydro-5-azido-5-deoxy-D-arabinitol (31).** Yield 0.10 g (10%); syrup;  $R_{\rm f}$  0.45 in 7:3 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  85.7, 81.0, 80.4 (C-4,3,2), 74.5 (C-1), 53.6 (C-5).

Anal. Calcd for C<sub>5</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub>: C, 37.74; H, 5.70; N, 26.40; O, 30.16. Found: C, 37.68; H, 5.79; N, 26.49.

**1,6-Diazido-1,6-dideoxy-D-mannitol (35).** According to GPII, the reaction was carried out at 130°C during 15 min. The crude product was chromatographed on silica gel (CH<sub>2</sub>Cl<sub>2</sub>). Compound **35** (0.60 g, 47%); colorless crystals; m.p.  $92-93^{\circ}C$ ;  $[\alpha]_{D}^{28}$  +18.4° (*c* 1.00 in MeOH),  $R_{\rm f}$  0.36 in 6:4 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (D<sub>2</sub>O):  $\delta$  72.1 (C-2,5), 71.9 (C-3,4), 56.4 (C-1,6); <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta$  4.78 (s, 4H, OH), 3.91 (m, 2H,  $J_{1,2}$ ,  $J_{5,6}$  6.0 Hz, H-2,5), 3.83 (d, 2H,  $J_{2,3}$ ,  $J_{4,5}$  9.0 Hz, H-3,4), 3.68 (dd, 2H,  $J_{1',2}$ ,  $J_{5,6'}$  2.3 Hz, H-1',6'), 3.52 (dd, 2H,  $J_{1,1'}$ ,  $J_{6,6'}$  13.0 Hz, H-1,6).

Anal. Calcd for  $C_6H_{12}N_6O_4$ : C, 31.04; H, 5.21; N, 36.19; O, 27.56. Found: C, 31.15; H, 5.33; N, 36.12.

**1,6-Diazido-1,6-dideoxy-3,4-***O***-sulfinyl-D-mannitol (37)**. According to GPII, the reaction was carried out at 90°C during 2 hr. The crude product was chromatographed on silica gel (CH<sub>2</sub>Cl<sub>2</sub>). Compound **37** (0.89 g, 58%); syrup;  $[\alpha]_D^{23} + 103.5^\circ$  (*c* 1.80 in CH<sub>2</sub>Cl<sub>2</sub>),  $R_f$  0.65 in 9 : 1 CH<sub>2</sub>Cl<sub>2</sub>-acetone; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  84.5 (C-4), 82.0 (C-3), 71.2 (C-5), 70.3 (C-2), 53.0 (C-1,6); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  4.92 (dd, 1H,  $J_{2,3}$  7.5 Hz,  $J_{3,4}$  5.0 Hz, H-3), 4.64 (dd, 1H,  $J_{4,5}$  7.0 Hz, H-4), 4.09 (m, 1H,  $J_{5,6}$  5.3,  $J_{5,6'}$  3.8 Hz, H-5), 3.79 (m, 1H,  $J_{1,2}$  3.5 Hz,  $J_{1',2}$  5.8 Hz, H-2), 3.62–3.42 (m, 4H, H-1,6).

Anal. Calcd for C<sub>6</sub>H<sub>10</sub>N<sub>6</sub>O<sub>5</sub>S: C, 25.90; H, 3.62; N, 30.20; O, 28.75; S, 11.52. Found: C, 25.78; H, 3.73; N, 30.33.

**1,6-Diazido-1,6-dideoxy-D-glucitol (41)**. According to GPII, the reaction was carried out at 130°C during 20 min. After evaporation of the solvent, the crude product was hydrolyzed by a catalytic amount of MeONa in 40 mL of MeOH at rt during one night under argon. The crude product was then neutralized by an acid resin Amberlyst 15 wet and filtered. The filtrate was evaporated under reduced pressure and chromatographed on silica gel (3 : 2 CH<sub>2</sub>Cl<sub>2</sub>–acetone). Compound **41** (0.66 g, 52%); colorless crystals; m.p. 62.5–63.5°C,  $[\alpha]_D^{19}$  +13.4° (*c* 0.70 in MeOH),  $R_f$  0.40 in 6 : 4 CH<sub>2</sub>Cl<sub>2</sub>–acetone; <sup>13</sup>C NMR (D<sub>2</sub>O):  $\delta$  74.1 (C-2), 73.5 (C-4), 72.1 (C-3), 71.3 (C-5), 55.9 (C-6), 55.4 (C-1); <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta$  4.79 (s, 4H, OH), 3.96 (m, 1H, H-2), 3.90 (m, 1H, H-5), 3.85 (dd, 1H,  $J_{2,3}$  6.0 Hz, H-3), 3.65 (dd, 1H,  $J_{3,4}$  1.7 Hz, H-4), 3.61 (dd, 1H,  $J_{5,6'}$  1.9 Hz, H-6'), 3.52 (dd, 1H,  $J_{1,1'}$  13.8,  $J_{1',2}$  3.3 Hz, H-1'), 3.50 (dd, 1H,  $J_{5,6}$  7.2 Hz,  $J_{6,6'}$  13.0 Hz, H-6), 3.47 (dd, 1H,  $J_{1,2}$  6.3 Hz, H-1).

Anal. Calcd for  $C_6H_{12}N_6O_4$ : C, 31.04; H, 5.21; N, 36.19; O, 27.56. Found: C, 31.12; H, 5.18; N, 36.26.

**1,6-Diazido-3,4-di**-*O*-benzyl-1,6-dideoxy-D-mannitol (45). According to GPII, the reaction was carried out at 90°C during 16 hr. The crude product was chromatographed on silica gel (9:1 hexane–EtOAc). Compound 45 (0.11 g, 10%); syrup;  $R_{\rm f}$  0.60 in 7:3



#### Synthesis of α,ω-Diazidoalditol Derivatives

hexane – EtOAc; <sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  140.5 (Ph-ipso), 130.5, 130.1, 129.7 (Ph), 81.4 (C-3,4), 76.7 (<u>CH<sub>2</sub>Ph</u>), 72.4 (C-2,5), 56.6 (C-1,6); <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  7.30 (m, 10H, Ph), 4.68 (m, 4H, CH<sub>2</sub>Ph), 3.98 (m, 2H,  $J_{2,3}$  7.5 Hz,  $J_{3,4}$  5.0 Hz, H-2,5), 3.88 (d, 2H,  $J_{2,3}$ ,  $J_{4,5}$  8.5 Hz, H-3,4), 3.55 (dd, 2H,  $J_{1,1'}$ ,  $J_{6,6'}$  12.6 Hz,  $J_{1',2}$ ,  $J_{5,6'}$  2.8 Hz, H-1',6'), 3.37 (dd, 2H,  $J_{1,2}$ ,  $J_{5,6}$  5.4 Hz, H-1,6).

Anal. Calcd for  $C_{20}H_{24}N_6O_4$ : C, 58.25; H, 5.83; N, 20.39. Found: C, 58.32; H, 5.85; N, 19.98.

**2,5-Anhydro-1-azido-3,4-di**-*O*-benzyl-1-deoxy-D-glucitol (47). According to GPII, the reaction was carried out at 90°C during 16 hr. The crude product was chromatographed on silica gel (9 : 1 hexane–EtOAc). Compound **47** (0.91 g, 89%); syrup;  $R_f$  0.43 in 7 : 3 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  137.5 (Ph-ipso), 128.6, 128.0, 127.7 (Ph), 83.9, 83.1, 82.4, 81.6 (C-4,5,3,2), 72.0, 71.9 (CH<sub>2</sub>Ph), 61.3 (C-6), 52.6 (C-1); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.34 (m, 10H, Ph), 4.52 (q, 4H, CH<sub>2</sub>Ph), 4.71–3.86 (4m, 6H, H-5,4,3,2,6,6'), 3.30 (dd, 2H, H-1,1'), 3.05 (m, 1H, OH).

Anal. Calcd for C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O<sub>4</sub>: C, 65.03; H, 6.28; N, 11.37; O, 17.32. Found: C, 64.93; H, 6.33; N, 11.29.

#### **General Procedure III for Acetylation (GPIII)**

After azidation, the crude product was directly dissolved in pyridine (20 mL) and an excess of acetic anhydride was added dropwise. After one night at room temperature, the solvent was evaporated under reduced pressure. The residue was dissolved in  $CH_2Cl_2$  (15 mL) and washed twice by 2 × 15 mL of HCl (0.1 M), 2 × 15 mL of saturated HCO<sub>3</sub>Na and 2 × 15 mL of water. The organic layer was evaporated and chromatographed on silica gel.

**2,3-Di-***O***-acetyl-1,4-diazido-1,4-dideoxyerythritol** (4). Reaction of crude product obtained from **2** according to GPII followed by GPIII gave **4** after chromatography on silica gel (8 : 2 hexane-AcOEt). Compound **4** (1.85 g, 88%); colorless crystals; m.p. 62–63°C;  $R_{\rm f}$  0.50 in 8 : 2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.4 (CO), 70.7 (C-2,3), 50.3 (C-1,4), 20.5 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.10 (m, 2H, H-2,3), 3.50 (dd, 2H,  $J_{1',2}$ ,  $J_{3,4'}$  5.4 Hz, H-1',4'), 3.40 (dd, 2H,  $J_{1,2}$ ,  $J_{3,4}$  2.9 Hz,  $J_{1,1'}$ ,  $J_{4,4'}$  13.4 Hz, H-1,4), 2.08 (s, 6H, CH<sub>3</sub>).

Anal. Calcd for  $C_8H_{12}N_6O_4$ : C, 37.50; H, 4.72; N, 32.80; O, 24.98. Found: C, 37.70; H, 4.80; N, 32.53.

**2,3,4-Tri-O-acetyl-1-azido-1-deoxy-D,L-erythritol** (6). Yield 0.71 g (65%); syrup;  $R_{\rm f}$  0.20 in 8 : 2 hexane – EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.9, 169.4, 169.3 (CO), 69.9 (C-3), 69.3 (C-2), 61.3 (C-4), 50.1 (C-1), 20.2 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.25 (m, 1H,  $J_{2,3}$  5.2 Hz, H-3), 5.00 (m, 1H, H-2), 4.25 (dd, 1H,  $J_{3,4'}$  3.3 Hz,  $J_{4,4'}$  12.2 Hz, H-4'), 4.10 (dd, 1H,  $J_{3,4}$  5.5 Hz, H-4), 3.32 (dd, 1H,  $J_{1,2}$  4.3,  $J_{1,1'}$  13.5 Hz, H-1'), 3.25 (dd, 1H,  $J_{1,2}$  8.8 Hz, H-1), 1.89, 1.87, 1.82 (3s, 9H, CH<sub>3</sub>).

Anal. Calcd for C<sub>10</sub>H<sub>15</sub>N<sub>3</sub>O<sub>6</sub>: C, 43.96; H, 5.53; N, 15.38; O, 35.13. Found: C, 43.63; H, 6.01; N, 15.53.

**2,3-Di-***O***-acetyl-1,4-diazido-1,4-dideoxy-D,L-threitol** (10). Reaction of crude product obtained from **8** according to GPII followed by GPIII gave **10** after chromato-graphy on silica gel (8 : 2 hexane–AcOEt). Compound **10** (1.99 g, 95%); syrup;  $R_{\rm f}$  0.42 in 8 : 2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.7 (CO), 70.4 (C-2,3), 50.3 (C-1,4),



	EPRINTS
--	---------

20.4 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.06 (m, 2H, H-2,3), 3.33 (dd, 2H,  $J_{1',2}$ ,  $J_{3,4'}$  4.5 Hz, H-1',4'), 3.26 (dd, 2H,  $J_{1,2}$ ,  $J_{3,4}$  7.2 Hz,  $J_{1,1'}$ ,  $J_{4,4'}$  13.3 Hz, H-1,4), 2.00, 1.97 (2s, 6H, CH<sub>3</sub>).

Anal. Calcd for  $C_8H_{12}N_6O_4$ : C, 37.50; H, 4.72; N, 32.80; O, 24.98. Found: C, 37.93; H, 4.62; N, 32.14.

**2,3,4-Tri-O-acetyl-1,5-diazido-1,5-dideoxyxylitol** (14). Reaction of crude product obtained from 12 according to GPII followed by GPIII gave 14 after chromatography on silica gel (9:1 hexane–AcOEt). 14 (1.62 g, 75%); colorless crystals; m.p. 54–55°C;  $R_{\rm f}$  0.40 in 8:2 hexane–AcOEt; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.8, 169.7 (CO), 69.9 (C-2,4), 69.7 (C-3), 50.7 (C-1,5), 20.5, 20.4 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.30 (t, 1H,  $J_{2,3}$ ,  $J_{3,4}$  5.2 Hz, H-3), 5.05 (m, 2H,  $J_{1,2}$ ,  $J_{4,5}$  5.8 Hz, H-2,4), 3.40 (dd, 2H,  $J_{1',2}$ ,  $J_{4,5'}$  4.1 Hz, H-1',5'), 3.32 (dd, 2H,  $J_{1,1'}$ ,  $J_{5,5'}$  13.2 Hz, H-1,5), 2.10, 2.00 (2s, 3H, 6H, CH<sub>3</sub>).

Anal. Calcd for  $C_{11}H_{16}N_6O_6$ : C, 40.25; H, 4.91; N, 25.60; O, 29.24. Found: C, 40.53; H, 4.61; N, 25.71.

**2,3,4,5-Tetra-O-acetyl-1-azido-1-deoxy-D,L-xylitol** (16). Yield 0.23 g (10%); syrup;  $R_{\rm f}$  0.60 in 6:4 hexane–AcOEt; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  170.0, 169.8 (CO), 69.8 (C-2), 69.2 (C-3), 68.5 (C-4), 61.6 (C-5), 50.1 (C-1), 20.3, 20.1, 19.9 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.30 (dd, 1H,  $J_{2,3}$  5.0,  $J_{3,4}$  5.3 Hz, H-3), 5.10 (ddd, 1H, H-4), 5.01 (ddd, 1H, H-2), 4.22 (dd, 1H,  $J_{4,5'}$  4.0 Hz,  $J_{5,5'}$  12.2 Hz, H-5'), 3.89 (dd, 1H,  $J_{4,5}$  6.3 Hz, H-5), 3.38 (dd, 1H,  $J_{1',2}$  4.3,  $J_{1,1'}$  13.5 Hz, H-1'), 3.33 (dd, 1H,  $J_{1,2}$  6.0 Hz, H-1), 2.00, 1.97, 1.95, 1.93 (4s, 12H, CH<sub>3</sub>).

Anal. Calcd for  $C_{13}H_{19}N_3O_8$ : C, 45.22; H, 5.55; N, 12.17; O, 37.07. Found: C, 45.33; H, 5.64; N, 12.51.

**2,3-Di-***O***-acetyl-1,4-anhydro-5-azido-5-deoxy-D,L-xylitol** (18). Yield 0.18 g (11%); syrup;  $R_{\rm f}$  0.35 in 8:2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  168.6 (CO), 77.4 (C-4), 76.3 (C-2), 75.2 (C-3), 70.7 (C-1), 48.7 (C-5), 19.7 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.00 (dd, 1H,  $J_{2,3}$  0.6 Hz,  $J_{3,4}$  3.6 Hz, H-3), 4.84 (m, 1H, H-2), 3.98 (dd, 1H, H-4), 3.85 (dd, 1H,  $J_{1',2}$  4.8 Hz,  $J_{1,1'}$  10.8 Hz, H-1'), 3.50 (dd, 1H,  $J_{1,2}$  2.0 Hz, H-1), 3.30 (m, 2H,  $J_{5,5'}$  11.9 Hz, H-5,5'), 1.89, 1.85 (2s, 12H, CH<sub>3</sub>).

Anal. Calcd for  $C_9H_{13}N_3O_5$ : C, 44.44; H, 5.39; N, 17.28; O, 32.89. Found: C, 44.53; H, 5.73; N, 17.62.

**2,3,4-Tri-O-acetyl-1,5-diazido-1,5-dideoxyribitol (22)**. Reaction of crude product obtained from **20** according to GPII followed by GPIII gave **22** after chromatography on silica gel (9 : 1 hexane–AcOEt). 1.25 g (58%); syrup;  $R_{\rm f}$  0.42 in 8 : 2 hexane–AcOEt; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.8, 169.3 (CO), 69.5 (C-2,4), 69.3 (C-3), 49.4 (C-1,5), 19.7, 19.6 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.23 (m, 2H, H-2,4), 5.13 (t, 1H,  $J_{2,3}$ ,  $J_{3,4}$  5.2 Hz, H-3), 3.48 (dd, 2H,  $J_{1',2}$ ,  $J_{4,5'}$  3.5 Hz, H-1',5'), 3.37 (dd, 2H,  $J_{1,2}$ ,  $J_{4,5}$  6.4 Hz,  $J_{1,1'}$ ,  $J_{5,5'}$  13.5 Hz, H-1,5), 2.07, 2.05 (2s, 6H, 3H, CH<sub>3</sub>).

Anal. Calcd for C<sub>11</sub>H<sub>16</sub>N<sub>6</sub>O<sub>6</sub>: C, 40.25; H, 4.91; N, 25.60; O, 29.24. Found: C, 40.63; H, 4.51; N, 25.65.

**2,3,4,5-Tetra-O-acetyl-1-azido-1-deoxy-D,L-ribitol** (24). Yield 0.23 g (10%); syrup;  $R_f$  0.63 in 6:4 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  170.2, 169.7, 169.5 (CO), 70.1 (C-2), 69.5 (C-3), 69.2 (C-4), 61.3 (C-5), 49.7 (C-1), 20.7, 20.6, 20.5 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.25 (ddd, 1H,  $J_{2,3}$  5.2 Hz,  $J_{3,4}$  5.3 Hz, H-3), 5.15 (m, 1H, H-4), 5.05 (m, 1H, H-2), 4.20 (dd, 1H,  $J_{4,5'}$  3.0,  $J_{5,5'}$  12.5 Hz, H-5'), 3.99 (dd, 1H,  $J_{4,5}$  6.0 Hz, H-5), 3.41 (dd, 1H,  $J_{1,2}$  3.5 Hz,  $J_{1,1'}$  13.2 Hz, H-1'), 3.30 (dd, 1H,  $J_{1,2}$  6.9 Hz, H-1), 2.05, 2.00 (2s, 6H, 6H, CH<sub>3</sub>).



Downloaded by [University of Leeds] at 09:22 21 August 2013

Anal. Calcd for C<sub>13</sub>H<sub>19</sub>N<sub>3</sub>O<sub>8</sub>: C, 45.22; H, 5.55; N, 12.17; O, 37.07. Found: C, 45.73; H, 5.51; N, 12.29.

**2,3-Di-***O***-acetyl-1,4-anhydro-5-azido-5-deoxy-D,L-ribitol** (26). Yield 0.21 g (13%); syrup; R<sub>f</sub> 0.35 in 8 : 2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.6, 169.5 (CO), 77.9 (C-4), 72.5 (C-2), 71.3 (C-3), 70.1 (C-1), 49.5 (C-5), 19.9, 19.8 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.05 (m, 1H,  $J_{2,3}$  5.5 Hz,  $J_{3,4}$  6.3 Hz, H-2), 4.95 (d, 1H, H-3), 3.90 (dd, 1H, H-4), 3.85 (dd, 1H,  $J_{1,2}$  5.0 Hz,  $J_{1,1'}$  10.2 Hz, H-1'), 3.60 (dd, 1H,  $J_{1,2}$  3.4 Hz, H-1), 3.45 (dd, 1H,  $J_{4,5'}$  5.6 Hz,  $J_{5,5'}$  12.5 Hz, H-5'), 3.33 (dd, 1H,  $J_{4,5}$  3.7 Hz, H-5), 1.85, 1.83 (2s, 6H, CH<sub>3</sub>).

Anal. Calcd for  $C_9H_{13}N_3O_5$ : C, 44.44; H, 5.39; N, 17.28; O, 32.89. Found: C, 44.48; H, 5.41; N, 17.35.

**2,3,4-Tri-O-acetyl-1,5-diazido-1,5-dideoxy-D-arabinitol** (**30**). Reaction of crude product obtained from **28** according to GPII followed by GPIII gave **30** after chromato-graphy on silica gel (9:1 hexane–EtOAc). Yield 1.12 g (52%); colorless crystals; m.p.  $29-30^{\circ}$ C,  $[\alpha]_{D}^{22} + 38.0^{\circ}$  (*c* 1.40 in CH<sub>2</sub>Cl<sub>2</sub>),  $R_{\rm f}$  0.41 in 8:2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.5, 169.4, 169.2 (CO), 69.5 (C-2), 68.9 (C-3), 68.7 (C-4), 50.7 (C-5), 50.3 (C-1), 20.0, 19.9, 19.8 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.31 (dd, 1H,  $J_{2,3}$  2.5 Hz,  $J_{3,4}$  8.5 Hz, H-3), 5.19 (m, 1H, H-4), 5.08 (m, 1H, H-2), 3.45 (dd, 1H,  $J_{1',2}$  3.3 Hz,  $J_{1,1'}$  13.6 Hz, H-1'), 3.33 (dd, 1H,  $J_{4,5'}$  5.2 Hz,  $J_{5,5'}$  13.0 Hz, H-5'), 3.29 (dd, 1H,  $J_{4,5}$  7.0 Hz, H-5), 3.24 (dd, 1H,  $J_{12}$  5.3 Hz, H-1), 2.06, 2.01 (2s, 6H, 3H, CH<sub>3</sub>).

Anal. Calcd for  $C_{11}H_{16}N_6O_6$ : C, 40.25; H, 4.91; N, 25.60; O, 29.24. Found: C, 40.88; H, 4.59; N, 25.65.

**2,3-Di-***O***-acetyl-1,4-anhydro-5-azido-5-deoxy-D-arabinitol** (**32**). Yield 0.24 g (15%); syrup;  $R_{\rm f}$  0.37 in 8:2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  168.6 (CO), 83.5 (C-4), 78.5 (C-3), 77.8 (C-3), 71.9 (C-1), 51.7 (C-5), 20.7 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.11 (m, 1H, H-2), 4.95 (dd, 1H,  $J_{2,3}$  0.4 Hz,  $J_{3,4}$  3.5 Hz, H-3), 3.99 (d, 2H,  $J_{1,2}$ ,  $J_{1',2}$  2.7 Hz, H-1,1'), 3.87 (ddd, 1H, H-4), 3.52 (dd, 1H,  $J_{4,5'}$  5.9 Hz,  $J_{5,5'}$  13.1 Hz, H-5'), 3.37 (dd, 1H,  $J_{4,5}$  3.2'Hz, H-5), 2.02 (s, 6H, CH<sub>3</sub>).

Anal. Calcd for C<sub>9</sub>H<sub>13</sub>N<sub>3</sub>O<sub>5</sub>: C, 44.44; H, 5.39; N, 17.28; O, 32.89. Found: C, 44.28; H, 5.62; N, 17.11.

**2,3,4,5-Tetra-***O***-acetyl-1,6-diazido-1,6-dideoxy-D-mannitol (36)**. Reaction of crude product obtained from **34** according to GPII followed by GPIII gave **36** after chromatography on silica gel (8 : 2 hexane–EtOAc). 1.10 g (50%); syrup;  $[\alpha] - D^{25} + 33.0^{\circ}$  (*c* 1.00 in CH<sub>2</sub>Cl<sub>2</sub>),  $R_{\rm f}$  0.39 in 7 : 3 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  168.8, 168.7 (CO), 67.5 (C-3,4), 67.1 (C-2,5), 49.7 (C-1,6), 19.7, 19.6 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.31 (d, 2H,  $J_{2,3}$ ,  $J_{4,5}$  8.1 Hz, H-3,4), 4.94 (m, 2H, H-2,5), 3.38 (dd, 2H,  $J_{1',2}$ ,  $J_{5,6'}$  3.2 Hz,  $J_{1,1'}$ ,  $J_{6,6'}$  13.5 Hz, H-1',6'), 3.18 (dd, 2H,  $J_{1,2}$ ,  $J_{5,6}$  5.6 Hz, H-1,6), 2.00 (s, 12H, CH<sub>3</sub>).

Anal. Calcd for C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>8</sub>: C, 42.00; H, 5.04; N, 20.99; O, 31.97. Found: C, 42.22; H, 5.14; N, 21.02.

**2,5-Di-***O***-acetyl-1,6-diazido-1,6-dideoxy-3,4-O-sulfinyl-D-mannitol** (38). Reaction of crude product obtained from 34 according GPII at 90°C followed by GPIII gave 38 after chromatography on silica gel (7 : 3 hexane–EtOAc). 1.23 g (62%); syrup;  $[\alpha]_D^{24}$  +86.3°(*c* 4.7 in CH<sub>2</sub>Cl<sub>2</sub>), *R*<sub>f</sub> 0.60 in 7 : 3 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.6 (CO), 81.3 (C-3), 81.2 (C-4), 71.9 (C-2), 70.8 (C-5), 50.2 (C-1), 49.5 (C-6), 20.6 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.20 (m, 1H, H-2), 5.00 (m, 1H, H-5), 4.86 (dd, 1H, *J*<sub>3,4</sub> 5.0 Hz, *J*<sub>4,5</sub> 4.6 Hz, H-4), 4.64 (dd, 1H, *J*<sub>2,3</sub> 7.0 Hz, H-3), 4.60 (dd, 1H, *J*<sub>1',2</sub> 3.4 Hz, *J*<sub>1,1'</sub>

ORDER		REPRINTS
-------	--	----------

15.0 Hz, H-1′), 3.50 (d, 1H, *J*<sub>1,2</sub> 4.8 Hz, H-1) 3.46 (d, 2H, *J*<sub>5,6</sub>, *J*<sub>5,6</sub>′ 4.5 Hz, H-6,6′), 2.07, 2.06 (2s, 6H, CH<sub>3</sub>).

Anal. Calcd for  $C_{10}H_{14}N_6O_7S$ : C, 33.15; H, 3.89; N, 23.20; O, 30.91; S, 8.85. Found: C, 33.52; H, 3.74; N, 22.95.

**2,3,4,5-Tetra-***O***-acetyl-1,6-diazido-1,6-dideoxy-D-glucitol** (**42**). Reaction of crude product obtained from **40** according to GPII followed by GPIII gave **42** after chromatography on silica gel (8 : 2 hexane – EtOAc). 1.16 g (53%); colorless crystals; m.p. 64–65°C,  $[\alpha]_{D}^{25}$  +4.01° (*c* 1.0 in CH<sub>2</sub>Cl<sub>2</sub>), *R*<sub>f</sub> 0.50 in 7 : 3 hexane – EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  168.8, 168.7 (CO), 69.2 (C-2), 68.2 (C-5), 68.0 (C-4), 67.5 (C-3), 49.5 (C-1), 49.2 (C-6); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  5.38 (dd, 1H, *J*<sub>2,3</sub> 7.0 Hz, *J*<sub>3,4</sub> 3.6 Hz, H-3), 5.29 (dd, 1H, *J*<sub>4,5</sub> 7.1 Hz, H-4), 5.03 (m, 1H, H-2), 4.97 (m, 1H, H-5), 3.49 (dd, 1H, *J*<sub>1,1</sub>' 14.0 Hz, *J*<sub>1',2</sub> 2.6 Hz, H-1'), 3.43 (dd, 1H, *J*<sub>5,6'</sub> 3.7 Hz, *J*<sub>6,6',2</sub> 13.4 Hz, H-6'), 3.38 (dd, 1H, *J*<sub>1,2</sub> 5.2 Hz, H-1), 3.30 (dd, 1H, *J*<sub>5,6</sub> 5.6 Hz, H-6), 2.08, 2.04, 2.03, 2.01 (4s, 12H, CH<sub>3</sub>).

Anal. Calcd for C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>8</sub>: C, 42.00; H, 5.04; N, 20.99; O, 31.97. Found: C, 42.33; H, 4.92; N, 20.77.

**2,3-Di-***O***-acetyl-1,6-diazido-3,4-di-***O***-benzyl-1,6-dideoxy-D-mannitol** (46). Reaction of crude product obtained from 44 according to GPII followed by GPIII gave 46 after chromatography on silica gel (9:1 hexane–EtOAc). 0.48 g (9%); syrup;  $R_f$  0.48 in 9:1 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  169.8 (CO), 139.9 (Ph-ipso), 130.0, 129.7, 129.5 (Ph), 76.5 (C-3,4), 71.8 (CH<sub>2</sub>Ph), 70.1 (C-2,5), 51.0 (C-1,6), 20.7 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.28 (m, 10H, Ph), 4.55 (m, 2H, H-2,5), 4.50 (m, 4H, CH<sub>2</sub>Ph), 4.01 (d, 2H,  $J_{2,3}$ ,  $J_{4,5}$  7.8 Hz, H-3,4), 3.35 (dd, 2H,  $J_{1,1'}$ ,  $J_{6,6'}$  13.1 Hz,  $J_{1',2}$ ,  $J_{5,6'}$  3.1 Hz, H-1',6'), 3.18 (dd, 2H,  $J_{1,2}$ ,  $J_{5,6}$  6.1'Hz, H-16), 2.01 (s, 6H, CH<sub>3</sub>).

Anal. Calcd for  $C_{24}H_{28}N_6O_6$ : C, 58.06; H, 5.68; N, 16.93; O, 19.33. Found: C, 58.13; H, 5.65; N, 16.73.

**1-O-Acetyl-2,5-anhydro-6-azido-3,4-di-O-benzyl-6-deoxy-D-glucitol (48)**. Yield 1.02 g (89%); syrup;  $[\alpha]_{D}^{25} + 49.3^{\circ}$  (*c* 1.0 in CH<sub>2</sub>Cl<sub>2</sub>),  $R_{\rm f}$  0.58 in 8 : 2 hexane–EtOAc; <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  170.7 (CO), 137.3 (C-ipso), 128.5, 128.0, 127.8 (Ph), 83.3 (C-4), 83.0 (C-5), 82.4 (C-3), 79.2 (C-2), 71.8 (CH<sub>2</sub> benzyl), 63.1 (C-1), 52.4 (C-6), 20.8 (CH<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.33 (m, 10H, Ph), 4.38–4.58 (m, 5H, CH<sub>2</sub>Ph, H-1'), 4.22–4.29 (m, 2H, H-2,1), 4.07 (dt, 1H,  $J_{1,2}$ ,  $J_{1',2}$ , 5.8 Hz,  $J_{2,3}$ , 2.9 Hz, H-5), 3.99 (dd, 1H,  $J_{3,4}$  1.0 Hz,  $J_{4,5}$ , 3.1 Hz, H-3), 3.91 (dd, 1H, H-4), 3.36 (d, 2H, H-6,6'), 2.03 (s, 3H, CH<sub>3</sub>).

Anal. Calcd for C<sub>22</sub>H<sub>25</sub>N<sub>3</sub>O<sub>5</sub>: C, 64.22; H, 6.12; N, 10.21; O, 19.44. Found: C, 64.33; H, 6.15; N, 10.44.

#### General Procedure for Azidation of α,ω-Dibromoalditols Derivatives

To a solution in DMF (8 mL) of crude product of  $\alpha,\omega$ -dibromoalditols obtained after acetylation from 0.2 g of unprotected alditols,<sup>[11]</sup> was added sodium azide (3 equiv.), and the mixture was stirred during 16 hr at 80°C. The crude product obtained was chromatographed on silica gel with 6:1 hexane–EtOAc as an eluant. The peracetylated  $\alpha,\omega$ -diazidoalditol derivatives **4** (*erythro*, 75%), **10** (D,L-*threo*, 76%), **14** (*xylo*, 64%), **22** (*ribo*, 50%), **30** (D-*arabino*, 70%), **36** (D-*manno*, 51%) et.**42** (D-gluco, 42%) were obtained.



ORDER		REPRINTS
-------	--	----------

#### ACKNOWLEDGMENT

The authors thank the Conseil Régional de Picardie for its financial support.

#### REFERENCES

- (a) Mancera, M.; Roffé, I.; Riva, M.; Silva, C.; Galbis, J.A. Synthesis of D-mannitol and L-iditol derivatives as monomers for the preparation of new regioregular AABBtype polyamides. Carbohydr. Res. 2002, *337*, 607–611; (b) Bird, T.B.; Black, W.A.P.; Dewar, E.T.; Hare, J.P. Polyamides containing carbohydrate residues. J. Chem. Soc. 1963, 1208–1212.
- Samoilova, O.I.; Seleverstova, I.A.; Dyatlova, N.M.; Yashunskii, V.G. Substances with complexing capability. XV. D-Manno-2,3,4,5-tetraoxy-1,6-diaminohexanetetraacetic acid. Zh. Obshch. Khim. 1973, 43, 365–369Chem. Abstr. 1973, 78, 148–165.
- 3. Chenera, B.; Boehm, J.C.; Dreyer, G.B. Synthesis of C2-symmetric and pseudosymmetric HIV-1 protease inhibitors from D-mannitol and D-arabitol. Bioorg. Med. Chem. Lett. **1991**, *1*, 219–222.
- 4. Dureault, A.; Greck, C.; Depezay, J.C. Diastereospecific synthesis of diaziridines from D-mannitol. Access to chiral  $\alpha$ -amino acids. Tetrahedron Lett. **1986**, 27, 4157–4160.
- (a) El Meslouti, A.; Beaupère, D.; Demailly, G.; Uzan, R. One-pot stereoselective synthesis of glycosyl azides via 1,2-cyclic sulfite. Tetrahedron Lett. 1994, 35, 3913–3916; (b) Guiller, A.; Gagnieu, C.H.; Pacheco, H. Substitution of sulfites of cyclic oxides by azide ion. Tetrahedron Lett. 1985, 26, 6343–6344; (c) Gyoung, Y.S.; Jeon, W.S. Nucleophilic substitution reactions of cyclic sulfites. J. Korean Chem. Soc. 1994, 38, 465–468; (d) Lorhay, B.B.; Ahuja, R.J. Synthesis of homochiral amino alcohols, aziridines and diamines via homochiral cyclic sulfites. J. Chem. Soc., Chem. Commun. 1991, 95–97; (e) Dubois, L.; Dodd, R.H. Stereocontrolled synthesis of aziridine-2-lactones from D-ribose and D-lyxose. Tetrahedron 1993, 49, 901–910.
- Van der Klein, P.A.M.; Filemon, W.; Veeneman, G.H.; Van der Marel, G.A.; Van Boom, J.H. Highly regioselective ring opening of five-membered cyclic sulfates with lithium azide: synthesis of azido sugars. J. Carbohydr. Chem. **1992**, *11*, 837–848.
- 7. Le Merrer, Y.; Dureault, A.; Greck, C.; Micas-Languin, D.; Gravier, C.; Depezay, J.C. Synthesis of diepoxides and diaziridines, precursors of enantiomerically pure  $\alpha$ -hydroxy and  $\alpha$ -amino aldehydes or acids, from D-mannitol. Heterocycles **1987**, *25*, 541–548.
- Larabi, M.L.; Fréchou, C.; Demailly, G.; Uzan, R. Direct synthesis of glycosyl azides. Tetrahedron Lett. 1994, 35, 2175–2178.
- (a) Le Merrer, Y.; Gauzy, L.; Gravier-Pelletier, C.; Depezay, J.C. Synthesis of C<sub>2</sub>-symmetric guanidino-sugars as potent inhibitors of glycosidases. Bioorg. Med. Chem. 2000, 8, 307–320; (b) Fitremann, J.; Duréault, A.; Depezay, J.C. 2,5-Disubstituted pyrrolidines from D-mannitol-derived *bis*-aziridines. Tetrahedron Lett. 1994, 35, 1201–1204.

ORDER		REPRINTS
-------	--	----------

- Nazhaoui, M.; Joly, J.P.; Kitane, S.; Berrada, M. 1,6-Dideoxy-D-mannitol-based 20crown-6 ethers: synthesis and influence of the substituents upon complexing properties toward phenylglycinium methyl esters. J. Chem. Soc. Perkin Trans 1 1998, 22, 3845–3850.
- 11. Crombez-Robert, C.; Benazza, M.; Fréchou, C.; Demailly, G. Efficient synthesis of  $\alpha, \omega$ -dibromodieoxyalditols as precursors for  $\alpha, \omega$ -dithioalkyladitols. Carbohydr. Res. **1997**, *303*, 359–365.
- Glacon, V.; Benazza, M.; Beaupère, D.; Demailly, G. Heterocyclisation of free or partially protected alditols via their bis-cyclic sulfate derivatives. Versatile synthesis of aza and thiodeoxyanhydroalditol with *erythro*, *threo*, *arabino*, *gulo*, *talo* or *manno* configuration. Tetrahedron Lett. 2000, 41, 5053–5056.
- Benazza, M.; Beaupère, D.; Uzan, R.; Demailly, G. Selective chlorination of pentitols. Carbohydr. Res. 1991, 218, 75–81.
- Moore, R.E.; Barchi, J.J., Jr.; Bartolini, G. Use of borate complexation in assigning relative stereochemistry of acyclic polyhydroxylated compounds. J. Org. Chem. 1985, 50, 374–379.
- Jurczak, J.; Bauer, T.; Chmielewski, M. A general approach to the synthesis of 2,3-di-O-protected derivatives of D-glyceraldehyde. Carbohydr. Res. 1987, 164, 493–498.
- Jung, M.E.; Kretschik, O. Enantiospecific total synthesis of L-2',3'-dideoxyisonucleosides via regioselective opening of optically active C2-symmetric 1,4-pentadiene bis-epoxide. J. Org. Chem. 1998, 63, 2975–2981.
- McCort, I.; Fort, S.; Duréault, A.; Depezay, J.C. Synthesis and evaluation as glycosidase inhibitors of 2,5-imino-D-glucitol and 1,5-imino-D-mannitol related derivatives. Bioorg. Med. Chem. 2000, 8, 135–143.
- 18. Baldwin, J.E. Rules for ring closure. J. Chem. Soc. Chem. Comm. 1976, 734-736.



## **Request Permission or Order Reprints Instantly!**

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/ Order Reprints" link below and follow the instructions. Visit the <u>U.S. Copyright Office</u> for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on <u>Fair Use in the Classroom</u>.

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our <u>Website</u> User Agreement for more details.

# **Request Permission/Order Reprints**

Reprints of this article can also be ordered at http://www.dekker.com/servlet/product/DOI/101081CAR120030470