

Design and Combinatorial Development of Shield-1 Peptide Mimetics Binding to Destabilized FKBP12

Daniel Madsen, Frederik Præstholt Jørgensen, Daniel Palmer, Milena Roux, Jakob V. Olsen, Mikael Bols, Sanne Schoffelen, Frederik Diness, and Morten Meldal

ACS Comb. Sci., **Just Accepted Manuscript** • DOI: 10.1021/acscombsci.9b00197 • Publication Date (Web): 06 Feb 2020

Downloaded from pubs.acs.org on February 10, 2020

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Design and Combinatorial Development of Shield-1 Peptide Mimetics Binding to Destabilized FKBP12

*Daniel Madsen,^{†‡} Frederik P. Jørgensen,[†] Daniel Palmer,^{†‡} Milena E. Roux,^{||} Jakob V. Olsen,^{||}
Mikael Bols,[†] Sanne Schoffelen,^{†‡} Frederik Diness,^{*†‡} Morten Meldal^{*†‡}*

[†]Department of Chemistry

[‡]Center for Evolutionary Chemical Biology,


^{||} Department of Biology

University of Copenhagen

Universitetsparken 5, 2100 Copenhagen, Denmark

E-mail: meldal@nano.ku.dk, fdi@chem.ku.dk

TOC – for table of contents use only:



1
2
3 ABSTRACT: Based on computational design, a focused one-bead one-compound library has been
4 prepared on microparticle encoded PEGA₁₉₀₀ beads consisting of small tripeptides with a triazole
5 capped *N*-terminal. The library was screened towards a double point-mutated version of the human
6 FKBP12 protein, known as the destabilizing domain (DD). Inspired by the decoded library hits,
7 unnatural peptide structures were screened in a novel on-bead assay, which was useful for a rapid
8 structure evaluation prior to off-bead resynthesis. Subsequently, a series of 19 compounds were
9 prepared and tested using a competitive fluorescence polarization assay, which led to the discovery
10 of peptide-ligands with low micromolar binding affinity towards the DD. The methodology
11 represents a rapid approach for identification of a novel structure scaffold, where the screening
12 and initial structure refinement was accomplished using small quantities of library building blocks.

13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28 KEYWORDS: Destabilizing domain, one-bead one-compound library, encoded beads, solid phase
29 synthesis.

30 31 32 33 34 INTRODUCTION

35
36
37 The engineered and double point-mutated version of the human FK506-binding protein, F36V-
38 L106P-FKBP12, is known as the destabilizing domain (DD).¹ This domain can confer its
39 instability to proteins-of-interests (POIs) when fused to it.^{2, 3} When expressed in cells, the DD and
40 the fusion protein will rapidly be degraded by the proteasome, resulting in low levels of the POI.¹
41 Upon stabilization of the DD by addition of a small ligand, the degradation is reduced which
42 induces a dose-dependent accumulation of the POI within the cell. Hereby, the level of gene
43 product can be chemically controlled, creating a powerful tool for interrogating biological systems.
44
45
46
47
48
49
50
51
52
53 The compound Shield-1 (Shld1, **Figure 1**) is a small molecule derived from the natural ligand
54 FK506 and has been shown to interact selectively with the DD.^{4, 5} The Shld1-DD system has

1
2
3 proven its vast utility in several biological systems, including living mice,⁶ plants,^{7, 8} parasites,^{9,}
4
5 ^{10, 11} mammalian cells,¹ and yeast.¹² Previous studies have reported K_i values of 7.5 nM from *in*
6
7 *vitro* experiments between shld1 and the DD.¹³ However, in plant systems genetically modified to
8
9 express DD-POI fusion proteins, high concentrations of shld1 was required for induce
10
11 accumulation of the destabilized complex.⁷ In studies with rice and wheat, treatment with 3-10 μ M
12
13 Shld1 solutions were required for achieving sufficient diffusion of the small Shld1 molecule
14
15 through the plant cuticle and subsequently stabilization of DD-POI.⁸ A recent study highlighted
16
17 the challenges of improving the plant cell penetrating properties of Shld1 derivatives, while
18
19 preserving sufficient potency of the modified compounds.¹³ In connection to this project,
20
21 alternative and novel stabilizing compounds with less demanding synthetic route were envisioned.
22
23
24 We describe in this study, a method for using computational design combined with a series of
25
26 encoded combinatorial synthesis and screening steps, as a rapid approach for developing new
27
28 peptide-based ligands that stabilizes the DD protein.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Illustration of the development process for small molecule peptide mimetics and their respectively attachment strategies. Prior to this study, DD-ligands such as SLF* and Shield-1 (Shld1) were derived by structure-based design originating from the natural product FK506. *Strategy 1* of the peptide mimetics involve standard solid-phase synthesis in the *C*- to *N*-direction, whereas in *strategy 2* the ligand is anchored via a carboxylate linker attached to a tyrosine side-chain.

The FKBP1s have been described as being “undruggable” for low molecular mass ligands with sufficient affinity¹⁴ and most synthetic ligands of the FKBP1s are derived from the natural product FK506 (**Figure 1**). Reported strategies include the use of macrocyclic peptide scaffolds^{15, 16} and structure-based design of the FK506-FKBP12 pharmacophore.¹⁷ Different synthetic approaches have been reported for the synthesis of Shld1-like structures, in which chiral reducing agents,¹⁸

chiral auxiliaries,¹⁹ addition of stoichiometric amount of chiral complexing agents,²⁰ and chromatographic separation of diastereomers,⁵ have been utilized to obtain the enantiopure compounds. Inspired by the structural insight of these previous studies,^{1, 5} we selected *in silico* for the design new scaffolds that mimic the binding of Shld1 to the DD. The scaffolds should be assembled from simple amino acid building blocks by solid phase peptide synthesis. The resulting initial structural design (*vide infra*) is presented in **Figure 1** and consists of tripeptides capped at the *N*-terminal with a triazole by CuAAC click chemistry.

We implemented and report here on a new on-bead binding assay using templated pre-folding of the labeled, disordered protein target for the evaluation of the target binding by the new scaffolds. The assay allowed rapid scaffold optimization. The solid-supported ligands could be synthesized in parallel using only 10 mg of resin per compound, thus enabling a rapid hit structure elucidation prior to off-bead resynthesis. The scaffold allowed synthesis of one-bead-one-compound (OBOC) libraries and comprised two strategies with different points of attachment to the support (**1** and **2**, **Figure 1**). The compound SLF*⁵ (**Figure 1**), which is closely related to shld1, has a carboxylic acid handle for attachment to the solid support, and was selected as a reference for studying the on-bead target interaction.

The screening of a focused combinatorial library, synthesized on optically encoded beads containing fluorescent microparticles,²¹ facilitated rapid identification of hit compounds through simple optical decoding. This library encoding approach is suitable for synthesis and screening of small to medium sized libraries (500-50000 beads) and permits, as do other bead based encoding technologies,^{22, 23, 24, 25} the identification of library members binding to the fluorescent target. The method relies on the fact that each bead in the library carries a unique distribution of microparticles as a code obtained by recording the microparticle coordinates that distinguish it from all other

beads in the library. The encoded library technology is particularly useful for bead-based libraries of compounds for which conventional mass-spectrometry methods are complicated and inefficient, as was the case in the presented study.²⁶

The structural insight gained from the on-bead assay assisted the design of 19 unnatural peptides which were screened in a competitive fluorescence polarization assay, and our findings has led to the discovery of novel and small unnatural peptides with low micromolar binding affinity towards the DD.

RESULTS AND DISCUSSION

With the aim of developing a core structure with modifiable moieties for combinatorial library development, a tripeptide with a triazole capped *N*-terminal was designed by computer modelling to adopt a similar binding pose to that of Shld1 in the DD binding site.⁵ The reference compound SLF* and an example of the computational designed ligand **1** is depicted in **Figure 2**. Modelling was performed with Molecular Operating Environment (MOE, CCG) using the crystal structure of SLF* bound to F36V-FKBP12 (PDB: 1BL4) as a starting point.⁵ Upon fixing the coordinates of the protein, the SLF* compound was replaced with peptide mimetics, soaked in a water droplet, and relaxed by energy minimization as described in detail in the supporting information. Modifications were explored on key features of the peptides during structural development. The pipecolic moiety is located in a groove on the active site of the protein and has in several studies shown to be important for binding to the protein.^{14, 17, 18}

,²⁷ A range of alternative ring sizes and substitution patterns were sampled during the modelling, but the pipecolic residue presented a superior fit to the pocket. Originating from an α -aminobutyric acid, the ethyl part adjacent to the triazole is designed to dock into the hydrophobic cavity formed after the F36V mutation,⁵ and various lengths of alkyl groups were investigated and selected in order to explore its capacity. Those alkyl groups providing favorable interactions in the pocket during modelling were included in subsequent library synthesis. For the remaining residues, interaction over the entire binding site was evaluated and structural components displaying appropriate contact with the protein were selected for combinatorial library synthesis.





Figure 2. Comparison of the binding modes of SLF* (A), and an example of peptide mimetic **1** (B) to the crystal structure of the SLF*-stabilized F36V-FKBP12 complex. In the modelling of the complex between F36V-FKBP12 and the peptide mimetic, key features of the ligand was

maintained while building the structure. The water-soaked complexes of F36V-FKBP12 with the peptide mimetic was subjected to several rounds simulated annealing (MOE, Amber 10, ETH; T: 430 K, 1 ns, 430-230 K, 1 ns, 230 K, 3 ns). In the first round, the coordinates of F36V-FKBP12 were fixed and in the subsequent rounds, the contact residues in F36V-FKBP12 were relaxed to allow an induced fit of F36V-FKBP12 and the ligand.

The design of the peptide mimetic scaffold allowed usage of the large pool of natural and unnatural amino acid building blocks in combination with the orthogonal CuAAC click chemistry to survey the pharmacophore interactions. In order to screen for optimal interaction with the protein for the immobilized substrates in the library, two different linker strategies were employed. As presented in **Scheme 1**, *Strategy 1* involve standard solid-phase peptide synthesis (SPPS) in the C- to N-direction. Whereas *Strategy 2* is anchored *via* a linker piece from the tyrosine sidechain, protruding from the other end of the compound and resembling the part of the SLF* used for immobilization of **18**. Presumably, attachment from this part of the peptide would exhibit a smaller degree of steric interaction with the DD. In order to initially verify the scaffold design and test the two linking strategies, 2-3 examples of the scaffold linked by each of the methods were synthesized on solid support (**Scheme 1**). The building block for the *strategy 2* linker was synthesized by coupling of **9** with benzylamine, followed by a substitution of ethyl 2-bromoacetate to give **10**. A Boc to Fmoc protection group exchange of **10** was performed, and after ester hydrolysis orthogonal to the Fmoc protection group,²⁸ **11** was formed in five steps with an overall yield of 79%, starting from Boc-protected L-tyrosine. A set of solid-supported compounds were then synthesized on a PEGA₁₉₀₀ resin (500-550 μm)²⁹ and subjected to a maltose binding protein (MBP) fused DD, which had been fluorescently labeled with the rhodamine-X fluorophore (MBP-DD-(ROX)). The MBP was fused to the DD to prevent aggregation of the destabilized protein and to increase the

overall polarity of the protein, hence reducing the non-specific binding. The unique swelling properties of the porous PEGA₁₉₀₀ resin in aqueous media enabled the large fusion protein to gain access to the bead interior.³⁰ As a reference compound for the initial test screen, a solid-supported version of the SLF* compound was synthesized by adapting known procedures (see supporting information).¹⁸



Scheme 1. Reagents and conditions: a) Fmoc-4-Amb-OH, TBTU, NEM, DMF, 2 h; b) Piperidine/DMF (1:4), 20 min; c) Fmoc-L-Phe-OH, TBTU, NEM, DMF, 2 h; d) Fmoc-L-Pip-OH,

1
2
3 TBTU, NEM, DMF, 2 h; e) Fmoc-D-Abu-OH, TBTU, NEM, DMF, 2 h; f) Imidazole-1-sulfonyl
4 azide hydrochloride, CuSO₄, K₂CO₃, H₂O, 16 h; g) Phenylacetylene, CuSO₄, ascorbic acid, t-
5 BuOH/H₂O (3:1), 16 h; h) Ac₂O/Pyridine/DMF (1:1:8), 30 min; i) 2-(3,4,5-
6 Trimethoxyphenyl)butanoic acid, TBTU, NEM, DMF, 2 h; j) Benzylamine, PyBOP, NEM, DMF,
7 1 h, 99%; k) ethyl-2-bromoacetate, Cs₂CO₃, DMF, 3 h, 86%; l) TFA/CH₂Cl₂ (1:1), 0°C to rt, 60
8 min; m) Fmoc-Cl, Na₂CO₃, Dioxane/H₂O (2:1), 0°C to rt, 1.5 h, 93% (2 steps); n) MgI₂, THF, mw
9 120°C, 15 min., quant; o) TBTU, NEM, DMF, 2 h.

10
11
12 The carboxylic acid part of the SLF* has been shown to have minor influence on the binding to
13 the DD protein and has previously been used as an anchor for solid-support while retaining its
14 interaction with the protein.³¹ To estimate any level of non-specific binding, three different control
15 tripeptides presenting negative, positive, and neutral charge (**19-21**, see **Figure 3**) were tested in
16 the on-bead binding assay. As presented in **Figure 3**, compounds from both attachment strategies
17 containing the 2-trimethoxyphenylbutyric acid building block showed significant binding, albeit
18 with lower affinity than the reference solid-supported SLF* (**18**) in the single concentration on-
19 bead screen. The acetylated compound **7** bound poorly to the protein, while the derivatives **6** and
20 **15** containing a phenyltriazole presented good binding, as measured by the fluorescence due to the
21 resin bound MBP-DD-(ROX). With a relatively high effective ligand concentration of
22 approximately 3.4 mM, a situation of on-bead binding saturation is presumably observed for strong
23 binding ligands. Indeed, compound **6** provided the same if not more fluorescence intensity than
24 resin bound SLF*. However, these results indicated that the triazole based mimetics of the 2-
25 trimethoxyphenylbutyric acid would be a good starting point for development of a combinatorial
26 library.




Figure 3. Single concentration binding tests of solid-supported ligands subjected to a fluorescently labeled DD protein (550 nM) for 16 h. The three tripeptides **19** (Ac-Glu-Phe-Gly-OH), **20** (Ac-Lys-Phe-Gly-OH), **21** (Ac-Ala-Phe-Gly-OH) were attached to the PEGA resin *via* the HMBA linker and used as controls for non-specific binding.

The main advantage of this approach was that the azido acid could be generated on bead from amino acid building blocks, hereby removing the need for introducing a stereocenter through enantioselective alkylation of a range of phenylacetic acid derivatives.^{19, 20} Due to anticipation of slightly better interaction, and ease of synthesis, *strategy 1* was selected for the development of the combinatorial library.

Library screening.

Using the split-and-mix approach, a library of in total 896 different compounds were synthesized on PEGA₁₉₀₀ beads (500-550 μ m) that were optically encoded by MicroParticleMatrix (MPM)-encoding (**Figure 4**).³² Encoding of the immobilized peptides was performed by recording three orthogonal images of every bead in each portion of the split process, to generate a microparticle coordinate matrix used as a barcode for the individual beads. The images were recorded using a

custom made MPM decoding instrument comprising a solid state laser, three CCD-cameras with tele centric optics and a carousel for individual bead handling.²¹ Subsequently, the isolated hit-beads were recorded and the 3-dimentional matrix represented by their microparticle coordinates were determined, allowing tracking through the synthetic history of the immobilized compound.²¹ Pipecolic acid and 4-(aminomethyl) benzoic acid were maintained as structural features during the library development. The remaining amino acid building blocks were varied and coupled using either coupling reagents TBTU or HATU. Subsequently, a solid-phase diazotransfer to the *N*-terminal amine was carried out to form the corresponding azide,³³ which in turn was reacted by the CuAAC click reaction using various alkynes (see supporting info Table S1 for the full list of library building blocks).




Figure 4. Development and screening of encoded one-bead one-compound library. (A) Using the split-and-mix approach, an 896-member library was synthesized on encoded PEGA₁₉₀₀ beads (500-550 μm) using the same conditions as presented in **scheme 1**. (B) and (C) bright field and

fluorescence image, respectively, of library beads after being subjected to a 1 μ M solution of shld1-stabilized MBP-DD-(ROX) for 48 h. The images presents a clear hit bead among inactive beads. (D) Typical encoded bead used in the present study. Three orthogonal images of the same bead are recorded and used in construction of the 3-dimentional matrix represented by the microparticle coordinates. These are used as a unique digital code to identify the structure of the compound immobilized on the bead. The code of the bead is recorded at each synthetic step and used for hit identification.


The structure variations of R, R₁, and R₂ were guided and chosen based on computational design as described above. The moieties on the R-position were primarily aliphatic and aromatic cyclic structures, whereas the R₁-position consisted of small linear and branched alkyl groups. Presumably, the enlargement of the hydrophobic cavity formed upon F36V mutation of FKBP12 could allow binding of longer alkyl groups, particularly in combination with the less bulky triazole moiety, as comparison to the reference trimethoxy phenylbutyric acid residue of SLF*. In order to induce the correct folding of the protein, the MBP-DD-(ROX) was preincubated for 30 minutes with a stoichiometric amount of Shld1 prior to the incubation of the library beads. A final concentration of 1 μ M MBP-DD-(ROX) was used and the beads were incubated for 48 hours to establish equilibrium between the individual compounds in a binding-competition assay. As presented in **Figure 4**, a clear difference in interaction with the protein among the solid-supported ligands was observed, which was visualized by recording the fluorescence intensity of the bead collections. The binding was quite selective and provided a number of bright beads on a background of many inactive beads. The identified hit-beads were manually sorted based on their apparent fluorescence intensity. The structures attached to the isolated encoded hit-beads were decoded giving a total of 16 different hit-structures out of 17 isolated beads (see supporting info


for all the hit structures).²¹ Nine of the hit compounds contained a 4-trifluoromethyl phenylalanine derivative as the second amino acid. In addition, 12 compounds had an ethyl group on the R₁-position, and primarily either a 4-methoxy phenyl or a cyclopropyl on the R₂-position.








































On-bead assay and structure evaluation.

In order to distinguish and verify the potency among the structures of the solid-supported hit compounds, a well-based on-bead binding assay was employed.³⁴ Guided by the results from the library screen, a set of immobilized substrates were resynthesized on PEGA₁₉₀₀ beads (125-200 μ m). The beads were singly sorted into a 96-well plate using a COMPAS bead sorter (Union Biometrica) and subjected to a dilution series of MBP-DD-(ROX). After 16 hours of incubation, the fluorescence intensity was measured on wide-field fluorescence microscope. The individual binding profile of the resynthesized peptide mimetics was compared to the solid-supported SLF* and graded in a scale from 1-4 (**Table 1**). In general, it was found that the most abundant substituents from the hits in the library screen, also proved to be the most potent in the bead assays. Compounds containing a 4-trifluoromethyl phenylalanine as the second amino acid, and an ethyl group on the R₁-position displayed the most promising binding profile. On the R₂-position, neither the phenyl group of **23** nor the 3-benzoic acid of **24**, were identified as hits during the screening, but were selected for comparison with the binding of **25**. These two compounds displayed a comparable binding profile to the anisole of **25**, which indicated that aromatic moieties on this position was preferred. Resynthesis of the peptide mimetics in the solid-supported screening format was realized on a low micromolar scale, requiring a minimal amount of the unnatural amino acid building-blocks, which in combination with the library screening gave an efficient and rapid structure evaluation.

Table 1. On-bead screening of compounds derived from the hits identified through optically encoded split-and-mix library technology.



\bullet = 

Cmp.	R	R ₁	R ₂	Potency ^a
18^b	-	-	-	++++
23				++++
24				++++
25				++++
26				+++
27				+++
28				++
29				+++
30				++
31				-
32				-
33				+
34				-
35				++


^a Binding profile of the peptide mimetics was compared to the solid-supported SLF* and their potency was graded in a scale from 1-4. ^b Immobilized SLF* (see **scheme 1**).



Off-bead peptide resynthesis and screening.



















The next step was to survey the non-linked scaffold for binding to the MBP-DD in solution. Based on the results obtained from the well-based on-bead assay, a set of 19 unnatural peptides were synthesized in solution or from solid-support (see supporting info). Using the same synthetic approach as presented in **Scheme 1**, the compounds containing a primary amide or a benzoic acid on the C-terminal of the peptides were synthesized from the Rink-amide linker and the 2-chloro trityl linker, respectively. The solid-supported compounds were cleaved upon acid treatment to yield the desired peptide mimetics in modest to excellent yields. The compounds with no handle for solid-support were synthesized in 3-4 steps in solution. For this, primary and secondary amines were coupled with a substituted Boc-protected L-phenylalanine derivative. Subsequently, the Boc-group was removed under acid treatment and the Boc-pipecolic acid was coupled using HBTU. In a subsequent one-pot protecting group removal and coupling procedure, an azido modified D- α -aminobutyric acid was introduced, and the final triazole was formed in a CuAAC reaction. Variation on the R₁-position (**Table 2**) included moieties of different size, polarity, and charge. The nineteen peptide mimetics were screened towards the MBP-DD using a recently described fluorescence polarization competition assay.¹³ Directed by the results from the on-bead assay, the unbound versions of **24** and **25** were synthesized and tested. The most promising of these compounds was the aminomethyl benzamide **37**, which showed a K_i of 21.5 μ M. It was found that omitting a R₁-substituent (**38**) reduced the activity fivefold, whereas enhancing the lipophilicity by changing the carboxylic acid to an ester or omitting the amide on the R₁-substituent led to loss of activity. Many of the resynthesized ligands showed poor solubility under the conditions of the assay, which is a crucial property for biological activity. This influenced the outcome of the fluorescence polarization competition assay and complicated the accurate assessment binding

constants. Reducing the lipophilicity of **36** revealed that changing to an aliphatic amide substituent (**40**), some activity could be achieved and that this was lost when switching to a lipophilic substituent (**49**). However, the effect appeared not only to be enhanced solubility, since benzoic acid derivatives (**45-47**) or shorter aliphatic amide substituent (**42-44**) in the R₁-position only led to inactive derivatives. Compound **40** exhibited a *K_i* of 33.9 μM and has a primary amide positioned in a similar distance from the pipercolic core as compound **37**, suggesting that a hydrogen bond donor in this position may be advantageous to the binding of the protein. Testing rather small moieties on the R₁-position in combination with or without changing the R- and R₂-substituents (**50-54**) interestingly revealed the two most potent compounds in the series (**51** and **53**) with *K_i*-values of 5.13 μM and 11.2 μM, respectively. Both compounds are among the most hydrophilic, but again this effect is not the only determining factor, as introducing the aliphatic amide substituent found the active derivative **40** in the R₁-position of **51** led to loss of activity (**41**). This underlines that further optimization of the scaffold has to be multi-dimensional, which exactly is the strength of combinatorial screening. However, the presented data demonstrates that the combination of computational design and combinatorial synthesis and screening is a powerful concept for developing new active scaffolds towards a given biological target.

Table 2. *K_i* values of resynthesized peptide mimetics **36-54**



Cmp	R	R ₁	R ₂	<i>K_i</i> (μM) ^a
Shld1	-	-	-	0.0075 ¹³
SLF*	-	-	-	0.0187 ¹³
36			CF ₃	>200

37		CF ₃	21.5 ± 4.0
38		H	CF ₃ 110 ± 109
39		H	CF ₃ >200
40		CF ₃	33.9 ± 7.6
41		CF ₃	>200
42		CF ₃	>200
43		CF ₃	>200
44		CF ₃	>200
45		CF ₃	>200
46		CF ₃	>200
47		NO ₂	>200
48		CF ₃	>200
49		CF ₃	>200
50		CF ₃	>200
51		CF ₃	5.13 ± 0.4
52		CF ₃	>200
53		NO ₂	11.2 ± 9.3
54		F	>200

^a A competitive fluorescence polarization assay was used to obtain the reported K_i values ± SD from three independent experiments.

Binding evaluation of peptide ligands.

The interactions between ligand **37** and F36V-FKBP12, involve two hydrophobic pockets (see **Figure 5A**). One for the ethyl group which interact with Ile-91, Gly-28, Val-36, Phe-99, Tyr 26 and Leu-97 of F36V-FKBP12, and one for the pipercolic acid, which is lined by Phe-46, Phe-99, Tyr-26, Ile-56 and Val-55, with Trp-59 positioned at the bottom of the pocket. In addition, the 4-trifluoromethyl-L-phenylalanine interact with Phe-46, and the aromatic ring of the amidomethylbenzamide residue interact with Ile-56. The bound **37** show a set of crucial hydrogen bonds. The C-terminal amide binds to the carbonyl group of Val-55 while the NH of the amidomethylbenzamide binds internally to the carbonyl of the pipercolic amide. This in turn binds amide NH of Ile-56. Tyr-82 forms a hydrogen bond to the deeply buried carbonyl of the triazole modified butyric acid, while the triazole forms a hydrogen bond to the NH of the Phe(CF₃) residue. Overall the fit of **37** is not as tight as that of SLF*.^{5, 13} The optimized interaction of the three peripheral aromatic rings seem to lift the core pipercolic amide and α -aminobutyric acid residue slightly from their binding pockets, since it cannot reach over the rim of the binding pocket. Hereby, interaction with the protein is compromised, and as visualized in **Figure 5C**, the SLF* is in significant closer contact with the protein in the deep hydrophobic binding pocket. The complex of F36V-FKBP12 with compound **51** presents an alternative binding mode in which the cyclopropane ring interacts with Ile-90, His-97, and Tyr-82 and the triazole binds perpendicularly to the NH of the Phe(CF₃). This redirects Tyr-82 to form a hydrogen bond to the carbonyl of Phe(CF₃), allowing optimal interaction of the morpholine residue with Ile-56. Surprisingly, the smaller compound **51** showed the highest binding affinity. However, since all peripheral residues of **51** are relatively small, they may allow optimal interaction of the core pipercolic amide and α -aminobutyric acid residues with the F36V-FKBP12. These observations suggest that the scaffold

could be even further improved by extending the core framework to position the peripheral aromatic residues at more favorable regions of the interaction interface, thus providing a better match of binding to the active site of the DD.




Figure 5. (A), (B), and (C) the binding modes of compound **37** and compound **51** (D) to the crystal structure of F36V-FKBP12 in the SLF*-stabilized complex presented by computational modeling. (A) Contact residues of interaction between **37** with F36V-FKBP12. (B) Hydrogen bond network of interacting residues between the complex **37** and F36V-FKBP12. C) The overlay of SLF* of the crystal structure (Green) and ligand **37** (grey) in the binding pocket of F36V-FKBP12. D) The alternate binding mode presented by compound **51**.

In summary, an 896 member one-bead one-compound library was synthesized on microparticle encoded PEGA₁₉₀₀ beads using the split-and-mix approach. The library was screened towards a fluorescently labeled version of the DD-protein and the screen resulted in 16 different structures, which were identified as hits and deconvoluted using in-house developed decoder equipment. General trends among the hits included the presence of a 4-trifluoromethyl-L-phenylalanine as the second amino acid and an ethyl-group on the R₁-position (**Figure 4**) originating from an α -D-aminobutyric acid. Guided by the library hits, solid supported peptide mimetics were resynthesized on small scale and screened in a novel on-bead binding format to confirm and distinguish the potency among the library hits. The structure information was then used for the synthesis of 19 peptide mimetics. By exploiting the fixed stereochemistry of the unnatural amino acids, the small peptide-like compounds were synthesized in few steps using either solution- or solid-phase synthesis. The peptide mimetics were tested in a competition fluorescence polarization assay and one compound with a low micromolar binding affinity towards the DD was identified. Analysis of the identified low potency ligand (**37**) indicates that binding may be partly compromised by the small displacement of the pipecolic and butyric acids caused by rim interaction of the peripheral residues. This may also add to the explanation, why a small compound like **51** bind better than **37**, while displaying less interaction. Our high-throughput

screening approach to identify a novel peptide-like structure scaffold for the DD has involved the combination of computational modeling, combinatorial library screening, and on-bead structure evaluation.

With a focus on improving the solubility of the peptide mimetics, the identified ligand **51** could serve as a starting point for development of unnatural peptides that mimic the Shld1-DD interaction. We envision that the present methodology can be applied to the synthesis and screening of larger peptide and non-peptide libraries, where the screening and initial structure refinement can be accomplished routinely and rapidly using small quantities of library building blocks in a high throughput manner using the MPM-encoding technology.

SUPPORTING INFORMATION

A full description of the materials and methods, including HPLC, NMR, and MS data can be found in the supporting info.

AUTHOR INFORMATION

Corresponding Authors. Professor Morten Meldal and Associate Professor Frederik Diness, CECB, Department of Chemistry, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark. Email: Meldal@nano.ku.dk, fdi@chem.ku.dk

Author Contributions. The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding Sources. University of Copenhagen – CECB-Lighthouse Grant 2013. Danish Council for Independent Research (DFF-4184-00019).

ACKNOWLEDGMENT

The University of Copenhagen has supported the present research through the UCPH – lighthouse program (CECB) and the free Research Council of Denmark.

ABBREVIATIONS

Ac: acetyl; Abu: α -aminobutyric acid; Amb: aminomethylbenzoic acid; Boc: *tert*-butoxycarbonyl; Comp: compound; DD: destabilizing domain; DMF: *N,N*-dimethylformamide; FKBP: FK506-binding protein; Fmoc: 9-fluorenylmethyloxycarbonyl; HATU: 1-[Bis(dimethylamino)methylene]-1*N*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxid hexafluorophosphate; HMBA: hydroxymethylbenzoic acid; MBP: maltose-binding protein; NEM: *N*-ethylmorpholine; NSB: non-specific binding; PEGA: polyethylene glycol dimethyl acrylamide; Pip: pipercolic acid; POI: protein of interest; PyBOP: benzotriazole-1-yl-oxy-tripyrrolidinophosphonium hexafluorophosphate; ROX: 5(6)-rhodamine-X; rt: room temperature; Shld1: Shield-1; SPPS: solid phase peptide synthesis; SPS: solid phase synthesis; TBTU: *N*-[(1*H*-benzotriazol-1-yl)(dimethylamino)methylene]-*N*-methylmethanaminium tetrafluoroborate; TFA: trifluoroacetic acid; THF: tetrahydrofuran.


REFERENCES

- (1) Banaszynski, L. A.; Chen, L. chun; Maynard-Smith, L. A.; Ooi, A. G. L.; Wandless, T. J. A Rapid, Reversible, and Tunable Method to Regulate Protein Function in Living Cells Using Synthetic Small Molecules. *Cell* **2006**, 126 (5), 995–1004.
- (2) Chu, B. W.; Banaszynski, L. A.; Chen, L. chun; Wandless, T. J. Recent Progress with FKBP-Derived Destabilizing Domains. *Bioorg. Med. Chem. Lett.* **2008**, 18 (22), 5941–5944.
- (3) Maynard-Smith, L. A.; Chen, L. C.; Banaszynski, L. A.; Ooi, A. G. L.; Wandless, T. J. A Directed Approach for Engineering Conditional Protein Stability Using Biologically Silent Small Molecules. *J. Biol. Chem.* **2007**, 282 (34), 24866–24872.
- (4) Grimley, J. S.; Chen, D. A.; Banaszynski, L. A.; Wandless, T. J. Synthesis and Analysis of Stabilizing Ligands for FKBP-Derived Destabilizing Domains. *Bioorg. Med. Chem. Lett.* **2008**, 18 (2), 759–761.
- (5) Clackson, T.; Yang, W.; Rozamus, L. W.; Hatada, M.; Amara, J. F.; Rollins, C. T.; Stevenson, L. F.; Magari, S. R.; Wood, S. A.; Courage, N. L.; Lu, X.; Cerasoli, F.; Gilman, M.; Holt, D. A. Redesigning an FKBP-Ligand Interface to Generate Chemical Dimerizers with Novel Specificity. *Proc. Natl. Acad. Sci.* **1998**, 95 (18), 10437–10442.
- (6) Banaszynski, L. A.; Sellmyer, M. A.; Contag, C. H.; Wandless, T. J.; Thorne, S. H. Chemical Control of Protein Stability and Function in Living Mice. *Nat. Med.* **2008**, 14 (10), 1123–1127.


- (7) Su, L.; Li, A.; Li, H.; Chu, C.; Qiu, J. L. Direct Modulation of Protein Level in Arabidopsis. *Mol. Plant* **2013**, *6* (5), 1711–1714.
- (8) Zhang, J.; Yin, K.; Sun, J.; Gao, J.; Du, Q.; Li, H.; Qiu, J. L. Direct and Tunable Modulation of Protein Levels in Rice and Wheat with a Synthetic Small Molecule. *Plant Biotechnol. J.* **2018**, *16* (2), 472–481.
- (9) Armstrong, C. M.; Goldberg, D. E. An FKBP Destabilization Domain Modulates Protein Levels in Plasmodium Falciparum. *Nat. Methods* **2007**, *4* (12), 1007–1009.
- (10) Brooks, C. F.; Johnsen, H.; van Dooren, G. G.; Muthalagi, M.; Lin, S. S.; Bohne, W.; Fischer, K.; Striepen, B. The Toxoplasma Apicoplast Phosphate Translocator Links Cytosolic and Apicoplast Metabolism and Is Essential for Parasite Survival. *Cell Host Microbe* **2010**, *7* (1), 62–73.
- (11) Madeira da Silva, L.; Owens, K. L.; Murta, S. M. F.; Beverley, S. M. Regulated Expression of the Leishmania Major Surface Virulence Factor Lipophosphoglycan Using Conditionally Destabilized Fusion Proteins. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106* (18), 7583–7588.
- (12) Rakhit, R.; Edwards, S. R.; Iwamoto, M.; Wandless, T. J. Evaluation of FKBP and DHFR Based Destabilizing Domains in Saccharomyces Cerevisiae. *Bioorg. Med. Chem. Lett.* **2011**, *21* (17), 4965–4968.
- (13) Jorgesen, F. P.; Madsen, D.; Meldal, M.; Olsen, J. V.; Petersen, M.; Granhøj, J.; Bols, M. Synthesis of Shld Derivatives, Their Binding the Destabilizing Domain and Influence on Protein Accumulation in Transgenic Plants. *J. Med. Chem.* **2019**, *62* (10), 5191–5216.
- (14) Dunyak, B. M.; Gestwicki, J. E. Peptidyl-Proline Isomerases (PPIases): Targets for Natural Products and Natural Product-Inspired Compounds. *J. Med. Chem.* **2016**, *59* (21), 9622–9644.
- (15) Wu, X.; Wang, L.; Han, Y.; Regan, N.; Li, P. K.; Villalona, M. A.; Hu, X.; Briesewitz, R.; Pei, D. Creating Diverse Target-Binding Surfaces on FKBP12: Synthesis and Evaluation of a Rapamycin Analogue Library. *ACS Comb. Sci.* **2011**, *13* (5), 486–495.
- (16) Guo, Z.; Hong, S. Y.; Wang, J.; Rehan, S.; Liu, W.; Peng, H.; Das, M.; Li, W.; Bhat, S.; Peiffer, B.; Ullman, B. R.; Tse, C. M.; Tarmakova, Z.; Schiene-Fischer, C.; Fischer, G.; Coe, I.; Paavilainen, V. O.; Sun, Z.; Liu, J. O. Rapamycin-Inspired Macrocycles with New Target Specificity. *Nat. Chem.* **2019**, *11* (3), 254–263.
- (17) Holt, D. A.; Luengo, J. I.; Yamashita, D. S.; Oh, H. J.; Konialian, A. L.; Yen, H. K.; Rozamus, L. W.; Brandt, M.; Bossard, M. J.; Levy, M. A.; Eggleston, D. S.; Liang, J.; Schultz, L. W.; Stout, T. J.; Clardy, J. Design, Synthesis, and Kinetic Evaluation of High-Affinity FKBP Ligands and the X-Ray Crystal Structures of Their Complexes with FKBP12. *J. Am. Chem. Soc.* **1993**, *115* (22), 9925–9938.
- (18) Keenan, T.; Yaeger, D. R.; Courage, N. L.; Rollins, C. T.; Pavone, M. E.; Rivera, V. M.; Yang, W.; Guo, T.; Amara, J. F.; Clackson, T.; Gilman, M.; Holt, D. A. Synthesis and Activity of Bivalent FKBP12 (1) Keenan, T.; Yaeger, D. R.; Courage, N. L.; Rollins, C. T.; Pavone, M. E.; Rivera, V. M.; Yang, W.; Guo, T.; Amara, J. F.; Clackson, T.; Gilman, M.; Holt, D. A. Synthesis and Activity of Bivalent FKBP12 Ligands. *Bioorg. Med. Chem.* **1998**, *6* (8), 1309–1335.
- (19) Gaali, S.; Kirschner, A.; Cuboni, S.; Hartmann, J.; Kozany, C.; Balsevich, G.; Namendorf,

- C.; Fernandez-Vizarra, P.; Sippel, C.; Zannas, A. S.; Draenert, R.; Binder, E. B.; Almeida, O. F. X.; Rühler, G.; Uhr, M.; Schmidt, M. V.; Touma, C.; Bracher, A.; Hausch, F. Selective Inhibitors of the FK506-Binding Protein 51 by Induced Fit. *Nat. Chem. Biol.* **2015**, *11* (1), 33–37.
- (20) Jørgensen, F. P.; Bols, M. An Inexpensive and Scalable Synthesis of Shld. *J. Org. Chem.* **2018**, *83* (11), 6050–6055.
- (21) Meldal, M.; Christensen, S. F. Microparticle Matrix Encoding of Beads. *Angew. Chem. Int. Ed.* **2010**, *49* (20), 3473–3476.
- (22) Farrer, R. A.; Copeland, G. T.; Previte, M. J. R.; Okamoto, M. M.; Miller, S. J.; Fourkas, J. T. Production, Analysis, and Application of Spatially Resolved Shells in Solid-Phase Polymer Spheres. *J. Am. Chem. Soc.* **2002**, *124* (9), 1994–2003.
- (23) Braeckmans, K.; De Smedt, S. C.; Roelant, C.; Leblans, M.; Pauwels, R.; Demeester, J. Encoding Microcarriers by Spatial Selective Photobleaching. *Nat. Mater.* **2003**, *2* (3), 169–174.
- (24) Battersby, B. J.; Bryant, D.; Meutermans, W.; Matthews, D.; Smythe, M. L.; Trau, M. Toward Larger Chemical Libraries: Encoding with Fluorescent Colloids in Combinatorial Chemistry. *J. Am. Chem. Soc.* **2000**, *122* (9), 2138–2139.
- (25) Fenniri, H.; Ding, L.; Ribbe, A. E.; Zyrianov, Y. Barcoded Resins: A New Concept for Polymer-Supported Combinatorial Library Self-Deconvolution. *J. Am. Chem. Soc.* **2001**, *123* (33), 8151–8152.
- (26) Kassel, D. B. Combinatorial Chemistry and Mass Spectrometry in the 21st Century Drug Discovery Laboratory. *Chem. Rev.* **2001**, *101* (2), 255–267.
- (27) Kolos, J. M.; Voll, A. M.; Bauder, M.; Hausch, F. FKBP Ligands—Where We Are and Where to Go? *Front. Pharmacol.* **2018**, *9*, 1–25.
- (28) Berthet, M.; Davanier, F.; Dujardin, G.; Martinez, J.; Parrot, I. MgI₂-Mediated Chemoselective Cleavage of Protecting Groups: An Alternative to Conventional Deprotection Methodologies. *Chem. Eur. J.* **2015**, *21* (31), 11014–11016.
- (29) Auzanneau, F. -I; Meldal, M.; Bock, K. Synthesis, Characterization and Biocompatibility of PEGA Resins. *J. Pept. Sci.* **1995**, *1* (1), 31–44.
- (30) Meldal, M. The One-Bead Two-Compound Assay for Solid Phase Screening of Combinatorial Libraries. *Biopolym. - Pept. Sci. Sect.* **2002**, *66* (2), 93–100.
- (31) Noblin, D. J.; Page, C. M.; Tae, H. S.; Gareiss, P. C.; Schneekloth, J. S.; Crews, C. M. A HaloTag-Based Small Molecule Microarray Screening Methodology with Increased Sensitivity and Multiplex Capabilities. *ACS Chem. Biol.* **2012**, *7* (12), 2055–2063.
- (32) Hu, H.; Nikitin, S. V.; Berthelsen, A. B.; Diness, F.; Schoffelen, S.; Meldal, M. Sustainable Flow Synthesis of Encoded Beads for Combinatorial Chemistry and Chemical Biology. *ACS Comb. Sci.* **2018**, *20* (8), 492–498.
- (33) Castro, V.; Blanco-Canosa, J. B.; Rodriguez, H.; Albericio, F. Imidazole-1-Sulfonyl Azide-Based Diazo-Transfer Reaction for the Preparation of Azido Solid Supports for Solid-Phase Synthesis. *ACS Comb. Sci.* **2013**, *15* (7), 331–334.

- (34) Li, M.; Hoeck, C.; Schoffelen, S.; Gotfredsen, C. H.; Meldal, M. Specific Electrostatic Molecular Recognition in Water. *Chem. Eur. J.* **2016**, 22 (21), 7206–7214.



ACS Paragon Plus Environment



Scheme 1. Reagents and conditions: a) Fmoc-4-Amb-OH, TBTU, NEM, DMF, 2 h; b) Piperidine/DMF (1:4), 20 min; c) Fmoc-L-Phe-OH, TBTU, NEM, DMF, 2 h; d) Fmoc-L-Pip-OH, TBTU, NEM, DMF, 2 h; e) Fmoc-D-Abu-OH, TBTU, NEM, DMF, 2 h; f) Imidazole-1-sulfonyl azide hydrochloride, CuSO₄, K₂CO₃, H₂O, 16 h; g) Phenylacetylene, CuSO₄, ascorbic acid, t-BuOH/H₂O (3:1), 16 h; h) Ac₂O/Pyridine/DMF (1:1:8), 30 min; i) 2-(3,4,5-Trimethoxyphenyl)butanoic acid, TBTU, NEM, DMF, 2 h; j) Benzylamine, PyBOP, NEM, DMF, 1 h, 99%; k) ethyl-2-bromoacetate, Cs₂CO₃, DMF, 3 h, 86%; l) TFA/CH₂Cl₂ (1:1), 0°C to rt, 60 min; m) Fmoc-Cl, Na₂CO₃, Dioxane/H₂O (2:1), 0°C to rt, 1.5 h, 93% (2 steps); n) MgI₂, THF, mw 120°C, 15 min., quant; o) TBTU, NEM, DMF, 2 h.

600x550mm (96 x 96 DPI)




Figure 2. Comparison of the binding modes of SLF* (A), and an example of peptide mimetic 1 (B) to the crystal structure of the SLF*-stabilized F36V-FKBP12 complex. In the modelling of the complex between F36V-FKBP12 and the peptide mimetic, key features of the ligand was maintained while building the structure. The water-soaked complexes of F36V-FKBP12 with the peptide mimetic was subjected to several rounds simulated annealing (MOE, Amber 10, ETH; T: 430 K, 1 ns, 430-230 K, 1 ns, 230 K, 3 ns). In the first round, the coordinates of F36V-FKBP12 were fixed and in the subsequent rounds, the contact residues in F36V-FKBP12 were relaxed to allow an induced fit of F36V-FKBP12 and the ligand.

550x360mm (96 x 96 DPI)




Figure 3. Single concentration binding tests of solid-supported ligands subjected to a fluorescently labeled DD protein (550 nM) for 16 h. The three tripeptides 19 (Ac-Glu-Phe-Gly-OH), 20 (Ac-Lys-Phe-Gly-OH), 21 (Ac-Ala-Phe-Gly-OH) were attached to the PEGA resin via the HMBA linker and used as controls for non-specific binding.

338x190mm (96 x 96 DPI)




Figure 4. Development and screening of encoded one-bead one-compound library. (A) Using the split-and-mix approach, an 896-member library was synthesized on encoded PEGA1900 beads (500-550 μm) using the same conditions as presented in scheme 1. (B) and (C) bright field and fluorescence image, respectively, of library beads after being subjected to a 1 μM solution of shld1-stabilized MBP-DD-(ROX) for 48 h. The images presents a clear hit bead among inactive beads. (D) Typical encoded bead used in the present study. Three orthogonal images of the same bead are recorded and used in construction of the 3-dimensional matrix represented by the microparticle coordinates. These are used as a unique digital code to identify the structure of the compound immobilized on the bead. The code of the bead is recorded at each synthetic step and used for hit identification.

500x379mm (96 x 96 DPI)





Figure 5. (A), (B), and (C) the binding modes of compound 37 and compound 51 (D) to the crystal structure of F36V-FKBP12 in the SLF*-stabilized complex presented by computational modeling. (A) Contact residues of interaction between 37 with F36V-FKBP12. (B) Hydrogen bond network of interacting residues between the complex 37 and F36V-FKBP12. C) The overlay of SLF* of the crystal structure (Green) and ligand 37 (grey) in the binding pocket of F36V-FKBP12. D) The alternate binding mode presented by compound 51

300x800mm (96 x 96 DPI)



Cmp.	R	R ₁	R ₂	Potency ^a
18 ^b	-	-	-	++++
23				++++
24				++++
25				++++
26				+++
27				+++
28				++
29				+++
30				++
31				-
32				-
33				+
34				-
35				++


^a Binding profile of the peptide mimetics was compared to the solid-supported SLF* and their potency was graded in a scale from 1-4. ^b Immobilized SLF* (see scheme 1).

Table 1. Synthesis and screening of compounds derived from the hits identified through split-and-mix library decoding.

300x819mm (96 x 96 DPI)

^a A competitive fluorescence polarization assay was used to obtain the reported K_i values \pm SD from three independent experiments.

300x1100mm (96 x 96 DPI)



539x237mm (43 x 41 DPI)