

Pd(0)-Catalyzed Cross-Coupling Reactions of Boron Derivatives with a Lactam-Derived *N*-Boc Enol Triflate

Ernesto G. Occhiato,* Andrea Trabocchi, and Antonio Guarna

Dipartimento di Chimica Organica “U. Schiff” and Centro di Studio sulla Chimica e la Struttura dei Composti Eterociclici e loro Applicazioni, C.N.R.; Università di Firenze, Via G. Capponi 9, I-50121 Firenze, Italy

labgua@chimorg.unifi.it

Received March 6, 2000

ABSTRACT




The cross-coupling reaction of 2-(1-alkenyl)-1,3,2-benzodioxaboroles, obtained from alkynes and catecholborane, and other boron derivatives with a lactam-derived *N*-Boc enol triflate occurred under very mild conditions in a THF–water medium employing $(\text{Ph}_3\text{P})_2\text{PdCl}_2$ as a catalyst, providing the corresponding 6-substituted *N*-Boc 3,4-dihydro-2*H*-pyridines in high yields.

In recent years, lactam-derived enol triflates have gained importance as intermediates in the synthesis of heterocyclic compounds. Hiemstra and Speckamp¹ have shown that these compounds can undergo palladium-mediated displacement of the triflate group with nucleophiles such as organotin and organozinc derivatives, as well as methoxycarbonylation reactions. Comins² and other authors^{3,4} have reported coupling reactions with organocuprates and organotin compounds, while Sonogashira-type cross-coupling reactions with monosubstituted acetylenes have been described by Isobe⁵ and, more recently, by Cha.⁶ Surprisingly, no examples of Suzuki-type reactions of lactam-derived enol triflates have been yet described, although the use of organoboron compounds as nucleophiles in Pd-catalyzed cross-coupling reactions with vinyl and aryl triflates or halides is a powerful methodology in organic synthesis.⁷

The possibility of forming new C–C bonds by Pd(0)-catalyzed coupling reactions of 2-(1-alkenyl)-1,3,2-benzo-

dioxaboroles and other boron derivatives with six-membered enol triflates such as **2** (Scheme 1), bearing a *N*-Boc protecting group, would certainly extend the utility and scope of this class of compounds⁸ in the preparation, and the further synthetic elaboration, of heterocyclic compounds.^{1,2,9} Moreover, the *N*-Boc protection provides a possible alternative

Scheme 1



a: LiHMDS, HMPA, PhNTf_2 , THF, -78 to 25°C , 15 h.
b: $(\text{Ph}_3\text{P})_2\text{PdCl}_2$, THF– H_2O , 2 M Na_2CO_3 , 40°C , 6–7 h.

- (1) (a) Luker, T.; Hiemstra, H.; Speckamp, W. N. *J. Org. Chem.* **1997**, 62, 8131–8140. (b) Luker, T.; Hiemstra, H.; Speckamp, W. N. *Tetrahedron Lett.* **1996**, 37, 8257–8260. (c) Bernabé, P.; Rutjes, F. P. J. T.; Hiemstra, H.; Speckamp, W. N. *Tetrahedron Lett.* **1996**, 37, 3561–3564.
- (2) Foti, C. J.; Comins, D. L. *J. Org. Chem.* **1995**, 60, 2656–2657.
- (3) Tsushima, K.; Hirade, T.; Hasegawa, H.; Murai, A. *Chem. Lett.* **1995**, 9, 801–802.
- (4) Beccalli, E. M.; Marchesini, A. *Tetrahedron* **1995**, 51, 2353–2362.
- (5) Okita, T.; Isobe, M. *Tetrahedron* **1995**, 51, 3737–3744.
- (6) Ha, J. D.; Lee, D.; Cha, J. K. *J. Org. Chem.* **1997**, 62, 4550–4551.
- (7) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, 95, 2457–2483.


to other electron-withdrawing protecting groups such as *N*-tosyl,¹ *N*-Bz,^{2,3} or *N*-Cbz,^{2,9} since it is removable under different conditions.

N-Boc enol triflate **2** was prepared in 75% yield by treatment of *N*-Boc lactam **1** with LiHMDS and *N*-phenyltriflimide, in the presence of HMPA.³ The reaction of **2** with 2-[(*E*)-1-hexenyl]-1,3,2-benzodioxaborole **3a**¹⁰ (Scheme 1) was first attempted under the classical conditions reported for the reaction of enol triflates with alkenylboronate, i.e., using (Ph₃P)₄Pd catalyst in dioxane at 85 °C and K₃PO₄ as a base.¹¹ However, the reaction did not occur at all, recovering after 5 h the unreacted starting material **2**. Instead, when we used (Ph₃P)₂PdCl₂ as a catalyst (5% mol) in a THF–water mixture and in the presence of Na₂CO₃ as a base (Scheme 1), the reaction proceeded very fast at 80 °C, affording **4a**¹² after 2 h in 81% yield and with maintenance of the double bond geometry; the reaction rate was slightly slower at 40 °C (82% yield after 6 h), while at room temperature it reached completion (82% yield) after 16 h.

We tested also the more hindered (*o*-tolyl)₂PdCl₂ for the reaction of **2** with **3a**, but the reaction was slower, furnishing **4a** in 16% yield after 48 h at room temperature.

Two other alkenylboronates, **3b,c**, prepared from the corresponding alkynes and catecholborane as reported,¹⁰ were used as nucleophiles (Scheme 1). The reactions, always carried out with (Ph₃P)₂PdCl₂, were successful in providing the corresponding products **4b,c**. After 6 h at 40 °C, the conversion was complete in both cases. Compound **4b**, obtained in 76% yield, maintained the *E* geometry of the double bond. This compound in particular was quite acid sensitive, and even when left in CDCl₃ solution it underwent ring opening to give **5** in a 6:1 ratio with **4b** after 48 h (Scheme 1).


For the introduction of an aryl group, we found convenient the use of aryldiethylboranes, 2-aryl-1,3,2-dioxaborinanes, and arylboronic acids, such as commercially available compounds **6a–c** (Scheme 2). Diethyl(3-pyridyl)borane **6a**



reacted fast with **2** under the usual conditions at 80 °C, providing **7** in 91% yield after 2 h. The reaction of boron derivative **6b** with **2** was complete after 3 h at 40 °C, affording 6-phenyl-substituted *N*-Boc 3,4-dihydro-2*H*-pyridine **8** in 85% yield. The same compound was obtained in 87% yield after 2 h at 40 °C using phenylboronic acid **6c**. In this case, we increased the volume of the aqueous 2 M Na₂CO₃ solution to avoid a possible ring opening of the

product, which we observed when performing the reaction under the usual conditions.

N-Boc-protected compounds **4a–c** are potential dienes for Diels–Alder reactions, as reported for closely related compounds in which the N atom bore a tosyl^{1a} or a Cbz group.⁹ We carried out the cycloaddition between **4a** and **4b** and *N*-phenylmaleimide **9** (2.5 equiv) (Scheme 3) in



refluxing benzene obtaining, after 2 h, *endo*-cycloadducts **10** (74%) and **11** (71%) in which migration of the double bond to the more substituted position occurred.⁹ The *endo* stereochemistry, easily assignable by the ¹H NMR coupling constant value of about 4.0 Hz between protons H_a and H_b (in cis relationship), is in accordance with the results obtained by Cha in the cycloaddition of *N*-phenylmaleimide with a similar *N*-Cbz-protected substrate.⁹

In conclusion, we have shown that the coupling reaction of alkenylboronates and other boron derivatives with six-membered lactam-derived *N*-Boc enol triflates is feasible under very mild conditions in THF–water employing (Ph₃P)₂PdCl₂ as a catalyst, which provides in high yields the corresponding 6-(1-alkenyl)- and 6-aryl-substituted *N*-Boc 3,4-dihydro-2*H*-pyridines. The former are useful dienes for Diels–Alder reactions in the construction of more complex heterocyclic structures.

Acknowledgment. We thank MURST, the University of Florence (COFIN 1998–2000), and Consiglio Nazionale delle Ricerche (CNR) for financial support. Dr. Dina Scarpi and Dr. Fabrizio Machetti are acknowledged for carrying out some preliminary experiments. Mr. Sandro Papaleo and Mrs. Brunella Innocenti are acknowledged for their technical support.

Supporting Information Available: Experimental procedures and characterization of compounds **4a–c**, **7–8**, **10**, and **11**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL005759U

(8) The extension of the methodology to five-membered enol triflates from *N*-acetylpyrrolidin-2-ones is not immediate due to the high instability and difficulty in the isolation of these compounds (see refs 1a,b and 2).

(9) Ha, J. D.; Kang, C. H.; Belmore, K. A.; Cha, J. K. *J. Org. Chem.* **1998**, 63, 3810–3811.

(10) (a) Brown, H. C.; Gupta, S. K. *J. Am. Chem. Soc.* **1975**, 97, 5249–5255. (b) Miyaura, N.; Suzuki, A. *Org. Synth.* **1990**, 68, 130–136.

(11) Oh-e, T.; Miyaura, N.; Suzuki, A. *J. Org. Chem.* **1993**, 58, 2201–2208.

(12) All new compounds were fully characterized and gave satisfactory elemental analysis.