


225. The Reaction between Thiosemicarbazones and Maleic Anhydride.

By JOHN M'LEAN and F. J. WILSON.


In continuation of previous work on thiosemicarbazones and thiazoles, the authors have shown that thiosemicarbazones and δ -substituted thiosemicarbazones combine with maleic anhydride to yield thiazole derivatives.

It was shown by Andreasch (*Monatsh.*, 1895, **16**, 789; 1897, **18**, 56) that thiourea reacted on heating with maleic or fumaric acid in aqueous solution, giving 2-imino-4-ketotetrahydrothiazole-5-acetic acid :




Diphenylthiourea reacted in a similar manner. Citraconic acid or its anhydride condensed similarly with thiourea or substituted thioureas.

In continuation of previous work (J., 1922, **121**, 870; 1923, **123**, 799; 1926, 2531; 1937, 556) we have found that the thiosemicarbazones of acetone, acetophenone, 3-methylcyclohexanone and benzaldehyde condense with maleic anhydride on heating in benzene solution, yielding derivatives of 2 : 4-diketotetrahydrothiazole-5-acetic acid :



We expected that hydrolysis of these compounds with dilute hydrochloric acid would give the hydrazone (I, $\text{CRR}' = \text{H}_2$), but the results were indefinite (cf. Stephen and Wilson, J., 1928, 1415). On heating with concentrated hydrochloric acid, however, 2 : 4-diketotetrahydrothiazole-2- α -phenylethylidenehydrazone-5-acetic acid gave acetophenone, hydrazine dihydrochloride, and 2 : 4-diketotetrahydrothiazole-5-acetic acid :



Hydrolysis with concentrated hydrochloric acid of compounds of type (I, $R'' = \text{Ph}$) produced resins and hydrazine dihydrochloride. However, 2:4-diketo-3-methyltetrahydrothiazole-2- α -phenylethylidenehydrazone-5-acetic acid (I; $R = R'' = \text{Me}$, $R' = \text{Ph}$) with concentrated hydrochloric acid gave acetophenone, hydrazine dihydrochloride, and 2:4-diketo-3-methyltetrahydrothiazole-5-acetic acid (II, $R'' = \text{Me}$) together with some resin.

EXPERIMENTAL.

2: 4-Diketotetrahydrothiazole-2-isopropylidenehydrazone-5-acetic acid (I; R = R' = Me, R'' = H) (from acetone-thiosemicarbazone in benzene-acetone, 15 minutes), m. p. 223° after recrystallisation from acetone-light petroleum (Found: C, 42.5; H, 4.9; N, 18.1. C₉H₁₁O₃N₃S requires C, 41.9; H, 4.8; N, 18.3%).

2: 4-Diketotetrahydrothiazole-2- α -phenylethylidenehydrazono-5-acetic acid (I; R = Me, R' = Ph, R'' = H) (from acetophenonethiosemicarbazone in benzene, 1 hour), m. p. 244°, from alcohol (Found: C, 54.2; H, 4.5; N, 14.4. $C_{13}H_{15}O_3N_3S$ requires C, 53.7; H, 4.5; N, 14.4%).

2 : 4-Diketotetrahydrothiazole-2-3'-methylcyclohexylidenehydrazone-5-acetic acid (I; RR' = C₆H₅Me, R'' = H) (from 3-methylcyclohexanonethiosemicarbazone in benzene, 20 minutes), m. p. 209°, from alcohol-benzene (Found : N, 14.9. C₁₅H₁₇ON₃S requires N, 14.8%).

2 : 4-Diketotetrahydrothiazole-2-benzylidenehydrazone-5-acetic acid (I; R = R' = H, R' = Ph) (from benzaldehydethiosemicarbazone in benzene-acetone, 20 minutes), m. p. 255°, from acetone (Found : N, 15.1, 15.3. $C_{12}H_{11}O_3N_3S$ requires N, 15.2%).

2 : 4-Diketo-3-phenyltetrahydrothiazole-2-isopropylidenehydrazine-5-acetic acid (I; R = R' = Me, R'' = Ph) (from acetone- δ -phenylthiosemicarbazone in benzene-acetone, 4 hours), m. p. 175—180°, from benzene-acetone (Found : N, 13.7. $C_{14}H_{15}O_3N_3S$ requires N, 13.8%).

2 : 4-Diketo-3-phenyltetrahydrothiazole-2- α -phenylethylidenehydrazone-5-acetic acid (I; R = Me, R' = R'' = Ph) (from acetophenone- δ -phenylthiosemicarbazone in toluene, 1 $\frac{1}{4}$ hours), m. p. 185°, from toluene (Found : N, 11.5. C₁₉H₁₇O₃N₃S requires N, 11.4%).

2 : 4-Diketo-3-phenyltetrahydrothiazole-2-cyclohexyldenehydrazone-5-acetic Acid (I; $\text{RR}' = \text{C}_6\text{H}_{10}$, $\text{R}'' = \text{Ph}$).—cycloHexanone-8-phenylthiosemicarbazone was prepared by refluxing the ketone and 8-phenylthiosemicarbazide in absolute alcohol on the water-bath for 2 hours; on cooling, the solution deposited the compound in needles, m. p. 114° after recrystallisation from light petroleum (Found : N, 17.1. $\text{C}_{15}\text{H}_{17}\text{N}_3\text{S}$ requires N, 17.0%). The thiazole, prepared from this phenylthiosemicarbazone in benzene (1 hour), was recrystallised from alcohol-benzene and melted at 218° (decomp.) (Found : N, 12.1. $\text{C}_{17}\text{H}_{19}\text{O}_2\text{N}_2\text{S}$ requires N, 12.2%).

2 : 4-Diketo-3-phenyltetrahydrothiazole-2-3'-methylcyclohexylidenehydrazone-5-acetic Acid (I; $RR' = C_6H_5Me$, $R'' = Ph$).—3-Methylcyclohexanone- δ -phenylthiosemicarbazone was prepared by refluxing equimolecular amounts of the ketone and δ -phenylthiosemicarbazide in absolute alcohol on the water-bath for 1 hour. On cooling and addition of a little water the compound separated, m. p. 139° after recrystallisation from benzene–light petroleum (Found : N, 15.9, 16.0. $C_{14}H_{19}N_3S$ requires N, 16.0%). The thiazole was prepared from this δ -phenylthiosemicarbazone in benzene (1 hour), and purified by addition of light petroleum to an alcoholic solution; it melted at 192° (Found : N, 11.7. $C_{18}H_{21}O_3N_3S$ requires N, 11.7%).

2 : 4-Diketo-3-phenyltetrahydrothiazole-2-benzylidenehydrazone-5-acetic acid (I; $R = H$, $R' = R'' = Ph$) (from benzaldehyde- δ -phenylthiosemicarbazone in benzene, $1\frac{1}{2}$ hours) melted at 215° after recrystallisation from alcohol (Found : N, 11.7. $C_{18}H_{17}O_3N_3S$ requires N, 11.9%).

2 : 4-Diketo-3-methyltetrahydrothiazole-2- α -phenylethylidenehydrazone-5-acetic Acid (I; $R = R'' = Me$, $R' = Ph$).—Acetophenone- δ -methylthiosemicarbazone was obtained by refluxing acetophenone with δ -methylthiosemicarbazide in absolute alcohol for $\frac{3}{4}$ hour. The compound separated on cooling and after recrystallisation from alcohol melted at 135° (Found : N, 20.2. $C_{10}H_{13}N_3S$ requires N, 20.3%). The thiazole was prepared from this substance in benzene (5 hours); some resinous matter separated during the reaction. It was recrystallised from aqueous alcohol and melted at 154° (Found : N, 13.7. $C_{14}H_{17}O_3N_3S$ requires N, 13.7%).

2 : 4-Diketo-3-methyltetrahydrothiazole-2-benzylidenehydrazone-5-acetic acid (I; $R = H$, $R' = Ph$, $R'' = Me$) (from benzaldehyde- δ -methylthiosemicarbazone in benzene, 4 hours) was recrystallised from acetone containing a little water and melted at 188° (Found : N, 14.7. $C_{13}H_{13}O_3N_3S$ requires N, 14.5%).

Hydrolysis Experiments.—2 : 4-Diketotetrahydrothiazole-2- α -phenylethylidenehydrazone-5-acetic acid was heated with concentrated hydrochloric acid for 30 minutes and then steam-distilled till no more acetophenone passed over. The cold solution was extracted ten times with ether; the extract on evaporation gave 2 : 4-diketotetrahydrothiazole-5-acetic acid, m. p. 170° after recrystallisation from acetone–petrol (Tambach, *Annalen*, 1894, **280**, 241) (Found : C, 34.5; H, 2.9; S, 18.6. Calc. : C, 34.3; H, 2.9; S, 18.4%). The aqueous portion on evaporation deposited hydrazine dihydrochloride, ammonium chloride, and a small quantity of resin.

2 : 4-Diketo-3-methyltetrahydrothiazole-2- α -phenylethylidenehydrazone-5-acetic acid was hydrolysed as above. The ethereal extract on evaporation left a syrup, which on scratching and standing solidified and after recrystallisation from benzene was identified as 2 : 4-diketo-3-methyltetrahydrothiazole-5-acetic acid in poor yield; it melted at 99° (Kallenberg, *Ber.*, 1923, **56**, 316) (Found : N, 7.5. Calc. : N, 7.4%). The aqueous portion on evaporation gave hydrazine dihydrochloride and a small amount of resinous matter.

We thank Imperial Chemical Industries, Ltd., for a research grant.

“FREELAND” CHEMICAL LABORATORIES,
THE ROYAL TECHNICAL COLLEGE, GLASGOW.

[Received, May 10th, 1939.]