LETTERS

TiCl₄ Promoted Formal [3 + 3] Cycloaddition of Cyclopropane 1,1-Diesters with Azides: Synthesis of Highly Functionalized Triazinines and Azetidines

Huan-Huan Zhang, Yong-Chun Luo,* Hua-Peng Wang, Wei Chen, and Peng-Fei Xu*

State Key Laboratory of Applied Organic Chemistry, College of Chemistry and Chemical Engineering, Lanzhou University, Lanzhou 730000, PR China

Supporting Information

ABSTRACT: A TiCl₄ promoted formal [3 + 3] cycloaddition of cyclopropane 1,1-diesters with azides has been developed for the synthesis of highly functionalized triazinines. Both stoichiometric and substoichiometric versions of this reaction were accomplished dependent on the choice of solvent. It is noteworthy that the corresponding products could be easily converted to biologically important azetidines by simple thermolysis.

Many heterocycles with NNN linkages in their cyclic arrangement embrace interesting biological activities.¹ Among them, 1,2,3-triazine represents a widely used lead structure with a multitude of interesting applications in numerous pharmacological fields. Various synthetic analogues of 1,2,3-triazine have been synthesized, and some of them have shown excellent pharmacological activity.² Combined with our interest in the cycloaddition of donor–acceptor cyclopropanes, we attempted to develop an efficient method for the straightforward synthesis of functionalized 1,4,5,6-tetrahydro-1,2,3-triazines, also named triazinines,³ from azides and cyclopropanes under mild conditions.

To date, several reactions between azides and cyclopropanes have been reported.⁴ Among which, Aubé and co-workers developed the reaction of alkyl azides with triethyl(1-methoxy-2,2-dimethyl-cyclopropoxy)silane to prepare a series of α amino- α' -diazomethyl ketone.^{4d,e} They speculated that a triazine intermediate was formed by [3 + 3] cycloaddition in this reaction but decomposed immediately (Figure 1). In addition, in several other reactions, some triazines were prepared from azides,^{3,5} but the synthesis of triazinines via the cycloaddition of cyclopropane with azides has not been reported.



Figure 1. Cycloaddition of azide with cyclopropanes.



Cyclopropane 1,1-diesters, a kind of versatile three-carbon zwitterionic synthons, have been widely used to assemble various cyclic skeletons by means of [3 + n] cycloaddition reactions promoted by Lewis acid.⁶ In these reactions, the scope and applicability of its [3 + 3] process are limited to a few 1,3-dipoles.^{7,8} So far, alkyl azides have not been employed in this [3 + 3] process. Herein, we report an efficient synthesis of triazinines derivatives by the formal [3 + 3] cycloaddition of azides with cyclopropane 1,1-diesters.

In our initial trials, benzyl azide 1a and dimethyl 2phenylcyclopropane-1,1-dicarboxylate 2a were employed to optimize the reaction conditions, and the results are summarized in Table 1. In order to activate 2a, a variety of common Lewis acids were attempted. At first, a stoichiometric

Table 1. Optimization of the Reaction Conditions

Ph へ		Lewis acid	Ph Y	COOMe COOMe
	^o Ph ^o COOMe	solvent	Ph~N ^N N	1
1a	2a		3aa	
entry ^a	Lewis acid (mol %)	solvent	time (h)	yield (%) ^c
1	SnCl ₄ (100)	CH_2Cl_2	16	complex
2	FeCl ₃ (100)	CH_2Cl_2	16	complex
3	$AlCl_3$ (100)	CH_2Cl_2	16	25
4^b	$TiCl_4$ (100)	CH_2Cl_2	2	89
5 ^b	$TiCl_4$ (50)	CH_2Cl_2	2	47
6^b	$TiCl_4$ (20)	HFIP	12	87
7^{b}	$TiCl_{4}$ (10)	HFIP	48	61

^{*a*}Reaction conditions: unless otherwise noted, the reactions were carried out under a N_2 atmosphere with 1a (0.5 mmol) and 2a (0.5 mmol) in 5 mL of solvent at rt and stirred for the indicated time. ^{*b*}The reaction was carried out at 0 °C to rt. ^{*c*}Isolated yield.

Received: August 13, 2014 Published: September 5, 2014 amount of Lewis acids was added and the reactions were carried out in dichloromethane. Using $SnCl_4$ and $FeCl_3$, the reaction mixture was very complex (entries 1, 2). Using $CeCl_3$ and $Ti({}^{i}PrO)_{4}$, desired product was not detected. However, when $AlCl_3$ was added, triazinine **3aa** was obtained in 25% yield (entry 3). This result encouraged us to continue the study. Fortunately, we found that 1.0 equiv of $TiCl_4$ enabled full conversion of the starting materials to **3aa** in 89% yield after 2 h (entry 4). Further screening of the reaction solvents demonstrated that this reaction is sensitive to the solvent. Compound **3aa** could not be formed when the reaction was conducted in 1,2-dichloroethane, chloroform or tetrachloromethane.

Furthermore, on the basis of above results, we continued our efforts to develop substoichiometric version of this reaction. At first, the amount of TiCl₄ was reduced to 50 mol %, but the yield of 3aa decreased dramatically (47%, entry 5). This result suggested that the generated product sequestered TiCl₄ in an unproductive manner and interrupted the circulation of TiCl₄. Thus, several other trifluoromethanesulfonate salts, such as $Sc(OTf)_3$, $Yb(OTf)_3$, $Zn(OTf)_2$, $In(OTf)_3$ were examined with 20 mol % catalyst loading. However, no reaction was observed even by prolonging the reaction time or raising the temperature. Similar result was obtained in the presence of Ni(ClO_4)₂. Right at that moment, TiCl₄ catalyzed intramolecular Schmidt reaction was developed by Aubé and co-workers.⁹ Using strong hydrogen-bond-donating solvent hexafluoro-2-propanol (HFIP) effectively overcame the inhabitation of product in catalysis. In this catalytic system, in situ-generated HCl from the reaction of HFIP with TiCl₄ is the catalytically active species. We were greatly inspired by this discovery and carried out our reaction in HFIP in the presence of 20 mol % TiCl₄. After 12 h, 3aa was obtained in 87% yield (entry 6). Reducing the catalyst loading to 10 mol %, the reaction became very slow. After 48 h, 3aa was obtained in 61% yield (entry 7). According to the above results, both stoichiometric and substoichiometric versions of this reaction have been developed. With the optimal conditions (entries 4, 6) in hand, we set forth to survey the substrate scope of this interesting reaction and demonstrate the utility of this process.

The scope of the reaction was investigated under stoichiometric and substoichiometric conditions, respectively. Various cyclopropane 1,1-diesters and azides were employed and the results are summarized in Table 2. The stoichiometric protocol was found to be tolerant to both electron-withdrawing and electron-donating substituents in cyclopropane 1,1-diesters, and all of them gave the corresponding triazinine derivatives 3aa-3bl as a single product in moderate to high yields. No significant difference in reactivity was observed between the benzyl azides 1a and decyl azides 1b in the reaction. The position of the substituent on the aryl group slightly influences the reactivity and sterically demanding ortho-substituted cyclopropanes always gave relatively lower yields of the desired product. For example, the reaction of *o*-ClC₆H₄- and *p*-ClC₆H₄substituted cyclopropanes 2d and 2g with 1a gave the corresponding 3ad and 3ag in 77% and 88% yield respectively, probably due to the larger steric effect relative to the latter (entries 4, 7). A vinyl group could also be used as substituent, offering 3al and 3bl in good yield (entries 12, 24). Unfortunately, when the aryl group on cyclopropane was replaced by methyl or hexyl group, the reactions did not work. To examine the feasibility of this method on a larger preparative scale, the model reaction leading to 3aa was also

Table 2. Investigation of the Reaction Scope

R ^{1_} 1a R ¹	N ₃ + = Bn		COOMe <u>TiCl₄</u>	$\begin{array}{c} \bullet R^2 \\ \bullet \\ R^1 \cdot N \\ \end{array}$		Me DMe
1b R ¹	= <i>n</i> -C ₁₀ H	21 2a-l		3	aa-bl	
					yield	$(\%)^a$
entry	1	2	\mathbb{R}^2	3	A^b	B^{c}
1	1a	2a	C ₆ H ₅	3aa	89	87
2	1a	2b	o-MeOC ₆ H ₄	3ab	75	57
3	1a	2c	o-MeC ₆ H ₄	3ac	79	80
4	1a	2d	o-ClC ₆ H ₄	3ad	77	49
5	1a	2e	p-MeOC ₆ H ₄	3ae	86	41
6	1a	2f	p-MeC ₆ H ₄	3af	89	86
7	1a	2g	p-ClC ₆ H ₄	3ag	88	85
8	1a	2h	p-BrC ₆ H ₄	3ah	87	74
9	1a	2i	p-FC ₆ H ₄	3ai	89	80
10	1a	2j	m-MeC ₆ H ₄	3aj	85	89
11	1a	2k	m-BrC ₆ H ₄	3ak	85	70
12	1a	21	$H_2C = CH$	3al	81	58
13	1b	2a	C ₆ H ₅	3ba	91	91
14	1b	2b	o-MeOC ₆ H ₄	3bb	78	74
15	1b	2c	o-MeC ₆ H ₄	3bc	84	81
16	1b	2d	o-ClC ₆ H ₄	3bd	83	83
17	1b	2e	<i>p</i> -MeOC ₆ H ₄	3be	85	87
18	1b	2f	p-MeC ₆ H ₄	3bf	90	88
19	1b	2g	p-ClC ₆ H ₄	3bg	87	87
20	1b	2h	p-BrC ₆ H ₄	3bh	88	84
21	1b	2i	p-FC ₆ H ₄	3bi	89	94
22	1b	2j	m-MeC ₆ H ₄	3bj	91	98
23	1b	2k	m-BrC ₆ H ₄	3bk	85	87
24	1b	21	H ₂ C=CH	3bl	86	61

^{*a*}Isolated yield. ^{*b*}Stoichiometric conditions: cyclopropane (0.5 mmol), azide (0.5 mmol), TiCl₄ (0.5 mmol, 100 mol %), CH_2Cl_2 (5 mL), 0 °C to rt, 2 h. ^{*c*}Substoichiometric conditions: cyclopropane (0.3 mmol), azide (0.3 mmol), TiCl₄ (0.06 mmol, 20 mol %), HFIP (3 mL), 0 °C to rt, 12 h. See Supporting Information for more details.

performed on a 5.0 mmol (1.17 g) scale. The reaction proceeded similarly to the small-scale experiment and provided **3aa** in 82% yield after 4 h. The molecular structure of **3ab** was unambiguously established by X-ray crystallographic analysis (see Supporting Information).¹⁰

When the reaction was conducted in HFIP with 20 mol % TiCl₄, benzyl azide generally gave lower yields than decyl azide probably due to the partial decomposition of benzyl azide under this conditions.¹¹ When vinyl substituted cyclopropane was employed, the yields reduced dramatically (entries 12, 24). This may be attributed to the electrophilic addition of carbocation to the vinyl group on cyclopropane. When *o*-MeOC₆H₄- and *p*-MeOC₆H₄-substituted cylcopropanes reacted with benzyl azide, the lower yields were obtained due to cyclodimerization of cyclopropanes (entries 2, 5).¹² Because of steric effect, *o*-ClC₆H₄- substituted cyclopropane also gave lower yield than *p*-ClC₆H₄- substituted ones (entries 4, 7).

An attractive feature of this cycloaddition reaction is the opportunity for ready access to *spiro*-triazinines. We were pleased to find that the [3 + 3] cycloaddition of cyclopropane 1,1-diester **2m**¹³ with **1b** under the stoichiometric conditions also proceeded well to give *spiro*-triazinine **3bm**, albeit with a modest yield (46%) (Scheme 1).

In an effort to gain knowledge on the mechanism of our reaction, the reaction between (S)-2a¹⁴ and 1b was carried out

Scheme 1. Synthesis of spiro-Triazinine



in the presence of 1.0 equiv of $TiCl_4$ and racemic mixture of **3ba** was obtained (Scheme 2). This indicated that the

Scheme 2. [3 + 3] Cycloaddition of Enantioenriched Cyclopropane

<i>n</i> -C ₁₀ H ₂₁ N ₃ +	Ph ^{```} COOMe COOMe	$\frac{\text{TiCl}_4 (100 \text{ mol }\%)}{\text{CH}_2\text{Cl}_2, 0 \text{ °C-rt}, n-\text{C}_1}$	Ph COOMe COOMe NOH21 N N N
1b	(S) -2a, 90% ee		<i>rac</i> - 3ba , yield: 94%

00011

mechanism of this cycloaddition is stepwise, rather than concerted process. In view of this, plausible reaction mechanism for these transformations is proposed in Scheme 3. Initially, under the activation of $TiCl_4$, the ring-opening of





cyclopropane 2 affords 1,3-zwitterionic intermediate A. Subsequently, a nucleophilic attack of an azide¹⁵ happens to form a new zwitterion B, which then undergoes an intramolecular nucleophilic attack to form the product 3.

As early as 1989, Dunkin and co-workers reported that the thermal decomposition of 1-methyl-1,2,3-benzotriazin-4(1H)one offered benzazetinone by loss of nitrogen.¹⁶ If our products decomposed in the same way, azetidines, a class of useful molecular,¹⁷ will be obtained. Therefore, the solution of 3aa in xylene was heated to reflux. After 4 h, 3aa was exhausted and 4aa was obtained in 46% yield (Scheme 4). In order to improve the yield, we changed the solvent to toluene or dimethyl sulfoxide to promote the reaction. But no expected increase of vield developed. Thermolysis of 3al, 3bc, 3bf and 3bh offered 4al, 4bc, 4bf and 4bh respectively in 40-52% yield. Among these reactions, imine 5 was obtained along with 4bh. This disclosed that the decomposition occurred via two pathways (Scheme 5). Under the thermolysis conditions, the cleavage of N-N bond affords intermediate C. Which undergoes an intramolecular nucleophilic substitution to furnish the desired azetidine 4 in path A or a Grob fragmentation to furnish the imine 5 in path B.

In conclusion, we have discovered the first $TiCl_4$ promoted [3 + 3] cycloaddition of 2-substituted cyclopropane-1,1-diesters with azides to devise complex triazinines with diverse substituent patterns in good yields. Both stoichiometric and substoichiometric versions of this reaction were investigated through judicious choice of solvent. Notably, this chemistry offers an efficient and practical strategy for the construction of

Letter





Scheme 5. Proposed Pathways of Thermolysis



medicinally valuable azetidines by means of simple thermolysis of the triazinine products. Further related studies utilizing this novel [3 + 3] cycloaddition for the development of other methodologies are currently underway in our laboratory.

ASSOCIATED CONTENT

Supporting Information

Experimental procedure and characterization data for all products, including ¹H NMR and ¹³C NMR spectra, X-ray structural information on **3ab** (CIF) and chiral HPLC chromatograms of (S)-**2a** and *rac*-**3ba**. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Authors

- *E-mail: luoych@lzu.edu.cn.
- *E-mail: xupf@lzu.edu.cn.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We are grateful for the NSFC (21202070, 21402073), Specialized Research Fund for the Doctoral Program of Higher Education (SRFDP 20110211120017), Gansu Provincial Natural Science Fund (2011GS04143) and the Fundamental Research Funds for the Central Universities (Lzujbky-2014-95 and Lzujbky-2014-65).

REFERENCES

(1) (a) Agalave, S. G.; Maujan, S. R.; Pore, V. S. Chem.—Asian J.
 2011, 6, 2696. (b) Xia, Y.; Liu, Y.; Wan, J.; Wang, M.; Rocchi, P.; Qu, F.; Iovanna, J. L.; Peng, L. J. Med. Chem. 2009, 52, 6083. (c) French, A. M.; Khan, M. T.; Scott, N. C. Br. J. Pharmacol. 1978, 63, 475. (d) Caliendo, G.; Fiorino, F.; Grieco, P.; Perissutti, E.; Santagada, V.; Meli, R.; Raso, G. M.; Zanesco, A.; Nucci, G. D. Eur. J. Med. Chem. 1999, 34, 1043. (e) Wanner, M. J.; Koomen, G.-J. J. Chem. Soc. Perkin Trans. 1 2002, 1877. (f) Larkin, J. M. G.; Hughes, S. A.; Beirne, D. A.; Patel, P. M.; Gibbens, I. M.; Bate, S. C.; Thomas, K.; Eisen, T. G.; Gore, M. E. Br. J. Cancer 2007, 96, 44. (g) Mueller, R.; Rachwal, S.; Varney, M. A.; Johnson, S. A.; Alisala, K.; Zhong, S.; Li, Y.-X.; Haroldsen, P.; Herbst, T.; Street, L. J. Bioorg. Med. Chem. Lett. 2011, 21, 7455.

(2) (a) Saravanan, J.; Mohan, S.; Roy, J. J. *Eur. J. Med. Chem.* **2010**, 45, 4365. (b) Kumar, R.; Singh, A. D.; Singh, J.; Singh, H.; Roy, R. K.; Chaudhary, A. *Mini-Rev. Med. Chem.* **2014**, *14*, 72.

(3) Schmidt, B. F.; Snyder, E. J.; Carroll, R. M.; Farnsworth, D. W.; Michejda, C. J.; Smith, R. H., Jr. J. Org. Chem. **1997**, *62*, 8660.

(4) (a) Mitani, M.; Tachizawa, O.; Takeuchi, H.; Koyama, K. J. Org. Chem. 1989, 54, 5397. (b) Wang, Y.-F.; Chiba, S. J. Am. Chem. Soc. 2009, 131, 12570. (c) Wang, Y.-F.; Toh, K. K.; Ng, E. P. J.; Chiba, S. J. Am. Chem. Soc. 2011, 133, 6411. (d) Desai, P.; Aubé, J. Org. Lett. 2000, 2, 1657. (e) Grecian, S.; Desai, P.; Mossman, C.; Poutsma, J. L.; Aubé, J. J. Org. Chem. 2007, 72, 9439.

(5) (a) Scadeng, O.; Ferguson, M. J.; West, F. G. Org. Lett. 2011, 13, 114. (b) Rostami, A.; Wang, Y.; Arif, A. M.; McDonald, R.; West, F. G. Org. Lett. 2007, 9, 703. (c) Pearson, W. H.; Fang, W.-H.; Kampf, J. W. J. Org. Chem. 1994, 59, 2682.

(6) For reviews of donor-acceptor cyclopropane, see: (a) Reissig, H.-U.; Zimmer, R. Chem. Rev. 2003, 103, 1151. (b) Yu, M.; Pagenkopf, B. L. Tetrahedron 2005, 61, 321. (c) De Simone, F.; Waser, J. Synthesis 2009, 20, 3353. (d) Campbell, M. J.; Johnson, J. S.; Parsons, A. T.; Pohlhaus, P. D.; Sanders, S. D. J. Org. Chem. 2010, 75, 6317. (e) Melnikov, M. Ya.; Budynina, E. M.; Ivanova, O. A.; Trushkov, I. V. Mendeleev Commun. 2011, 21, 293. (f) Tang, P.; Qin, Y. Synthesis 2012, 44, 2969. (g) Schneider, T. F.; Kaschel, J.; Werz, D. B. Angew. Chem., Int. Ed. 2014, 53, 5504.

(7) Selected examples for nitrones: (a) Sapeta, K.; Kerr, M. A. J. Org. Chem. 2007, 72, 8597. (b) Yang, H.-B.; Shi, M. Org. Biomol. Chem. 2012, 10, 8236. (c) Gorbacheva, E. O.; Tabolin, A. A.; Novikov, R. A.; Khomutova, Y. A.; Nelyubina, Y. V.; Tomilov, Y. V.; Ioffe, S. L. Org. Lett. 2013, 15, 350.

(8) (a) Sapeta, K.; Kerr, M. A. Org. Lett. 2009, 11, 2081. (b) Lebold, T. P.; Leduc, A. B.; Kerr, M. A. Org. Lett. 2009, 11, 3770. (c) Leduc, A. B.; Lebold, T. P.; Kerr, M. A. J. Org. Chem. 2009, 74, 8414. (d) Perreault, C.; Goudreau, S. R.; Zimmer, L. E.; Charette, A. B. Org. Lett. 2008, 10, 689. (e) Zhou, Y.-Y.; Li, J.; Ling, L.; Liao, S.-H.; Sun, X.-L.; Li, Y.-X.; Wang, L.-J.; Tang, Y. Angew. Chem., Int. Ed. 2013, 52, 1452. (f) Sathishkannan, G.; Srinivasan, K. Chem. Commun. 2014, 50, 4062. (g) Talukdar, R.; Tiwari, D. P.; Saha, A.; Ghorai, M. K. Org. Lett. 2014, 16, 3954.

(9) Motiwala, H. F.; Fehl, C.; Li, S.-W.; Hirt, E.; Porubsky, P.; Aubé, J. J. Am. Chem. Soc. **2013**, 135, 9000.

(10) Crystal data for **3ab** has been deposited in CCDC as deposition number 959858. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre *via* www.ccdc.com. According to the crystal structure and the ¹H NMR spectra, the conformations of triazinines **3** may be different in solid and solution state. See Supporting Information for more details.

(11) Schildknegt, K.; Agrios, K. A.; Aubé, J. *Tetrahedron Lett.* **1998**, 39, 7687.

(12) Ivanova, O. A.; Budynina, E. M.; Chagarovskiy, A. O.; Trushkov, I. V.; Melnikov, M. Y. *J. Org. Chem.* **2011**, *76*, 8852.

(13) Carson, C. A.; Kerr, M. A. Org. Lett. **2009**, *11*, 777.

(14) Davies, H. M. L.; Bruzinski, P. R.; Lake, D. H.; Kong, N.; Fall,
 M. J. J. Am. Chem. Soc. 1996, 118, 6897.

(15) (a) Pearson, W. H.; Schkeryantz, J. S. *Tetrahedron Lett.* **1992**, 33, 5291. (b) Pearson, W. H.; Walavalkar, R.; Schkeryantz, J. S.; Fang,

W.-K.; Blickensdorf, J. D. J. Am. Chem. Soc. 1993, 115, 10183.
(16) Dunkin, I. R.; Lynch, M. A.; Withnall, R.; Boulton, A. J.;

Henderson, N. J. Chem. Soc., Chem. Commun. 1989, 1777.

(17) Brandi, A.; Cicchi, S.; Cordero, F. M. Chem. Rev. 2008, 108, 3988.