

Characterization of 14,15-Epoxyeicosatrienoyl-Sulfonamides as 14,15-Epoxyeicosatrienoic Acid Agonists: Use for Studies of Metabolism and Ligand Binding

Wenqi Yang, Blythe B. Holmes, V. Raj Gopal, R. V. Krishna Kishore, Bhavani Sangras, Xiu-Yu Yi, J. R. Falck, and William B. Campbell

Department of Pharmacology and Toxicology, Medical College of Wisconsin, Milwaukee, Wisconsin (W.Y., B.B.H., X.-Y.Y., W.B.C.); and Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, Texas (V.R.G., R.V.K.K., B.S., J.R.F.)

Received January 10, 2007; accepted February 26, 2007

ABSTRACT

Epoxyeicosatrienoic acids (EETs) are cytochrome P450 epoxygenase metabolites of arachidonic acid. EETs mediate numerous biological functions. In coronary arteries, they regulate vascular tone by the activation of smooth muscle large-conductance, calcium-activated potassium (BK_{Ca}) channels to cause hyperpolarization and relaxation. We developed a series of 14,15-EET agonists, 14,15-EET-phenylidosulfonamide (14,15-EET-PISA), 14,15-EET-biotinsulfonamide (14,15-EET-BSA), and 14,15-EET-benzoyldihydrocinnamide-sulfonamide (14,15-EET-BZDC-SA) as tools to characterize 14,15-EET metabolism and binding. Agonist activities of these analogs were characterized in precontracted bovine coronary arterial rings. All three analogs induced concentration-dependent relaxation and were equipotent with 14,15-EET. Relaxations to these analogs were inhibited by the BK_{Ca} channel blocker iberiotoxin (100 nM), the 14,15-EET antagonist 14,15-epoxyeicosa-5(Z)-enoyl-

methylsulfonamide (10 μ M), and abolished by 20 mM extracellular K^+ . 14,15-EET-PISA is metabolized to 14,15-dihydroxyeicosatrienoyl-PISA by soluble epoxide hydrolase in bovine coronary arteries and U937 cells but not U937 cell membrane fractions. 14,15-EET- $P^{125}I$ SA binding to human U937 cell membranes was time-dependent, concentration-dependent, and saturable. The specific binding reached equilibrium by 15 min at 4°C and remained unchanged up to 30 min. The estimated K_d and B_{max} were 148.3 ± 36.4 nM and 3.3 ± 0.5 pmol/mg protein, respectively. These data suggest that 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA are full 14,15-EET agonists. 14,15-EET- $P^{125}I$ SA is a new radiolabeled tool to study EET metabolism and binding. Our results also provide preliminary evidence that EETs exert their biological effect through a membrane binding site/receptor.

Epoxyeicosatrienoic acids (EETs) are cytochrome P450 epoxygenase metabolites of arachidonic acid. There are four regioisomers of EET: 5,6-, 8,9-, 11,12-, and 14,15-EET (Capdevila et al., 1981; Rosolowsky and Campbell, 1996; Nithipatikom et al., 2001). They play key roles in regulating vascular tone and homeostasis. Previous studies from our lab and other groups have characterized EETs as endothelium-de-

rived hyperpolarizing factors in the coronary circulation (Rosolowsky and Campbell, 1993; Fulton et al., 1995; Campbell et al., 1996; Fisslthaler et al., 1999). Vascular endothelium synthesizes and releases EETs in response to bradykinin, acetylcholine, arachidonic acid, flow, or cyclic stretch (Pinto et al., 1987; Campbell et al., 1996; Rosolowsky and Campbell, 1996; Fisslthaler et al., 1999; Gauthier et al., 2005; Huang et al., 2005). EETs then activate smooth muscle membrane large-conductance, calcium-activated potassium (BK_{Ca}) channels to cause hyperpolarization and vascular relaxation (Campbell et al., 1996; Gauthier et al., 2005). In addition, EETs are involved in other biological functions

This work was supported by the National Institutes of Health (Grants HL-51055 and GM-31278) and by the Robert A. Welch Foundation.

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.
doi:10.1124/jpet.107.119651.

ABBREVIATIONS: EET, epoxyeicosatrienoic acid; BK_{Ca} , large-conductance, calcium-activated potassium channel; sEH, soluble epoxide hydrolase; DHET, dihydroxyeicosatrienoic acid; 14,15-EET-PISA, 14,15-EET-phenylidosulfonamide; 14,15-EET-BSA, 14,15-EET-biotinsulfonamide; 14,15-EET-BZDC-SA, 14,15-EET-benzoyldihydrocinnamide-sulfonamide; THF, tetrahydrofuran; NHS, *N*-hydroxysuccinimide; TLC, thin layer chromatography; HPLC, high-performance liquid chromatography; IBTX, iberiotoxin; 14,15-EEZE-mSA, 14,15-epoxyeicosa-5(Z)-enoyl-methylsulfonamide; HBSS, Hanks' balanced salt solution; BCA, bovine coronary artery; CDU, 1-cyclohexyl-3-dodecylurea; MS, mass spectrometry; 14,15-EET-mSA, 14,15-EET-methylsulfonamide; U46619, 9,11-dideoxy-9 α ,11 α -methanoepoxy-prostaglandin $F_{2\alpha}$; HMPA, hexamethyl phosphoramide; DMF, *N,N*-dimethylformamide.

including anti-inflammation (Node et al., 1999), antipyresis (Kozak et al., 2000), angiogenesis (Michaelis and Fleming, 2006), and fibrinolysis (Node et al., 2001). Mechanisms involved in EET bioactivities include activation of K^+ channels (Campbell et al., 1996; Yang et al., 2005), G-protein ADP-ribosylation (Li et al., 1999), protein kinase A, and tyrosine kinase pathways (Node et al., 2001; Michaelis and Fleming, 2006). To maintain full agonist vascular activity of 14,15-EET in bovine coronary arteries, key structural components including the carbon-1 (C-1) carboxyl group, the 8,9-double bond, the 14(*S*),15(*R*)-cis epoxide group, and a 14-carbon distance between the carboxyl and the epoxide group are required (Falck et al., 2003a; Gauthier et al., 2004). Likewise, specific structural components are required for 11,12-EET-mediated inhibition of vascular cell adhesion molecule-1 expression (Falck et al., 2003b). Such structure-activity relationships indicate a possible lipid-protein binding event that requires precise conformation of the EET molecules. Numerous studies have indicated that the EET action site may be the plasma membrane. 11,12-EET activates coronary smooth muscle BK_{Ca} channels in inside-out patches where a small portion of plasma membrane was excised (Li and Campbell, 1997). In aortic smooth muscle cells, a membrane-impermeable 14,15-EET derivative was equipotent as 14,15-EET in inhibiting aromatase activity, indicating that the EET acts on the cell surface (Snyder et al., 2002). Wong et al. (1993, 2000) reported protein binding sites in both intact guinea pig monocytes and mononuclear cell membranes by radioligand binding studies using [3H]14,15-EET. [3H]14,15-EET binding to guinea pig monocytes was sensitive to mild protease treatment, which suggests a protein identity of the 14,15-EET binding site. However, a membrane receptor/binding protein for EETs has not been purified or cloned. Tools to identify the putative EET binding protein/receptor are needed.

EETs are rapidly taken up into cells when applied exogenously (Snyder et al., 2002). Whether this uptake is through simple diffusion or a protein transporter is unclear. EETs are actively metabolized by multiple enzyme pathways. The major pathways are hydrolysis of the epoxide groups by soluble epoxide hydrolase (sEH) to the corresponding vicinal-dihydroxyeicosatrienoic acids (DHETs), β -oxidation that results in chain-shortened products, and esterification into membrane phosphoglycerolipids (Spector et al., 2004). These studies were done with ^{14}C - or 3H -labeled EETs synthesized from radiolabeled arachidonic acid. At present, no commercial company markets radiolabeled EETs. The synthesis of radiolabeled EETs from radiolabeled arachidonic acid is inefficient and costly because: 1) yields cannot exceed 50%, or diepoxides are formed; and 2) chemical methods produce all four EET regioisomers in varying amounts (Corey et al., 1979; Rosolowsky and Campbell, 1993). Thus, there is a need to produce radiolabeled EETs efficiently, cheaply and with high specific activity for studies of EET transport, metabolism, and binding.

To serve these purposes, we developed and characterized a series of stable 14,15-EET agonists, 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA (see Fig. 2). These analogs contain an *N*-acylsulfonamide group instead of a terminal carboxylic acid. The *N*-acylsulfonamide group has a similar pK_a as a carboxyl group and obviates β -oxidation and phospholipid esterification (Backes and Ellman, 1994; Snyder

et al., 2002). For 14,15-EET-PISA, a benzyl ring that may be iodinated with [^{125}I] or [^{127}I] is attached to the sulfonamide group. For 14,15-EET-BSA and 14,15-EET-BZDC-SA, a biotin or BZDC moiety is covalently attached to the sulfonamide group through a benzyl linker. We characterized 14,15-EET- $P^{125}ISA$ as a radioactive tracer to study EET metabolism and as a radioligand for studying the binding kinetics of the putative EET receptor/binding site. It was used as a model for studying the metabolism of other EET-sulfonamides.

Materials and Methods

Preparation of 14,15-EET-PISA, 14,15-EET-BZDC-SA, and 14,15-EET-BSA

Preparation of 14,15-EET-PISA (5) (Fig. 1, Scheme 1). *Synthesis of N-hydroxysuccinimide-ester 2.* *N*-Hydroxysuccinimide (115 mg, 1 mmol), dried azeotropically using anhydrous toluene and then under high vacuum for 2 h, was added to a solution of 14,15-EET (1) (132 mg, 0.412 mmol) in dry THF (8 ml) at room temperature under an argon atmosphere followed by *N,N'*-dicyclohexylcarbodiimide (94 mg, 0.453 mmol). After stirring overnight, the solvent was removed in vacuo, and the residue was purified by SiO_2 column chromatography eluting with hexanes/EtOAc (70:30) to give *N*-hydroxysuccinimide (NHS)-ester 2 (151 mg, 88%). TLC: $R_f \sim 0.30$ (50% EtOAc/hexanes); 1H NMR ($CDCl_3$, 300 MHz) δ 0.92 (t, $J = 7.0$ Hz, 3H), 1.32–1.60 (m, 8H), 1.82 (quintet, $J = 7.3$ Hz, 2H), 2.17–2.22 (m, 3H), 2.37–2.45 (m, 1H), 2.62 (t, $J = 7.3$ Hz, 2H), 2.80–2.88 (m, 8H), 2.92–2.99 (m, 2H), 5.34–5.54 (m, 6H).

Synthesis of stannane 3. A mixture of 3-bromobenzenesulfonamide (Sigma-Aldrich Co., St. Louis, MO; 1 g, 4.23 mmol), tetrakis(triphenylphosphine)palladium(0) (0.24 g, 0.209 mmol), and bis(tributyltin) (4.3 g, 7.42 mmol) was stirred at room temperature in a sealed pressure tube under an argon atmosphere in anhydrous CH_3CN (25 ml) and toluene (5 ml) for 10 min, then warmed to 95°C. After 48 h, the reaction mixture was cooled to room temperature, concentrated in vacuo, and the residue was purified by silica gel column chromatography to yield 3 (870 mg, 51%). TLC: $R_f \sim 0.32$ (30% EtOAc/hexanes); 1H NMR ($CDCl_3$, 400 MHz) δ 0.88 (t, $J = 7.4$ Hz, 9H), 1.11 (app t, $J = 7.4$ Hz, 6H), 1.25–1.36 (m, 6H), 1.42–1.58 (m, 6H), 5.08 (s, 2H), 7.42–7.47 (m, 1H), 7.66 (dd, $J_{H,H} = 8.3$ Hz, $J_{Sn,H} = 34.5$ Hz, 1H), 7.85 (dd, $J = 1.8$, 8.4 Hz, 1H), 8.02 (dd, $J_{H,H} = 1.8$ Hz, $J_{Sn,H} = 34.5$ Hz, 1H).

Synthesis of sulfonamide 4. A solution of 3-(tri-*n*-butylstannanyl)-benzenesulfonamide (3) (80 mg, 0.179 mmol) and 4-(dimethylamino)pyridine (20 mg, 0.157 mmol) in dry HMPA (1.5 ml) was cannulated into a solution of NHS-ester 2 (60 mg, 0.143 mmol) in HMPA (2.5 ml) under an argon atmosphere. After stirring for 10 min at room temperature, the reaction mixture was warmed to 65°C and allowed to stir for 16 h, then quenched by the addition of water (10 ml). The mixture was extracted with EtOAc (3×15 ml), and the combined extracts were washed with water, brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by SiO_2 column chromatography to give stannane 4 (38 mg, 36%). TLC: $R_f \sim 0.52$ (30% EtOAc/hexanes); 1H NMR ($CDCl_3$, 300 MHz) δ 0.86–0.93 (m, 12H), 1.11 (app t, $J = 7.4$ Hz, 6H), 1.26–1.39 (m, 10H), 1.47–1.68 (m, 12H), 2.04 (dd, $J = 7.0$, 7.0 Hz, 2H), 2.17–2.26 (m, 3H), 2.43–2.51 (m, 1H), 2.69–2.91 (m, 4H), 3.02–3.12 (m, 2H), 5.20–5.61 (m, 6H), 7.40–7.48 (m, 1H), 7.70 (dd, $J_{H,H} = 8.4$, $J_{Sn,H} = 34.5$ Hz, 1H), 7.96 (dd, $J = 1.8$, 8.4 Hz, 1H), 8.08 (dd, $J_{H,H} = 1.8$ Hz, $J_{Sn,H} = 34.5$ Hz, 1H), 8.99 (br s, 1H).

Synthesis of iodide 5. A solution of sodium iodide (40 mg, 0.027 mmol) in water (1.0 ml) was added to a 0°C solution of stannane 4 (20 mg, 0.027 mmol) in 95% EtOH (4 ml) under an argon atmosphere followed by H_2O_2 (30% aq. soln, 24 μ l, 0.213 mmol) and peracetic acid (32% in AcOH, 26 μ l, 0.213 mmol). The reaction mixture was




Fig. 1. Synthesis of 14,15-EET-sulfonamides in Schemes 1, 2, and 3.

warmed to room temperature. After 2 h, the reaction mixture was quenched with saturated aqueous sodium thiosulfate (1 ml) and extracted with EtOAc (3×15 ml). The combined organic extracts were washed with water, brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by silica gel column chromatography to give 5 (9 mg, 58%). TLC: $R_f \sim 0.32$ (30% EtOH/hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 0.92 (t, $J = 7$ Hz, 3H), 1.32–1.69 (m, 1H), 2.04 (dd, $J = 7.0, 7.0$ Hz, 2H), 2.18–2.26 (m, 3H), 2.46–2.54 (m, 1H), 2.70–2.94 (m, 4H), 3.06–3.17 (m, 2H), 5.20–5.60 (m, 6H), 7.27 (dd, $J = 8.4, 8.4$ Hz, 1H), 7.95 (dd, $J = 1.8, 8.4$ Hz, 1H), 7.96 (dd, $J = 1.8, 8.4$ Hz, 1H), 8.05 (dd, $J = 1.8, 8.4$ Hz, 1H), 8.33 (dd, $J = 1.8, 1.8$ Hz, 1H), 9.37 (br s, 1H).

Preparation of 14,15-EET-BZDC-SA (8) (Fig. 1, Scheme 2). *Synthesis of aniline 6.* *n*-Butyl lithium (0.21 ml of a 2.5 M solution in hexanes, 0.539 mmol) was added slowly to a -78°C solution of 4-aminobenzenesulfonamide (93 mg, 0.539 mmol) in anhydrous THF (8 ml) and HMPA (2 ml) under an argon atmosphere. After 5 min, a solution of NHS-ester 2 (150 mg, 0.359 mmol) in THF (4 ml) was added at -78°C , and then the mixture was slowly warmed to room temperature. After stirring overnight, the reaction mixture was quenched with saturated aq. NH_4Cl (5 ml) and extracted with EtOAc (3×20 ml). The combined organic extracts were washed with water, brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by SiO_2 column chromatography to give 6 (120 mg, 71%). TLC: $R_f \sim 0.34$ (50% EtOAc/hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 0.92 (t, $J = 7.0$ Hz, 3H), 1.32–1.60 (m, 8H), 1.63 (quintet, $J = 7.3$ Hz, 2H), 2.03 (dd, $J = 7.3, 7.3$ Hz, 2H), 2.18–2.26 (m, 2H), 2.42–2.49 (m, 1H), 2.71–2.89 (m, 4H), 3.01–3.07 (m, 2H), 4.31 (br s, 2H), 5.25–5.54 (m, 6H), 6.65 (d, $J = 8.6$ Hz, 2H), 7.80 (d, $J = 8.6$ Hz, 2H), 9.01 (br s, 1H).

Synthesis of BZDC 8. A solution of 4-(benzoyl)benzenepropanoic acid NHS ester (Kort et al., 2000) (7) (22.3 mg, 0.063 mmol) and Et_3N (13 μl , 0.126 mmol) in DMF (2 ml) was added dropwise to a solution of 6 (20 mg, 0.042 mmol) in anhydrous DMF (2 ml) followed by 4-(dimethylamino)pyridine (7.7 mg, 0.063 mmol) under an argon atmosphere. After 48 h, the reaction mixture was quenched with

water (3 ml), extracted with EtOAc (3×15 ml), and the combined organic extracts were washed with water, brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by silica gel column chromatography to yield 8 (13 mg, 30%). TLC: $R_f \sim 0.42$ (20% EtOAc/ CH_2Cl_2); ^1H NMR (CDCl_3 , 400 MHz) δ 0.90 (t, $J = 7.0$ Hz, 3H), 1.31–1.62 (m, 8H), 1.63 (quintet, $J = 7.3$ Hz, 2H), 2.41–2.53 (m, 1H), 2.71–2.92 (m, 6H), 3.02–3.20 (m, 4H), 5.20–5.60 (m, 6H), 7.34 (d, $J = 8.2$ Hz, 2H), 7.48 (t, $J = 7.6$ Hz, 2H), 7.56–7.68 (m, 3H), 7.73 (d, $J = 7.9$ Hz, 2H), 7.78 (d, $J = 7.9$ Hz, 2H), 7.88 (d, $J = 8.8$ Hz, 2H), 8.21 (br s, 1H), 9.35 (br s, 1H).

Preparation of 14,15-EET-BSA (10) (Fig. 1, Scheme 3). *Synthesis of acid 9.* 4-Carboxybenzenesulfonamide (Aldrich Chemical Co., Milwaukee, WI; 130 mg, 0.358 mmol), dried azeotropically with anhydrous benzene and then dried under high vacuum for 2 h, was dissolved in anhydrous THF (6 ml) and HMPA (1.5 ml) and cooled to -78°C under an argon atmosphere. To this, *n*-butyl lithium (0.57 ml of a 2.5 M solution in hexanes, 0.862 mmol) was added, and after 10 min, a solution of 2 (180 mg, 0.431 mmol) in THF (4 ml) was slowly added. The reaction mixture was then slowly warmed to room temperature. After stirring overnight, the reaction mixture was quenched with saturated aqueous NH_4Cl solution (15 ml), extracted with EtOAc (3×30 ml), and the combined organic extracts were washed with water, brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by silica gel column chromatography to give acid 9 (85 mg, 71%). TLC: $R_f \sim 0.23$ (50% EtOAc/hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 0.91 (t, $J = 7.0$ Hz, 3H), 1.34–1.60 (m, 8H), 2.02–2.12 (m, 2H), 2.26 (t, $J = 8.5$ Hz, 2H), 2.42–2.49 (m, 1H), 2.71–2.89 (m, 4H), 3.07–3.16 (m, 2H), 5.23–5.57 (m, 6H), 8.14 (d, $J = 8.4$ Hz, 2H), 8.23 (d, $J = 8.6$ Hz, 2H).

Synthesis of biotin 10. To a -78°C solution of 9 (50 mg, 0.099 mmol) in dry CH_2Cl_2 (3 ml) under an argon atmosphere were added 1-methylpiperidine (Aldrich; 12 μl , 0.099 mmol) followed by isobutyl chloroformate (Aldrich; 13 μl , 0.099 mmol). After stirring for 45 min at the same temperature, the reaction mixture was warmed to -25°C , stirred for another 30 min, and then a solution of biotin hydrazide (Sigma-Aldrich Co.; 100 mg, 0.39 mmol) in DMF (1 ml,

prewarmed to $\sim 80^\circ\text{C}$ to dissolve) was added. The reaction mixture was slowly warmed to room temperature over 1 h. After stirring at room temperature overnight, the reaction mixture was diluted with CH_2Cl_2 (10 ml), filtered, washed with water, and concentrated in vacuo. The residue was purified by SiO_2 PTLC to give 10 (15 mg, 38%) and recovered starting material (30 mg). TLC: $R_f \sim 0.45$ (10% $\text{MeOH}/\text{CH}_2\text{Cl}_2$); ^1H NMR (CDCl_3 , 300 MHz) δ 0.90 (t, $J = 6.7$ Hz, 3H), 1.28–1.42 (m, 6H), 1.43–1.84 (m, 10H), 1.99–2.10 (m, 2H), 2.25 (apparent t, $J = 7.0$ Hz, 2H), 2.28–2.33 (m, 2H), 2.36 (apparent t, $J = 6.8$ Hz, 2H), 2.68–2.78 (m, 4H), 2.84 (apparent t, $J = 5.8$ Hz, 2H), 2.91–3.01 (m, 2H), 3.18–3.28 (m, 1H), 3.32 (br s, 1H, D_2O exchangeable), 3.90 (d, $J = 6.4$ Hz, 1H, D_2O exchangeable), 4.30–4.65 (m, 1H), 4.48–4.54 (m, 1H), 5.26–5.40 (m, 4H), 5.42–5.60 (m, 2H), 8.04 (d, $J = 8.5$ Hz, 2H), 8.10 (d, $J = 8.5$ Hz, 2H).

Synthesis of 14,15-EET- P^{125}ISA and 14,15-DHET- P^{125}ISA

To synthesize 14,15-EET- P^{125}ISA , carrier-free Na^{125}I (2 μl , pH 9–10, 200 μCi , 17.4 Ci/mg) (GE Healthcare, Arlington Heights, IL) was mixed with NaI (1 μl , 1.08 mg in 10 ml H_2O). 14,15-EET-stannane (4) (10 μg in 43 μl of ethanol) was added to the NaI solution followed by H_2O_2 (10.67 nmol in 1.25 μl of H_2O) and peracetic acid (2.69 nmol in 1.68 μl of acetic acid) at 4°C (Fig. 1, scheme 1). The reaction mixture was warmed to room temperature and allowed to react for 45 min. On quenching with saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (250 μl), the aqueous layer was extracted with ethyl acetate (2×300 μl). The combined organic extracts were washed with H_2O (300 μl) and 0.9% NaCl (300 μl) and dried under N_2 . The residues were purified by reverse-phase high-performance liquid chromatography (HPLC) as described previously (Campbell et al., 1996). The extracts were redissolved in 100 μl of acetonitrile/glacial acetic acid (999:1) and 100 μl of distilled H_2O and resolved on a Nucleosil C18 reverse-phase column (5 μm , 4.6×250 mm; Phenomenex, Torrance, CA). Solvent A was distilled H_2O , and solvent B was acetonitrile/glacial acetic acid (999:1). A linear gradient from 50% solvent B in solvent A to 94% solvent B within 40 min was used at a flow rate of 1 ml/min. Column effluent corresponding to synthetic 14,15-EET-PISA was collected, extracted, and dried under N_2 .

To synthesize 14,15-DHET- P^{125}ISA , 14,15-EET- P^{125}ISA was dissolved in HEPES buffer containing 10 mM HEPES, 150 mM NaCl, 5 mM KCl, 1.8 mM CaCl_2 , 1 mM MgCl_2 , and 5.5 mM glucose, pH 7.4. Recombinant human sEH (35 μg) was added and incubated for 30 min at 37°C . The residues were extracted with ethyl acetate and purified by HPLC as described above. Specific activities of 14,15-EET- P^{125}ISA and 14,15-DHET- P^{125}ISA were 257.5 Ci/mmol.

Vascular Reactivity of Bovine Coronary Arteries

Bovine hearts were purchased from a local slaughterhouse. The left anterior descending coronary artery was dissected and cleaned of connective tissue. Arteries of 2-mm diameter were cut into rings (3-mm width) and suspended on a pair of stainless hooks in a 6-ml water-jacketed organ chamber in Krebs' buffer consisting of 119 mM NaCl, 4.8 mM KCl, 24 mM NaHCO_3 , 0.2 mM KH_2PO_4 , 0.2 mM MgSO_4 , 11 mM glucose, 0.02 mM EDTA, and 3.2 mM CaCl_2 . The buffer was equilibrated with 95% O_2 -5% CO_2 and maintained at 37°C . Tension was continuously recorded as described previously (Campbell et al., 1996). In brief, submaximal concentrations of the thromboxane-mimetic U46619 (10–20 nM; Cayman Chemical Co., Ann Arbor, MI) were administered to contract the vessels to 50 to 75% of KCl-induced contraction. Cumulative concentrations of 14,15-EET, 14,15-EET-PISA, 14,15-EET-BSA, or 14,15-EET-BZDC-SA were added to the chamber. In some studies, the arteries were incubated with ibuprofen (IBTX; 100 nM; Sigma-Aldrich Co.), 14,15-epoxyeicosa-5(Z)-enoyl-methylsulfonamide (14,15-EEZE-mSA; 10 μM), or vehicle for 10 min before U46619 contraction (Gauthier et al., 2003). In the high extracellular potassium ($[\text{K}^+]_o$) studies, K^+ was increased to 20 mM by substitution of Na^+ . Results are expressed as percent relaxation with 100% relaxation representing basal tension.

Culture of U937 Cells and Membrane Preparation

U937 cells were cultured in suspension in RPMI 1640 medium (Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum (Hyclone, South Logan, UT), 25 mM HEPES, 2 mM L-glutamine, 1 mM sodium pyruvate, 100 U/ml penicillin, 100 $\mu\text{g}/\text{ml}$ streptomycin, and 0.25 $\mu\text{g}/\text{ml}$ amphotericin B. The cultures were maintained at a density of 5 to 10×10^5 cells/ml at 37°C in a humidified atmosphere of 5% CO_2 in air. Membrane extraction was modified from previously reported protocol (Wong et al., 1993). Cells were collected by centrifugation at 1000g for 2 min and washed twice with Hanks' balanced salt solution (HBSS) containing 5 mM CaCl_2 , 5 mM MgCl_2 , and protease inhibitor cocktail (Roche Diagnostics, Mannheim, Germany). The pellet was resuspended with HBSS (without CaCl_2 or MgCl_2) containing protease inhibitors and sonicated for 20-s bursts on ice five times at setting 4. The homogenate was centrifuged at 1000g for 10 min at 4°C to remove unbroken cells. The supernatant was then centrifuged at 150,000g for 45 min at 4°C . The pellet represents the membrane fraction and was resuspended in binding buffer consisting of 10 mM HEPES, 5 mM CaCl_2 , 5 mM MgCl_2 , and 5 mM EDTA, pH 7.4.

Bovine Coronary Artery Homogenate Preparation

Bovine coronary arteries (BCAs) were dissected as described above, cut into small pieces, and homogenized with a glass homogenizer in binding buffer containing miconazole (10 μM ; Sigma-Aldrich Co.) and protease inhibitor cocktail (Roche) at 4°C . Unbroken tissue and cell debris were removed by centrifugation at 6000g for 5 min at 4°C . The supernatant represents the BCA homogenate that contains membrane and cytosolic fractions.

Tissue Incubation of 14,15-EET- P^{125}ISA

14,15-EET- P^{125}ISA was incubated and extracted using the following protocols: 1) 14,15-EET- P^{125}ISA (30–33 nM) was incubated with 500 μg of BCA homogenate in binding buffer for 1 h at 30°C . The metabolites were extracted by ethyl acetate. In some incubations, 1 μM 1-cyclohexyl-3-dodecylurea (CDU), an sEH inhibitor (Morisseau et al., 2002), was included. 2) 14,15-EET-PISA (10 μM), with tracer amounts of 14,15-EET- P^{125}ISA , was incubated with BCA rings (1 g) in Krebs' buffer bubbled with 95% O_2 -5% CO_2 for 15 min at 37°C . The metabolites were harvested by solid-phase extraction using the BondElut C_{18} octadecylsilica column (Varian Inc., Harbor City, CA) as described previously (Nithipatikom et al., 2001). 3) 14,15-EET- P^{125}ISA (35 nM) was incubated with U937 cells (800,000 cells) in 200 μl of HBSS for 15 min at 37°C . The metabolites were extracted by liquid-liquid extraction using 500 μl of chloroform/methanol (1:2) followed by the addition of 167 μl of chloroform and 150 μl of H_2O . The organic phase was collected and dried under N_2 . 4) 14,15-EET- P^{125}ISA (1 nM) was incubated with 50 μg of U937 cell membrane proteins in 200 μl of binding buffer for 15 min at 4°C . In some studies, incubations were performed with buffer alone or boiled membranes. The metabolites were extracted by the same liquid-liquid extraction protocol as for intact U937 cell incubations. The extracts were then analyzed by reverse-phase HPLC as described above. The column effluent was collected in 0.5-ml fractions, and radioactivity was measured by a Packard Cobra-II Auto-Gamma counter (PerkinElmer Life and Analytical Sciences, Boston, MA).

Mass Spectrometry Analysis

To identify the 14,15-EET-PISA metabolite, 14,15-EET-PISA was incubated with BCA homogenate and extracted as described above. The HPLC effluent (21 min) containing the 14,15-EET-PISA metabolite was collected, extracted with cyclohexane/ethyl acetate (50:50), and further analyzed by mass spectrometry (MS). The MS analysis used an Agilent 1100 MSD mass spectrometer (Agilent Technologies, Palo Alto, CA) with electrospray ionization. The residues were introduced into the electrospray chamber by flow injection. Solvent A was water, and solvent B was acetonitrile with both solvents containing

0.005% glacial acetic acid. Fifty percent of each solvent was used at a flow rate of 0.2 ml/min, and data were collected for 3 min. Detection was in the negative ion mode. Results are expressed as relative abundance, with the most abundant ion set as 100%.

Radioligand Binding

U937 membranes (50 μ g) were incubated with 14,15-EET- P^{125} ISA in the presence of 20 μ M 14,15-EEZE-mSA (nonspecific binding) or an equivalent amount of ethanol (total binding) at 4°C with shaking. For the time course study, 1 nM 14,15-EET- P^{125} ISA was incubated at various times (1–30 min). For the concentration-dependent study, increasing concentrations of 14,15-EET- P^{125} ISA (0.1–150 nM) were incubated for 15 min. Incubations (total volume, 200 μ l) were performed in binding buffer and were terminated by filtration through GF/A glass filter paper on a Brandel harvester (Brandel Inc., Gaithersburg, MD). Filters were washed four times with 5 ml of ice-cold binding buffer and counted by a Packard Cobra-II Auto-Gamma counter. Specific binding was defined as total binding minus nonspecific binding. The K_d (equilibrium dissociation constant) and B_{max} (maximal binding site density) values are determined from saturation studies using nonlinear regression to fit the data to the single-site binding equation using Prism software (GraphPad Inc., San Diego, LA).

Statistical Analysis

Data are expressed as means \pm S.E.M. Statistical evaluation was performed by a one-way analysis of variance followed by the Student-Newman-Keuls multiple comparison test when significant differences were present. $P < 0.05$ was considered statistically significant.

Results

Previously, we characterized 14,15-EET-methylsulfonamide (14,15-EET-mSA; structure shown in Fig. 2) as a full agonist of 14,15-EET (Falck et al., 2003a). The *N*-acetylmethylsulfonamide group has a similar pK_a as the carboxyl group (Backes and Ellman, 1994) and is a commonly used substitute for a carboxyl group (Imig et al., 2001; Snyder et al., 2002). Based on this knowledge, we developed 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA, which contain an iodine, biotin, and BZDC group, respectively, attaching to the sulfonamide group through a benzyl linker (Fig. 2).

In bovine coronary arteries precontracted with U46619, 14,15-EET causes concentration-dependent relaxation with an EC_{50} of 1 μ M and a maximum relaxation averaging 80 to 94% at 10 μ M. Likewise, 14,15-EET-PISA, 14,15-EET-BSA,




Fig. 2. Structures of 14,15-EET and 14,15-EET-sulfonamides.

and 14,15-EET-BZDC-SA relaxed these arteries with an EC_{50} of 1 μ M and maximal relaxation of 84.5 ± 7.5 , 89.6 ± 3.9 , and $92.9 \pm 5.0\%$ at 10 μ M, respectively (Fig. 3A). We tested the effect of K^+ channel inhibition on these analog-mediated relaxations. Increasing extracellular K^+ from 4 to 20 mM eliminated relaxations to 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA (maximum relaxation = 7.8 ± 12.8 , 8.6 ± 6.3 , and $4.5 \pm 3.2\%$, respectively, Fig. 3, B–D). The BK_{Ca} channel blocker, IBTX (100 nM), markedly attenuated the 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA-induced relaxations (maximum relaxation =




Fig. 3. Effects of 14,15-EET and 14,15-EET-sulfonamides on vascular tone of U46619 precontracted bovine coronary arteries. A, relaxations to 14,15-EET, 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA. B to D, effects of the BK_{Ca} channel inhibitor, IBTX (100 nM); the EET antagonist, 14,15-EEZE-mSA (10 μ M); or increased extracellular K^+ (20 mM) on 14,15-EET-PISA (B), 14,15-EET-BSA (C), and 14,15-EET-BZDC-SA (D) relaxations. *, significantly different from control, $P < 0.05$.

39.3 ± 12.8 , 20.8 ± 2.6 , and $17.3 \pm 6.1\%$, respectively, Fig. 3, B–D). We next examined the effect of 14,15-EEZE-mSA, a 14,15-EET antagonist (Gauthier et al., 2003), on analog-mediated relaxation. Pretreatment with 14,15-EEZE-mSA ($10 \mu\text{M}$) inhibited 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA relaxations (maximal relaxation = 23.5 ± 9.1 , 21.4 ± 13.1 , and $34.7 \pm 8.6\%$, Fig. 3, B–D). These results demonstrate that 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA caused vascular relaxation by mechanisms similar to 14,15-EET, i.e., they cause relaxation by activating BK_{Ca} channels, and the relaxations are inhibited by the 14,15-EET antagonist.

14,15-EET is metabolized through three major pathways including β -oxidation, epoxide hydration, and phospholipid esterification. 14,15-EET-mSA, the parent compound of 14,15-EET-PISA, does not undergo β -oxidation or incorporation into membrane lipids (Snyder et al., 2002). It is reasonable to predict that 14,15-EET-mSA derived analogs are likewise not metabolized by these pathways. However, the 14,15-epoxide group is susceptible to hydration. We used 14,15-EET- P^{125}ISA as a radioactive tracer to study its metabolism in BCA homogenates. 14,15-EET- P^{125}ISA was incubated with buffer or BCA homogenate in the presence and absence of $1 \mu\text{M}$ CDU for 1 h at 30°C . The metabolites were extracted and resolved on reverse-phase HPLC. With buffer, 94.8% of radioactivity was extracted into the organic phase. The HPLC chromatogram showed one single peak at 29 min (peak 29), which comigrates with the synthetic 14,15-EET-PISA standard (Fig. 4A). After 30-min incubation with BCA homogenate at 30°C , an additional more polar metabolite migrating at 21 min (peak 21) was produced. The radioactivity ratio of peak 21 to peak 29 was 1.27 (Fig. 4B). In the

presence of $1 \mu\text{M}$ CDU, an sEH inhibitor, the production of peak 21 was inhibited. The peak 21/peak 29 ratio was reduced to 0.22 (Fig. 4C). Similar metabolism was observed with 90 min incubation and similar inhibition was obtained using another sEH inhibitor, adamantyl dodecanoic acid urea ($1 \mu\text{M}$) (data not shown). This suggests that peak 21 is likely the sEH metabolite of 14,15-EET-PISA, 14,15-DHET-PISA. To identify this metabolite, peak 21 was collected, re-extracted, and analyzed by negative electrospray ionization MS. The most abundant ion in the mass spectrum of this peak was at m/z 602.1 (M-H) (Fig. 4E). This indicated a molecular weight of 603, which is consistent with the molecular weight of 14,15-DHET-PISA. Thus, peak 21 was identified as 14,15-DHET-PISA. We then tested 14,15-EET- P^{125}ISA metabolism by bovine coronary arterial rings using conditions under which the vascular activities were tested. After a 15-min incubation at 37°C , the majority of radioactivity eluted at 29 min, and a small portion of radioactivity was converted into peak 21. The peak 21/peak 29 ratio was 0.18 (Fig. 4D). These data indicate that 14,15-EET-PISA is hydrolyzed to 14,15-DHET-PISA by sEH in both BCA homogenate and arterial rings. The arterial rings showed less conversion. There was no evidence of the metabolism of 14,15-EET-PISA by β -oxidation.

Similar results were obtained when 14,15-EET- P^{125}ISA was incubated with U937 cells for 15 min at 37°C (Fig. 5A). The major metabolite eluted at 29 min, and a small radioactive product eluted at 21 min. Only the 29-min radioactive metabolite was observed when the cells were incubated at 4°C (data not shown). Thus, U937 cells metabolized 14,15-EET-PISA to 14,15-DHET-PISA, as did coronary arteries. Next, we investigated the stability of 14,15-EET- P^{125}ISA




Fig. 4. Metabolism of 14,15-EET- P^{125}ISA by BCA homogenate and arterial rings. A to C, HPLC chromatograms of extracted 14,15-EET- P^{125}ISA metabolites after incubation with buffer, BCA homogenate ($500 \mu\text{g}$) in the presence or absence of CDU ($1 \mu\text{M}$) for 30 min at 30°C , respectively. D, HPLC chromatogram of extracted 14,15-EET- P^{125}ISA metabolites after incubation with arterial rings for 15 min at 37°C . E, Negative ion electrospray ionization MS spectrum of 14,15-EET-PISA metabolite (peak 21).




Fig. 5. Metabolism of 14,15-EET- P^{125} ISA by U937 cells and U937 membrane protein. A, 14,15-EET- P^{125} ISA (35 nM) was incubated with intact U937 cells (800,000 cells) for 15 min at 37°C. B to D, 14,15-EET- P^{125} ISA (1 nM) was incubated with buffer (B), boiled U937 membrane protein (C), or U937 membrane protein (50 μ g, D) for 15 min at 4°C. HPLC chromatograms of extracted 14,15-EET- P^{125} ISA metabolites from the incubations were shown.

under the experimental conditions of radioligand binding. After incubation with U937 membranes at 4°C for 15 min, the majority of the radioactivity was extracted with an organic solvent, and the extraction efficiency was $93.6 \pm 1.0\%$ ($n = 3$). When 14,15-EET- P^{125} ISA was incubated with boiled and control U937 membranes, the percentage of radioactivity in the organic phase was 95.8 ± 0.1 and $95.6 \pm 0.2\%$, respectively ($n = 3$). The HPLC chromatograms showed a single peak eluting at 29 min (Fig. 5, B–D). These results indicate that 14,15-EET- P^{125} ISA is metabolically stable under the conditions for radioligand binding.

To determine whether 14,15-EET- P^{125} ISA is a radioligand useful for the characterization of 14,15-EET binding, we used membrane fractions from U937 cells, a cell line that was reported to contain an EET binding site (Wong et al., 1997). To test whether 14,15-EET- P^{125} ISA binds to U937 membrane in a time-dependent manner, 1 nM 14,15-EET- P^{125} ISA was incubated with U937 membranes (50 μ g of protein) at 4°C for 1 to 30 min. The binding of 14,15-EET- P^{125} ISA to U937 membranes was rapid and specific. Equilibrium was reached in 15 min and




Fig. 6. 14,15-EET- P^{125} ISA binding to U937 membrane protein. A, time course of 14,15-EET- P^{125} ISA binding. 14,15-EET- P^{125} ISA (1 nM) was incubated with 50 μ g of protein at 4°C for various time (1–30 min). $n = 2$. B, concentration-dependent 14,15-EET- P^{125} ISA binding. Increasing concentrations of 14,15-EET- P^{125} ISA (0.1–150 nM) were incubated with 50 μ g of protein at 4°C for 15 min. $n = 2$ –7. C, 14,15-DHET- P^{125} ISA binding to U937 membranes. Increasing concentrations of 14,15-DHET- P^{125} ISA (2.5–100 nM) were incubated with 50 μ g of protein at 4°C for 15 min. $n = 2$ –3.

remained stable until 30 min (Fig. 6A). Then, we tested the saturability of the specific 14,15-EET- P^{125} ISA binding. Various concentrations of 14,15-EET- P^{125} ISA (0.1–150 nM) were incubated with U937 membrane (50 μ g of protein) at 4°C for 15 min. 14,15-EET- P^{125} ISA-specific binding was concentration-dependent and reached an apparent plateau at 100 nM (Fig. 6B). The data were fit to a one-site saturation model, and the predicted K_d and B_{max} were 148.3 ± 36.4 nM and 3.3 ± 0.5 pmol/mg protein, respectively. To confirm the specificity of this binding site to 14,15-EET-PISA, the binding of 14,15-DHET- P^{125} ISA to U937 membranes was determined. Specific binding of 14,15-DHET- P^{125} ISA to U937 membranes was very low with a maximum binding of 0.3 pmol/mg protein (Fig. 6C). The data poorly fit with saturation models. This suggests that 14,15-DHET-PISA does not specifically bind to U937 membranes. Therefore, the 14,15-epoxide group is required for 14,15-EET- P^{125} ISA binding to U937 membranes. Taken together, binding of 14,15-EET- P^{125} ISA to U937 membranes is specific, time-dependent, concentration-dependent, and saturable.

Discussion

Our present study developed and characterized a series of stable 14,15-EET-mSA derivatives, 14,15-EET-PISA, 14,15-

EET-BSA, and 14,15-EET-BZDC-SA, as tools to study EET transport, metabolism, and binding. The analogs caused concentration-dependent relaxation of bovine coronary arteries and retained full agonist potency and activity compared with 14,15-EET. Attachment of a bulky group at the C1 of 14,15-EET did not alter its vascular activity. The relaxations to the analogs were blocked by inhibition of K^+ efflux by 20 mM extracellular K^+ and BK_{Ca} channel blockade by iberiotoxin. These data suggest that these agonists, like 14,15-EET, mediate vascular relaxation by activation of BK_{Ca} channels and membrane hyperpolarization. 14,15-EEZE-mSA, a known 14,15-EET antagonist, inhibited the relaxations (Gauthier et al., 2003). It is likely that these agonists bind to the same binding site or receptor as 14,15-EET and act through the similar cellular mechanisms as 14,15-EET.

EET-mediated activation of vascular smooth muscle BK_{Ca} channels requires active $G\alpha$ protein and may involve ADP-ribosylation of $G\alpha$ (Li and Campbell, 1997; Li et al., 1999). Likewise, $G\alpha$ is required for 11,12-EET-mediated activation of BK_{Ca} channels expressed in human embryonic kidney 293 cells (Fukao et al., 2001). Because G proteins characteristically couple to membrane receptors, it is likely that EETs initiate their signaling events by binding to a G protein-coupled receptor. Node et al. (2001) showed that 11,12-EET stimulated GTP-binding activity of $G\alpha$ in human saphenous vein endothelial cells and activated endothelial t-PA gene expression through a cAMP-dependent mechanism. Taken together, this suggests that EET may mediate their cellular events through a G protein-coupled receptor signaling pathway. However, characterization and purification of such a membrane receptor/binding site have been limited due to lack of proper tools. In addition, the mechanisms of EETs uptake and secretion by cells are unclear. Many eicosanoids such as prostaglandins (Kanai et al., 1995), leukotrienes (Leier et al., 1994), and arachidonic acid (Krischer et al., 1998) have been described to enter and leave cells through transporter-facilitated diffusion. It is unknown whether a similar transporter is present for EETs.

We designed 14,15-EET-mSA derivatives as tools to study EET transport, metabolism, and binding. Each agonist contains a functional group, i.e., a biotin, a photolinking reagent, or a radioiodinated group. With aid of convenient avidin conjugates such as agarose beads and fluorophore conjugates, 14,15-EET-BSA can be used to pull down an EET binding protein or image 14,15-EET binding sites. Upon ultraviolet radiation, 14,15-EET-BZDC-SA cross-links with molecules in close proximity. This reaction may add a tag to the target molecule, which may be detected by mass spectrometric analysis. 14,15-EET-PISA can be radiolabeled with [^{125}I] to provide a new radioactive 14,15-EET agonist. Radioactive EET analogs are valuable tools to study their metabolism and binding.

Currently, radiolabeled 14,15-EET is made by epoxidation of ^{14}C - or 3H -labeled arachidonic acid, which results in all four EET regioisomers and less than 25% production of 14,15-EET (Corey et al., 1979; Rosolowsky and Campbell, 1993). In contrast, radioiodination of 14,15-EET- $P^{125}ISA$ from 14,15-EET-stannane is specific and efficient. In addition, [^{125}I] has higher specific activity than [3H], which results in increased sensitivity for radioligand binding assays. Therefore, 14,15-EET- $P^{125}ISA$ is a radioactive 14,15-EET

agonist with high specific activity, low production cost, and efficient yields.

β -Oxidation and phospholipid esterification of 14,15-EET require the C-1 carboxyl group. Other routes of 14,15-EET metabolism include sEH hydration, cytochrome P450 ω -hydroxylation (Cowart et al., 2002), and glutathione conjugation (Spearman et al., 1985) and are independent of the C-1 carboxyl group. Therefore, 14,15-EET- $P^{125}ISA$ is a useful tool to study C1-carboxyl-independent metabolism of 14,15-EET. Like 14,15-EET, 14,15-EET-PISA is hydrolyzed into its corresponding 14,15-DHET-PISA. This metabolism is effectively inhibited by sEH inhibitors. Among the tested tissues, the hydration was most active in BCA homogenate where sEH is exposed, markedly less active in BCA rings and U937 cells where sEH is intracellular and less accessible, and absent in U937 membrane fractions where sEH is not localized. These data indicate that 14,15-EET-PISA is actively metabolized into 14,15-DHET-PISA by sEH.

Previously in the human leukemic monocyte lymphoma cell line U937 cells, a 14,15-EET binding site was characterized by whole-cell radioligand binding using [3H]14,15-EET. The K_d and B_{max} were 13.84 ± 2.58 nM and 3.54 ± 0.28 pmol/ 10^6 cells, respectively (Wong et al., 1997). We used U937 cell membrane fraction to test 14,15-EET- $P^{125}ISA$ binding. 14,15-EET- $P^{125}ISA$ remained stable and unchanged when it was incubated with U937 membrane proteins at 4°C for 15 min. In agreement with the [3H]14,15-EET whole-cell binding data, 14,15-EET- $P^{125}ISA$ bound to U937 membranes in a time-dependent, concentration-dependent, and saturable manner. Interestingly, the binding of 14,15-EET- $P^{125}ISA$ to U937 membranes was with a lower affinity and lower receptor density. 14,15-EET- $P^{125}ISA$ binding to U937 membranes reached equilibrium by 15 min and remained unchanged to 30 min at 4°C. The K_d was 148.3 ± 36.4 nM, which was 10-fold higher than that observed with [3H]14,15-EET in whole cells. The B_{max} of 14,15-EET- $P^{125}ISA$ binding to U937 membranes was 3.3 ± 0.5 pmol/mg protein, which was lower than that observed in whole cells. The reason for these discrepant data could be due to esterification of [3H]14,15-EET into phospholipids or conjugation to cellular components. This could contribute to the observed binding in whole cells, whereas these processes are absent or less active in membrane fractions. It is also possible that intracellular components maintain the binding sites in a conformation that favors binding to the ligand. This intracellular factor is missing in extracted membranes, which may change the receptor conformation to a ligand-unfavorable form. In other words, a smaller portion of the receptors in the membrane fractions than whole cell is available for the ligand, which results in a decreased receptor density. In addition, ligands differ in affinity. The addition of the sulfonamide group to C-1 may reduce the affinity for the binding site. The variability between U937 cell lines could also account for the discrepancies. The 14,15-EET- $P^{125}ISA$ binding requires the 14,15-epoxide because its hydrolyzed product, 14,15-DHET- $P^{125}ISA$, did not bind to U937 membranes. Taken together, our binding data validate the use of 14,15-EET- $P^{125}ISA$ as a radiolabeled tool to study 14,15-EET binding.

EETs are key regulators of vascular tone and active mediators in pathophysiologic conditions such as anti-inflammation, fibrinolysis, angiogenesis, and ischemic reperfusion protection (Seubert et al., 2004; Nithipatikom et al., 2006).

However, whether they initiate their signaling through binding to specific protein(s)/receptor(s) is unclear. 14,15-EET-PISA, 14,15-EET-BSA, and 14,15-EET-BZDC-SA are 14,15-EET agonists that serve as tools to identify and purify EET binding protein(s). 14,15-EET-P¹²⁵ISA, in particular, is a radioactive tracer to study EET metabolism and a radioligand to characterize EET binding kinetics.

Acknowledgments

We thank Dr. Bruce Hammock for the kind gift of recombinant sEH and sEH inhibitors, Dr. Kathy Gauthier for review of the manuscript, and Gretchen Barg for secretarial assistance.

References

- Backes BJ and Ellman JA (1994) Carbon-carbon bond-forming methods on solid support: utilization of Kenner's "safety-catch" linker. *J Am Chem Soc* **116**:11171–11172.
- Campbell WB, Gebremedhin D, Pratt PF, and Harder DR (1996) Identification of epoxyeicosatrienoic acids as endothelium-derived hyperpolarizing factors. *Circ Res* **78**:415–423.
- Capdevila J, Parkhill L, Chacos N, Okita R, Masters BS, and Estabrook RW (1981) The oxidative metabolism of arachidonic acid by purified cytochromes P-450. *Biochem Biophys Res Commun* **101**:1357–1363.
- Corey EJ, Niwa H, and Falck JR (1979) Selective epoxidation of eicosa-cis-5,8,11,14-tetraenoic (arachidonic) acid and eicosa-cis-8,11,14-trienoic acid. *J Am Chem Soc* **101**:586–587.
- Cowart LA, Wei S, Hsu MH, Johnson EF, Krishna MU, Falck JR, and Capdevila JH (2002) The CYP4A isoforms hydroxylate epoxyeicosatrienoic acids to form high affinity peroxisome proliferator-activated receptor ligands. *J Biol Chem* **277**:35105–35112.
- Falck JR, Krishna UM, Reddy YK, Kumar PS, Reddy KM, Hittner SB, Deeter C, Sharma KK, Gauthier KM, and Campbell WB (2003a) Comparison of vasodilatory properties of 14,15-EET analogs: structural requirements for dilation. *Am J Physiol* **284**:H337–H349.
- Falck JR, Reddy LM, Reddy YK, Bondlela M, Krishna UM, Ji Y, Sun J, and Liao JK (2003b) 11,12-epoxyeicosatrienoic acid (11,12-EET): structural determinants for inhibition of TNF- α -induced VCAM-1 expression. *Bioorg Med Chem Lett* **13**:4011–4014.
- Fisslthaler B, Popp R, Kiss L, Potente M, Harder DR, Fleming I, and Busse R (1999) Cytochrome P450 2C is an EDHF synthase in coronary arteries. *Nature (Lond)* **401**:493–497.
- Fukao M, Mason HS, Kenyon JL, Horowitz B, and Keef KD (2001) Regulation of BK(Ca) channels expressed in human embryonic kidney 293 cells by epoxyeicosatrienoic acid. *Mol Pharmacol* **59**:16–23.
- Fulton D, Mahboubi K, McGiff JC, and Quilley J (1995) Cytochrome P450-dependent effects of bradykinin in the rat heart. *Br J Pharmacol* **114**:99–102.
- Gauthier KM, Edwards EM, Falck JR, Reddy DS, and Campbell WB (2005) 14,15-epoxyeicosatrienoic acid represents a transferable endothelium-dependent relaxing factor in bovine coronary arteries. *Hypertension* **45**:666–671.
- Gauthier KM, Falck JR, Reddy LM, and Campbell WB (2004) 14,15-EET analogs: characterization of structural requirements for agonist and antagonist activity in bovine coronary arteries. *Pharmacol Res* **49**:515–524.
- Gauthier KM, Jagadeesh SG, Falck JR, and Campbell WB (2003) 14,15-epoxyeicosatrienoic acid (14,15-EET) is a 14,15- and 5,6-EET antagonist in bovine coronary arteries. *Hypertension* **42**:555–561.
- Huang A, Sun D, Jacobson A, Carroll MA, Falck JR, and Kaley G (2005) Epoxyeicosatrienoic acids are released to mediate shear stress-dependent hyperpolarization of arteriolar smooth muscle. *Circ Res* **96**:376–383.
- Imig JD, Falck JR, Wei S, and Capdevila JH (2001) Epoxygenase metabolites contribute to nitric oxide-independent afferent arteriolar vasodilation in response to bradykinin. *J Vasc Res* **38**:247–255.
- Kanai N, Lu R, Satriano JA, Bao Y, Wolkoff AW, and Schuster VL (1995) Identification and characterization of a prostaglandin transporter. *Science (Wash DC)* **268**:866–869.
- Kort MD, Luijendijk J, Marel GAVd, and Boom JHV (2000) Synthesis of photoaffinity derivatives of adenophostin A. *Eur J Org Chem* **2000**:3085–3092.
- Kozak W, Kluger MJ, Kozak A, Wachulec M, and Dokladny K (2000) Role of cytochrome P-450 in endogenous antipyreresis. *Am J Physiol* **279**:R455–R460.
- Krischer SM, Eisenmann M, and Mueller MJ (1998) Transport of arachidonic acid across the neutrophil plasma membrane via a protein-facilitated mechanism. *Biochemistry* **37**:12884–12891.
- Leier I, Jedlitschky G, Buchholz U, Cole SP, Deeley RG, and Keppler D (1994) The MRP gene encodes an ATP-dependent export pump for leukotriene C4 and structurally related conjugates. *J Biol Chem* **269**:27807–27810.
- Li PL and Campbell WB (1997) Epoxyeicosatrienoic acids activate K⁺ channels in coronary smooth muscle through a guanine nucleotide binding protein. *Circ Res* **80**:877–884.
- Li PL, Chen CL, Bortell R, and Campbell WB (1999) 11,12-Epoxyeicosatrienoic acid stimulates endogenous mono-ADP-ribosylation in bovine coronary arterial smooth muscle. *Circ Res* **85**:349–356.
- Michaelis UR and Fleming I (2006) From endothelium-derived hyperpolarizing factor (EDHF) to angiogenesis: epoxyeicosatrienoic acids (EETs) and cell signaling. *Pharmacol Ther* **111**:584–595.
- Morisseau C, Goodrow MH, Newman JW, Wheelock CE, Dowdy DL, and Hammock BD (2002) Structural refinement of inhibitors of urea-based soluble epoxide hydrolases. *Biochem Pharmacol* **63**:1599–1608.
- Nithipatikom K, Grall AJ, Holmes BB, Harder DR, Falck JR, and Campbell WB (2001) Liquid chromatographic-electrospray ionization-mass spectrometric analysis of cytochrome P450 metabolites of arachidonic acid. *Anal Biochem* **298**:327–336.
- Nithipatikom K, Moore JM, Isbell MA, Falck JR, and Gross GJ (2006) Epoxyeicosatrienoic acids in cardioprotection: ischemic versus reperfusion injury. *Am J Physiol* **291**:H537–H542.
- Node K, Huo Y, Ruan X, Yang B, Spiecker M, Ley K, Zeldin DC, and Liao JK (1999) Anti-inflammatory properties of cytochrome P450 epoxygenase-derived eicosanoids. *Science (Wash DC)* **285**:1276–1279.
- Node K, Ruan XL, Dai J, Yang SX, Graham L, Zeldin DC, and Liao JK (2001) Activation of G α s mediates induction of tissue-type plasminogen activator gene transcription by epoxyeicosatrienoic acids. *J Biol Chem* **276**:15983–15989.
- Pinto A, Abraham NG, and Mullane KM (1987) Arachidonic acid-induced endothelial-dependent relaxations of canine coronary arteries: contribution of a cytochrome P-450-dependent pathway. *J Pharmacol Exp Ther* **240**:856–863.
- Rosolowsky M and Campbell WB (1993) Role of PGI₂ and epoxyeicosatrienoic acids in relaxation of bovine coronary arteries to arachidonic acid. *Am J Physiol* **264**:H327–H335.
- Rosolowsky M and Campbell WB (1996) Synthesis of hydroxyeicosatetraenoic (HETEs) and epoxyeicosatrienoic acids (EETs) by cultured bovine coronary artery endothelial cells. *Biochim Biophys Acta* **1299**:267–277.
- Seubert J, Yang B, Bradbury JA, Graves J, Degraff LM, Gabel S, Gooch R, Foley J, Newman J, Mao L, et al. (2004) Enhanced postischemic functional recovery in CYP2J2 transgenic hearts involves mitochondrial ATP-sensitive K⁺ channels and p42/p44 MAPK pathway. *Circ Res* **95**:506–514.
- Snyder GD, Krishna UM, Falck JR, and Spector AA (2002) Evidence for a membrane site of action for 14,15-EET on expression of aromatase in vascular smooth muscle. *Am J Physiol* **283**:H1936–H1942.
- Spearman ME, Prough RA, Estabrook RW, Falck JR, Manna S, Leibman KC, Murphy RC, and Capdevila J (1985) Novel glutathione conjugates formed from epoxyeicosatrienoic acids (EETs). *Arch Biochem Biophys* **242**:225–230.
- Spector AA, Fang X, Snyder GD, and Weintraub NL (2004) Epoxyeicosatrienoic acids (EETs): metabolism and biochemical function. *Prog Lipid Res* **43**:55–90.
- Wong PY, Lai PS, and Falck JR (2000) Mechanism and signal transduction of 14 (R), 15 (S)-epoxyeicosatrienoic acid (14,15-EET) binding in guinea pig monocytes. *Prostaglandins Other Lipid Mediat* **62**:321–333.
- Wong PY, Lai PS, Shen SY, Belosludtsev YY, and Falck JR (1997) Post-receptor signal transduction and regulation of 14(R),15(S)-epoxyeicosatrienoic acid (14,15-EET) binding in U-937 cells. *J Lipid Mediat Cell Signal* **16**:155–169.
- Wong PY, Lin KT, Yan YT, Ahern D, Iles J, Shen SY, Bhatt RK, and Falck JR (1993) 14(R),15(S)-epoxyeicosatrienoic acid (14(R),15(S)-EET) receptor in guinea pig mononuclear cell membranes. *J Lipid Mediat* **6**:199–208.
- Yang W, Gauthier KM, Reddy LM, Sangras B, Sharma KK, Nithipatikom K, Falck JR, and Campbell WB (2005) Stable 5,6-epoxyeicosatrienoic acid analog relaxes coronary arteries through potassium channel activation. *Hypertension* **45**:681–686.

Address correspondence to: Dr. William B. Campbell, Department of Pharmacology and Toxicology, Medical College of Wisconsin, 8701 Watertown Plank Road, Milwaukee, WI 53226. E-mail: wbcamp@mcw.edu