

## Isopentenyl Diphosphate Isomerase. Mechanism-Based Inhibition by Diene Analogues of Isopentenyl Diphosphate and Dimethylallyl Diphosphate

Zheng Wu,<sup>‡</sup> Johan Wouters,<sup>§</sup> and C. Dale Poulter<sup>\*,‡</sup>

Contribution from the Department of Chemistry, University of Utah, Salt Lake City, UT 84112, Laboratoire de Chimie Biologique Structurale, 61 Rue de Bruxelles, Namur, Belgium, and Institut de Recherches Wiame, Campus Ceria, Bruxelles, Belgium

Received September 8, 2005; E-mail: poulter@chemistry.utah.edu

Abstract: Isopentenyl diphosphate isomerase (IDI) catalyzes the interconversion of isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP). This is an essential step in the mevalonate entry into the isoprenoid biosynthetic pathway. The isomerization catalyzed by type I IDI involves protonation of the carbon-carbon double bond in IPP or DMAPP to form a tertiary carbocation, followed by deprotonation. Diene analogues for DMAPP (E-2-OPP and Z-2-OPP) and IPP (4-OPP) were synthesized and found to be potent active-site-directed irreversible inhibitors of the enzyme. X-ray analysis of the E-I complex between Escherichia coli IDI and 4-OPP reveals the presence of two isomers that differ in the stereochemistry of the newly formed C3-C4 double bond in the hydrocarbon chain of the inhibitor. In both adducts C5 of the inhibitor is joined to the sulfur of C67. In these structures the methyl group formed upon protonation of the diene moiety in 4-OPP is located near E116, implicating that residue in the protonation step.

The isomerization of isopentenyl diphosphate (IPP) to dimethylallyl diphosphate (DMAPP) is a crucial activation step in isoprenoid biosynthesis, during which IPP is converted to its highly electrophilic isomer. These two molecules are the building blocks used to construct over 35 000 isoprenoid compounds found in nature. In Archaea, Eukaryota, and some Bacteria, IPP is synthesized from acetyl-CoA by the mevalonate (MVA) pathway and is the exclusive product from the phosphorylation and decarboxylation of the key intermediate mevalonic acid.<sup>1,2</sup> IPP is required for biosynthesis of DMAPP in the MVA pathway, and IPP isomerase is essential.<sup>3</sup> In most bacteria and plant chloroplasts, IPP and DMAPP are synthesized from pyruvate and D-glyceraldehyde phosphate by the methylerythritol phosphate (MEP) pathway.<sup>4</sup> The final step produces both compounds during the reduction of hydroxydimethylallyl diphosphate. IPP isomerase activity is often found in organisms that utilize the MEP pathway, but the enzyme is not essential.

Two convergently evolved IPP isomerases are known.<sup>5,6</sup> The type I IPP isomerase was discovered in the late 1950s<sup>7</sup> and is found in Eukaryota and Bacteria. The enzyme requires a divalent metal and catalyzes the antarafacial isomerization of IPP and DMAPP<sup>8</sup> by a protonation-deprotonation mechanism<sup>9</sup> (see

Scheme 1. Protonation-Deprotonation Mechanism for Isomerization of IPP and DMAPP

$$PPO \xrightarrow{H}_{H} \xrightarrow{H^{*}}_{HB^{*}} PPO \xrightarrow{H}_{H} \xrightarrow{B}_{H} \xrightarrow{H^{*}}_{H} PPO \xrightarrow{H}_{H} \xrightarrow{H}_{H} \xrightarrow{H^{*}}_{H} PPO \xrightarrow{H}_{H} \xrightarrow{H}_{H} \xrightarrow{H}_{H} PPO \xrightarrow{H}_{H} \xrightarrow{H}_{H}$$

Scheme 1). During the reaction, the pro-R hydrogen (blue) is removed from C(2) of IPP, and a proton from solvent (red) is added to the re-face at C4 of the double bond to generate the E-methyl group at C(3).

Type II IPP isomerase is found in Archaea and in Bacteria.<sup>10</sup> Typically, a given bacterium has only one form of IPP isomerase, but, in at least one instance, the type I and type II enzymes coexist.<sup>11,12</sup> Type II IPP isomerase requires reduced flavin and divalent metal cofactors for activity. The stereochemistry of the reaction is similar to that catalyzed by the type I enzyme,<sup>13,14</sup> although the facial selectivity for proton addition at C(4) has not yet been determined.

- (5) Anderson, M. S.; Muehlbacher, M.; Street, I. P.; Proffitt, J.; Poulter, C. D. J. Biol. Chem. 1989, 264, 19169-1917
- (6) Kandea, K.; Kuzuyama, T.; Takagi, M.; Hayakawa, Y.; Seto, H. Proc. Natl. Acad. Sci. U.S.A. 2001, 98, 932-937. (7)
- Agranoff, B. W.; Eggerer, H.; Henning, U.; Lynen, F. J. Am. Chem. Soc. **1959**, 81, 1254–55. (8) Clifford, K.; Cornforth, J. W.; Mallaby, R.; Philips, R. T. Chem. Commun.
- 1971, 1599-1600.
- (9) Muehlbacher, M.; Poulter, C. D. *Biochemistry* 1988, 27, 7315–7328.
   (10) Rohdich, F.; Bacher, A.; Eisenreich, W. *Bioorg. Chem.* 2004, *32*, 292–
- 308. (11) Kuzuyama, T.; Seto, H. Nat. Prod. Rep. 2003, 20, 171-183.
- (12) Laupitz, R.; Hecht, S.; Amslinger, S.; Zepeck, F.; Kaiser, J.; Richter, G.; Schramek, N.; Steinbacher, S.; Huber, R.; Arigoni, D.; Bacher, A.; Eisenreich, W.; Rohdich, R. Eur. J. Biochem. 2004, 271, 2658-2669.
- (13) Laupitz, R.; Grawert, T.; Rieder, C.; Zepeck, F.; Bacher, A.; Arigoni, D.; Rohdich. F.; Eisenreich, W. *Chem. Biodiversity* **2004**, *1*, 1367–1376.
  (14) Barkley, S. J.; Desai, S. V.; Poulter, C. D. Org. Lett. **2004**, *6*, 5019–5021.

<sup>&</sup>lt;sup>‡</sup> University of Utah.

<sup>&</sup>lt;sup>§</sup> Laboratoire de Chimie Biologique Structurale and Institut de Recherches Wiame.

<sup>(1)</sup> Poulter, C. D.; Rilling, H. Biosynthesis of Isoprenoid Compounds; Wiley: New York, 1981. (2) Koyama, T.; Ogura, K. Comprehensive Natural Product Chemistry;

Pergmon: Oxford, 1999 (3) Mayer, M. P.; Hahn, F. M.; Stillman, D. J.; Poulter, C. D. Yeast 1992, 8,

<sup>743-748</sup> 

<sup>(4)</sup> Rodríguez-Concepción, M.; Boronat, A. Plant Physiol. 2002, 130, 1079-1089

Several lines of evidence support the protonation-deprotonation mechanism shown in Scheme 1 for type I IPP isomerase. These include proton exchange experiments,<sup>15</sup> substituent effects on the rate of the reaction,<sup>9,16</sup> tight binding of transition-state analogues,9 and irreversible inhibition by mechanism-based inhibitors.9 Covalent modification of active-site residues by the irreversible inhibitors and site-directed mutagenesis of the modified amino acids implicate two amino acids, C67 and E116 (Escherichia coli numbering), in the proton-transfer steps.<sup>17</sup> X-ray analysis of E. coli IPP isomerase shows that these amino acids are located on opposite sides of the active site.<sup>18</sup> E116 is also part of a hexacoordinate binding site for a divalent metal. In the structure of the enzyme containing 2-(N,N-dimethylamino)ethyl diphosphate, a transition-state analogue for the putative tertiary cationic intermediate where C(3) is replaced by a positively charged N-H ammonium moiety, one of the carboxylate oxygens in the E116 side chain is coordinated to the  $H-N^+$  unit and the other to the divalent metal. In another structure, where the enzyme was inactivated with the epoxide derivative of IPP, the oxirane ring had been opened to give a primary alcohol at C(4) with concomitant formation of a thioether bond between C(3) of the inhibitor and the sulfhydryl moiety of C67.<sup>17</sup> The X-ray structures of the enzyme•inhibitor complexes also contained a second divalent metal, which was coordinated to nonbridging oxygens at P(1) and P(2) of the diphosphate moiety, the side-chain carboxylate of E87, and the amide carbonyl oxygen in C67.

Replacement of the active-site cysteine in yeast IPP isomerase by serine produced a modest 2-fold increase in  $K_{\rm M}$  but reduced  $k_{\rm cat}$  by ~10<sup>4</sup>.<sup>17</sup> The related alanine mutant was inactive. Likewise, substitution of the active-site glutamate with glutamine or valine gave inactive proteins. Interestingly, an X-ray structure of the C67A mutant of *E. coli* IPP isomerase, which had been treated with an epoxy analogue of IPP, showed that the protein was covalently modified.<sup>19</sup> In this case the oxirane ring had opened to form an ester linkage between C(3) of the inhibitor and the side-chain carboxylate moiety of E116. Thus, the "inactive" protein retained the ability to activate and open the oxirane ring, suggesting a role for E116 in the protonation step of the isomerization reaction.

Diene analogues of oxidosqualene have been used to intercept carbocationic intermediates as allylic cations, which react with active-site nucleophiles and abort the cascade of cyclization reactions leading to the tetracyclic skeleton polycyclic triterpenes.<sup>20,21</sup> We thought that similar analogues of IPP and DMAPP might be potent mechanism-based inhibitors of IPP isomerase by creating an electrophilic center in the substrate analogues that is susceptible to alkylation. We now report the synthesis of two diene analogues of DMAPP and a diene analogue of IPP, their irreversible inactivation of type I IPP isomerase from *Schizosaccharomyces pombe*, and an X-ray

- (16) Reardon, J. E.; Abeles, R. H. *Biochemistry* 1986, 25, 5609–5616.
   (17) Street, I. P.; Coffman, H. R.; Baker, J. A.; Poulter, C. D *Biochemistry* 1994,
- Wouters, J.; Oudjama, Y.; Barkley, S. J.; Tricot, C.; Stalon, V.; Droogmans, L.; Poulter, C. D. *J. Biol. Chem.* **2003**, 278, 11903–11908.
   Wouters, J.; Oudiama, Y.; Stalon, V.; Droogmans, L.; Poulter, C. D.
- (19) Wouters, J.; Oudjama, Y.; Stalon, V.; Droogmans, L.; Poulter, C. D. *Proteins* **2004**, *54*, 216–221.
   (20) Xiao, X.-Y.; Prestwich, G. D. J. Am. Chem. Soc. **1991**, *113*, 9673–9674.
- (20) Xiao, X.-Y.; Prestwich, G. D. J. Am. Chem. Soc. 1991, 113, 9673–9674.
   (21) Ceruti, M.; Rocco, F.; Viola, F.; Balliano, G.; Milla, P.; Arpicco, S.; Cattel, L. J. Med. Chem. 1998, 41, 540–554.

structure of *E. coli* IPP isomerase covalently modified by the IPP analogue that provides insights about the protonation step.

## **Experimental Section**

(*E*)-3-Methyl-2,4-pentadien-1-ol (E-2-OH). A solution of 1.0 g (10.4 mmol) of (*E*)-3-methyl-2-penten-4-yn-1-ol (E-1-OH) in 10.0 mL of anhydrous ethyl alcohol was reduced using 0.148 g of Lindlar catalyst. The catalyst was removed by filtration, solvent was removed by rotary evaporation, and the resulting yellow oil was purified by flash chromatography on silica gel (3/2 hexane/diethyl ether) followed by distillation at 50–55 °C (Kugelrohr, 3 mmHg) (lit.<sup>22</sup> bp 74 °C at 14 mmHg) to give 0.801 g (77%) of a colorless oil:<sup>22,23</sup> <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>)  $\delta$  0.37 (1H, s (broad)), 1.53 (3H, s), 4.40 (2H, dd,  $J_{\rm H,H} = 6.6$  Hz), 4.96 (1H, d,  $J_{\rm H,H} = 10.5$  Hz), 5.09 (1H, dd,  $J_{\rm H,H} = 17.4$ , Hz), 5.52 (1H, d,  $J_{\rm H,H} = 6.6$  Hz), 6.33 40 (1H, dd,  $J_{\rm H,H} = 17.4$ , 10.8 Hz); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  12.2, 59.4, 112.6, 132.4, 135.8, 141.7.

(*E*)-5-Bromo-3-methylpenta-1,3-diene (E-2-Br). To a solution of 45 mg (0.42 mmol) of E-2-OH in CH<sub>2</sub>Cl<sub>2</sub> were added 182 mg (0.55 mmol) of CBr<sub>4</sub> and 144 mg (0.55 mmol) of triphenylphosphine. After 3 h at room temperature, pentane was added, the resulting suspension was filtered, and solvent was removed at reduced pressure. The residue was purified by distillation at 40 °C (Kugelrohr, 3 mmHg) to give 49 mg (62%) of a colorless oil:<sup>24</sup> <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.85 (3H, d, *J*<sub>H,H</sub> = 1.5 Hz), 4.13 (2H, d, *J*<sub>H,H</sub> = 8.4 Hz), 5.12 (1H, d, *J*<sub>H,H</sub> = 10.8 Hz), 5.30 (1H, d, *J*<sub>H,H</sub> = 17.1 Hz), 5.38 (1H, d, *J*<sub>H,H</sub> = 10.8 Hz), 5.78 (1H, dd, *J*<sub>H,H</sub> = 17.4, 10.8 Hz).

(*Z*)-3-Methyl-2,4-pentadien-1-ol (*Z*-2-OH). Following the procedure described for **E-2-OH**, 2.36 g (24.5 mmol) of (*Z*)-3-methyl-2-pentene-4-yn-1-ol (*Z*-1-OH) was reduced using 0.40 g of Lindlar catalyst to give 1.67 g (75%) of a colorless oil:<sup>22,23</sup> <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>)  $\delta$  0.76 (1H, s (broad)), 1.72 (3H, s), 4.21 (2H, t, *J*<sub>H,H</sub> = 6.6 Hz), 5.01 (1H, d, *J*<sub>H,H</sub> = 11.1 Hz), 5.13 (1H, dd, *J*<sub>H,H</sub> = 17.3, 1.0 Hz), 5.57 (1H, d, *J*<sub>H,H</sub> = 6.6 Hz), 6.68 (1H, dd, *J*<sub>H,H</sub> = 17.4, 10.8 Hz); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  20.1, 58.5, 115.2, 130.6, 134.0, 134.6.

(**Z**)-5-Chloro-3-methylpenta-1,3-diene (**Z**-2-Cl). To a solution of 0.367 g (2.75 mmol) of NCS in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> at -30 °C was added 1.86.4 mg (3.0 mmol) of dimethyl sulfide. The mixture was allowed to warm to 0 °C for 5 min and then cooled to -40 °C before 246 mg (2.51 mmol) of **Z**-2-OH was added. The suspension was allowed to warm to 0 °C and stirred for 2 h. The resulting solution was washed with saturated NaCl, solvent was removed at reduced pressure, and the resulting yellow oil was distilled at 35 °C (Kugelrohr, 4 mmHg) to give 0.175 g (60%) of a colorless oil:<sup>25 1</sup>H NMR (NMR (CDCl<sub>3</sub>)  $\delta$  1.72 (3H, s) 4.21 (2H, d,  $J_{\rm H,H} = 6.6$  Hz), 5.01 (1H, d,  $J_{\rm H,H} = 11.1$  Hz), 5.31 (1H, dd,  $J_{\rm H,H} = 17.3$ , 1.0 Hz), 5.57 (1H, d,  $J_{\rm H,H} = 6.6$  Hz), 6.68 (1H, dd,  $J_{\rm H,H} = 17.4$ , 10.8 Hz); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  20.1, 58.5, 115.2, 130.6, 134.0, 134.6.

**3-Methylene-4-penten-1-ol (4-OH).** Hexane was removed from 8.4 mL (17 mmol) of a solution of butyllithium (2.0 M in hexane) before a solution containing 170 mg (16.8 mmol) of diisopropylamine, 1.68 mL of (1.68 mmol) of potassium *tert*-butoxide (1.0 M in THF), and 1.50 g (16.8 mmol) of 4-methylenetetrahydropyran (**3**) in 20 mL of THF (precooled at -75 °C) was added. After 2 h at -50 °C, the reaction was quenched with saturated potassium dihydrogen phosphate and extracted with 30/70 diethyl ether/hexane. Solvent was removed at reduced pressure (bp 68–71 °C at 22 mmHg) to give 0.75 g (50%) of a colorless oil:<sup>23</sup> <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.80 (1H, s (broad)), 2.52 (2H, t,  $J_{\rm H,H} = 6.6$  Hz), 3.77 (2H, dt,  $J_{\rm H,H} = 6.3$  Hz), 509–5.30 (4H, m), 6.40 (1H, ddd,  $J_{\rm H,H} = 18.0$ , 10.5, 0.6 Hz); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  34.8, 61.1, 114.0, 117.8, 138.5, 142.8.

- (23) Margot, C.; Rizzolio, M.; Scholosser, M. Tetrahedron 1990, 46, 2411– 2424.
   (24) Clark K B.; Culshaw P. N.; Griller, D.; Lossing, F. P.; Martinbo Simoes
- (24) Clark, K. B.; Culshaw, P. N.; Griller, D.; Lossing, F. P.; Martinho Simoes, J. A.; Walter, J. C. J. Org. Chem. **1991**, 56, 5535-5539.
  (25) Mayr, H.; Heilman, W. Tetrahedron **1986**, 42, 6657-6662.

<sup>(15)</sup> Street, I. P.; Christensen, D. J.; Poulter, C. D. J. Am. Chem. Soc. 1990, 112, 8577–8578.

<sup>(22)</sup> Burger, B. V.; Plessis, du E.; Garbers, C. F.; Pachler, K. G. R. J. Chem. Soc., Perkin Trans. 1 1973, 584–587.

3-Methylene-4-pentenyl Tosylate (4-OTs). To a solution of 93 mg (0.50 mmol) of p-toluenesulfonyl chloride and 73 mg (0.60 mmol) of 4-(N,N-dimethylamino)pyridine (DMAP) in 4 mL of CH2Cl2 was added 47 mg (0.48 mmol) of 4-OH. After 2 h, the mixture was poured into hexane, and the resulting precipitates were removed by filtration. Solvent was removed at reduced pressure, and the residue was chromatographed on silica gel (40/60 diethyl ether/hexane) to give 113 mg (92%) of a clear colorless oil: IR (neat) 3100, 3043, 3015, 2959, 2917, 1650, 1603, 1358, 1309, 1293, 1230, 1188, 1176, 1096, 970 cm<sup>-1</sup>; UV (CHCl<sub>3</sub>)  $\lambda_{max}$  222 ( $\epsilon$  18 600), 197 ( $\epsilon$  13 800) nm; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.45 (3H, s), 2.59 (2H, td,  $J_{\rm H,H}$  = 7.2, 0.9 Hz), 4.14 (2H, t,  $J_{\rm H,H}$  = 7.2 Hz), 5.0–5.15 (4H, m), 6.29 (1H, dd,  $J_{H,H} = 17.6$ , 10.5 Hz), 7.35  $(1H, d, J_{H,H} = 8.4 \text{ Hz}), 7.79 (1H, d, J_{H,H} = 8.4 \text{ Hz}); {}^{13}\text{C NMR} (CDCl_3)$ δ 21.9, 31.1, 68.9, 114.0, 118.6, 128.0, 129.9, 130.0, 133.2, 138.0, 140.7, 144.9; HRMS (EI at 70 eV) calcd for C13H16O3S 252.0816, found 252.0797.

General Procedure for Synthesis of Diphosphates. The procedure of Davission and co-workers<sup>26</sup> was followed with the exception that HPLC was used in the final purification step instead of flash chromatography on cellulose. To a solution of 1.5-3.0 equiv of tris-(tetra-n-butylammonium) hydrogen pyrophosphate in CH<sub>3</sub>CN was added a solution of tosylate, chloride, or bromide (0.25-0.50 mM) in CH<sub>3</sub>CN. The resulting mixture was allowed to stir for 2-3 h at room temperature. Solvent was removed at reduced pressure, and the resulting residue was dissolved in a minimal volume of 25 mM NH<sub>4</sub>HCO<sub>3</sub> (ionexchange buffer). The clear solution was passed through DOWEX AG 50W-X8 (100-200 mesh, ammonium form) that had been equilibrated with two column volumes of the ion exchange buffer at a flow rate of one column volume/15 min. The resulting clear solution was concentrated by rotary evaporation and lyophilized. The residue was then transferred to a centrifuge tube and dissolved in 100 mM NH<sub>4</sub>HCO<sub>3</sub>. The resulting solution was extracted two or three times with a 1:1 mixture of CH<sub>3</sub>CN and 2-propanol. CH<sub>3</sub>CN and 2-propanol were removed at reduced pressure, and the aqueous layer was lyophilized. The residue was by HPLC, and the purified diphosphate was stored at −78 °C.

(E)-3-Methyl-2,4-pentadienyl Diphosphate (E-2-OPP). A solution of 90 mg (0.92 mmol) of E-2-Br in CH<sub>3</sub>CN was treated with 2.74 g (3.00 mmol) of tris(tetra-n-butylammonium) hydrogen pyrophosphate. The ammonium form of the diphosphate was purified by HPLC on an Asahipak ODP-90 HPLC column with gradient elution (solvent A =100 mM NH<sub>4</sub>HCO<sub>3</sub>, solvent B = CH<sub>3</sub>CN; 0-20 min (A) 20-35 min 1:1 (A/B), 35-45 min, 1:1 (A/B)) to give 120 mg (42%) of a white solid: UV (25 mM NH<sub>4</sub>HCO<sub>3</sub>) λ<sub>max</sub> 228 (ε 14 800), 197 (ε 10 600) nm; <sup>1</sup>H NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  1.60 (3H, s), 4.40 (2H, dd,  $J_{\rm H,H} = J_{\rm H,P} = 6.6$  Hz), 4.94 (1H, d,  $J_{\rm H,H} = 11.1$  Hz), 5.26 (1H, d,  $J_{\rm H,H}$ = 17.7 Hz), 5.53 (1H, t,  $J_{H,H}$  = 6.6 Hz), 6.29 (1H, dd,  $J_{H,H}$  = 17.7, 10.8 Hz); <sup>13</sup>C NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  14.1, 65.3 (d, J<sub>C,P</sub> = 5.2 Hz), 112.7, 129.9 (d,  $J_{C,P} = 7.9$  Hz), 140.9, 143.2; <sup>31</sup>P NMR (D<sub>2</sub>O/ ND<sub>4</sub>OD, pH = 8.0)  $\delta$  -1.90 (1P, d,  $J_{P,P}$  = 22 Hz), -4.76 (1P, d,  $J_{P,P}$ = 22 Hz); HRMS ((–)-FAB) calcd for  $C_6H_{11}P_2O_7$  256.9980, found 256.9966.

(Z)-3-Methyl-2,4-pentadienyl Diphosphate (Z-2-OPP). Following the procedure described for E-2-OPP, 63 mg (0.54 mmol) of Z-2-CI was treated with 0.901 g (1.0 mmol) of tris(tetra-*n*-butylammonium) hydrogen pyrophosphate to give 87 mg (52%) of a white solid: UV (25 mM NH<sub>4</sub>HCO<sub>3</sub>)  $\lambda_{max}$  229 ( $\epsilon$  8900), 196 ( $\epsilon$  6800) nm; <sup>1</sup>H NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  1.72 (3H, s), 4.44 (2H, dd,  $J_{H,H} = J_{H,P} =$ 7.0 Hz), 5.13 (1H, d,  $J_{H,H} = 11.1$  Hz), 5.25 (1H, d,  $J_{H,H} = 17.4$  Hz), 5.50 (1H, t,  $J_{H,H} = 7.3$  Hz), 6.71 (1H, dd,  $J_{H,H} = 17.2$ , 10.9 Hz); <sup>13</sup>C NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  21.6, 64.2 (d,  $J_{C,P} = 5.2$  Hz), 119.3, 127.7 2 (d,  $J_{C,P} = 7.8$  Hz), 135.6, 140.1; <sup>31</sup>P NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  -1.83 (1P, d,  $J_{P,P} = 21$  Hz, P1), -4.83 (1P, d,  $J_{P,P} = 21$  Hz, P2); HRMS ((-)-FAB) calcd for C<sub>6</sub>H<sub>11</sub>P<sub>2</sub>O<sub>7</sub> 256.9980, found 256.9959. **3-Methylene-4-pentenyl Diphosphate (4-OPP).** Following the procedure described for **E-2-OPP**, 230 mg (1.10 mmol) of **4-OTs** was treated with 2.98 g (3.30 mmol) of tris(tetra-*n*-butyl) ammonium pyrophosphate to give 147 mg (43%) of a white solid: UV (25 mM NH<sub>4</sub>HCO<sub>3</sub>)  $\lambda_{max}$  224 ( $\epsilon$  16 600) nm; <sup>1</sup>H NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  2.51 (1H, dd,  $J_{H,H}$  = 18.0, 10.5 Hz), 3.98 (2H, dd,  $J_{H,H}$  = 7.2 Hz), 5.03–5.31 (4H, m), 6.37 (1H, dd,  $J_{L,H}$  = 18.0, 10.5 Hz); <sup>13</sup>C NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  34.5 (d,  $J_{C,P}$  = 5.6 Hz), 67.7 (d,  $J_{C,P}$  = 4.8 Hz), 116.9, 120.5, 141.1, 145.3; <sup>31</sup>P NMR (D<sub>2</sub>O/ND<sub>4</sub>OD, pH = 8.0)  $\delta$  –0.46 (1P, d,  $J_{P,P}$  = 22 Hz), -4.63 (1P, d,  $J_{P,P}$  = 22 Hz); HRMS ((–)-FAB) calcd for C<sub>6</sub>H<sub>11</sub>P<sub>2</sub>O<sub>7</sub> 256.9980, found 256.9972.

Assay for IPP Isomerase. *Schizosaccharomyces pombe*<sup>27</sup> and *Escherichia coli*<sup>28</sup> IPP isomerase were obtained as described previously. The activity of the enzymes was measured by the acid-liability protocol<sup>9</sup> in a total volume 200  $\mu$ L of 50 mM HEPES buffer, pH 7.0, containing 10 mM MgCl<sub>2</sub>, 200 mM KCl, 0.5 mM DTT, 1 mg/mL BSA, and 400  $\mu$ M [1-<sup>14</sup>C]IPP (10  $\mu$ Ci/ $\mu$ mol). The mixture was equilibrated at 37 °C and initiated by addition of 10  $\mu$ L of a solution of enzyme in assay buffer. Addition of a His<sub>6</sub> tag to *E. coli* IPP isomerase did not significantly alter its kinetic properties.

Time-dependent inactivation of the enzyme was measured as follows. Assay buffer containing varying concentrations of inhibitor was equilibrated at 37 °C before a 20-fold higher concentration of enzyme than required for the standard assay was added to a final volume of 200  $\mu$ L. Samples (10  $\mu$ L) were removed at different times and added to assay buffer containing 400  $\mu$ M [<sup>14</sup>C]IPP (10  $\mu$ Ci/ $\mu$ mol) to a final volume of 200  $\mu$ L.

Crystallization of E. coli IPP isomerase. Crystals of recombinant (C-terminal His-tagged) E. coli IPP isomerase were grown at room temperature by the hanging drop method as described by Oudjama et al.28 Briefly, protein (10 mg/mL) was equilibrated against a reservoir containing PEG2000 (16%), ammonium sulfate (100 mM), and MnCl<sub>2</sub> (10 mM) buffered with Tris/maleate at pH 5.5. Complexes with 4-OPP were obtained by soaking crystals of the enzyme in a 25 mM solution of the inhibitor in Tris/maleate (100 mM, pH 5.5), PEG2000 (16%), ammonium sulfate (100 mM), MnCl<sub>2</sub> (10 mM), and glycerol (25%). These conditions were identical to those used to crystallize the wildtype enzyme. After soaking, a crystal was flash frozen and diffraction data were collected with a Mar345 imaging plate system from Marresearch equipped with Osmic optics and running on an FR591 rotating anode generator. Diffraction data were processed with the MarFLM suite. In the presence of metal, the protein crystallizes, with two molecules in the asymmetric unit, in orthorhombic space group  $P2_12_12_1$  with cell parameters a = 69.3, b = 72.6, and and c = 92.5 Å. A data set was recorded at 2.05 Å resolution. Refinement was performed with the program SHELXL97 using the structure of unliganded IPP isomerase (PDB reference code: 1hxz) as starting model.

## Results

Synthesis of Diene Analogues of IPP and DMAPP. The syntheses of DMAPP analogues E-2-OPP and Z-2-OPP and IPP analogue 4-OPP are shown in Scheme 2. The E- and Z-isomers of diene analogues for DMAPP (E-2-OH and Z-2-OH) were prepared from commercially available isomeric acetylenic alcohols E-1-OH and Z-1-OH by selective hydrogenation of the triple bond using Lindlar catalyst. Initially both alcohols were converted to the corresponding bromides. However, the yield for the Z-isomer was substantially lower, and the less reactive chloride was prepared instead. The halides were phosphorylated by treatment with tris(tetra-*n*-butylammonium)-hydrogen pyrophosphate to give E- and Z-2-OPP. The diene

<sup>(26)</sup> Davisson, V. J.; Woodside, A. B.; Neal, T. R.; Stremler, K. E.; Muehlbacher, M.; Poulter, C. D. J. Org. Chem. 1986, 51, 4768–4779.

<sup>(27)</sup> Hahn, F. M.; Poulter, C. D. J. Biol. Chem. 1995, 270, 11298-11303.

<sup>(28)</sup> Oudjama, Y.; Derbecq, V.; Sainz, G.; Clantin, B.; Tricot, C.; Stalon, V.; Villeret, V.; Droogmans, L. Acta Crystallogr. Sect. D, Biol. Crystallogr. 2001, 57, 287–288.

Scheme 2. Synthesis of Diphosphates E-2-OPP, Z-2-OPP, and 4-OPP



analogue for IPP was synthesized from 4-methylene tetrahydropyran. The pyran ring was opened upon treatment with lithium diisopropyl amide (LDA)/potassium *tert*-butoxide. Alcohol **4-OH** was converted to the corresponding tosylate, followed by treatment with inorganic pyrophosphate to give **4-OPP**.

**Kinetic Studies.** Preliminary experiments indicated that S. pombe and E. coli IPP isomerase was irreversibly inactivated upon incubation with the diene analogues of IPP and DMAPP. The kinetic constants for inactivation of the S. pombe enzyme were measured by incubation of the protein with varying concentrations of the inhibitors. Samples were removed over a period of 30 min and diluted 20-fold into buffer containing [14C]IPP, and the activity of the enzyme was measured. In a control experiment, the enzyme retained >97% of its activity over a period of 1 h when incubated under similar conditions without addition of an inhibitor. In each case concentrationdependent first-order inactivation was observed (Figure 1). The rate constants for inactivation  $(k_{inact})$  were determined from a semilogarithmic plot of residual enzyme activity versus time. The rate constants for inhibition at saturating levels of the inhibitors  $(k_{\rm I})$  and the inhibitor constants  $(K_{\rm I})$  were determined from replots of  $k_{inact}$  versus the reciprocal of the inhibitor concentrations. The values for  $k_{\rm I}$  and  $K_{\rm I}$  are given in Table 1. All of the diene analogues were potent inhibitors of the enzyme. **E-2-OPP** was the most efficient inhibitor as measured by  $k_{\rm I}$  $K_{\rm I}$ . Although **Z-2-OPP** was ~10 times more reactive than **E-2-OPP**, its  $K_{\rm I}$  was substantially higher. **4-OPP** was only slightly less efficient than E-2-OPP. Both of the DMAPP analogues were alternate substrates for chain elongation by avian FPP synthase, with steady-state kinetic constants similar to those for DMAPP (data not shown); however, neither compound inactivated the enzyme.

**Crystallographic Structure of the** *E. coli* **4-OPP-IPP Isomerase Complex.** The quality of the crystallographic data and the refinement are summarized in Table 2. Unambiguous electron density in the initial Fourier difference  $(F_o - F_c)$  maps (contoured at  $3\sigma$ ) confirmed the presence of the inhibitor within the active site of the protein. Simulated-annealing omit maps of the refined models confirmed the placement on the inhibitor with a distinct geometry for both molecules in the asymmetric unit. Electron density is observed consistent with formation of a covalent bond between the sulfur of Cys67 in the active site and C5 in **4-OPP** and a new carbon–carbon double bond between C3 and C4. Interestingly, in one protein molecule the stereochemistry of the double bond in the covalently attached diene inhibitor is Z, while in the other it is E. The position of

**Table 1.** Kinetic Constants for Irreversible Inhibition of *S. pombe* 

inhibitor	<i>k</i> <sub>l</sub> (s <sup>-1</sup> )	<i>K</i> ι (μM)	$k_{\rm I}/K_{\rm I} ({\rm s}^{-1}\mu{\rm M}^{-1})$
E-2-OPP	$17 \pm 1.2$	$0.91\pm0.08$	19
Z-2-OPP	$186 \pm 11$	$84 \pm 7$	2.2
4-OPP	$43 \pm 5.1$	$3.2 \pm 0.2$	13

the diphosphate group in **4-OPP** is similar in both geometries. A second metal binds to nonbridging oxygens in the diphosphate moiety, and the stabilizing roles of Lys21, Arg51, Lys55, Glu83, and Glu87 are confirmed.



*Figure 1.* Plots of ln[% inactivation] versus time for incubation of IPP isomerase with **E-2-OPP**, **Z-2-OPP**, and **4-OPP**. Panel A, [**E-2-OPP**] = 0 ( $\bigcirc$ ), 0.4 ( $\bigcirc$ ), 0.5 ( $\square$ ), 0.7 ( $\blacksquare$ ), 1.0 ( $\triangle$ ), and 2.5 ( $\triangle$ )  $\mu$ M. Panel B, [**Z-2-OPP**] = 0 ( $\bigcirc$ ), 15 ( $\bigcirc$ ), 20 ( $\square$ ), 30 ( $\blacksquare$ ), 40 ( $\triangle$ ), and 50 ( $\triangle$ )  $\mu$ M. Panel C, [**4-OPP**] = 0 ( $\bigcirc$ ), 10 ( $\bigcirc$ ), 12 ( $\square$ ), 15 ( $\blacksquare$ ), 20 ( $\triangle$ ), and 25 ( $\triangle$ )  $\mu$ M.

Table 2. Statistics of Data Collection and Structure Refinement for IPP Isomerase-4-OPP

Crystal Data	
space group	$P2_{1}2_{1}2_{1}$
cell dimensions (Å)	68.633
	70.681
	90.773
subunits per asymmetric unit	2
Data Set	
wavelength (Å)	1.54179
highest resolution (Å)	2.03
total reflections	55557
unique reflections	26732
observed reflections $(I > 2\sigma(I))$	23693
completeness $(\%)^a$	93.3 (93.2)
$R_{\text{merge}}(\%)^a$	8.0 (27.5)
$I/\sigma(I)$	6.2
Refinement	
resolution range (Å)	10.0 - 2.0
number of protein atoms	2831
number of ligand atoms	52
number of water molecules	85
$R_{\rm cryst}$ (observed/all data) (%)	18.8/22.1
$R_{\text{free}}^{b}(\%)$	23.9
rms deviations	
bond lengths (Å)	0.007
bond angles (Å)	0.022
Ramachadran Plot <sup>c</sup>	
core region (%)	95.8
inner core region (%)	72.8
-	

<sup>a</sup> Statistics for the highest resolution shell (2.08-1.97 Å) are given in parentheses. <sup>b</sup> Free test subset represents 10% of total unique reflections. As defined by SHELXPRO,<sup>29</sup> Gly and Pro residues excluded.

## Discussion

The substrates for chain elongation,<sup>30,31</sup> isomerization,<sup>9</sup> and cyclization reactions<sup>31</sup> in isoprenoid metabolism are processed by reactions that proceed through discrete carbocationic intermediates. A common feature of these reactions is the addition of an electrophile  $(R_3^+)$  to a carbon–carbon double bond to generate a tertiary cation, as illustrated in Scheme 3. While the carbocationic intermediates are sufficiently electrophilic to alkylate any of the nucleophiles typically found in enzyme active sites, enzymes that process carbocations typically do not suffer inactivation during catalysis. However, if the structure of the intermediate is altered in a manner that changes the location of the positive charge in the active site, alkylation is commonly seen. For example, diene analogues of oxidosqualene ( $R_2 =$ -CH=CH<sub>2</sub>), which generate allylic cations during cyclization, irreversible inactivate oxidosqualene cyclase<sup>20</sup> and squalenehopene cyclase<sup>21</sup> during turnover. We reasoned that the IPP and





DMAPP diene analogues 2-E-OPP, 2-Z-OPP, and 4-OPP were likely candidates for irreversible inhibition of IPP isomerase by a related mechanism. In this case, an allylic cation is generated by protonation of the diene in an active site known to contain carboxylate and thiol nucleophiles. All three dienes were potent time-dependent irreversible inhibitors of S. pombe and E. coli IPP isomerase.

The isomeric forms of covalently bound 4-OPP are shown in Figure 2. The structures mostly differ in the stereochemistry of the C3-C4 double bond and the conformation of the hydrocarbon chain of the bound analogue. The most likely mechanism for inactivation involves protonation of the methylene group in 4-OPP to give isomeric allylic cations, which span C3, C4, and C5, followed by alkylation of the thiol group in C67 (Scheme 4). Since the barrier to rotation about the C3-C4 partial double bond in the allylic cation should exceed 10 kcal/mol,<sup>32</sup> we conclude that 4-OPP is bound in s-cis and s-trans conformations in the E·I complex prior to protonation.



Figure 2. Structures of the isomeric IPP isomerase-4-OPP complexes.

Scheme 4. Mechanism for Inhibition of Type I IPP Isomerase by **4-OPP** 



One of the carboxyl oxygens in the side chain of E116 is coordinated with a tightly bound divalent metal ion in type I IPP isomerase. The other carboxyl oxygen points toward the substrate in the binding pocket. The distance between the noncoordinated oxygen and the newly formed methyl group in the Z-adduct is 2.80 Å, and thus E116 is the only amino acid that could protonate the diene to give the structure that was seen. In the E-adduct, the methyl group is rotated to increase the distance to 3.98 Å, but again the non-coordinated oxygen of E116 is the only obvious source for the proton. This oxygen is also strongly hydrogen-bonded to the positively charged nitrogen in the tightly bound transition-state analogue 2-(N,N-dimethy-

- (30) Erickson, H.; Poulter, C. D. J. Am. Chem. Soc. 2003, 23, 6885-6888.
   (31) Liang, P.-H.; Ko, T.-P.; Wang, A. Eur. J. Biochem. 2002, 269, 3339-3354
- (32) Allinger, N. L.; Siefert, J. H. J. Am. Chem. Soc. 1975, 97, 752-760.

<sup>(29)</sup> Sheldrick, G. M.; Schneider, T. R. Methods Enzymol. 1997, 277, 319-343.

lamino)ethyl diphosphate.<sup>18</sup> Thus, E116 is a likely active-site acid in the protonation—deprotonation mechanism for type I IPP isomerase. Coordination with the divalent metal would enhance the acidity of the carboxyl group, although the state of protonation of that residue in the resting enzyme has not been established. In addition, the tertiary carbocationic intermediate would be stabilized by the carboxylate moiety following the protonation step.

In summary, diene analogues of IPP and DMAPP are potent active-site-directed irreversible inhibitors of type I IPP isomerase. During the inhibition reaction, the IPP analogue **4-OPP** is protonated at its C3 methylene to generate an allylic cation that alkylates the thiol group in C67 to form a primary thioether. An X-ray structure of inactivated *E. coli* IPP isomerase shows that the methyl group generated by protonation is located near E116, thereby implicating that residue in the protonation step of the isomerization reaction.

Acknowledgment. This project was supported by NIH Grant GM 25521 (C.D.P.) and an IISN Grant from the FNRS (Belgium). The authors thank Y. Oudjama and V. Durisotti for purification of the *E. coli* enzyme and crystallization assays.

JA056187H