LETTERS

Rhenium-Catalyzed Synthesis of 1,3-Diiminoisoindolines via Insertion of Carbodiimides into a C–H Bond of Aromatic and Heteroaromatic Imidates

Zijia Wang,[†] Shunsuke Sueki,^{†,‡,§} Motomu Kanai,[†] and Yoichiro Kuninobu^{*,†,‡}

[†]Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan [‡]CREST, Japan Science and Technology Agency (JST), 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Supporting Information

ABSTRACT: The rhenium-catalyzed synthesis of 1,3-diiminoisoindolines and their related compounds from aromatic or heteroaromatic imidates and carbodiimides are reported via C–H bond activation. This reaction is the first example of a transition-metal-catalyzed insertion of carbodiimides into an aromatic or heteroaromatic C–H bond and a novel method for synthesizing 1,3-diiminoisoindolines and their related compounds. Unsymmetrical 1,3-diiminoisoindolines were easily obtained using this method. The reaction proceeded in good to excellent yield using a variety of substrates.



1,3-Diiminoisoindolines are diimidated phthalimides and partial structures of phthalocyanines and their metal complexes. Phthalocyanines and their metal complexes are useful as organic functional materials, such as pigments, organic electroluminescence, and organic field effect transistors.¹ In addition, 1,3-diiminoisoindolines and their derivatives exhibit several biological activities as a complement component 3a antagonist and antimalarial candidates.² Synthetic reactions of 1,3-diiminoisoindolines, however, are still rare.³ In addition, synthesis of unsymmetrical 1,3-diiminoisoindolines using previously reported synthetic methods remains difficult.

We considered a new synthetic method of 1,3-diiminoisoindolines via C-H bond activation (Scheme 1): (1) a 1,3-

Scheme 1. Retrosynthetic Route of 1,3-Diiminoisoindolines



diiminoisoindoline is obtained by intramolecular nucleophilic cyclization of intermediate A and elimination of methanol; (2) intermediate A is formed by the insertion of a carbodiimide into a rhenium–carbon bond of intermediate B; and (3) intermediate B is generated by oxidative addition of an aromatic imidate to a rhenium catalyst (C–H bond activation).

Transition-metal-catalyzed C-H insertion reactions of unsaturated molecules have recently received considerable attention, and many reactions using several types of transition metal catalysts have been developed.⁴ To the best of our knowledge, however, there are no examples of transition-metalcatalyzed insertion of carbodiimides into an aromatic C–H bond. We reported new synthetic reactions of cyclic compounds by the insertion of unsaturated molecules into a C–H bond of various substrates and successive intramolecular nucleophilic cyclization under rhenium catalysis.^{5–7} We then considered that synthesis of 1,3-diiminoisoindolines, shown in Scheme 1, could be achieved using a rhenium catalyst. Herein we report the rhenium-catalyzed synthesis of 1,3-diiminoisoin-dolines from aromatic imidates and carbodiimides.

We began our studies with several transition metal complexes and salts using aromatic imidate **1a** and N_rN' -diisopropylcarbodiimide (**2a**) as model substrates. A rhenium complex, $\text{Re}_2(\text{CO})_{10}$, exhibited high catalytic activity. Treatment of **1a** with **2a** in the presence of the $\text{Re}_2(\text{CO})_{10}$ catalyst in chlorobenzene at 170 °C for 24 h gave 1,3-diiminoisoindoline **3a** in 99% yield (Scheme 2). Several other transition metal complexes promoted the annulation reaction, but the yields of **3a** were low.⁸ The stereochemistry of the imino groups was





Received: April 8, 2016

Organic Letters

determined by nuclear Overhauser effect experiments using (E)-N-((E)-5-bromo-2-isopropyl-3-(isopropylimino)-1-isoindolinylidene)-1-propanamine **3A** (see Supporting Information for details). Interestingly, the desired reaction proceeded in advance of the consumption of **2a** by the addition of methanol (byproduct) to **2a**.⁹ This reaction is the first example of transition-metal-catalyzed insertion of a carbodiimide into a C– H bond of aromatic and heteroaromatic compounds and a novel example of the synthesis of an unsymmetric 1,3diiminoisoindoline derivative.

We then investigated the substrate scope of aromatic and heteroaromatic imidates (Scheme 3). The desired reaction





proceeded, and 1,3-diiminoisoindoline **3b** was obtained in 85% yield when an aromatic imidate with a phenyl group on the nitrogen, **1b**, was used as a substrate. In the case of *N*-octyl aromatic imidates bearing an electron-donating or -withdrawing group, **1c**-**1g**, the corresponding 1,3-diiminoisoindolines **3c**-**3g** were afforded in 96%-99% yield. In these substrates, the functional groups remained unchanged during the reactions. A mixture of two regioisomers, **3h** and **3h**', was formed when an aromatic imidate with a methyl group at the *meta*-position, **1h**, was used as a substrate. In this reaction, the major product **3h** was produced by the reaction at the sterically less hindered site. The desired reaction also proceeded regioselectively when

naphthyl imidate 1i and heteroaromatic imidates 1j and 1k were used as substrates.

Next, we examined the substrate scope of carbodiimides (Scheme 4). Dicyclohexylcarbodiimide (2b) afforded the

Scheme 4. Reactions between Aromatic Imidate 1a and Several Carbodiimides 2^a



corresponding 1,3-diiminoisoindoline 3l in 90% yield. Carbodiimides bearing primary alkyl groups, 2c-2e, also produced 1,3-diiminoisoindolines 3m-3o in 82%-94% yield without loss of the functional groups. The corresponding 1,3-diiminoisoindolines 3p and 3q were obtained in good yield using dibenzylic carbodiimides as substrates. Interestingly, the regioselectivity was completely controlled in the reaction between 1a and unsymmetric carbodiimide 2h, and a single product 3r was produced. A diaryl carbodiimide 2i also gave the corresponding 1,3-diiminoisoindoline 3s in 83% yield.

A proposed mechanism for the formation of 1,3diiminoisoindolines 3 is shown in Scheme 5: (1) oxidative addition of aromatic imidate 1 to a rhenium catalyst (C–H bond activation); (2) insertion of carbodiimide 2 into the formed rhenium–carbon bond; (3) stereoinversion of a C==N double bond to avoid the steric repulsion between substituents R^3 and R^4 , and intramolecular nucleophilic cyclization; and (4) reductive elimination and elimination of methanol to give 1,3diiminoisoindoline 3 and regenerate the rhenium catalyst. As described above, the stereochemistry of product 3 was determined by nuclear Overhauser effect experiments. In addition, the formation of methanol was detected by ¹H NMR of the crude mixture. Scheme 5. Proposed Mechanism for the Formation of 1,3-Diiminoisoindolines 3



The annulation reaction proceeded in excellent yield, even in gram scale (Scheme 6). An annulation reaction of 1.09 g of aromatic imidate 1a with carbodiimide 2a gave 1.28 g of the desired product 3a in 85% yield.

Scheme 6. Gram-Scale Synthesis of 1,3-Diiminoisoindoline



A double annulation reaction also proceeded in excellent yield (Scheme 7). Treatment of aromatic diimidate 4 with *N*,*N*-

Scheme 7. Double Annulation Reaction



diisopropylcarbodiimide (2a) in the presence of the $\text{Re}_2(\text{CO})_{10}$ catalyst in chlorobenzene at 170 °C for 24 h gave 1,3diiminoisoindoline 5 in 97% yield as a single product. The reaction proceeded highly regioselectively, and only product 5 was obtained, whereas a regioisomer could be formed.

In summary, we successfully applied rhenium catalysis to synthesize 1,3-diiminoisoindolines and their related compounds from aromatic or heteroaromatic imidates and carbodiimides via C–H bond activation. The desired reaction proceeded in advance of the consumption of carbodiimides by the addition of methanol (byproduct) to carbodiimides. To the best of our knowledge, this is the first example of a transition-metalcatalyzed C–H insertion of carbodiimides, and this reaction is a novel method for synthesizing 1,3-diiminoisoindolines and their related compounds. Introduction of different substituents on the imino groups of 1,3-diiminoisoindolines by previously reported synthetic methods remains difficult. From this point of view, the present method is useful for synthesizing unsymmetrical 1,3-diiminoisoindolines. The reaction proceeded in good to excellent yield using a variety of substrates, even in gram scale. Double annulation also proceeded in excellent yield to give the desired product as a single product. We hope that the present reaction will become a useful method to synthesize 1,3-diiminoisoindolines and their related compounds.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.or-glett.6b01012.

Experimental procedures and compound characterization data (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: kuninobu@mol.f.u-tokyo.ac.jp.

Present Address

[§]Department of Chemistry, Graduate School of Science and Engineering, Tokyo Metropolitan University, 1-1 Minami Osawa, Hachioji, Tokyo, 192-0397, Japan.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported in part by CREST from JST and Grant-in-Aid for Scientific Research (B) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

REFERENCES

(1) (a) Tang, Q.; Li, H.; He, M.; Hu, W.; Liu, C.; Chen, K.; Wang, C.; Liu, Y.; Zhu, D. Adv. Mater. 2006, 18, 65. (b) Chaidogiannos, G.; Petraki, F.; Glezos, N.; Kennou, S.; Nešpůrek, S. Appl. Phys. A: Mater. Sci. Process. 2009, 96, 763. (c) Armstrong, N. R.; Wang, W.; Alloway, D. M.; Placencia, D.; Ratcliff, E.; Brumbach, M. Macromol. Rapid Commun. 2009, 30, 717. (d) Würthner, F.; Kaiser, T. E.; Saha-Möller, C. R. Angew. Chem., Int. Ed. 2011, 50, 3376.

(2) (a) Grant, E. B.; Guiadeen, D.; Singer, M.; Argentieri, D.; Hlasta, D. J.; Wachter, M. Bioorg. Med. Chem. Lett. 2001, 11, 2817.
(b) Mombelli, P.; Witschel, M. C.; van Zijl, A. W.; Geist, J. G.; Rottmann, M.; Freymond, C.; Roehl, F.; Kaiser, M.; Illarionova, V.; Fischer, M.; Siepe, I.; Schweizer, W. B.; Brun, R.; Diederich, F. ChemMedChem 2012, 7, 151.

(3) (a) Siegl, W. O. J. Org. Chem. 1977, 42, 1872. (b) Sato, R.; Senzaki, T.; Shikazaki, Y.; Goto, T.; Saito, M. Chem. Lett. 1984, 13, 1423. (c) Forsyth, T. P.; Williams, D. B. G.; Montalban, A. G.; Stern, C. L.; Barrett, A. G. M.; Hoffman, B. M. J. Org. Chem. 1998, 63, 331.
(d) Meder, M. B.; Gade, L. H. Eur. J. Inorg. Chem. 2004, 2004, 2716.
(e) Petrova, T. D.; Platonov, V. E. Russ. J. Org. Chem. 2005, 41, 218.
(f) Biitseva, A. V.; Hordiyenko, O. V.; Sukach, V. A.; Vovk, M. V.; Pichugin, K. A.; Konovalova, I. S.; Shishkin, O. V. Monatsh. Chem. 2008, 139, 939.

(4) For selected recent reviews, see: (a) Colby, D. A.; Bergman, R. G.; Ellman, J. A. Chem. Rev. 2010, 110, 624. (b) Satoh, T.; Miura, M. Chem. - Eur. J. 2010, 16, 11212. (c) Kuninobu, Y.; Takai, K. Chem. Rev. 2011, 111, 1938. (d) Song, G.; Wang, F.; Li, X. Chem. Soc. Rev. 2012, 41, 3651. (e) Wang, C. Synlett 2013, 24, 1606. (f) Rouquet, G.; Chatani, N. Angew. Chem., Int. Ed. 2013, 52, 11726. (g) Zhang, M.;

Zhang, Y.; Jie, X.; Zhao, H.; Li, G.; Su, W. Org. Chem. Front. **2014**, *1*, 843. (h) Tsurugi, H.; Yamamoto, K.; Nagae, H.; Kaneko, H.; Mashima, K. Dalton Trans. **2014**, 43, 2331. (i) Liu, B.; Hu, F.; Shi, B.-F. ACS Catal. **2015**, *5*, 1863.

(5) For examples of rhenium-catalyzed C(sp²)-H transformations via insertion of unsaturated molecules into a C(sp²)-H bond and successive intramolecular nucleophilic cyclization, see: (a) Kuninobu, Y.; Kawata, A.; Takai, K. J. Am. Chem. Soc. 2005, 127, 13498.
(b) Kuninobu, Y.; Tokunaga, Y.; Kawata, A.; Takai, K. J. Am. Chem. Soc. 2006, 128, 202. (c) Kuninobu, Y.; Nishina, Y.; Shouho, M.; Takai, K. Angew. Chem., Int. Ed. 2006, 45, 2766. (d) Kuninobu, Y.; Nishina, Y.; Nakagawa, C.; Takai, K. J. Am. Chem. Soc. 2006, 128, 12376.
(e) Kuninobu, Y.; Nishina, Y.; Matsuki, T.; Takai, K. J. Am. Chem. Soc. 2008, 130, 14062. (f) Kuninobu, Y.; Matsuki, T.; Takai, K. Org. Lett. 2010, 12, 2948. (g) Kuninobu, Y.; Yu, P.; Takai, K. Org. Lett. 2010, 12, 4274. (h) Sueki, S.; Guo, Y.; Kanai, M.; Kuninobu, Y. Angew. Chem., Int. Ed. 2013, 52, 11879. (i) Geng, X.; Wang, C. Org. Lett. 2015, 17, 2434. (j) Jin, X.; Yang, X.; Yang, Y.; Wang, C. Org. Chem. Front. 2016, 3, 268.

(6) For examples of rhenium-catalyzed C(sp²)-H transformations, see: (a) Kuninobu, Y.; Tokunaga, Y.; Takai, K. Chem. Lett. 2007, 36, 872. (b) Kuninobu, Y.; Kikuchi, K.; Tokunaga, Y.; Nishina, Y.; Takai, K. Tetrahedron 2008, 64, 5974. (c) Kuninobu, Y.; Fujii, Y.; Matsuki, T.; Nishina, Y.; Takai, K. Org. Lett. 2009, 11, 2711. (d) Kuninobu, Y.; Nakahara, T.; Yu, P.; Takai, K. J. Organomet. Chem. 2011, 696, 348. (e) Kuninobu, Y.; Ohta, K.; Takai, K. Chem. Commun. 2011, 47, 10791. (f) Xia, D.; Wang, Y.; Du, Z.; Zheng, Q.-Y.; Wang, C. Org. Lett. 2012, 14, 588. (g) Tang, Q.; Xia, D.; Jin, X.; Zhang, Q.; Sun, X.-Q.; Wang, C. J. Am. Chem. Soc. 2013, 135, 4628. (h) Geng, X.; Wang, C. Org. Biomol. Chem. 2015, 13, 7619.

(7) For several examples of transition-metal-catalyzed $C(sp^2)$ -H transformations via insertion of unsaturated molecules into a $C(sp^2)$ -H bond and successive intramolecular nucleophilic cyclization, see: (a) Tsuchikama, K.; Kasagawa, K.; Endo, K.; Shibata, T. Synlett 2010, 2010, 97. (b) Sun, Z.-M.; Chen, S.-P.; Zhao, P. Chem. - Eur. J. 2010, 16, 2619. (c) Tran, D. N.; Cramer, N. Angew. Chem., Int. Ed. 2010, 49, 8181. (d) Patureau, F. W.; Besset, T.; Kuhl, N.; Glorius, F. J. Am. Chem. Soc. 2011, 133, 2154. (e) Lian, Y.; Bergman, R. G.; Ellman, J. A. Chem. Sci. 2012, 3, 3088. (f) Li, B.-J.; Wang, H.-Y.; Zhu, Q.-L.; Shi, Z.-J. Angew. Chem., Int. Ed. 2012, 51, 3948. (g) Hou, W.; Zhou, B.; Yang, Y.; Feng, H.; Li, Y. Org. Lett. 2013, 15, 1814. (h) Tan, P. W.; Juwaini, N. A. B.; Seayad, J. Org. Lett. 2013, 15, 5166. (i) Qi, Z.; Wang, M.; Li, X. Org. Lett. 2013, 15, 5440. (j) Hung, C.-H.; Gandeepan, P.; Cheng, C.-H. ChemCatChem 2014, 6, 2692. (k) Rodríguez, A.; Albert, J.; Ariza, X.; Garcia, J.; Granell, J.; Farràs, J.; La Mela, A.; Nicolás, E. J. Org. Chem. 2014, 79, 9578. (1) De Sarkar, S.; Ackermann, L. Chem. -Eur. J. 2014, 20, 13932. (m) Liu, W.; Zell, D.; John, M.; Ackermann, L. Angew. Chem., Int. Ed. 2015, 54, 4092. (n) Sueki, S.; Wang, Z.; Kuninobu, Y. Org. Lett. 2016, 18, 304. For a review, see: (o) Zhang, Z.-S.; Chen, K.; Shi, Z.-J. Chem. Sci. 2014, 5, 2146.

(8) For the results of reactions between **1b** and **2a** in toluene at 150 °C: ReBr(CO)₅, 26%; [ReBr(CO)₃(thf)]₂, 30%; Re₂(CO)₁₀, 33%; [Rh(OH)(cod)]₂/*rac*-BINAP, 8%; [Cp*IrCl₂]₂, 16%. The desired reaction did not proceed using the following transition metal complexes and salts: $Mn_2(CO)_{10}$, $MnBr(CO)_5$, $Ru_3(CO)_{12}$, $RuH_2(CO)(PPh_3)_3$, [RuCl₂(*p*-cymene)]₂, RhCl(PPh_3)₃, [Cp*RhCl₂]₂, [Ir(OMe)(cod)]₂/dtbpy, Pd(OAc)₂, and Cu(OAc)₂.

(9) (a) Inoue, Y.; Taguchi, M.; Hashimoto, H. Synthesis **1986**, *1986*, 332. (b) Aresta, M.; Dibenedetto, A.; Fracchiolla, E.; Giannoccaro, P.; Pastore, C.; Pápai, I.; Schubert, G. J. Org. Chem. **2005**, *70*, 6177.