


This article was downloaded by: [Aston University]

On: 24 January 2014, At: 10:52

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

An Efficient and Mild Deprotection of 1,3-Oxathiolanes to Carbonyl Compounds Using the Superoxide Ion

Satish Kumar Singh^a & Krishna Nand Singh^a

^a Department of Chemistry, Faculty of Science ,
Banaras Hindu University , Varanasi, India

Published online: 04 Sep 2009.

To cite this article: Satish Kumar Singh & Krishna Nand Singh (2009) An Efficient and Mild Deprotection of 1,3-Oxathiolanes to Carbonyl Compounds Using the Superoxide Ion, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 184:9, 2339-2343

To link to this article: <http://dx.doi.org/10.1080/10426500802465850>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or

indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

An Efficient and Mild Deprotection of 1,3-Oxathiolanes to Carbonyl Compounds Using the Superoxide Ion

Satish Kumar Singh and Krishna Nand Singh

Department of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi, India

An efficient deprotection of 1,3-oxathiolanes to carbonyl compounds has been achieved under the mild reaction conditions of tetraethylammonium superoxide in an aprotic medium at room temperature.

Keywords Deprotection; 1,3-oxathiolane; phase-transfer catalyst; superoxide ion

INTRODUCTION













Protection and deprotection are common features of multistep organic synthesis. Among the various functional groups, the protection of the carbonyl group using 1,3-oxathiolanes has attracted a great deal of interest owing to their remarkable applications. They can be used as acyl carbanion equivalents¹ for carbon–carbon bond forming reactions. The utility of chiral 1,3-oxathiolanes for the synthesis of optically active tertiary alcohols having a carbonyl functionality at the α -position has been nicely demonstrated in organic synthesis.^{2,3} Moreover, the use of oxathiolanes is much more convenient than *O,O*-acetals or *S,S*-acetals. Although a large number of methods has been reported for the deprotection of 1,3-dithiolanes,⁴ only a few are known for the deprotection of 1,3-oxathiolanes.⁵ Usual and current methods for the deprotection of 1,3-oxathiolanes employ ammonium tribromide,⁶ IBX- β -cyclodextrin,⁷ BNBS,⁸ isoamyl nitrite,⁹ chloramine-T,¹⁰ TMSOTf,¹¹ NCS-AgNO₃,¹² NBS,¹³ and Amberlys-15/glyoxylic acid.¹⁴

Received 30 July 2008; accepted 10 September 2008.


The authors are thankful to UGC, New Delhi, for financial support.

Address Correspondence to K. N. Singh, Department of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi 221005, India. E-mail: knsinghbhu@yahoo.co. in

TABLE I Deprotection of 1,3-Oxathiolanes with *In Situ* Generated Tetraethylammonium Superoxide

Entry	Substrate	Time (h)	Yield (%) ^a
1	 1a	1.5	83
2	 1b	2.0	81
3	 1c	2.0	79
4	 1d	2.5	77
5	 1e	2.0	80
6	 1f	2.5	76
7	 1g	2.5	73
8	 1h	2.75	84
9	 1i	3.0	91
10	 1j	3.0	90
11	 1k	3.0	88
12	 1l	3.0	65

^aYields are pure isolated product.



SCHEME 1 Superoxide-induced deprotection of 1,3-oxathiolanes.

RESULTS AND DISCUSSION

In light of the above, and as a part of our ongoing research on the use of the superoxide ion¹⁵ in organic synthesis, we report herein a mild and efficient method to restore carbonyl compounds from 1,3-oxathiolanes using in situ generated tetraethylammonium superoxide in dry DMF at room temperature (Scheme 1).

A wide variety of structurally different 1,3-oxathiolanes (**1a-1l**) were prepared by the reaction of the appropriate carbonyl compound with 2-mercaptoethanol in the presence of Montmorillonite K-10 clay¹⁶ (**1a-1d**, **1f**, **1j-1l**) and PAS¹⁷ (**1e**, **1g**, **1h**, **1i**) as a catalyst in dichloromethane. The reaction conditions were then optimized for the transformation of 1,3-oxathiolanes to the parent carbonyl compounds. As an outcome, a mixture of substrate, potassium superoxide, and tetraethylammonium bromide in the molar ratio 1:2:1 in dry DMF was found to give the best result. Subsequently various 1,3-oxathiolanes (**1a-1l**) were smoothly converted to their parent carbonyl compounds (**2a-2l**) under the optimized conditions at room temperature (Table I). The products **2a-2h** were accompanied by the formation of the corresponding acids, albeit in low yield (10–15%).

CONCLUSION

The present findings demonstrate the role of the superoxide ion as a convenient and safe deprotecting agent for 1,3-oxathiolanes to the parent carbonyl compounds.

EXPERIMENTAL

Potassium superoxide and tetraethylammonium bromide were produced from E. Merck, Germany and were used as received. Dry *N,N*-dimethylformamide (DMF), from Aldrich, USA, was stored over molecular sieves (4Å) prior to use. The other reagents and solvents were of A.R. grade. All the protected and deprotected compounds were fully characterized by their IR, ¹H NMR, and ¹³C NMR spectra. IR spectra

were recorded on a JASCO FT/IR-5300 spectrophotometer. NMR spectra were run on a JEOL AL300 FTNMR spectrometer; chemical shifts are given in δ ppm, relative to TMS as the internal reference. All the products were purified by column chromatography.

REACTION OF IN SITU GENERATED TETRAETHYLAMMONIUM SUPEROXIDE WITH 1,3-OXATHIOLANES (1a–1l): GENERAL PROCEDURE

A mixture of potassium superoxide (1.13 g, 16 mmol) and tetraethylammonium bromide (1.68 g, 8 mmol) (weight under a nitrogen atmosphere using an atmosbag) in dry DMF (25 mL) was stirred for 15 min, and then the substrate 1,3-oxathiolane (**1a–1l**) (8 mmol) was added. After the reaction was over (1.5 to 3 h) as indicated by TLC, a cold brine solution (10 mL) was introduced to decompose the unreacted potassium superoxide, followed by the addition of saturated sodium hydrogen carbonate solution (10 mL). The reaction mixture was extracted with diethyl ether (3 \times 15 mL). The combined organic extract was dried over anhydrous Na₂SO₄, filtered, and evaporated to give the crude product (**2a–2l**), which was purified by column chromatography on silica gel using *n*-hexane:ethyl acetate (9:1) as an eluent. The aqueous phase was acidified with hydrochloric acid and extracted with diethyl ether (3 \times 10 mL) to isolate the acidic product.

REFERENCES

- [1] (a) E. L. Eliel and S. Morris-Natschke, *J. Am. Chem. Soc.*, **106**, 2937 (1984); (b) K. Fuji, M. Udea, K. Sumi, K. Kaziwara, E. Fujita, T. Iwashita, and I. Miura, *J. Org. Chem.*, **50**, 657 (1985).
- [2] J. E. Lynch and E. L. Eliel, *J. Am. Chem. Soc.*, **106**, 2943 (1984).
- [3] K. Utimoto, A. Nakamura, and S. Matsubara, *J. Am. Chem. Soc.*, **112**, 8189 (1990).
- [4] T. W. Greene and P. G. M. Wuts, *Protective groups in organic synthesis*, 3rd ed., (John Wiley and Sons, Inc., New York, 1999), pp. 329–344.
- [5] T. W. Greene and P. G. M. Wuts, *Protective groups in organic synthesis*, 3rd ed., (John Wiley and Sons, Inc., New York, 1999), pp. 346–347.
- [6] E. Mondel, P. R. Sahu, G. Bose, and A. T. Khan, *Tetrahedron Lett.*, **43**, 2843 (2002).
- [7] M. S. Reddy, M. Narender, A. Mahesh, Y. V. D. Nageswar, and K. R. Rao, *Synth. Commun.*, **36**, 3771 (2006).
- [8] R. Ghorbani-Vaghei and A. Khazaei, *Phosphorus, Sulfur, and Silicon*, **179**, 1787 (2004).
- [9] K. Fuji, K. Ichikawa, and E. Fujita, *Tetrahedron Lett.*, **19**, 3561 (1978).
- [10] D. W. Emerson and H. Wynberg, *Tetrahedron Lett.*, **12**, 3445 (1971).
- [11] (a) T. Ravindranathan, S. P. Chavan, and S. W. Dantale, *Tetrahedron Lett.*, **36**, 2285 (1995); (b) T. Ravindranathan, S. P. Chavan, R. B. Tejawani, and J. P. Varghese, *J. Chem. Soc., Chem. Commun.*, 1750 (1991); (c) T. Ravindranathan, S. P. Chavan,

- J. P. Varghese, S. W. Dantale, and R. B. Tejawani, *J. Chem. Soc., Chem. Commun.*, 1937 (1994).
- [12] (a) S. V. Frye and E. L. Eliel, *Tetrahedron Lett.*, **26**, 3907 (1985); (b) K. Nishide, K. Yokota, D. Nakamura, T. Sumiya, M. Node, M. Ueda, and K. Fuji, *Tetrahedron Lett.*, **34**, 3425 (1993).
- [13] B. Karimi, H. Seradj, and M. H. Tabaei, *Synlett*, 1798 (2000).
- [14] S. P. Chavan, P. Soni, and S. K. Kamat, *Synlett*, 1251 (2001).
- [15] (a) K. N. Singh, *Synth. Commun.*, **37**, 2651 (2007); (b) R. S. Raghuvanshi and K. N. Singh, *Synth. Commun.*, **37**, 1371 (2007); (c) R. S. Raghuvanshi and K. N. Singh, *Synth. Commun.*, **36**, 3075 (2006); (d) S. Singh and K. N. Singh, *Synth. Commun.*, **35**, 2597 (2005); (e) S. Singh, M. Verma, and K. N. Singh, *Synth. Commun.*, **34**, 4471 (2004).
- [16] S. Gogoi, J. C. Borah, and N. C. Barua, *Synlett*, 1592 (2004).
- [17] A. Majee, S. K. Kundu, and S. Islam, *Synth. Commun.*, **36**, 3767 (2006).