This article was downloaded by: [University of Chicago Library] On: 30 November 2014, At: 05:16 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lcar20</u>

SYNTHESIS OF VINYL $\alpha\text{-}D\text{-}GLUCOPYRANOSIDES$ FROM MIXED ACETAL GLYCOSIDES

Jocelyn Edathil^a, Joan Nguyen^a, Kevin Hughes^a, Walter J. Boyko^a & Robert M Giuliano^b

^a Department of Chemistry, Villanova University, Villanova, PA, 19085, U.S.A.

^b Department of Chemistry, Villanova University, Villanova, PA, 19085, U.S.A. Published online: 11 Dec 2006.

To cite this article: Jocelyn Edathil , Joan Nguyen , Kevin Hughes , Walter J. Boyko & Robert M Giuliano (2001) SYNTHESIS OF VINYL α -D-GLUCOPYRANOSIDES FROM MIXED ACETAL GLYCOSIDES , Journal of Carbohydrate Chemistry, 20:1, 81-85, DOI: <u>10.1081/CAR-100102545</u>

To link to this article: http://dx.doi.org/10.1081/CAR-100102545

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

J. CARBOHYDRATE CHEMISTRY, 20(1), 81–85 (2001)

COMMUNICATION

SYNTHESIS OF VINYL α -D-GLUCOPYRANOSIDES FROM MIXED ACETAL GLYCOSIDES¹

Jocelyn Edathil, Joan Nguyen, Kevin Hughes, Walter J. Boyko, and Robert M. Giuliano^{*}

Department of Chemistry, Villanova University, Villanova, PA 19085

Vinyl D-glucopyranosides are useful intermediates in a wide variety of applications in synthetic carbohydrate chemistry. They have been utilized as chiral auxiliaries in the inverse-electron-demand cycloaddition with isoquinolinium salts, giving homochiral tetralins (Bradsher cycloaddition),² and as precursors to enantiomerically pure cyclobutanols by [2+2] cycloaddition with ketenes.³ Vinyl glycosides that are unsaturated at C2-C3 undergo thermal Claisen rearrangement to give C-3 branched glycal derivatives.⁴ Isopropenyl⁵ and butenyl glycosides have been used as glycosyl donors in oligosaccharide synthesis.⁶ Vinyl glycosides and mixed acetal glycosides have been studied as substrates for glycosidases.^{7,8} In more recent studies, vinylated sugars, in which the vinyl group is attached to nonanomeric hydroxyl groups, have been utilized in the synthesis of cyclooctanoic mimetics of carbohydrates,⁹ as precursors to *C*-glycosides,¹⁰ and as intermediates in the synthesis of β -mannosides by intramolecular tethering and delivery.¹¹

Vinyl glycosides are most often synthesized from glucopyranosides with an unblocked anomeric hydroxyl group by transvinylation with mercuric acetate¹² or from glycosyl halides by nucleophilic displacement with bis(acylmethyl)mercury reagents.¹³

Elimination reactions of 2-(phenylselenyl)ethyl glycosides¹⁴ and 2-(trimethylammonium)-ethyl glycosides,¹² and photolysis of 4-oxopentyl glycosides by Norrish Type II reactions also give vinyl glycosides.¹⁵ Our studies of hetero-Diels-Alder reactions in carbohydrate synthesis required access to vinyl D-glucopyranosides. A mercury-free preparation of vinyl ethers derived from chiral alcohols has been reported,¹⁶ in which treatment of mixed acetals with trimethylsilyl trifluoromethane sulfonate and an amine (Gassman method)¹⁷ gave good yields of vinyl ethers. We were intrigued by the possibility of synthesizing vinyl glycosides by the Gassman method, since the required monosaccharides that

ORDER		REPRINTS
-------	--	----------

are unblocked at the anomeric center are readily available with both ether and ester protecting groups on the remaining hydroxyl groups. Treatment of these starting materials with vinyl ethers in the presence of an acid catalyst would provide the mixed acetal glycosides needed for the elimination reaction.¹⁸ Mixed acetal glycosides are also available by the reaction of TMS glycosides and dialkyl acetals in the presence of TMS-triflate.¹⁹ Herein, we describe the synthesis of vinyl and isopropenyl α -D-glucopyranosides by the Gassman method.

Both 2,3,4,6-tetra-*O*-benzyl-D-glucopyranoside and its corresponding tetra-*O*-acetyl derivative,²⁰ **1a** and **1b**, were treated with ethyl vinyl ether in dichloromethane in the presence of PPTS to give the mixed acetal α -D-glucopyranosides **2a**⁷ and **2b**⁸ in 85-90% yields as a 1:1 mixture of diastereomers that were not separated. The major products (~9:1) were the α -anomers in each case. Reduction of the quantity of acid catalyst and the use of no more than five equivalents of vinyl ether gave clean product that could be used directly in the elimination step (see Experimental).



Treatment of the benzyl-protected mixed acetal glycoside (2b) with trimethylsilyl trifluoromethanesulfonate and triethylamine gave the vinyl α -D-glucopyranoside 3^{12} in 68% yield after purification by flash chromatography. Products resulting from ring-opening, glycals formed by elimination of the anomeric substituent, or the TMS glycoside 4 that would result from an alternative elimination were not detected. The regioselectivity observed is consistent with that reported by Dujardin, Rossignol, and Brown,¹⁶ and it is assumed to result from complexation of the trimethylsilyl cation with the ethoxy group oxygen of the mixed acetal, rather than the oxygen at C-1.







VINYL α-D-GLUCOPYRANOSIDES

Downloaded by [University of Chicago Library] at 05:16 30 November 2014

The synthesis of isopropenyl 2,3,4,6-tetra-*O*-benzyl- α -D-glucopyranoside by the Gassman method was also successful. Treatment of the mixed acetal glycoside **5** (prepared from **1b** using 2-methoxypropene) gave a product mixture containing a 2:1 ratio of isopropenyl glycoside **6**⁵ to TMS glycoside **4**⁸, from which **6** was isolated in 54% yield.



The preparation of acetate ester-protected vinyl glycosides from mixed acetal glycosides **2a** by this procedure gave poor yields (20%) of vinyl glycoside, with decomposition being observed. Benzoate esters were not attempted, but these may be more suitable if competing enolization in the acetates is responsible for the low yields. In an example of the use of this method for the vinylation of other sites on the pyranose ring, the 3-*O*-vinyl ether **8** was obtained in 40% yield from mixed acetal **7**, which was readily prepared from methyl 4,6-*O*-benzylidene-2-deoxy- α -D-*ribo*-hexopyranoside.²¹

In summary, a mercury-free synthesis of vinyl and isopropenyl glycosides was developed based on the Gassman method. Vinylation of hydroxyl groups at other positions also appears to be feasible. Further studies to test the scope of the reaction, the optimization of experimental conditions, and applications of the vinylated sugars are in progress.



EXPERIMENTAL

Preparation of Mixed Acetal Glycosides 2a, 2b, 5, and 7. A solution of the alcohol (1 mmol), vinyl ether (ethyl vinyl ether or 2-methoxypropene for 5, 5 mmol), and PPTS (10 mg) in dry dichloromethane (20 mL) was stirred at room temperature until TLC (1:3 ethyl acetate-hexanes) showed that starting material was consumed (1-3 h). Solid sodium carbonate was added and after stirring for several minutes, the mixture was filtered and the filtrate concentrated under reduced pressure to give syrupy mixed acetal glycosides $2a^7$ and $2b^8$ in 85-90% yields, which were used without further purification. Compound 5 had $[\alpha]_D + 60^\circ$



Marcel Dekker, Inc.

270 Madison Avenue, New York, New York 10016

83

ORDER		REPRINTS
-------	--	----------

(*c* 1.1, chloroform); HRMS (ES+) Calcd. for $C_{38}H_{44}O_7Na$ [M+Na]: 635.3009. Found: 635.2985. Compound **7**: Anal. Calcd for $C_{18}H_{26}O_6$: C, 63.89; H, 7.74. Found: C, 63.55; H, 7.53.

Preparation of Vinyl Glycosides 3, 6 and Vinyl Ether 8. To a solution of mixed acetal glycoside (1 mmol) in dichloromethane (3 mL) cooled to 0 °C was added triethylamine (1.6 equiv) followed by TMS- triflate (1.3 equiv). The reaction mixture was allowed to warm to room temperature and then stirred overnight. Additional triethylamine (0.4 equiv) and TMS-triflate (0.3 equiv) was added and stirring was continued until TLC (1:3 ethyl acetate-hexanes) showed the consumption of the starting material, usually another 18-24 h. Vinyl glycosides had slightly higher R_f values than mixed acetal glycosides, for example 0.38 and 0.54 for **5** and **6**, respectively (1:3 ethyl acetate-hexanes). Dilute sodium hydroxide solution (2 mL) was added slowly to the reaction, after which it was diluted with dichloromethane (20 mL). The organic phase was washed with saturated sodium chloride solution, dried (Na₂SO₄) and concentrated. Vinyl glycosides 3^{12} and 6^{5} were purified by flash chromatography using 10% ethyl acetate-hexanes. The same procedure was used for the preparation of 3-O-vinyl sugar 8: Rf 0.20 (1:4 ethyl acetate/hexanes); α]_D +104° (*c* 1.0, chloroform); Anal. Calcd for C₁₆H₂₀O₅: C, 65.74; H, 6.90. Found: C, 65.43; H, 6.89.

ACKNOWLEDGEMENTS

The authors thank the Howard Hughes Medical Institute and the Chemistry Department of Villanova University for financial support of this research. Support for Jocelyn Edathil from the Villanova University Honors Program is also gratefully acknowledged.

REFERENCES

- 1. Presented at the 20th International Carbohydrate Symposium, Hamburg, Germany; Aug 27-Sept 1, 2000.
- Choudhury, A.; Franck, R. W.; Gupta, R. B. Cycloaddition of Isoquinolinium Salts: Homochiral Tetralins via Dienophiles Bearing Chiral Auxiliaries. Tetrahedron Lett. 1989, 37, 4921-4924.
- Ganz I.; Kunz, H. Carbohydrates as Chiral Auxiliaries. [2+2] Cycloadditions of Ketenes to Enol Ethers. Synthesis 1994, 1353-1358.
- de Raadt, A.; Ferrier, R. J. Syntheses and Reactions of Saturated and 2,3-Unsaturated Vinyl and 1'-Substituted Vinyl Glycosides. Carbohydr. Res. 1991, 216, 93-107.
- Marra, A.; Esnault, J.; Veyrieres, A.; Sinaÿ, P. Isopropenyl Glycosides and Cogeners as Novel Classes of Glycosyl Donors: Theme and Variations. J. Am. Chem. Soc. 1992, 114, 6354-6360.
- (a) Boons, G. -J.; Heskamp, B.; Hout, F. Vinyl Glycosides in Oligosaccharide Synthesis: A Strategy for the Preparation of Trisaccharide Libraries Based on Latent-Active Glycosylation. Angew. Chem. Int. Ed. Engl. 1996, 35, 2845-2847. (b) Boons, G.

Copyright @ Marcel Dekker, Inc. All rights reserved





VINYL α-D-GLUCOPYRANOSIDES

- -J.; Isles, S. Vinyl Glycosides in Oligosaccharide Synthesis. 2. The Use of Allyl and Vinyl Glycosides in Oligosaccharide Synthesis. J. Org. Chem. **1996**, *61*, 4262-4271.
- Dettinger, H. -M.; Lehman, J.; Wallenfals, K. Glycosides for Testing Glycosides: Vinyl, 1-Ethoxylethyl, and 1-Ethoxybut-3-enyl D-glucopyranosides. Carbohydr. Res. 1980, 87, 63-70.
- Tietze, L. F.; Fischer, R.; Guder, H. J.; Goerlach, A.; Neumann, M.; Krach, T. Synthesis of Acetal-α-Glucosides. A Stereoselective Entry into a New Class of Compounds. Carbohydr. Res. 1987, 164, 177-194.
- Wang, W.; Zhang, Y; Zhou, H.; Sinaÿ, P. Stereoselective Synthesis of Cyclooctano Mimetics of β-L-Glucopyranoside. In Abstracts of Papers, 20th International Carbohydrate Symposium, Hamburg, Germany; Aug 27-Sept 1, 2000; Thiem, J., Ed.; LCI Publishers, Hamburg, 2000, B-247.
- Godage, H. Y.; Fairbanks, A. J. Stereoselective Synthesis of C-Glycosides via Tebbe Methylenation and Claisen Rearrangement. Tetrahedron Lett. 2000, 41, 7589.
- Godage, H. Y.; Fairbanks, A. J. Stereoselective Synthesis of α-Glucosides and β-Mannosides via Temporary Molecular Tethering. In Abstracts of Papers, 20th International Carbohydrate Symposium, Hamburg, Germany, Aug 27-Sept 1, 2000; Thiem, J., Ed.; LCI Publishers, Hamburg, 2000, B-287.
- Perrine, T. D.; Glaudemans, C. P. J.; Ness, R. K.; Kyle, J.; Fletcher, Jr. H. G. Syntheses with Partially Benzylated Sugars. VII. The Anomeric Vinyl D-Glucopyranosides. J. Org. Chem., **1967**, *32*, 664-669 (1967).
- de Raadt, A.; Ferrier, R. J. A Simple, Efficient Synthesis of Vinyl β-D-Glucopyranosides. J. Chem. Soc., Chem. Commun. 1987, 1009-1010.
- 14. Rollin, P.; Verez Bencomo, V.; Sinaÿ, P. Use of Selenium in Carbohydrate Chemistry: Formation of Vinyl Glycosides. Synthesis, **1984**, 134-135.
- 15. Cottier, L.; Remy, G.; Descotes, G. Photochemical Synthesis of *O*-Vinyl Glycosides and Their Transformation into *C*-Branched Sugars. Synthesis **1979**, 711-712.
- Dujardin, G.; Rossignol, S.; Brown, E. Efficient Mercury-free Preparation of Vinyl and Isopropenyl Ethers of Chiral Secondary Alcohols and α-Hydroxyesters. Tetrahedron Lett. **1995**, *36*, 1653-1656.
- Gassman. P.; Burns, S. J.; Pfister, K. B. Synthesis of Cyclic and Acyclic Enol Ethers (Vinyl Ethers). J. Org. Chem., 1993, 58, 1449-1457.
- Tietze, L. F.; Logers, M. Stereoselective Synthesis of Acetal-β-glucosides by Reaction of 2,3,4,6-Tetra-O-acetyl-β-D-glucopyranose with Enol Ethers. Liebigs Ann. Chem. 1990, 261-265.
- Tietze, L. F.; Goerlach, A.; Beller, M. Synthesis of Glycoconjugates of Acetal-Glycosides with Lysine and Tripeptides for Selective Cancer Therapy. Liebigs Ann. Chem. 1988, 565-577.
- Zhang, J.; Kovacs, P. An Alternative Method for Regioselective, Anomeric Deacylation of Fully Acetylated Carbohydrates. J.Carbohydr. Chem. 1999, 18, 461-469.
- 21. Rosenthal, A.; Catsoulacos, P. Synthesis of Branched-Chain Sugars by the Wittig Reaction Can. J. Chem. **1968**, *46*, 2868-2872.

Received September 19, 2000 Accepted December 22, 2000

Downloaded by [University of Chicago Library] at 05:16 30 November 2014



85

Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the <u>U.S. Copyright Office</u> for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on <u>Fair Use in the Classroom</u>.

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our <u>Website</u> <u>User Agreement</u> for more details.

Order now!

Reprints of this article can also be ordered at http://www.dekker.com/servlet/product/DOI/101081CAR100102545