Facile Barton–McCombie Deoxygenation of Alcohols with Tetrabutylammonium Peroxydisulfate and Formate Ion

Hee Sock Park, Hee Yoon Lee, and Yong Hae Kim*

Center for Molecular Design and Synthesis, Department of Chemistry, KAIST, Daejeon 305-701, Republic of Korea

kimyh@kaist.ac.kr

Received April 21, 2005

ABSTRACT



A new method for efficient radical deoxygenation of alcohols is described for preparing bulk chemicals avoiding scale-up problems. Treatment of various thiocarbonyl derivatives with $(Bu_4N)_2S_2O_8$ and HCO₂Na in DMF afforded the corresponding deoxygenated products in excellent yields. The deoxygenation appears to be initiated by the transfer of a single electron to thiocarbonyl derivatives from $CO_2^{\bullet-}$ rather than from SO4^{•-}.

Radical deoxygenation of aliphatic alcohols is useful and important in organic synthesis. One of the most well-known methods is described by Barton and co-workers.¹ It utilizes the combination of the radicophilic nature of the thiocarbonyl group of thionocarbonates and xanthates and Bu₃SnH.² Although Bu₃SnH is the reagent most commonly used for effecting this reduction, it has some drawbacks related to the toxic organotin compounds.³

Considerable efforts have been made to devise more acceptable replacements such as silanes,⁴ dialkyl phosphates,⁵ and hypophosphorous acid and its salts⁵ instead of the tin hydride. Despite these efforts and in consideration of the prices of reagents, the control in radical reactions, and the simplicity of product separation, new methods for efficient and cost-effective deoxygenation processes are still in

demand especially for preparing bulk chemicals avoiding scale-up problems.

Among the known deoxygenation reactions, suitable hydrogen atom sources are inevitably used in combination with proper radical initiators such as AIBN and peroxides. The explosive property of AIBN and peroxides is another drawback of this kind of radical reaction. Thus the reaction control is also difficult in large-scale synthesis.⁶

It is reported that ethyl carbon radical generated from Et_3B —air interacts with the thiocarbonyl group in xanthate and drives the deoxygenation, although yields are somewhat low.^{7a} Additionally, undecyl carbon radicals arising from the thermolysis of the lauroyl peroxide were used for the same reaction.^{7b} Formation of carbon dioxide radical anion (CO₂•⁻) is reported by the reaction of the one-electron oxidant ammonium peroxydisulfate and HCO₂Na.⁸

In our earlier work, tetrabutylammonium peroxydisulfate $((Bu_4N)_2S_2O_8)$ was readily prepared by mixing 2 equiv of tetrabutylammonium hydrogen sulfate and potassium per-

Vol. 7, No. 15

<u>3187–3190</u>

⁽¹⁾ Barton, D. H. R.; McCombie, S. W. J. Chem. Soc., Perkin Trans. 1 1975, 1574.

⁽²⁾ Barton, D. H. R.; Ozbalik, N. *Phosphorus, Sulfur Silicon Relat. Elem.* **1989**, *43*, 349 and references therein.

^{(3) (}a) Graham, A. E.; Thomas, A. V.; Yang, R. J. Org. Chem. 2000, 65, 2583.

^{(4) (}a) Chatgilialoglu, C. Acc. Chem. Res. **1992**, 25, 188 and references therein. (b) Barton, D. H. R.; Jang, D. O.; Jaszberenyi, J. Cs. Tetrahedron **1993**, 49, 7193. (c) Cole, S. J.; Kirwan, J. N.; Roberts, B. P.; Willis, C. R. J. Chem. Soc., Perkin Trans. 1 **1991**, 103. (d) Togo, H.; Matsubayashi, S.; Yamazaki, O.; Yokoyama, M. J. Org. Chem. **2000**, 65, 2816.

^{(5) (}a) Barton, D. H. R.; Jang, D. O.; Jaszberenyi, J. Cs. J. Org. Chem. **1993**, 58, 6838. (b) Sato, T.; Koga, H.; Tsuzuki, K. *Heterocycles* **1996**, 42, 499.

⁽⁶⁾ Wehrstedt, K. D.; Knorr, A.; Schuurman, P. J. Loss Prev. Process Ind. 2003, 16, 523.

^{(7) (}a) Barton, D. H. R.; Jang, D. O.; Jaszberenyi, J. Cs. *Tetrahedron Lett.* **1990**, *31*, 3991. (b) Quiclet-Sire, B.; Zard, S. Z. *Tetrahedron Lett.* **1998**, *39*, 9435.

^{(8) (}a) Janzen, E. G.; Towner, R. A.; Brauer, M. Free Radical Res. Commun. **1988**, 4, 359. (b) Hu, C.-M.; Qing, F.-L. J. Org. Chem. **1991**, 56, 6348.

sulfate in water followed by extraction with methylene dichloride.9 In contrast to metal or ammonium peroxydisulfate, (Bu₄N)₂S₂O₈ is readily soluble in organic solvents such as acetonitrile, DMF, acetone, CH₂Cl₂, CHCl₃, and ethyl ether. Radicophilic cleavage of thiocarbonyl derivatives has been examined using (Bu₄N)₂S₂O₈ and HCO₂Na as a new alkyl radical generator.

Compound 4 is regarded to be a candidate as a therapeutic Alzheimer's disease drug that protects neurons from Ab toxicity and is in the preclinical stage as a hepatoprotective drug.¹⁰ In our new alternative process for improving the purification step, a key step in the synthesis of 4 involved the conversion of **3** to **4** (Scheme 1).



Barton-McCombie deoxygenation of 3 with Bu₃SnH in toluene at 110 °C gives 4 in 76% yields. Purification steps have been needed to remove residual toxic organotin byproducts in order to meet regulatory stipulations. This procedure affects product isolation.

Furthermore the reaction of 1 using another method,¹¹ using less toxic diphenylsilane and AIBN in toluene at 110 °C, gave 4 in 70% yields after a deprotection reaction with catalytic HCl in methanol.

Substrate 3 was reacted with (Bu₄N)₂S₂O₈/HCO₂Na (3 equiv/6 equiv) in DMF at 50 °C for 4 h to afford 4 in excellent yield of 95%. In the absence of HCO₂Na or $(Bu_4N)_2S_2O_8$, no product 4 was obtained: starting material 3 was recovered. This result indicates that both $(Bu_4N)_2S_2O_8$ and HCO₂Na are necessary for this deoxygenation.

(11) Barton, D. H. R.; Jang, D. O.; Jaszberenyi, J. Cs. Synlett 1991, 435.

In connection with these results, various alcohols were subjected to this optimized deoxygenation protocol. The results obtained are shown in Table 1. As shown, several

 Table 1. Deoxygenation of Primary, Secondary, and Tertiary
 Alcohols with (Bu₄N)₂S₂O₈ (3 equiv)/HCO₂Na (6 equiv) in DMF

entry	substrate	temp (°C)	time (min)	product	yield (%) ^a
1	5a	50	30	5b	100
2	6a	65	15	6b	86
3	7a	40	60	7b	96
4	8a	75	30	8b	100
5	9b	65	15	9c	95
6	10b	65	15	10e	75
7	10b	65	15	10e	98^b
8	10c	65	20	10e	90^b
9	10d	65	15	10e	97^b
10	11b	65	15	11c	65
11	11b	65	15	11c	75^b
12	12b	65	15	12d	65^b
13	12b	65	15	12d	73^c
14	12c	65	15	12d	75^c
15	13	60	60	14	95
16	15a	65	120	16a	95^d
17	15a	80	20	16a	95^e
18	15a	80	10	16a	95
19	15b	65	15	16b	90
20	17	65	60	18	86^b

^a Isolated yields obtained by silica gel column chromatography. ^b 8 equiv Na₂CO₃ was added. ^c 8 equiv Et₃N was added. ^d DMSO. ^e 3 equiv (HCO₂) 2Cu.

different primary, secondary, and tertiary alcohols were successively deoxygenated in high to excellent yields.

The S-methyl xanthate **9b** was prepared from **9a**,¹² which was obtained from the reduction of 7α -acetyl-6,14-endoethanotetrahydrothebaine¹³ using DIBAL-H. As an opioid derivative, **9c**¹⁴ was obtained in excellent yield. Substrates 10b,¹⁵ 10c,¹⁶ and 10d¹⁷ were prepared from glucofuranose derivative 10a. Compounds 11b and 12b were prepared from 11a¹⁸ and 12a,¹⁹ respectively. Their 3-deoxy sugar derivatives 10e¹⁵ and 11c¹⁸ and 5-deoxy sugar derivative 12d²⁰ obtained are useful intermediates in pharmaceutical synthesis.

Of special note, **12d** was obtained by the indirect deoxygenation of 12b. Because cyclic 3,5-O-thionocarbonate 12c

- (17) Dentey, R. W., Haldy, D. G., Dewis, S. W., Realinead, R. S.,
 Rushworth, W. I. J. Chem. Soc. C. **1969**, 826.
 (15) Iacono, S.; Rasmussen, J. R. Org. Synth. **1986**, 64, 57.
 (16) Tormo, J.; Fu, G. C. Org. Synth. **2002**, 78, 239.
 (17) Hagiwara, H.; Ohtsubo, S.; Kato, M. Tetrahedron **1997**, 53, 2415.
- (18) Nicolaou, K. C.; Daines, R. A.; Uenishi, J.; Li, W. S.; Papahatjis, D. P.; Chakraborty, T. K. J. Am. Chem. Soc. 1988, 110, 4672.
- (19) Suhara, Y.; Nihei, K.-i.; Kurihara, M.; Kittaka, A.; Yamaguchi, K.;

^{(9) (}a) Jung, J. C.; Choi, H. C.; Kim, Y. H. Tetrahedron Lett. 1993, 34, 3581. (b) Hwang, J. P.; Yang, S. G.; Kim, Y. H. Chem. Commun. 1997, 1355.

^{(10) (}a) Mook-Jung, I.; Shin, J.-E.; Yun, S. H.; Huh, K.; Koh, J. Y.; Park, H. K.; Jew, S.-S.; Jung, M. W. *J. Neurosci. Res.* **1999**, *58*, 417. (b) Jew, S.-S.; Yoo, C.-h.; Lim, D.-y.; Kim, H.; Mook-Jung, I.; Jung, M. W.; Choi, H.; Jung, Y.-h.; Kim, H.; Park, H. K. Bioorg. Med. Chem. Lett. 2000, 10.119.

^{(12) (}a) Bentley, K. W.; Hardy, D. G.; Meek, B. J. Am. Chem. Soc. 1967, 89. 3273. (b) The ¹H NMR spectral data of **9a** is reported: Fulmor, W.; Lancaster, J. E.; Morton, G. O.; Brown, J. J.; Howell, C. F.; Nora, C. T.; Hardy, R. A., Jr. J. Am. Chem. Soc. 1967, 89, 3322.

⁽¹³⁾ Bentley, K. W.; Hardy, D. G. J. Am. Chem. Soc. 1967, 89, 3267. (14) Bentley, K. W.; Hardy, D. G.; Lewis, J. W.; Readhead, M. J.;

Fujishima, T.; Konno, K.; Miyata, N.; Takayama, H. J. Org. Chem. 2001, 66, 8760. (20) Shah, R. H.; Schaeffer, H. J.; Murray, D. H. J. Pharm. Sci. 1965,

^{54, 15.}



was deoxygenated also to give **12d** (entry 14), it can be concluded that the reaction of **12b** goes through the cyclic thionocarbonate intermediate **12c**. Since deoxygenation of a primary alcohol is somewhat difficult, it can be an alternative deoxygenation method for primary alcohols. In these conversions, triethylamine or sodium carbonate was used as a base additive for diminishing the deprotection of products and reactants.

It is noteworthy that deoxygenation of 13^{21} gave the coupled product 14^{22} (entry 15). The diphenylmethyl radicals arising from the radical-induced C-O bond cleavage events may be too stabilized to abstract a hydrogen atom from the

formate ion (or other possible sources) and end up dimerizing (Scheme 2). The diphenylmethyl radicals do not provide



sufficient steric inhibition to radical-radical combination reactions, which is required for persistence.²³

Deoxygenation of **15a** and **15b**,²¹ which possess labile benzylic hydrogens, gave styrene derivatives **16a**²⁴ and **16b**,²⁵ respectively. Furthermore, bis-xanthate **17**^{4b} was deoxygenated to give the dideoxygenated product **18**^{4b} having a carbon–carbon double bond.

Besides sodium formate as a hydrogen source, copper formate and ammonium formate were tested with **10b**. The reactions were monitored by TLC analysis. A copper formate system did not give any deoxygenated product **10e**: starting material was recovered. The system of ammonium formate gave a low yield of **10e** (8%).

It can be postulated that alkyl radical C (Scheme 3) abstracts a hydrogen atom from formate ion, tetrabutylam-



monium ion, or solvent to form the deoxygenated product. To investigate the hydrogen source of products, **19** (Table 2) was synthesized by a conventional method¹⁵ using CD_3I instead of CH_3I .

The deoxygenation of **10b**, **10c**, and **19** has been examined in the presence of DCO₂Na or HCO₂Na in various solvents (Table 2). The amount of atom % deuterium incorporation of **20** was fully characterized by ¹H NMR.²⁶ None of the

⁽²¹⁾ Cristol, S. J.; Seapy, D. G. J. Org. Chem. 1982, 47, 132.

⁽²²⁾ Wakui, H.; Kawasaki, S.; Satoh, T.; Miura, M.; Nomura, M. J. Am. Chem. Soc. **2004**, *126*, 8658.

⁽²³⁾ Hirano, T.; Li, W.; Abrams, L.; Krusic, P. J.; Ottaviani, M. F.; Turro, N. J. J. Am. Chem. Soc. **1999**, *121*, 7170.

⁽²⁴⁾ Strassburg, R. W.; Gregg, R. A.; Walling, C. J. Am. Chem. Soc. 1947, 69, 2141.

⁽²⁵⁾ Cho, S.; Kang, S.; Keum, G.; Kang, S. B.; Han, S.-Y.; Kim, Y. J. Org. Chem. 2003, 68, 180.

⁽²⁶⁾ It has been reported that *N*-phenylthioxocarbamate of **10a** reacted with Bu₃SnD/AIBN to give **20**- β D in 85% yield by attacking of deuterium on the less hindered β -face selectively. Oba, M.; Nishiyama, K. *Tetrahedron* **1994**, *50*, 10193.

Table 2. Deoxygenation of **10b**, **10c**, and **19** with $(Bu_4N)_2S_2O_8$ (3 equiv), HCO₂Na or DCO₂Na (6 equiv), and Na₂CO₃ (8 equiv) in 0.05 M Solvent at 65 °C for 15 min



entry	substrate	conditions	ratio of βD/βH in 20 ^a	yield (%) ^b
1	10b	DCO ₂ Na, DMF	22/78	98
2	10b	$DCO_2Na, DMF-d_7$	20/80	98
3	10b	DCO_2Na , $DMSO$	17/83	97
4	10c	DCO ₂ Na, DMF	47/53	98
5	10c	DCO ₂ Na, DMSO	91/9	90^c
6	19	HCO ₂ Na, DMF	0/100	98

 a Determined by $^1\!{\rm H}$ NMR. b Isolated yields after silica gel column chromatography. c The reaction time was 30 min.

substrates (10b, 10c, and 19) give any 3α -deuterated product under the reaction conditions. Thus deuteration occurred at the 3-position in a β -face selective manner.

When **10b** reacted with $(Bu_4N)_2S_2O_8/DCO_2Na$ in DMF and DMF- d_7 , the atom % D incorporation of **20** reached 22% and 20%, respectively (entries 1 and 2). In the case of **10c** the atom % D incorporation of **20** was 47% in DMF. Interestingly, when DMSO was used as a solvent the deuteration using $(Bu_4N)_2S_2O_8/DCO_2Na$ occurred with 91 atom % D incorporation at the selective position. This result can be applied to obtain the deuterated 3-deoxy furanose derivatives.²⁷ Since an undeuterated $(Bu_4N)_2S_2O_8/HCO_2Na$ system did not give deuterated product **20** and quantitative yield of **10e** (98%, entry 6) was obtained, one of the hydrogen sources must be the hydrogen of HCO₂Na.

A plausible mechanism can be suggested from previously known results of tin hydride reduction.^{7a} The deoxygenation appears to be initiated by the transfer of a single electron to thiocarbonyl derivatives from $CO_2^{\bullet-}$ rather than from $SO_4^{\bullet-}$. The combination of the radicophilic nature of thiocarbonyl derivatives and $CO_2^{\bullet-}$ might be utilized to form alkyl radical **C** (Scheme 3).

In summary, an efficient and versatile deoxygenation method has been developed via the thiocarbonyl derivatives of alcohols such as xanthates, phenyl thionocarbonates, and thiocarbonylimidazolides. This new process may be generally available in the deoxygenation of alcohols bearing acid-sensitive functionalities. Of particular note, in the case of **10c** (phenyl thionocarbonate) the (Bu₄N)₂S₂O₈/DCO₂Na deoxygenation system in DMSO can provide the deuterated furanose derivatives at C3 position.

Supporting Information Available: General procedures, experimental procedures, characterization data, and copies of ¹H and/or ¹³C NMR spectra for compounds **1**, **3**, **4**, **5a**,**b**, **6a**,**b**, **7a**,**b**, **8a**,**b**, **9a**-**c**, **10b**-**e**, **11b**,**c**, **12b**-**d**, and **13**-**20**. This material is available free of charge via the Internet at http://pubs.acs.org.

OL050886H

⁽²⁷⁾ For recent review, see: Foldesi, A.; Trifonova, A.; Kundu, M. K.; Chattopadhyaya, J. *Nucleotides, Nucleosides Nucleic Acids* 2000, 19, 1615.