

that the C_1 -H of the *cis* compound has 3–5 Hz of a J value but that of the *trans* 10–12 Hz. The fact that the signal¹²⁾ due to the C_1 -H of these products (12) appears at δ 3.70 as a doublet having about 4 Hz of a J value allowed us to assign the configuration of them to be *cis*.

Refluxing these products (12) with chloral in $CHCl_3$ afforded 1,2,3,4-tetrahydro-1-naphthylformamides¹¹⁾ (13) in 83.0% [13a: mp 182.5–185°] and 87.0% [13b: mp 161–162°] yields. Dehydrogenation of these formamides (13) with DDQ gave the fully aromatized formamides¹¹⁾ (14) in 84.9% [14a: mp 217–219°] and 80.6% [14b: mp 211–215°] yields.

Bischler-Napieralski reaction of the trimethoxy-formamide (14a) with $POCl_3$ in xylene afforded red needles, mp 281–284° (lit.^{5a)} mp 275–276°) in 42.4% yield, which was identified with an authentic sample of sanguirubine⁵⁾ (3) chloride.

On the other hand, treatment of the pentamethoxy-formamide (14b) with $POCl_3$ in CH_3CN gave orange needles, mp 140–145° (dec.) (lit.^{5a)} mp 163–164°) in 60.8% yield, which was identified with an authentic sample of sanguilutine⁵⁾ (4) chloride. This material was also characterized as ϕ -cyanide (15), mp 228–232° (lit.^{5a)} mp 232–233°).

Acknowledgement We wish to thank Prof. J. Slavík, Purkyně University, for the gift of the samples of sanguirubine chloride and sanguilutine chloride.

Faculty of Pharmaceutical Sciences,
Chiba University, 1-33, Yayoi-cho,
Chiba, 260, Japan

HISASHI ISHII
TOSHIKO WATANABE¹³⁾
TSUTOMU ISHIKAWA

Received July 15, 1978

13) Present address: School of Pharmaceutical Science, Toho University, 2-2-1, Miyama, Funabashi, 274 Japan.

[Chem. Pharm. Bull.]
26(10)3254–3256(1978)

UDC 547.786.057 : 547.232.04

Synthesis of Isoxazole Derivatives by Means of O-Acylation of Aliphatic Nitro Compounds in the Presence of Acetylenic Compounds¹⁾

O-Acylation of some aliphatic nitro compounds or their alkali metal salts with acid chlorides in N,N-dimethylacetamide in the presence of acetylenic compounds afforded isoxazole derivatives in fairly good yield. The reaction is shown to arise from 1,3-dipolar cycloaddition of acetylenic compounds to the nitrile oxide intermediates which would result from the fragmentation of the initially formed O-acylated product.

Keywords—isoxazole; aliphatic nitro compounds; nitrile oxide; acetylenic compounds; dipolar aprotic solvent; O-acylation; fragmentation; 1,3-dipolar cycloaddition

Several examples have been reported of the O-acylation of aliphatic nitro compounds in which stable nitronic carboxylic anhydrides (mixed anhydrides) are formed from secondary nitroalkanes with acylating agents, while in the case of primary nitroalkanes, the correspond-

1) The Synthetic Reactions of Aliphatic Nitro Compounds. Part XIII: Presented at the 37th Annual Meeting of the Chemical Society of Japan, Yokohama, April, 1978: Abstracts, II, p. 731: Part XII of this series: E. Kaji, H. Kohno, and S. Zen, *Bull. Chem. Soc. Jpn.*, **50**, 928 (1977).

ing mixed anhydrides have not been isolated and the rearrangement of initially formed mixed anhydrides to hydroxamic acid esters has been observed.^{2,3)}

Our current interest in alkylation of nitronate ambident anions in a dipolar aprotic solvent,⁴⁾ also directed towards acylation of aliphatic nitronate anions in such a solvent. We have found that the reaction of primary nitro compounds such as nitroethane, phenylnitromethane, and methyl nitroacetate with acetyl chloride in N,N-dimethylacetamide (DMA) at room temperature gave 3,4-disubstituted furoxanes (3: R¹=CH₃, C₆H₅, or COOCH₃)⁵⁾ characterized by physical and spectroscopic identification. The mechanism of formation of the furoxanes appears to involve dimerization of nitrile oxides as intermediates which would result from the fragmentation of initially formed mixed anhydrides (1) as shown in Chart 1.




Chart 1

To support above suggestion and hoping this acylation to an isoxazole synthesis, we attempted to trap the nitrile oxides (2) with several acetylenic compounds as dipolarophiles.⁶⁾

Typical experimental conditions are exemplified as follows (Method-A): to a stirred solution of 300 mg (4 mmol) of nitroethane in 15 ml of anhydrous DMA, was added 4 ml of 1 N sodium methoxide in methanol. The mixture was cooled to 5° in an ice-bath, 0.28 ml (4 mmol) of acetyl chloride and 0.49 ml (4 mmol) of dimethyl acetylenedicarboxylate were added. After the addition had been completed the mixture was stirred for about 16 hr at room temperature, then 60 ml of ice-water was added. The resulting mixture was extracted with benzene. The extract was concentrated, and the resultant syrup was chromatographed on silica gel with hexane-ethyl acetate (3:1) as developer, giving 595 mg of 3-methyl-4,5-bis-(methoxycarbonyl)isoxazole (4a): mp 31–32° (methanol-water) in 75% yield. The infrared (IR), proton magnetic resonance (PMR) spectra, and elemental analysis of the product (4a)

- 2) A.T. Nielsen, "The Chemistry of the Nitro and Nitroso Groups," Part I, ed. by H. Feuer, Interscience Publishers, New York, 1969, pp. 461–468, and literatures cited therein.
- 3) A. McKillop and R.J. Kobylecki, *Tetrahedron*, **30**, 1365 (1974).
- 4) E. Kaji, A. Igarashi, and S. Zen, *Bull. Chem. Soc. Jpn.*, **49**, 3181 (1976) and our preceding papers.
- 5) J.V.R. Kaufman and J.P. Picard, *Chem. Rev.*, **59**, 429 (1959).
- 6) A closely related work has been shown that reaction of primary nitroalkanes with isocyanate and a catalytic amount of triethylamine in the presence of acetylenic compounds gives isoxazoles, in which the intermediate is also believed to be nitrile oxide. (T. Mukaiyama and T. Hoshino, *J. Am. Chem. Soc.* **82**, 5339 (1960); Some applications of the above reaction were reported: G.B. Bachman and L.E. Strom, *J. Org. Chem.*, **28**, 1150 (1963); J.E. McMurry, "Organic Syntheses," vol. 53, ed. by A. Brossi, John Wiley and Sons, Inc., New York, 1973, p. 59.)

TABLE I. Isoxazole Derivatives synthesized by Means of O-Acylation of Aliphatic Nitro Compounds with Acetyl Chloride

Compounds 4	R ¹	R ²	R ³	mp (°C) bp (°C/Torr)	Yield (%)	
					Method-A	Method-B
a	CH ₃	CO ₂ CH ₃	CO ₂ CH ₃	31—32 ^{a)}	75	44
b	CH ₃	H	CH ₂ OAc	65—73/2 ^{b)}	39	36
c	CH ₃	H	CO ₂ C ₂ H ₅	27—28 ^{c)}	54	45
d	CH ₃	H	C ₆ H ₅	65—66 ^{d)}	36	23
e	C ₆ H ₅	CO ₂ CH ₃	CO ₂ CH ₃	63.5—64.5 ^{e)}	76	71
f	C ₆ H ₅	H	CH ₂ OAc	37—38 ^{f)}	57	44
g	C ₆ H ₅	H	CO ₂ C ₂ H ₅	45—46 ^{g)}	63	53
h	C ₆ H ₅	H	C ₆ H ₅	141—143 ^{h)}	47	44
i	CO ₂ CH ₃	CO ₂ CH ₃	CO ₂ CH ₃	101—102 ⁱ⁾	20	17

a) Reported bp 74—76° (0.05 Torr).⁸⁾

b) Anal. Calcd. for C₇H₉NO₃: C, 54.19; H, 5.85; N, 9.03. Found: C, 54.54; H, 5.49; N, 9.25. MS *m/e*: 155 (M⁺). PMR (CDCl₃) δ: 2.21 (3H, singlet, COCH₃), 2.30 (3H, singlet, CH₃), 5.11 (2H, singlet, CH₂), 6.16 (1H, singlet, =CH).

c) Reported mp 27°. ⁷⁾

d) Reported mp 67°. ⁸⁾

e) Reported mp 62—63°. ⁹⁾

f) mp is not shown in Ref. 9.

g) Reported mp 47°. ¹⁰⁾

h) Reported mp 141°. ¹¹⁾

i) Anal. Calcd. for C₉H₉NO₃: C, 44.45; H, 3.73; N, 5.76. Found: C, 44.42; H, 3.66; N, 5.60. MS *m/e*: 243 (M⁺). PMR (CDCl₃) δ: 3.94 (3H, singlet, CO₂CH₃), 3.97 (6H, singlet, CO₂CH₃); Corresponding tris(ethoxycarbonyl)isoxazole has been reported in Ref. 12.

gave satisfactory results. Table I shows physical constants and analytical data of isoxazole derivatives synthesized by above manner. Another method employed sodium ethanenitronate (Method-B) also gave the isoxazole (4a) in relatively low yield (44%) as shown in Table I. Application of a phase transfer catalyst, *e.g.*, 18-crown-6 in benzene or triethylbenzylammonium chloride in benzene–water resulted in 31 and 14% yield respectively for 4a.

When benzoyl chloride (one equivalent) was used as an acylating agent for methoxycarbonylmethanenitronate, methyl 2-chloro-2-benzoyloxyiminoacetate (5) was isolated as a by-product; mp 125.5—126.5° (methanol–water), 33% yield, IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 1770 (ester C=O), 1740 (C₆H₅C=O), 1608 (C=N), PMR (CDCl₃) δ: 4.00 (3H, singlet, ester CH₃), 7.50—8.22 (5H, multiplet, C₆H₅), Anal. Calcd. for C₁₀H₈ClNO₄: C, 49.69; H, 3.31; N, 5.80. Found: C, 49.28; H, 3.32; N, 5.74. This compound (5) would be formed by addition reaction of excess benzoyl chloride to the nitrile oxide.

It should be noted that this method of the synthesis of isoxazoles is relatively facile and mild comparing with already reported ones.^{3,10)} Furthermore, as using olefinic dipolarophiles such as dimethyl fumarate, methyl acrylate and ethyl β-nitrocinnamate, some isoxazoline derivatives could be obtained; further details of these cases are now in progress.

Acknowledgement One of the authors (E. K.) wishes to thank Matsunaga Science Foundation for financial support.

School of Pharmaceutical Sciences
Kitasato University
Shirokane, Minato-ku, Tokyo 108, Japan

EISUKE KAJI
KAZUHO HARADA
SHONOSUKE ZEN

Received August 18, 1978

- 7) C. Musante, *Gazz. Chim. Ital.*, **70**, 685 (1940).
- 8) G. Casnati and A. Ricca, *Tetrahedron Lett.*, **1967**, 327.
- 9) P. Battioni, L. Vo-Quang, J.C. Raymond, and Y. Vo-Quang, *C.R. Acad. Sci. Paris (C)*, **t.271**, 1468 (1970).
- 10) J.E. Franz and H.K. Pearl, *J. Org. Chem.*, **41**, 1296 (1976).
- 11) G. Labbe and G. Mathys, *J. Org. Chem.*, **39**, 1221 (1974).
- 12) L. Panizzi, *Gazz. Chim. Ital.*, **70**, 119 (1940).