GDCh



Hypervalent Iodine

How to cite:

International Edition:doi.org/10.1002/anie.202007439German Edition:doi.org/10.1002/ange.202007439

An α-Cyclopropanation of Carbonyl Derivatives by Oxidative Umpolung

Adriano Bauer, Giovanni Di Mauro⁺, Jing Li⁺, and Nuno Maulide^{*}

In memory of Kilian Muñiz

Abstract: The reactivity of iodine(III) reagents towards nucleophiles is often associated with umpolung and cationic mechanisms. Herein, we report a general process converting a range of ketone derivatives into α -cyclopropanated ketones by oxidative umpolung. Mechanistic investigation and careful characterization of side products revealed that the reaction follows an unexpected pathway and suggests the intermediacy of non-classical carbocations.

he importance of the carbonyl functionality has been identified already at the dawn of modern organic chemistry in the 19th century and remains at a cardinal point of chemical synthesis. With the advent of the umpolung approach, developed by Seebach and Corey, an important paradigm shift was provided, delivering a systematic perspective to overcome the limitations of the natural polarity of the carbonyls (Scheme 1 a-I).^[1,2] The inversion of the polarity through derivatization was showcased first by dithiane chemistry, where a thicketal could be effectively used as C1 nucleophile (Scheme 1 a-II).^[1,3] Some umpolung reagents such as the cyanide anion^[4,5] or thiazolium based NHCs^[6] (both react as C1 nucleophiles and derive formally from formic acid) can transfer this umpolung-reactivity through catalysis (Scheme 1 a-III).^[7] A third approach to reverse the polarity of a given functional group consist in its oxidation or reduction by an external or internal reagent (Scheme 1a-IV).^[8] Ketones and their derivatives for instance are commonly employed in oxidative umpolung reactions. Classically promoted by toxic elements such as Hg(II),^[9,10] Tl(III),^[9,11] Pb(IV),^[9,12,13] or Se(IV)^[14] more modern variants rely on Bi(V),^[15] *N*-oxides,^[16-19] Mn(III),^[20,21] halosuccinimides,^[22] sulfoxides,^[23] and on iodine(III).^[24-29]

[*] A. Bauer, G. Di Mauro,^[+] N. Maulide
Institute of Organic Chemistry, University of Vienna
Währinger Strasse 38, 1090 Vienna (Austria)
E-mail: Nuno.Maulide@univie.ac.at
Homepage: https://organicsynthesis.univie.ac.at/
J. Li^[+]
Department of Chemistry, Tohoku University
Aoba-ku, 980-8578 Sendai (Japan)

The α -functionalization of certain ketones through oxidative umpolung with iodine(III) was pioneered already in the 1960s,^[30,31] has been extensively investigated in the 1980s,^[32–36] and gained further attention in more recent years.^[37–41] It is believed that the reaction involves an enolonium species (compound **8**, Scheme 1 a).^[42,43] This highly electrophilic intermediate can react with a variety of different nucleophiles. Such oxidations are often promoted by acids^[32] or less frequently by bases,^[44] with enol ethers^[32,41] or active methylene compounds^[37,38] being commonly employed.

In many cases however, the enolonium species does not directly react with an external nucleophile. Instead, skeletal rearrangements are observed leading to stabilized carbocationic intermediates. Net 1,2-phenyl migrations for instance which proceed *via* a phenonium intermediate^[45,46] are frequently witnessed (Scheme 1b).^[47-49] This illustrates that α -umpolung chemistry of carbonyl compounds can be leveraged to a new avenue for carbocationic rearrangements which no



Scheme 1. a) Natural polarity and types of umpolung. b) α -Arylation by C-C bond activation. c) This work: oxidative α -cyclopropanation.

Angew. Chem. Int. Ed. 2020, 59, 1-6

 $\textcircled{\sc c}$ 2020 The Authors. Published by Wiley-VCH GmbH

These are not the final page numbers!

^{[&}lt;sup>+</sup>] These authors contributed equally to this work.

Supporting information and the ORCID identification number(s) for

the author(s) of this article can be found under: https://doi.org/10.1002/anie.202007439.

^{© 2020} The Authors. Published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

longer require halide/leaving group abstractions^[50,51] or protonation of carbon–carbon multiple bonds.^[52–54] Building on these developments, we herein report a new protocol for cyclopropanation by exploiting the propensity of certain enolonium species to engage in carbocationic rearrangements.

Inspired by the aforementioned reports, we started to investigate the enolonium species, which is prone to rearrangement reactions. The allylic silylenol ether **14a** was chosen as the first substrate since homoallylic cations are known to undergo fast rearrangements.^[55–60] When treated with activated iodosobenzene at low temperature, fast formation of the cyclopropane **16aa** was observed (Scheme 2a). The product was isolated in more than 70% yield as a single diastereoisomer (*trans*). The only other product detected was the diastereomerically pure cyclobutane **20aa**. The regiochemical outcome of the formation of **20aa** was surprising, since a classical Prins-type attack of the double bond should lead to a 1,3-substituted cyclobutane product. This result urged us to investigate the mechanism further by isotope labeling.

When the substrate **14a**, marked with a ¹³C-label at the terminal olefin, was employed the product was observed to be a 1:1 mixture of isotopomers (Scheme 2b). Similar results were obtained by deuterium labeling (Scheme 2c). However, here a slight imbalance of the two isotopologues was observed, in favor of the endocyclically labeled compound



Scheme 2. a) Preliminary result. b) ¹³C labelling study. c) Deuterium labelling study. The slight imbalance in isotopologue-ratio can be accounted for by a secondary isotope effect. d) The *trans* cyclopropane can be synthesized similarly from a cyclopropyl silylenol ether. e) Other appended double bonds do not participate during the oxidation event.

 $[\boldsymbol{D}_2]\textbf{-14aa'},$ which is suggestive of a secondary kinetic isotope effect. $^{[88]}$

Both results are consistent with a highly fluxional intermediate and suggest the intermediacy of a non-classical cyclopropylcarbinyl cation at least as a transient species. Two additional experiments further ascertained our assumption. The β -cyclopropyl silylenol ether **21** led to the formation of the same product, likely by the formation of the same cationic intermediate (Scheme 2 d). Furthermore, when an additional CH₂-spacer was installed between the olefin and the silyl enolether (Scheme 2 e–**22**) cyclization was completely absent.^[89] Instead, a simple α -functionalization took place showcasing that the homoallylic double bond does not participate in a general Prins-type nucleophilic attack.

Based on our findings and previously reported experiments we propose the following mechanism for the cyclopropanation of **14a** (Scheme 3): First the metastable enolonium species **24** (or **25** respectively) is formed which interacts with the nucleophilic alkene in close proximity. Under extrusion of PhI a non-classical C_s symmetric cyclopropyl carbinyl cation is formed (**26**), which may be in equilibrium with two enantiomeric classical carbocations (**27/28**).^[90] Those can be trapped to yield the observed *trans* cyclopropane. The equilibrium is highly dynamic and thus other carbocations may contribute. However, the cyclobutonium **29/30** might benefit from carbonyl bridging, explaining the formation and the stereochemical outcome observed for side product **20 aa**.

Although homoallylic rearrangements involving nonclassical carbocations have been studied previously, they typically lead to inseparable mixtures of products and are perceived as synthetically unappealing.^[55–60]

As shown in Scheme 4, a broad variety of substrates could engage in this oxidative cyclopropanation in good yields and with high diastereoselectivities. A range of aromatic rings, ranging from electron-poor to electron-rich, was tolerated with no large effect on yield and/or selectivity. The aromatic ring itself is not required for high yields or selectivities since



Scheme 3. Proposed mechanism for the cyclopropanation of linear silylenol ethers.

www.angewandte.org

2

© 2020 The Authors. Published by Wiley-VCH GmbH

Angew. Chem. Int. Ed. 2020, 59, 1-6

K These are not the final page numbers!





Scheme 4. Product scope of the oxidative α -cyclopropanation of linear ketone derivatives. All yields refer to pure, isolated material. Reactions conducted on a 0.2 mmol scale, see Supporting Information for reaction conditions.

aliphatic ketones gave comparable results (**16ia–16ja**). Moreover, as the precursor of **16ja** was employed as a 1:1 *E:Z* mixture (see Supporting Information for details), we believe that the geometrical purity of the starting material is inconsequential to the reaction outcome.^[91]

Interestingly, trisubstituted alkenes did not give the expected mesylate product but the corresponding alcohol instead (**16 lb**, **16 mb**). Although *trans* selectivity for the cyclopropanation was excellent and yields were typically higher, the reaction showed no selectivity over a third stereogenic center in the cyclopropylcarbinyl position (**16 mb**), possibly a result of the intermediacy of a highly stabilized tertiary cyclopropylcarbinyl cation.^[61]

The syntheses of cyclopropanes are often mediated by metals,^[62-64] but metal free approaches are also known^[65,66] such as the Johnson–Corey–Chaykovsky reaction.^[67,68] Several reviews have been published.^[69–71] Certain α -cyclopropyl ketones can be generated from similar starting materials by epoxidizing the alkene using NBS under strongly basic conditions, followed by deprotonation of the ketone.^[72] However, those approaches suffer from reproducibility issues^[73] and are limited to the formation of the alcohol.^[92] Herein, we were able to show that the oxidative umpolung approach allows the decoration of the cyclopropylcarbonyl

cation with a variety of nucleophiles in good yields and exclusive *trans* selectivities using trimethylsilyl trifluoromethanesulfonate (TMSOTf) as the activator (Scheme 5). Halides, such as chloride or iodide (**16ad–16ae**), but also oxygen-(**16ah**) and sulfur-nucleophiles (**16ag**) were found to be competent reaction partners. We were also pleased to isolate the fluorinated product **16ac** in very good yield and excellent diastereoselectivity.

At this stage, we were intrigued by the possibility of alternative rearrangements using other silvlenol ether scaffolds. Because our experiments indicate the presence of the non-classical cyclopropylcarbinyl carbocation, we were wondering if an enolonium species reminiscent to the 2-norbornyl cation would lead to similar results (Scheme 6). The 2norbornyl cation has been part of one of the most famous (and long) debates in organic chemistry.^[74-76] When compound 17b was submitted to similar reaction conditions again a cyclopropanated product was observed in good yield: the nortricyclene 19b (Scheme 6).^[77-80] Nortricyclene formation is often observed when non-classical 2-norbornyl carbocations are generated (compound 31). This is believed to result from a γ -elimination of the non-classical cation,^[78–82] which can be interpreted as a metastable protonated cyclopropane. It is also interesting to note that this cation has not only been generated by halide abstraction of a 2-halonorbornyl precursor but also by protonation of nortricyclene.^[83] This reaction is applicable to a broad array of different ketones. Importantly, the hypothetical side-product 32 and the rearranged product 33 were mostly absent from the reaction mixture.

Furthermore, we were pleased to find that 2-norbornyl amides undergo oxidative cyclopropanation through amide umpolung chemistry using 2-iodo pyridine, trifluoromethanesulfonic anhydride, and pyridine-*N*-oxide (PNO) subsequently.^[84–87] Even though a pyridine base is used for this approach, elimination towards the norbornene derivative **36** represents only a minor reaction pathway (Scheme 7). This result showcases the aforementioned tenet of non-classical



Scheme 5. Nucleophile scope of the oxidative α -cyclopropanation of linear ketone derivatives. All yields refer to the pure, isolated *trans* isomer, unless otherwise stated. Reactions conducted on a 0.2 mmol scale, see Supporting Information for reaction conditions.

© 2020 The Authors. Published by Wiley-VCH GmbH

GmbH www.angewandte.org These are not the final page numbers!





Scheme 6. Ketone scope for the cyclopropanation *via* non-classical 2norbornyl cations. All yields refer to pure, isolated material. Reactions conducted on a 0.2 mmol scale, see Supporting Information for reaction conditions.



Scheme 7. Nortricyclene synthesis by amide activation. All yields refer to pure, isolated material. Reactions conducted on a 0.2 mmol scale, see Supporting Information for reaction conditions.

carbocations, whereby different precursors can yield the same product.

In conclusion, we have demonstrated that 2-electron oxidative umpolung is a valuable concept for the metal-free construction of cyclopropanes. Our results strongly suggest the intermediacy of non-classical carbocations and rely on the unusual stability of the intermediates involved.

Acknowledgements

This work was supported by the FWF (Project P32206) and the European Research Council Horizon 2020 (ERC CoG 682002 VINCAT). We thank the University of Vienna for continued support of our research programs. Dr. H. Kählig (University of Vienna) is acknowledged for expert assistance with NMR experiments and Ing. Alexander Roller (FH) for single-crystal X-ray diffraction studies.

Conflict of interest

The authors declare no conflict of interest.

Keywords: amide activation \cdot hypervalent iodine \cdot non-classical carbocation \cdot umpolung

- [1] D. Seebach, E. J. Corey, J. Org. Chem. 1975, 40, 231–237.
- [2] D. Seebach, Angew. Chem. Int. Ed. Engl. 1979, 18, 239–258; Angew. Chem. 1979, 91, 259–278.
- [3] E. J. Corey, D. Seebach, Angew. Chem. Int. Ed. Engl. 1965, 4, 1075–1077; Angew. Chem. 1965, 77, 1134–1135.
- [4] N. Zinin, Ann. Pharm. 1839, 31, 329-332.
- [5] H. Stetter, M. Schreckenberg, Angew. Chem. Int. Ed. Engl. 1973, 12, 81–81; Angew. Chem. 1973, 85, 89–89.
- [6] H. Stetter, Angew. Chem. Int. Ed. Engl. 1976, 15, 639-647; Angew. Chem. 1976, 88, 695-704.
- [7] X. Bugaut, F. Glorius, Chem. Soc. Rev. 2012, 41, 3511.
- [8] O. Miyata, T. Miyoshi, M. Ueda, ARKIVOC 2012, 60.
- [9] D. J. Rawlinson, G. Sosnovsky, *Synthesis* **1973**, 567–603.
- [10] W. Treibs, M. Weissenfels, Chem. Ber. 1960, 93, 1374-1379.
- [11] M. E. Kuehne, T. J. Giacobbe, J. Org. Chem. 1968, 33, 3359– 3369.
- [12] S. Moon, H. Bohm, J. Org. Chem. 1972, 37, 4338-4340.
- [13] R. Criegee, K. Klonk, Justus Liebigs Ann. Chem. 1949, 564, 1–9.
- [14] H. L. Riley, J. F. Morley, N. A. C. Friend, J. Chem. Soc. 1932, 1875.
- [15] D. H. R. Barton, J.-C. Blazejewski, B. Charpiot, J.-P. Finet, W. B. Motherwell, M. T. B. Papoula, S. P. Stanforth, J. Chem. Soc. Perkin Trans. 1 1985, 2667.
- [16] A. Hall, K. Jones, T. Jones, N. Killeen, R. Pörzig, P. Taylor, S. Yau, N. Tomkinson, *Synlett* 2006, 3435–3438.
- [17] N. Takeda, M. Furuishi, Y. Nishijima, E. Futaki, M. Ueda, T. Shinada, O. Miyata, Org. Biomol. Chem. 2018, 16, 8940–8943.
- [18] T. Miyoshi, T. Miyakawa, M. Ueda, O. Miyata, Angew. Chem. Int. Ed. 2011, 50, 928–931; Angew. Chem. 2011, 123, 958–961.
- [19] T. Miyoshi, S. Sato, H. Tanaka, C. Hasegawa, M. Ueda, O. Miyata, *Tetrahedron Lett.* 2012, 53, 4188–4191.
- [20] C. Tanyeli, Ç. Iyigün, Tetrahedron 2003, 59, 7135-7139.
- [21] N. K. Dunlap, M. R. Sabol, D. S. Watt, *Tetrahedron Lett.* 1984, 25, 5839–5842.
- [22] Y. Wei, S. Lin, H. Xue, F. Liang, B. Zhao, Org. Lett. 2012, 14, 712-715.
- [23] X. Huang, N. Maulide, J. Am. Chem. Soc. 2011, 133, 8510-8513.
- [24] A. J.-D. Lauriers, C. Y. Legault, Asian J. Org. Chem. 2016, 5, 1078–1099.
- [25] A. Flores, E. Cots, J. Bergès, K. Muñiz, Adv. Synth. Catal. 2019, 361, 2–25.
- [26] A. Yoshimura, V. V. Zhdankin, Chem. Rev. 2016, 116, 3328– 3435.
- [27] T. Dohi, Y. Kita, *Iodine Chemistry and Applications*, Wiley, Hoboken, **2014**, pp. 103–157.
- [28] V. V. Zhdankin, ARKIVOC 2009, 1.
- [29] V. V. Zhdankin, *Hypervalent Iodine Chemistry*, Wiley, Chichester, **2014**.
- [30] E. Zbiral, M. Rasberger, Tetrahedron 1969, 25, 1871-1874.
- [31] F. Mizukami, M. Ando, T. Tanaka, J. Imamura, Bull. Chem. Soc. Jpn. 1978, 51, 335–336.
- [32] R. M. Moriarty, O. Prakash, M. P. Duncan, R. K. Vaid, H. A. Musallam, J. Org. Chem. 1987, 52, 150–153.

www.angewandte.org

© 2020 The Authors. Published by Wiley-VCH GmbH

These are not the final page numbers!

- [33] G.F. Koser, A.G. Relenyi, A.N. Kalos, L. Rebrovic, R.H. Wettach, J. Org. Chem. 1982, 47, 2487–2489.
- [34] V. V. Zhdankin, M. Mullikin, R. Tykwinski, B. Berglund, R. Caple, N. S. Zefirov, A. S. Koz'min, J. Org. Chem. 1989, 54, 2605-2608.
- [35] N. S. Zefirov, V. V. Zhdankin, Y. V. Dan'kov, V. D. Sorokin, V. N. Semerikov, A. S. Koz'min, R. Caple, B. A. Berglund, *Tetrahedron Lett.* **1986**, *27*, 3971–3974.
- [36] R. Moriarty, O. Prakash, M. P. Duncan, J. Chem. Soc. Perkin Trans. 1 1987, 53, 559–561.
- [37] O. S. Shneider, E. Pisarevsky, P. Fristrup, A. M. Szpilman, Org. Lett. 2015, 17, 282–285.
- [38] M. J. Galligan, R. Akula, H. Ibrahim, Org. Lett. 2014, 16, 600-603.
- [39] A. H. Abazid, B. J. Nachtsheim, Angew. Chem. Int. Ed. 2020, 59, 1479–1484; Angew. Chem. 2020, 132, 1495–1500.
- [40] A.-A. Guilbault, B. Basdevant, V. Wanie, C. Y. Legault, J. Org. Chem. 2012, 77, 11283–11295.
- [41] K. N. Parida, G. K. Pathe, S. Maksymenko, A. M. Szpilman, *Beilstein J. Org. Chem.* 2018, 14, 992–997.
- [42] S. Arava, J. N. Kumar, S. Maksymenko, M. A. Iron, K. N. Parida, P. Fristrup, A. M. Szpilman, *Angew. Chem. Int. Ed.* **2017**, *56*, 2599–2603; *Angew. Chem.* **2017**, *129*, 2643–2647.
- [43] S. Beaulieu, C. Y. Legault, Chem. Eur. J. 2015, 21, 11206-11211.
- [44] R. M. Moriarty, H. Hu, S. C. Gupta, *Tetrahedron Lett.* 1981, 22, 1283–1286.
- [45] K. Freudenberg, Sitzungsber. Heidelb. Akad. Wiss. Math.-Naturwiss. 1927, 10, 3.
- [46] D. J. Cram, J. Am. Chem. Soc. 1949, 71, 3863-3870.
- [47] J. Li, A. Bauer, G. Di Mauro, N. Maulide, Angew. Chem. Int. Ed. 2019, 58, 9816–9819; Angew. Chem. 2019, 131, 9921–9924.
- [48] S. M. Banik, J. W. Medley, E. N. Jacobsen, *Science* 2016, 353, 51– 54.
- [49] U. Farid, F. Malmedy, R. Claveau, L. Albers, T. Wirth, Angew. Chem. Int. Ed. 2013, 52, 7018–7022; Angew. Chem. 2013, 125, 7156–7160.
- [50] C. Pardo, M. Charpentier-Morize, J. Chem. Soc. Chem. Commun. 1982, 1037–1039.
- [51] J. P. Bégué, M. Charpentier-Morize, Acc. Chem. Res. 1980, 13, 207–212.
- [52] A. Bauer, N. Maulide, Org. Lett. 2018, 20, 1461-1464.
- [53] A. Bauer, N. Maulide, *Tetrahedron* **2018**, *74*, 6883–6889.
- [54] E. Stefan, R. E. Taylor, Tetrahedron Lett. 2015, 56, 3416-3419.
- [55] J. D. Roberts, R. H. Mazur, J. Am. Chem. Soc. 1951, 73, 2509– 2520.
- [56] J. D. Roberts, W. Bennett, R. Armstrong, J. Am. Chem. Soc. 1950, 72, 3329–3333.
- [57] D. N. Kevill, M. A. Abduljaber, J. Org. Chem. 2000, 65, 2548– 2554.
- [58] P. C. Myhre, G. G. Webb, C. S. Yannoni, J. Am. Chem. Soc. 1990, 112, 8992–8994.
- [59] G. A. Olah, G. K. Surya Prakash, G. Rasul, J. Am. Chem. Soc. 2008, 130, 9168–9172.
- [60] N. J. Demjanov, M. Luschnikov, J. Russ. Phys. Chem. Soc. 1903, 35, 26–42.
- [61] G. A. Olah, K. K. Laali, Q. Wang, G. K. S. Prakash, Onium Ions, Wiley, New York, 1998.
- [62] E. J. T. Phipps, T. Rovis, J. Am. Chem. Soc. 2019, 141, 6807– 6811.
- [63] H. E. Simmons, R. D. Smith, J. Am. Chem. Soc. 1958, 80, 5323 5324.
- [64] H. Ikeda, K. Nishi, H. Tsurugi, K. Mashima, Chem. Sci. 2020, 11, 3604–3609.

- [65] C. D. Papageorgiou, M. A. Cubillo de Dios, S. V. Ley, M. J. Gaunt, Angew. Chem. Int. Ed. 2004, 43, 4641–4644; Angew. Chem. 2004, 116, 4741–4744.
- [66] J. Lee, D. Ko, H. Park, E. J. Yoo, Chem. Sci. 2020, 11, 1672-1676.
- [67] A. W. Johnson, R. B. LaCount, J. Am. Chem. Soc. 1961, 83, 417– 423.
- [68] E. J. Corey, M. Chaykovsky, J. Am. Chem. Soc. 1965, 87, 1353– 1364.
- [69] C. Ebner, E. M. Carreira, Chem. Rev. 2017, 117, 11651-11679.
- [70] H. Pellissier, *Tetrahedron* **2008**, *64*, 7041–7095.
- [71] E. M. D. Allouche, A. B. Charette, *Synthesis* **2019**, *51*, 3947–3963.
- [72] L. Dechoux, M. Ebel, L. Jung, J. F. Stambach, *Tetrahedron Lett.* 1993, 34, 7405–7408.
- [73] J. Cossy, N. Blanchard, C. Meyer, Eur. J. Org. Chem. 2001, 339– 348.
- [74] T. P. Nevell, E. de Salas, C. L. Wilson, J. Chem. Soc. 1939, 1188– 1199.
- [75] S. Winstein, D. S. Trifan, J. Am. Chem. Soc. 1949, 71, 2953-2953.
- [76] H. C. Brown, *The Nonclassical Ion Problem*, Springer US, Boston, 1977.
- [77] S. Winstein, E. Clippinger, R. Howe, E. Vogelfanger, J. Am. Chem. Soc. 1965, 87, 376–377.
- [78] K. Alder, H. K. Schäfer, H. Esser, H. Krieger, R. Reubke, Justus Liebigs Ann. Chem. 1955, 593, 23–35.
- [79] K. Alder, F. Brochhagen, Chem. Ber. 1954, 87, 167-179.
- [81] C. B. Kelly, A. M. Colthart, B. D. Constant, S. R. Corning, L. N. E. Dubois, J. T. Genovese, J. L. Radziewicz, E. M. Sletten, K. R. Whitaker, L. J. Tilley, *Org. Lett.* **2011**, *13*, 1646–1649.
- [82] X. Creary, A. Heffron, G. Going, M. Prado, J. Org. Chem. 2015, 80, 1781–1788.
- [83] G. A. Olah, Acc. Chem. Res. 1976, 9, 41-52.
- [84] C. R. Gonçalves, M. Lemmerer, C. J. Teskey, P. Adler, D. Kaiser, B. Maryasin, L. González, N. Maulide, *J. Am. Chem. Soc.* 2019, 141, 18437–18443.
- [85] D. Kaiser, C. J. Teskey, P. Adler, N. Maulide, J. Am. Chem. Soc. 2017, 139, 16040-16043.
- [86] A. Bauer, N. Maulide, Chem. Sci. 2019, 10, 9836-9840.
- [87] R. Da Costa, M. Gillard, J. B. Falmagne, L. Ghosez, J. Am. Chem. Soc. 1979, 101, 4381–4383.
- [88] Rehybridization during the nucleophilic attack on one of the two enantiotopic carbon atoms in the achiral non-classical carbocation might cause a delay in the formation of the exocyclically labeled compound.
- [89] The conditions later used for the nucleophile scope (Scheme 5) were employed here.
- [90] We do not know if either **26/27** or **28** (or both) are minima on the potential energy surface.
- [91] When the central methylene group of the silylenol ether substrate was decorated with two methyl groups, only traces of the desired cyclopropane were observed (crude ¹H-NMR analysis). It is possible that a [1,2]-methyl shift took place, although no products could be isolated from the complex mixture.
- [92] Please note that the assignment of the obtained diastereoisomer was incorrect in reference [62]. In reference [63] strong evidence that the *trans* (and not the *cis*) isomer is obtained is provided.

Manuscript received: May 23, 2020

Revised manuscript received: June 11, 2020 Version of record online:

www.angewandte.org

GDCh



Communications



Communications



An impossible triangle? Iodine(III) promotes the α -cyclopropanation of simple ketone derivatives. Mechanistic studies reveal that an unusual intermediate might be involved.

6 www.angewandte.org

 $\textcircled{\sc c}$ 2020 The Authors. Published by Wiley-VCH GmbH

Angew. Chem. Int. Ed. 2020, 59, 1-6

These are not the final page numbers!