

Direct Catalytic Anti-Markovnikov Hydroetherification of Alkenols

David S. Hamilton and David A. Nicewicz*

Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599-3290, United States


Supporting Information

ABSTRACT: A direct intramolecular anti-Markovnikov hydroetherification reaction of alkenols is described. By employing catalytic quantities of commercially available 9-mesityl-10-methylacridinium perchlorate and 2-phenyl-malononitrile as a redox-cycling source of a H-atom, we report the anti-Markovnikov hydroetherification of alkenes with complete regioselectivity. In addition, we present results demonstrating that this novel catalytic system can be applied to the anti-Markovnikov hydrolactonization of alkenoic acids.


The development of catalytic protocols for the direct addition of heteroatom nucleophiles to alkenes has been an area of intense study over the past two decades.¹ The vast majority of these methods give rise to primarily Markovnikov-type addition products. Given the challenges associated with reversal of innate alkene polarization, there are comparatively few methods that allow for the direct anti-Markovnikov addition of nucleophiles to olefins.^{1,2} Although Hartwig^{3,4} and Grubbs⁵ have demonstrated transition metal catalyst systems for the anti-Markovnikov addition of amines and water, respectively, to alkenes, success has been limited to terminal styrenes. We were drawn to the possibility that single electron oxidation of olefins to their respective cation radicals could provide a basis to develop a general catalyst system for a range of heteroatom nucleophiles with unactivated alkenes.^{6–9} Herein, we report the direct intramolecular anti-Markovnikov addition of alcohols to alkenes via a unique two-component single electron photoredox system. This transformation provides a reactivity profile that complements traditional Markovnikov-based Brønsted acid catalyzed reactions (Scheme 1).^{10,11}

Seminal work from Arnold¹² and Gassman^{13,14} provided the first evidence for cation radical-mediated anti-Markovnikov reactivity. Arnold further characterized the initial nucleophile-cation radical adduct as the three-membered intermediate **8** by density functional theory calculations.¹⁵ It is likely that the observed anti-Markovnikov selectivity results from the rupture of the weaker of the two C–X bonds, giving rise to the more stable radical intermediate **9** (Scheme 2). Additionally, Gassman and Arnold have each demonstrated that single electron photooxidants can serve as effective single electron oxidants to access reactive olefin cation radicals (**7**). To date, however, this method remains significantly limited in scope and requires nearly stoichiometric quantities of the photooxidant that can often result in oxidant incorporation into the reaction products as well as undesired side reactivity.¹⁶ Truly catalytic

Scheme 1. Divergent Regioselectivity in Alkene Addition Reactions



Scheme 2. Anti-Markovnikov Reactivity of Cation Radicals



photosensitized anti-Markovnikov alcohol additions are limited to 1,1-diarylethylenes.^{17–19}

After analysis of this body of literature, we proposed that the critical step preventing the development of catalytic protocols was the fate of putative radical intermediate **9**. We hypothesized that using an alternative class of photooxidants might enable a general approach to this problem. We recognized that good candidates for a single electron redox catalyst should (i) exhibit nearly complete redox reversibility, (ii) be capable of oxidizing alkenes in the range +1.0 to +2.0 V, and (iii) be positively charged to minimize unproductive back electron transfer to **7** via minimization of Coulombic attraction in the reduced (neutral) form of the catalyst.

Reports of commercially available 9-mesityl-10-methylacridinium perchlorate (**2**), first employed by Fukuzumi et al., drew our attention as a photooxidant for this application.²⁰ Given the

Received: September 28, 2012

Published: October 31, 2012

acridinium moiety's strong absorption band in the visible region ($\lambda = 430$ nm), high excited state oxidizing power ($E_{1/2}^{\text{red}*} = +2.06$ V vs SCE),²¹ and utility in a number of reported transformations relying on single electron transfer pathways,^{22,23} we predicted that cation radicals could be conveniently generated from an electronically diverse range of alkenes. Additionally, the reduced form of the acridinium catalyst (**11**) is a moderate single electron reductant ($E_{1/2}^{\text{ox}} = -0.57$ V vs SCE)²¹ that we presumed would be capable of return electron transfer to radical intermediate **12**.

As a starting point, we focused on the development of a catalytic system for the direct intramolecular anti-Markovnikov hydroetherification of alkenols.^{2,24} To date, Mizuno has reported the only known direct anti-Markovnikov hydroetherification reaction of alkenols which is believed to proceed via exciplex formation and is limited to diphenylethylene alkenes.¹⁸ To begin, we subjected alkenol **4** to 5 mol % of catalyst **2** in degassed 1,2-dichloroethane (DCE) under irradiation with 450 nm LEDs. The anti-Markovnikov adduct, tetrahydrofuran **6**, was obtained, albeit in low yields (36% yield, Table 1, entry 1), though significantly higher than when

Table 1. Catalyst Optimization Studies^a

entry	conditions	conversion ^b	yield ^b	6:5 ^b
1	standard conditions	83%	36%	>20:1
2 ^{c,d}	0.2 equiv of 9,10-dicyanoanthracene instead of 2	21%	5%	>20:1
3 ^{c,e}	0.5 equiv of 1-cyanonaphthalene instead of 2	47%	15%	>20:1
4	with 0.5 equiv of <i>N</i> -hydroxyphthalimide	48%	41%	>20:1
5	with 0.5 equiv of 9-phenylfluorene	78%	51%	>20:1
6	with 0.5 equiv of PhCH(CN) ₂ (3)	89%	73%	>20:1
7 ^f	no photooxidant	<5%	<5%	—
8 ^f	no light	<5%	<5%	—
9 ^{f,g}	Ru(bpy) ₃ Cl ₂ instead of 2	<5%	<5%	—

^aReactions irradiated with a 15 W 450 nm LED flood lamp.

^bDetermined by ¹H NMR analysis. ^cIrradiated with 10 × 8 W T5 fluorescent bulbs (output > 290 nm). ^dBenzene as solvent. ^eMeCN as solvent with 0.5 equiv of biphenyl. ^fWith 0.5 equiv of **3**. ^gWith 1.0 equiv of methyl viologen.

employing the cyanoarene photooxidants (entries 2 and 3) used by Mizuno and Gassman in their pioneering studies. No trace of the Markovnikov adduct (**5**) was observed and conversion of the starting alkenol was relatively high (83%), but yields were significantly diminished by extensive unidentifiable byproduct formation that likely arose from competing radical processes.

After evaluation of a number of known single electron photooxidants failed to afford synthetically useful yields of the desired adduct, we felt that a distinctly different approach to this problem was required. Speculating that the reduction of radical **9** was still limiting reactivity, we hypothesized that employing a H-atom donor could facilitate this process while simultaneously serving as a single electron redox mediator.

Potential H-atom donors were selected on the basis of their respective homolytic bond dissociation energies (BDE). To



ensure exothermic H-atom transfer, we limited our survey of potential H-atom redox catalysts to moieties possessing R–H bonds with BDE <90 kcal/mol (Table 1, entries 3–5).²⁵ Though 0.5 equiv of either *N*-hydroxyphthalimide (BDE = 87 kcal/mol, entry 3) or 9-phenylfluorene (BDE = 74 kcal/mol, entry 4) gave modest increases in reaction efficiency, we were pleased to find that 2-phenylmalononitrile (**3**, BDE = 77 kcal/mol) furnished anti-Markovnikov adduct **6** in 73% yield (entry 5) with no trace of the undesired Markovnikov regioisomer. Further control experiments demonstrate that both the acridinium photocatalyst and light are necessary for reactivity (entries 6, 7).²⁶ The utility of the acridinium catalyst as a single electron photooxidant is underscored when compared directly with the frequently employed Ru(bpy)₃²⁺,²⁷ which failed to give any of the desired product (entry 8). This result demonstrates the advantage of acridinium catalysts as visible light single electron photooxidants and should allow for greater latitude in potential substrates with alkenes possessing oxidation potentials ranging up to +2.0 V.

Our mechanistic hypothesis outlined in Scheme 3 proposes that, following H-atom transfer from **3**, the resulting radical **13** could serve as an oxidant for radical **11**, regenerating the ground state photooxidation catalyst (**2**). Following this redox event, proton transfer would regenerate the H-atom donor (**3**) and furnish the desired product.

Having identified a viable catalyst system, we investigated the scope of the intramolecular anti-Markovnikov hydroalkoxylation of alkenols (Table 2). Electronically distinct styrenes (entries 1–3) ranging from electron-rich (4-(MeO)C₆H₄, entry 1; 80% yield) to electron-deficient (4-ClC₆H₄, entry 2; 60% yield) provided good yields of the desired 5-*exo* adducts. Additionally, Thorpe–Ingold assistance is not required in the backbone of the molecule, as the substrate in entry 3, which lacked the geminal dimethyl substituent, gave nearly identical levels of reaction efficiency (82% yield) as in entry 1 (80% yield). Furthermore, the mild reaction conditions are highlighted in entry 6, where a silyl-protected alcohol remains unperturbed by the cyclization conditions. A gram-scale reaction of the alkenol in entry 4 produced the expected tetrahydrofuran product in 77% isolated yield. Though long reaction times are required for most substrates, significantly shorter reaction times are possible by increasing the amount of **3** employed.²⁸

In addition to the formal 5-*exo* cyclization mode (entries 1–6), other ring closure types were possible. The alkenol in entry 8 underwent 6-*exo* cyclization to furnish the anticipated disubstituted tetrahydropyran adduct in 68% yield and 2.5:1 d.r. Treatment of β -citronellol to the catalyst conditions resulted in 7-*exo* cyclization in modest but reproducible yields (46% yield, 1.2:1 d.r., entry 9). The reactions in entries 8 and 9 required 2.0 equiv of PhCH(CN)₂ to avoid longer reaction times. Given their high oxidation potentials, monosubstituted alkenes are inaccessible by this catalyst system; however, expansion of the substrate scope will be a focus of future efforts through catalyst development.

It is particularly noteworthy that all of the reactions in Table 2 furnished the anti-Markovnikov hydroalkoxylation adducts exclusively. To emphasize the unique regioselectivity of this process, a direct comparison of alkene reactivity with cation radicals or Brønsted acids is depicted in eqs 1 and 2. Alkenol **14** is known to undergo Brønsted acid assisted Markovnikov hydroetherification to furnish tetrahydropyran **15**, while tetrahydrofuran **16** is obtained exclusively using our catalytic



that this unique approach to managing open-shell pathways holds promise to develop additional anti-Markovnikov hydrofunctionalization reactions. Studies further investigating this transformation as well as other heteroatom nucleophiles in this context are currently underway.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental procedures and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

nicewicz@unc.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors are grateful to J. Roberts for aid in obtaining the cyclic voltammograms of the alkenol substrates and C. Price for the preparation of some of these substrates. The project described was supported by The University of North Carolina at Chapel Hill, Award No. R01 GM098340 from the National Institute of General Medical Sciences and an Eli Lilly New Faculty Award.

■ REFERENCES

- (1) For comprehensive reviews on catalytic functionalization of olefins, see: (a) Beller, M.; Seavad, J.; Tillack, A.; Jiao, H. *Angew. Chem., Int. Ed.* **2004**, *43*, 3368. (b) Hintermann, L. *Top. Organomet. Chem.* **2010**, *31*, 123. (c) Müller, T. E.; Hultsch, K. C.; Yus, M.; Foubelo, F.; Tada, M. *Chem. Rev.* **2008**, *108*, 3795.
- (2) Catalytic anti-Markovnikov additions of nucleophiles to olefins have been described as one of the “top 10 challenges for catalysis.” See: Haggin, J. *Chem. Eng. News* **1993**, *71*, 23.
- (3) Utsunomiya, M.; Kuwano, R.; Kawatsura, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2003**, *125*, 5608.

- (4) Utsunomiya, M.; Kuwano, R.; Kawatsura, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2004**, *126*, 2702.
- (5) Dong, G.; Teo, P.; Wickens, Z. K.; Grubbs, R. H. *Science* **2011**, *333*, 1609.
- (6) For a review on reactivity patterns of cation-radicals, see: Schmittel, M.; Burghart, A. *Angew. Chem., Int. Ed.* **1997**, *36*, 2550.
- (7) For electrochemical intramolecular etherifications mediated by cation radicals, see: (a) Sutterer, A.; Moeller, K. D. *J. Am. Chem. Soc.* **2000**, *122*, 5636. (b) Moeller, K. D. *Synlett* **2009**, *8*, 1208.
- (8) For hydroetherifications of alkene cation radicals generated by heterolysis of β -substituted radicals, see: (a) Crich, D.; Ranganathan, K.; Neelamkavil, S.; Huang, X. *J. Am. Chem. Soc.* **2003**, *125*, 7942. (b) Crich, D.; Ranganathan, K. *J. Am. Chem. Soc.* **2005**, *127*, 9924. (c) Crich, D.; Shirai, M.; Brebion, F.; Rumthao, S. *Tetrahedron* **2006**, *62*, 6501. (d) Crich, D.; Brebion, F.; Suk, D. H. *Top. Curr. Chem.* **2006**, *263*, 1.
- (9) For a largely theoretical perspective on intramolecular nucleophilic capture of cation radicals by tethered alcohols, see: Roth, H. D.; Herberich, T.; Sauers, R. R.; Wang, H. *Tetrahedron* **2006**, *62*, 6471.
- (10) Jeong, Y.; Kim, D.; Choi, Y.; Ryu, J. *Org. Biomol. Chem.* **2011**, *9*, 374–378.
- (11) Smith, M. B.; March, J. *March's Advanced Organic Chemistry*; John Wiley and Sons: New York, 2001; p 993.
- (12) Neuteufel, R. A.; Arnold, D. R. *J. Am. Chem. Soc.* **1973**, *95*, 4080.
- (13) Gassman, P. G.; Bottorff, K. J. *Tetrahedron Lett.* **1987**, *28*, 5449.
- (14) Gassman, P. G.; Bottorff, K. J. *J. Am. Chem. Soc.* **1987**, *109*, 7547.
- (15) Arnold, D. R.; Chan, M. S. W.; McManus, K. A. *Can. J. Chem.* **1996**, *74*, 2143.
- (16) McManus, K. A.; Arnold, D. A. *Can. J. Chem.* **1995**, *73*, 2158.
- (17) Mizuno, K.; Nakanishi, I.; Ichinose, N.; Otsuji, Y. *Chem. Lett.* **1989**, *41*, 1095.
- (18) Mizuno, K.; Tamai, T.; Nishiyama, T.; Tani, K.; Sawasaki, M.; Otsuji, Y. *Angew. Chem., Int. Ed.* **1994**, *33*, 2113.
- (19) Asaoka, S.; Kitazawa, T.; Wada, T.; Inoue, Y. *J. Am. Chem. Soc.* **1999**, *121*, 8486.
- (20) Fukuzumi, S.; Kotani, H.; Ohkubo, K.; Ogo, S.; Tkachenko, N. V.; Lemmetyinen, H. *J. Am. Chem. Soc.* **2004**, *126*, 1600.
- (21) Ohkubo, K.; Mizushima, K.; Iwata, R.; Souma, K.; Suzuki, S.; Fukuzumi, S. *Chem. Commun.* **2010**, *46*, 601.
- (22) Kotani, H.; Ohkubo, K.; Fukuzumi, S. *J. Am. Chem. Soc.* **2004**, *126*, 15999.
- (23) Ohkubo, K.; Nanjo, T.; Fukuzumi, S. *Org. Lett.* **2005**, *7*, 4265.
- (24) Sanford, M. S.; Groves, J. T. *Angew. Chem., Int. Ed.* **2004**, *43*, 588.
- (25) Please see the Supporting Information for a more extensive list of the H-atom donors that were evaluated.
- (26) The BF₄[−] salt of catalyst **2** can be employed without significant variation in yields.
- (27) For reviews, see: (a) Yoon, T. P.; Ischay, M. A.; Du, J. *Nat. Chem.* **2010**, *2*, 527. (b) Narayanam, J. M. R.; Stephenson, C. R. J. *Chem. Soc. Rev.* **2011**, *40*, 102.
- (28) Please see the Supporting Information for details.
- (29) Seitz, M.; Reiser, O. *Curr. Opin. Chem. Biol.* **2005**, *9*, 285.