View Article Online View Journal



# Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: X. Wei, L. Wang, S. Du, L. Wu and Q. Liu, *Org. Biomol. Chem.*, 2016, DOI: 10.1039/C5OB02121A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/obc

## RSCPublishing

### COMMUNICATION

Journal Name

Cite this: DOI: 10.1039/xoxxooooox

## Visible-Light Photoredox Intramolecular Difluoroacetamidation: Facile Synthesis of 3,3-Difluoro-2-Oxindoles from Bromodifluoroacetamides

Received ooth January 2012, Accepted ooth January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Published on 08 January 2016. Downloaded by UNIVERSITY OF NEBRASKA on 11/01/2016 12:54:13.

Xiao-Jing Wei, <sup>a</sup> Lin Wang, <sup>a</sup> Shao-Fu Du, <sup>a</sup> Li-Zhu Wu<sup>b</sup> and Qiang Liu<sup>a</sup>\*

We report an operationally simple, visible-light-driven protocol for intramolecular C-H difluoroacetamidation of arenes for the synthesis of biological relevant 3, 3-difluoro-2oxindoles at room temperature. Using fac-Ir(ppy)<sub>3</sub> as a photocatalyst and a 3W blue LED as a light source, an array of difluoroxindoles were prepared from rapidly available tertiary aryl bromodifluoroacetamides in moderate to excellent yields.

Fluorine-containing organic molecules can be extensively found in a number of biologically active natural products, pharmaceutically relevant candidates, agrochemical reagents, and functional materials.<sup>1</sup> It is an important and general strategy to introduce fluorine and fluorinated substitutes in the small molecule for structure-based drug development and pharmaceutical research. The fluorinated functional groups change their physiochemical properties significantly due to the enhanced chemical, physical, and metabolic stability of the fluorinated moiety.<sup>2</sup> Accordingly, there has been great interest and substantial effort in the synthesis of fluorine-containing small molecules.<sup>3</sup> Especially, various synthetic methods



Fig. 1. Examples of biologically significant molecules containing 3, 3-difluoro-2-oxindoles.

for the introduction of the gem-difluoromethylene group (CF<sub>2</sub>) into organic compounds that have been well developed are consider as useful synthesis strategy for modification of biologically active compounds.<sup>4</sup>

Derivatives of oxindoles are an extremely important class of compounds due to their frequent appearance in a variety of naturally occurring compounds and their wide range of biological activities. Such as these kinds of compounds have been shown to display antimicrobial, antiviral, anticancer, and antiinflammatory activities.<sup>5</sup> The 3, 3-difluoro-2-oxindole ring system are of significant biological importance as bioisoteric analogues for biological studies in development of potential medicinal agents. Up to now, numerous natural compounds containing the 3, 3-difluoro-2-oxindoles ring moieties revealed potential utilities in biomedical studies.<sup>6</sup> Thus, search for efficiently synthetic methodologies to prepare these substrates are very meaningful for the detailed studies of their therapeutic application.<sup>7</sup> Generally, these types of difluoroxindoles could be obtained via nucleophilic fluorination of the corresponding isatin derivatives with diethylaminosulfur trifluoride (DAST) or Deoxofluor,<sup>8a-8c</sup> or by electrophilic fluorination of indoles with Selectfluor or NFSI.<sup>8d</sup> It is also possible to access difluoroxindoles via Cu-mediated intramolecular cyclization of iododifluoroacetamides in moderate yield.<sup>9a</sup> Very recently, Buchwald et al. used Pa-catalyzed C-H difluoroalkylation and BrettPhos as crucial ligand to construct difluoroxindoles derivatives.9b These valuable protocols have proven to be efficient for the preparation this kind of compounds. However, instability of the requisite reagents and bad to modest functionalgroup tolerance, expensive fluorination reagents, the production of a large excess of wastes from reaction systems, the synthesis difficulty of the required starting materials, as well as the requirement for rather high reaction temperature et al. remain impediments of these approaches. Therefore the development of more efficient and practical strategies in these areas is still highly desirable.



Scheme 1. Different protocols for the syntheses of 3, 3-difluoro-2-oxindoles.

For the past few years, photocatalysis reactions, especially the use of visible light as a nearly non-polluting energy source to enable sustainable organic synthesis have attracted great interest.<sup>10</sup> Visiblelight photocatalysis reactions avoided the disadvantage of highenergy ultraviolet radiation, and as the high transparency of most organic compounds in visible-light region, it is helpful to minimize side reactions which is often associated with the employment of UV light. Photoredox catalysis with transitional metal complexes has emerged as a powerful tool for redox reactions due to its ability to cause efficient photo-induced electron transfer processes under mild conditions. For instance, fac-Ir(ppy)<sub>3</sub>, an Ir polypyridine complexe with high reducibility in the excited state has been well demonstrated by some valuable work that generated carbon-centered radicals from organohalides.<sup>11</sup> During the course of our investigations on the applications of photoredox catalysis in organic synthesis, 11h,12 we found a promising approach for synthesizing an array of substituted difluoroacetamidated arenes via intermolecular radical addition to arenes.<sup>128</sup> We speculated that this strategy might also be effective for intramolecular aminocarbonyldifluoromethyl radical providina addition to arenes to afford biologically relevant 3,3-difluoro-2oxindoles.13

Herein, We started our investigation by exploring the reaction of 2bromo-2,2-difluoro-*N*-methyl-*N*-phenylacetamide (**1a**) in various photochemical conditions toward the synthesis of 3,3-difluoro-2oxindoles. It was encouraging to see that Table 1. Optimization of reaction conditions



<sup>*a*</sup> Reaction conditions: a mixture of **1a** (0.20 mmol), base (0.24 mmol, 1.2 equiv), and photocatalyst (0.5 mol %) in solvent was irradiated with a 3 W blue LEDs at room temperature for 24 h <sup>*b*</sup> Isolated yield. <sup>*c*</sup> Irradiated without *fac*-Ir(ppy)<sub>3</sub>. <sup>*d*</sup> The reaction was conducted in dark. <sup>*e*</sup> N.D. = not detected. DMF = *N*, *N*- dimethylformamide

the aimed product 2a was obtained in 58% yield after 24 hours' irradiation (blue LEDs,  $\lambda$  = 450 nm) at room temperature when the reaction was performed in N,N-dimethylformamide (DMF) with Ru(bpy)<sub>3</sub>Cl<sub>2</sub>·6H<sub>2</sub>O as photocatalyst and Na<sub>2</sub>CO<sub>3</sub> as base (Table 1, entry 1). Several photocatalysts were applied to this reaction to replace Ru(bpy)<sub>3</sub>Cl<sub>2</sub>·6H<sub>2</sub>O (Table 1, entries 2-5). It was found that when Ru(phen)<sub>3</sub>Cl<sub>2</sub> was applied in this reaction conditions, the reaction yields rose to 62% (Table 1, entry 2), but when Re[phen(CO)<sub>3</sub>]NCS or organic photocatalyst TBA-eosinY was tested in the reaction (Table 1, entries 2-3), just trace amount of desired product was found. When fac-Ir(ppy)<sub>3</sub> was used in the reaction, the yields rose to 91% dramatically. Therefore we choose fac-Ir(ppy)<sub>3</sub> as the ideal photocatalyst. Other different bases including Na<sub>2</sub>HPO<sub>4</sub>, NaHCO<sub>3</sub>,  $K_2CO_3$ ,  $K_2HPO_4$  were then tested (Table 1, entries 6-10), and the best result was achieved when K<sub>2</sub>HPO<sub>4</sub> was introduced to the system. In this case we obtained the desired product 2a in 93% isolated yield. In addition, when the reaction was carried out in the absecne of fac-Ir(ppy)<sub>3</sub> or in the dark (Table 1, entry 11 and 12), no conversion could be observed, which indicated that the reaction was indeed a visiblelight-driven photoredox process.

With the optimized condition in hand, we examined the substrate scope of this visible-light-driven synthesis of 3,3-difluoro-2-oxindoles. A broad array of substituted 2-bromo-2,2-difluoro-*N*-phenylacetamides can efficiently convert to the corresponding 3,3-difluoro-2-oxindoles derivatives in moderate to excellent yields (Table 2, entries 1-14). When the electron-withdrawing groups such as fluoro, ethoxycarbonyl and methoxycarbonyl which located at *para* positions of the phenylacetamide, the aimed product **2b**, **2c**, **2d** can be produced

Table2 Scope survey of visible-light catalyzed synthesis of 3,3-difluoro-2-oxindoles.<sup>a, v</sup>

Published on 08 January 2016. Downloaded by UNIVERSITY OF NEBRASKA on 11/01/2016 12:54:13.

Journal Name





<sup>*a*</sup> Reaction conditions: a mixture of **1a** (0.20 mmol, 1.0 equiv),  $K_2$ HPO<sub>4</sub> (0.24 mmol, 1.2 equiv) and photocatalyst *fac*-Ir(ppy)<sub>3</sub> (0.5 mol %) in DMF (3 mL) was irradiated with a 3 W blue LEDs at room temperature for 24 h <sup>*b*</sup> Yield of isolated product.

smoothly in excellent yields (Table 2, entries 2-4). Moreover, this kind of reaction was successfully applied to the 2-bromo-2,2-difluoro-*N*-phenylacetamides with electron-donating groups at the arenes (Table 2, entries 5 and 6). The substrate with di-substituted aniline **1g** was also compatible with this new visible-light-driven cyclization approach, although the reaction proceeded less efficiently than mono-substituted

cases (Table 2, entry 7). In general, arenes bearing electron-withdrawing substituents afforded higher yields than those with electron-donating groups. The lower transformation of 2-bromo-2,2-difluoro-*N*-phenylacetamides with electron-donating groups may be attributed to the inefficient electron transfer between the excited photocatalyst and the substrates.

Next, we examined 2-bromo-2,2-difluoroacetamides with different amine moieties instead of *N*-methylaniline. The reactions of 2-bromo-2,2-difluoro-*N*-methyl-*N*-(naphthalen-2-yl)acetamide (1h) generated 2h as the only product (Table 2, entry 8). When tetrahydrobenzoazepine (1i) and tetrahydroquinoline (1j) were employed as the amine moiety of the substrate, tricyclic products 2i and 2j were formed in 79% and 92% yields, respectively (Table 2, entries 9 and 10). In addition, 2-bromo-2,2difluoro-*N*-phenyl-acetamides possessing *N*-protecting groups such as phenyl(1k), cyclohexyl(1l), 2-cyanoethyl (1m) and 2-hydroxyethyl (1n) also resulted in satisfactory yields, highlighting the potential of the method in organic synthesis.



Scheme 3. Mechanism investigation

Published on 08 January 2016. Downloaded by UNIVERSITY OF NEBRASKA on 11/01/2016 12:54:13.

In order to understand the reaction mechanism and to verify the generation of the 2,2-difluoro-*N*-methyl-*N*-phenylacetamide radical **3** at the initial step of the reaction, the radical inhibitors such as BHT and TEMPO were introduced into the photochemical system, and no 3,3-difluoro-2-oxindole **2a** was detected in both cases. To our delight, the adduct of 2,2-difluoro-*N*-methyl-*N*-phenylacetamide radical **3** with the radical inhibitors were detected by ESI-HRMS, thus providing straightforward evidence of the formation of the radical **3** (Scheme 3).

Based on the radical trapped experimental results and our previous work in this field,<sup>12</sup> we proposed a possible mechanism. Visible-light irradiation of the photoredox catalyst *fac*-Ir<sup>III</sup>(ppy)<sub>3</sub> produced the excited <sup>3</sup>[fac-lr<sup>III</sup>(ppy)<sub>3</sub>]\*, which can reduce 2-bromo-2,2difluorophenylacetamide 1a by single electron transfer (SET). The SET procedure generates electron-deficient radical **3** along with fac-Ir<sup>IV</sup>(ppy)<sub>3</sub>. Then intramolecular  $\pi$ -addition of the radical **3** produces a cyclohexadiene radical 4. The  $\alpha$ -proton of radical 4 is highly acidic<sup>14</sup> because it is activated by both the incorporation of the 2,5cyclohexadienyl radical moiety into a larger aromatic system and the negative inductive effect of difluoroacetamide. Therefore. deprotonation of 4 should be occurred efficiently and give radical anion 5 with the aid of K<sub>2</sub>HPO<sub>4</sub> . Finally, SET from radical anion 5 to fac-Ir<sup>IV</sup>(ppy)<sub>3</sub> affords the desired product **2a** and regenerates the photocatalyst.

In summary, we have developed a catalytic and efficient method for the construction of  $_{3,3}$ -difluoro-2-oxindoles *via* visible light photocatalysis for the first time. The highlight of this photo-reaction is the operational simplicity, avoidance of



#### Scheme 4. Proposed Mechanism Cycle

expensive fluorination reagent, complex ligand and high temperature. Moreover, this method is also an efficient method for the C-H difluoroalkylation with readily available starting materials.

We thank the financial support from NSFC (No. 21172102), the Ministry of Science and Technology of China (973 Program, 2013CB834804), and the Fundamental Research Funds for the Central Universities (Izujbky-2013-47).

#### Notes and references

<sup>*a*</sup> State Key Laboratory of Applied Organic Chemistry, Lanzhou University, 222 South Tianshui Road, Lanzhou 730000, P. R. China. Email: <u>liuqiang@lzu.edu.cn</u>

<sup>b</sup> Key Laboratory of Photochemical Conversion and Optoelectronic Materials, Technical Institute of Physics and Chemistry, University of Chinese Academy of Sciences, Haidian District, Beijing, 100190, P. R. China.

† Electronic Supplementary Information (ESI) available: Experimental procedures, characterization data, and copies of NMR spectra. See DOI: 10.1039/c000000x/

- (a) K. Müller, C. Faeh and F. Diederich, *Science*, 2007, **317**, 1881; (b)
  S. Purser, P. R. Moore, S. Swallow and V. Gouverneur, *Chem. Soc. Rev.*, 2008, **37**, 320; (c) V. D. Romanenko and V. P. Kukhar, *Chem. Rev.*, 2006, **106**, 3868; (d) T. Furuya, A. S. Kamlet and T. Ritter, *Nature*, 2011, **473**, 470; (e) K. L. Kirk, *Org. Process Res. Dev.*, 2008, **12**, 305; (f) R. Banks, B. Smart and J. Tatlow, *Organofluorine Chemistry: Principles and Commercial Applications* (Eds. R. E. Banks, B. E. Smart, J. C. Tatlow), Plenum Press, New York, 1994.
- (a) D. J. Adams and J. H. Clark, *Chem. Soc. Rev.*, 1999, 28, 225; (b)
  W. K. Hagmann, *Med. Chem.*, 2008, 51, 4359; (c) C. Hollingworth and V. Gouverneur, *Chem. Commun.*, 2012, 48, 2929; (d) F. Babudri, G. M. Farinola, F. Naso and R. Ragni, *Chem. Commun.*, 2007, 43, 1003; (b) O. A.Tomashenko and V. V. Grushin, *Chem. Rev.*, 2011, 111, 4475; (c) W. K. Hagmann, *J. Med. Chem.*, 2008, 51, 4359.
- (a) C. Alonso, E. M. de Marigorta, G. Rubiales and F. Palacios, *Chem. Rev.*, 2015, **115**, 1847; (b) J. Charpentier, N. Früh and A. Togni, *Chem. Rev.*, 2015, **115**, 650; (c) C. Ni, M. Hu and J. Hu, *Chem. Rev.*, 2015, **115**, 765; (d) X. Liu, C. Xu, M. Wang and Q. Liu, *Chem. Rev.*, 2015, **115**, 683; (e) (f) D. J. Adams and J. H. Clark, *J. Org. Chem.*,

Journal Name

2000, 65, 1456; (g) C. Chen, L. Chu, F.-L. Qing, J. Am. Chem. Soc., 2012, 134, 12454; (h) O. Marrec, T. Billard, J. P. Vors, S. Pazenok and B. R. Langlois, Adv. Synth. Catal., 2010, 352, 2831; (i) K. N. Hojczyk, P. Feng, C. Zhan and M.-Y. Ngai, Angew. Chem., Int. Ed., 2014, 53, 14559; (j) B. Bayarmagnai, C. Matheis, K. Jouvin and L. J. Goossen, Angew. Chem., Int. Ed., 2015, 54, 5753; (k) C. Huang, T. Liang, S. Harada, E. Lee and T. Ritter J. Am. Chem. Soc., 2012, 134, 6548; (1) Q. Qi, Q. Shen and L. Lu, J. Am. Chem. Soc., 2012, 134, 6548; (m) Q. Lefebvre, R. Pluta and M. Rueping, Chem. Commun., 2015, 51, 4394; (n) K. Fujikawa, Y. Fujioka, A. Kobayashi and H. Amii, Org. Lett., 2011, 13, 5560; (o) J. Wu, Y. Gu, X. Leng and Q. Shen, Angew. Chem., Int. Ed., 2015, 54, 7648; (p) T. Liang, C. N. Neumann and T. Ritter, Angew. Chem., Int. Ed., 2013, 52, 8214; (q) J. A. Kalow and A. G. Doyle, J. Am. Chem. Soc., 2010, 132, 3268; (r) Y. Ye, S. D. Schimler, P. S. Hanley and M. S. Sanford, J. Am. Chem. Soc., 2013, 135, 16292.

- (a) J. Hu and W. Zhang, Chem. Commun., 2009, 48, 7465; (b) Q.-Q. Min, Z. Yin, Z. Feng, W.-H. Guo and X. Zhang, J. Am. Chem. Soc., 2014, 136, 1230; (c) P. S. Fier and J. F. Hartwig, J. Am. Chem. Soc., 2012, 134, 5524; (d) M. Fedoryńsk, Chem. Rev., 2003, 103, 1099; (e) W. R. Dolbier, Jr. and M. A. Battiste, Chem. Rev., 2003, 103, 1071; (f) L. Li, F. Wang, C. Ni and J. Hu, Angew. Chem., Int. Ed., 2013, 52, 12390; (g) G. K. S. Prakash, S. K. Ganesh, J.-P. Jones, A. Kulkarni, K. Masood, J. K. Swabeck and G. A. Olah, Angew. Chem., Int. Ed., 2012, 51, 12090; (h) J. Wu, Y. Gu, X. Leng and Q. Shen, Angew. Chem., Int. Ed., 2015, 54, 7648; (i) X. Lin and Z. Weng, Org. Biomol. Chem., 2015, 13, 3432; (j) J.-Y. Wang, Y.-M. Su, F. Yin, Y. Bao, X. Zhang, Y.-M. Xu and X.-S. Wang Chem.Commun., 2014, 50, 4108; (k) X.-Y. Deng, J.-H. Lin, J. Zheng and J.-C. Xiao, Chem. Commun., 2015, 51, 8805.
- (a) A. B. Dounay and L. E. Overman, *Chem. Rev.*, 2003, **103**, 2945;
  (b) C. V. Galliford and K. A. Scheidt, *Angew. Chem., Int. Ed.*, 2007, **46**, 8748;
   (c) K. Shen, X. Liu, L. Lin and X. Feng, *Chem. Sci.*, 2012, **3**, 327;
   (d) B. M. Trost and M. K. Brennan, *Synthesis*, 2009, 3003;
   (e) C. V. Galliford and K. A. Scheidt, *Angew. Chem.*, 2007, **119**, 8902; *Angew. Chem. Int. Ed.*, 2007, **46**, 8748;
   (f) R. Dalpozzo, G. Bartoli, and G. Bencivenni, *Chem. Soc. Rev.*, 2012, **41**, 7247;
   (g) G. S. Singh and Z. Y. Desta, *Chem. Rev.*, 2012, **112**, 6104;
   (h) F. Zhou, Y.-L. Liu and J. Zhou, *Adv. Synth. Catal.*, 2010, **352**, 1381;
   (i) Vine, K. L.; Matesic, L.; Locke, J. M.; Ranson, M.; Skropeta, D. *Anti-Cancer Agents Med. Chem.*, 2009, **9**, 397.
- (a) N. Zhou, A. M. Polozov, M. O'Connell, J. Burgeson, P. Yu, W. Zeller, J. Zhang, E. Onua, J. Ramirez, G. A. Palsdottir, G. V. Halldorsdottir, T. Andresson, A. S. Kiselyov, M. Gurney and J. Singh, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 2658; (b) A. K. Podichetty, A. Faust, K. Kopka, S. Wagner, O. Schober, M. Schäfers, G. Haufe, *Bioorg. Med. Chem.*, 2009, **17**, 2680; (c) S. S. Nigudkar, K. J. Stine and A. V. Demchenko, *J. Am. Chem. Soc.*, 2014, **136**, 921.
- (a) Y. Ohtsuka and T. Yamakawa, *Tetrahedron*, 2011, **67**, 2323; (b)
  L. A. McAllister, R. A. McCormick, K. M. James, S. Brand, N. Willetts and D. J. Procter, *Chem. Eur. J.*, 2007, **13**, 1032; (c) J. C. Torres, S. J. Garden, A. C. Pinto, F. S. Q. da Silva and N. Boechat, *Tetrahedron*, 1999, **55**, 1881.
- (a) R. P. Singh, U. Majumder and J. M. Shreeve, J. Org. Chem., 2001, 66, 6263; (b) W. J. Middleton and E. M. Bingham, J. Org. Chem., 1980, 45, 2883; (c) T. Umemoto, R.P. Singh, Y. Xu and N. Saito, J.

*Am. Chem. Soc.*, 2010, **132**, 18199; (d) Y. H. Lim, Q. Ong, H. A. Duong, T. M. Nguyen and C. W. Johannes, *Org. Lett.*, 2012, **14**, 5676.

- (a) J. Zhu, W. Zhang, L. Zhang, J. Liu, J. Zheng and J. Hu, J. Org. Chem., 2010, 75, 5505; (b) S.-L. Shi and S. L. Buchwald, Angew. Chem., 2015, 127, 1666; Angew. Chem. Int. Ed., 2015, 54, 1646.
- For reviews on visible light photoredox catalysis, see: (a) J. Xuan and W.-J. Xiao, Angew. Chem., Int. Ed., 2012, **51**, 6828; (b) M. N. Hopkinson, B. Sahoo, J. L. Li and F. Glorius, Chem. Eur. J., 2014, **20**, 3874; (c) L. Shi and W. Xia, Chem. Soc. Rev., 2012, **41**, 7687; (d) K. Zeitler, Angew. Chem., Int. Ed., 2009, **48**, 9875; (e) Y. Xi, H. Yi and A. Lei, Org. Biomol. Chem., 2013, **11**, 2387; (f) T. P. Yoon, M. A. Ischay and J. Du, Nat. Chem., 2010, **2**, 527; (g) J. M. R. Narayanam, and C. R. J. Stephenson, Chem. Soc. Rev., 2011, **40**, 102; (h) D. P. Hari and B. König, Angew. Chem. Int. Ed., 2013, **52**, 4734; (i) C. K. Prier, D. A. Rankic and D. W. C. Macmillian, Chem. Rev., 2013, **113**, 5322.
- For selected examples of the application of *fac*-Ir(ppy)<sub>3</sub> in visible-light photocatalysis, see: (a) A. McNally, C. K. Prier and D. W. C. MacMillan, *Science*, 2011, **334**, 1114; (b) Y. Yasu, T. Koike and M. Akita, *Angew. Chem., Int. Ed.*, 2012, **51**, 9567; (c) H. Jiang, Y. Cheng, R. Wang, M. Zheng, Y. Zhang and S. Yu, *Angew. Chem., Int. Ed.*, 2013, **52**, 13289; (d) P. Xu, A. Abdukader, K. Hu, Y. Cheng and C. Zhu, *Chem. Commun.*, 2014, **50**, 2308; (e) Q. Liu, H. Yi, J. Liu, Y. Yang, X. Zhang, Z. Zeng and A. Lei, *Chem. Eur. J.*, 2013, **19**, 5120; (f) W.-J. Yoo, T. Tsukamoto and S. Kobayashi, *Org. Lett.*, 2015, **17**, 3640; (g) Q. Wang, J. Huang and L. Zhou, *Adv. Synth. Catal.*, 2015, **357**, 2479; (h) X.-J. Wei, D.-T. Yang, L. Wang, T. Song, L.-Z. Wu and Q. Liu, *Org. Lett.*, 2013, **15**, 6054.
- (a) L. Wang, X.-J. Wei, W.-L. Jia, J.-J. Zhong, L.-Z. Wu and Q. Liu, Org. Lett., 2014, 16, 5842; (b) L. Wang, X.-J. Wei, W.-L. Lei, H. Chen, L.-Z. Wu and Q. Liu, Chem. Commun., 2014, 50, 15916.
- For selected examples on photoredox-catalyzed difluoromethylation, see: (a) C.-J. Wallentin, J. D. Nguyen, P. Finkbeiner and C. R. J. Stephenson, J. Am. Chem. Soc., 2012, 134, 8875; (b) X. Sun and S. Yu, Org. Lett., 2014, 16, 2938; (c) W. Fu, M. Zhu, G. Zou, C. Xu, Z. Wang and B. Ji, J. Org. Chem., 2015, 80, 4766; (d) C. Yu, N. Iqbal, S. Park and E. J. Cho, Chem. Commun., 2014, 50, 12884; (e) J. Jung, E. Kim, Y. You and E. J. Cho, Adv. Synth. Catal., 2014, 365, 2741; (f) P. Xu, K. Hu, Z. Gu, Y. Cheng and C. Zhu, Chem. Commun., 2015, 51, 7222; (g) W. Li, X. Zhu, H. Mao, Z. Tang and C. Zhu, Chem. Commun., 2014, 50, 7521; (h) Y. M. Su, Y. Hou, F. Yin, Y. M. Xu, Y. Li, X. Zheng and X. S. Wang, Org. Lett., 2014, 16, 2958; (i) X.-J. Tang, C. S. Thomoson and W. R. Dolbier, Org. Lett., 2014, 16, 4594; (j) X.-J. Tang and W. R. Dolbier, Angew. Chem., Int. Ed., 2015, 54, 4246.
- B. Zhang, C. Mück-Lichtenfeld, C. G. Daniliuc and A. Studer, *Angew. Chem., Int. Ed.* 2013, **52**, 10792.