


The Formation of Allyl Sulphides by Phenylthio-migration: Control by Silicon †

By Ian Fleming,* Ian Paterson, and Andrew Pearce, University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW


When γ -silyl- β -phenylthio-alcohols are treated with acid, the strategically placed silyl group encourages the re-arrangement of the phenylthio-group, both from a secondary migration origin to a secondary migration terminus, and from a secondary migration origin to a tertiary migration terminus (4) \rightarrow (6). Geraniol/nerol (12) and linalool (14) have been synthesised from a common intermediate (11) using this type of reaction. Phenylthio-migration from a tertiary migration origin (17) \rightarrow (3) can be controlled to a limited extent by a suitably placed silyl group, but it is easier to achieve direct β -elimination of the silyl and phenylthio-groups (17) \rightarrow (18).

BROWNBRIDGE and Warren¹ have shown that phenylthio-groups migrate in an acid-catalysed reaction of the type (1) \rightarrow (2) + (3) to give allyl sulphides. Rearrangement only takes place, however, when it is 'downhill,' *i.e.* when the phenylthio-group goes from a secondary migration origin to a primary migration terminus, or from a tertiary migration origin to either a secondary or a primary migration terminus. In a preliminary communication² we described how we extended the scope of this easy route to allyl sulphides by incorporating in the starting materials a trimethylsilyl group, so placed that it encouraged rearrangement of a phenylthio-group from a secondary migration origin to a secondary migration terminus and even 'uphill' from a secondary migration origin to a tertiary migration

makes it possible to rearrange them in daylight in high yield to their (more-substituted) allylic isomers (7).^{1,4} Thus our work, taken with Warren's, makes it possible to synthesise a wide range of allyl sulphides: of the



SCHEME 1 (i) TsOH-C₆H₆ reflux (secondary alcohols) or TsOH-MeCN-20 °C (tertiary alcohols); (ii) sunlight-CCl₄, 20 °C



terminus. In this paper we describe the experimental details, and illustrate the power of the method by syntheses of geraniol/nerol and linalool.

RESULTS AND DISCUSSION

Scheme 1 summarises the reactions used to establish that the silyl group encourages rearrangement in cases where rearrangement would not otherwise have taken place. Thus the compounds analogous to (4a) and (4b), but without the silyl group, gave no rearrangement on treatment with acid;¹ in contrast (4a) and (4b) themselves gave high yields of the rearranged allyl sulphides (6a) and (6b), respectively. Tertiary alcohols similar to (4c) and (4d), but without a silyl group, were known³ to undergo simple dehydration on treatment with acid; in contrast, (4c) and (4d) rearranged to give the allyl sulphides (6c) and (6d). The success of these reactions is probably a result of the easy loss of the silyl group from an intermediate episulphonium ion (5). Each of the allyl sulphides (6) has a substitution pattern which

* No reprints available.

eighteen possible substitution patterns (ignoring geometrical isomers), we have, between us, synthesised examples of eleven. Since allyl sulphides can easily be converted with allylic rearrangement to allyl alcohols,⁵ the same wide range of allyl alcohols is also made available by this route.

Two well-known natural products, geraniol (12) and linalool (14), are allyl alcohols which belong in this category. To test the power of our method we have synthesised these compounds, the former accompanied inevitably by its stereoisomer, nerol. The synthesis is shown in Scheme 2, and is noteworthy only in that we had some difficulty finding the best conditions for the key step (10) \rightarrow (11) (see Experimental section). One cause of the relatively low yield appeared to be the participation of the remote double bond [(10) \rightarrow (15) \rightarrow (16)]. The best conditions found were probably not optimal, but did lead to the allyl sulphide (11) in 58% isolated yield. The oxidation of this sulphide gave a mixture of the corresponding diastereoisomeric sulphoxides. These compounds had already been prepared in a different way by Evans and his co-workers,⁶ who had converted them to geraniol/nerol (12) (9:1; 55%). We got geraniol/nerol in a ratio of 67:33 in 64% yield; the different ratio is almost certainly a consequence of our having a different ratio of the diastereoisomeric sulphoxides. Evans's route, although simpler than ours, did not go by way of the sulphide (11). Since the

(3a) and (20) in the ratio 2 : 5, indicating that to some extent a proton has been lost rather than the silyl group, both in the initial reaction and possibly in the fluoride-catalysed reaction too. In our experience this is rare, but it is not unknown.¹⁰ The best conditions for the rearrangement in this case were to use a large excess of lithium chloride and to do the reaction in hexamethylphosphoramide, when the allyl sulphide (3a) was produced (n.m.r.) in 60% yield.

Syntheses of the Compounds used in this Work.—The syntheses are shown in Scheme 4 and were not remarkable. The only seriously low yields were in the steps (23)→(24) and (23)→(25), which must be a consequence of the ease of 'enolisation,' since the nitrile (27) gave a good yield of the aldehyde (28). The low yield of the ketone (25) was avoided by the (later) synthesis of the same ketone (9) shown in Scheme 2. The mixture of diastereoisomers of (4a) was rich in one isomer (*ca.* 4 : 1) when prepared from (24) and in the other isomer (*ca.* 1 : 4) when prepared from (25). However, the subsequent reaction (4a)→(6a) was the same with alcohol from either source.

EXPERIMENTAL

Rearrangements (4)→(6)→(7).—**3-Phenylthiobut-1-ene** (6a). The alcohol (4a) (113 mg, either diastereoisomer) in dry benzene (5 ml) was heated under reflux (light excluded) with toluene-*p*-sulphonic acid monohydrate (85 mg) for 5 min. The mixture was cooled, poured into sodium carbonate solution (50 ml, saturated), and the aqueous layer extracted with ether (2 × 10 ml). The combined organic layers were dried (Na₂SO₄) and evaporated *in vacuo*, all procedures being carried out as far as possible with light excluded. Preparative thin layer chromatography (t.l.c.) (silica gel, benzene) gave 3-(phenylthio)but-1-ene¹¹ (6a) (68 mg, 93%) as an oil; ν_{\max} 1 640 (C=C) and 1 580 cm⁻¹ (aromatic); $\tau(\text{CCl}_4)$ 2.5—2.9 (5 H, m, SPh), 4.28 (1 H, ddd, *J* 7, 9, and 18 Hz, CH=CH_AH_B), 4.89—5.34 (2 H, m, CH=CH_AH_B), 6.35 (1 H, quintet, *J* 7 Hz, CHSPh), and 8.64 (3 H, d, *J* 7 Hz, Me); *m/e* 164 (100%, M⁺), 149 (10, M — Me), and 110 (67, PhSH).

1-(Phenylthio)but-2-ene (7a). The olefin (6a) (50 mg) in carbon tetrachloride (1 ml) was exposed to bright sunlight for 6 h. The solution was evaporated *in vacuo* to give a 1 : 4 mixture of *cis*- and *trans*-1-(phenylthio)but-2-ene¹² (7a) (49 mg, 98%) as an oil (Found: C, 72.9; H, 7.5; S, 19.2. Calc. for C₁₀H₁₂S: C, 73.1; H, 7.4; S, 19.5%); ν_{\max} (CCl₄) 1 630 (C=O) and 1 580 cm⁻¹ (aromatic); $\tau(\text{CCl}_4)$ 2.6—3.0 (5 H, m, SPh), 5.3—5.7 (2 H, m, vinyl), 6.5—6.7 (2 H, m, CH₂), and 8.3—8.7 (3 H, m, Me); *m/e* 164 (31%, M⁺), 149 (10, M — Me), and 110 (100, PhSH).

3-Phenyl-3-(phenylthio)propene (6b). The alcohol (4b) (158 mg, a mixture of diastereoisomers) was stirred at room temperature with toluene-*p*-sulphonic acid monohydrate (98 mg) in dry benzene (10 ml) for 1 h in the absence of light. Work-up as for (6a) above gave 3-phenyl-3-(phenylthio)propene (6b) (110 mg, 99%) as an oil (Found: C, 79.3; H, 6.44; S, 14.2. C₁₅H₁₄S requires C, 79.4; H, 6.23; S, 14.2%); ν_{\max} (CCl₄) 1 635 (C=C) and 1 580 cm⁻¹ (aromatic); $\tau(\text{CCl}_4)$ 2.6—2.9 (10 H, m, PhS + Ph), 3.94 (1 H, ddd, *J* 8, 10, and 18 Hz, CH=CH_AH_B), 4.99 (1 H, d, *J* 10 Hz, CH=CH_AH_B), 5.03 (1 H, d, *J* 18 Hz, CH_AH_B), and 5.31 (1 H, d,

J 8 Hz, CHSPh); *m/e* 226 (62%, M⁺), 117 (100, M — PhS), and 110 (20, PhSH).

1-Phenyl-3-(phenylthio)propene (7b). The olefin (6b) (80 mg) in carbon tetrachloride (1 ml) was exposed to ordinary daylight over a period of two days. The solution was evaporated *in vacuo* and the solid crystallised from ethanol to give *trans*-1-phenyl-3-(phenylthio)propene (7b) (65 mg, 80%), m.p. 76—78 °C (lit.¹¹ 78 °C); ν_{\max} (CCl₄) 1 630 (C=C) and 1 580 cm⁻¹ (aromatic); $\tau(\text{CCl}_4)$ 2.6—3.0 (10 H, m, PhS + Ph), 3.62 (1 H, s, *J* 16 Hz, PhCH=CH), 3.70—4.00 (1 H, m, PhSCH₂CH=), and 6.35 (2 H, d, *J* 6 Hz, PhSCH₂); *m/e* 226 (41%, M⁺), 117 (30, M — PhS) and 110 (100, PhSH).

3,3-Dimethyl-3-(phenylthio)propene (6c). The alcohol (4c) (134 mg) in acetonitrile (2 ml, distilled from phosphorus pentaoxide) was added at room temperature to a stirred solution of toluene-*p*-sulphonic acid monohydrate (48 mg) in acetonitrile (10 ml, distilled as above) in the absence of light and the mixture maintained at room temperature for 30 min. Work-up as for (6a) above gave 3,3-dimethyl-3-(phenylthio)propene (6c)¹³ (86 mg, 96%) as an oil, *R*_F 0.7; ν_{\max} (CCl₄) 1 640 cm⁻¹ (C=C); $\tau(\text{CCl}_4)$ 2.5—2.9 (5 H, m, PhS), 4.08 (1 H, dd, *J* 11 and 18 Hz, CH=CH₂), 6.16 (1 H, dd, *J* 11 and 2 Hz, *cis*-CH=CH_AH_B), 6.34 (1 H, dd, *J* 18 and 2 Hz, *trans*-CH=CH_AH_B), and 8.65 (6 H, s, Me); *m/e* 178 (10%, M⁺), 110 (100, PhSH), and 109 (47, PhS).

3-Methyl-1-(phenylthio)but-2-ene (7c).—The olefin (6c) (86 mg) in carbon tetrachloride (1 ml) was exposed to bright sunlight for 12 h. The solution was evaporated *in vacuo* and t.l.c. (silica gel, benzene) gave 3-methyl-1-(phenylthio)but-2-ene (7c)¹⁴ (80 mg, 93%), as an oil, *R*_F 0.7; ν_{\max} (CCl₄) 1 635 cm⁻¹ (C=C); $\tau(\text{CCl}_4)$ 2.6—3.0 (5 H, m, PhS), 4.64—4.76 (1 H, m, vinyl), 6.66 (2 H, d, *J* 8 Hz, CH₂), and 8.29, 8.43 (total 6 H, 2 s, Me_A and Me_B); *m/e* 178 (100%, M⁺), 110 (84, PhSH) and 109 (28, PhS).

3-Ethyl-3-(phenylthio)pentene (6d). The alcohol (4d) (104 mg, 0.35 mmol) in dry acetonitrile (3 ml) was added, in the dark, to a stirred solution of toluene-*p*-sulphonic acid monohydrate (50 mg) in dry acetonitrile (10 ml) and the mixture kept for 5 h at room temperature. Work-up as for (6a) above gave 3-ethyl-3-(phenylthio)pentene (6d) (61 mg, 83%), *R*_F 0.8; ν_{\max} (film) 1 640 cm⁻¹ (C=C); $\tau(\text{CCl}_4)$ 2.35—2.85 (5 H, m, Ph), 4.15 (1 H, dd, *J* 11 and 17 Hz, CH=CH₂), 5.03 (1 H, dd, *J* 11 and 2 Hz, *cis*-CH=CH_AH_B), 5.4 (1 H, dd, *J* 17 and 2 Hz, *trans*-CH=CH_AH_B), 8.14—8.7 (4 H, m, CH₂Me), and 8.75—9.11 (6 H, m, CH₂Me) (Found: M⁺, 206.113 l. C₁₃H₁₈S requires M, 206.112 9); *m/e* 206 (9%), 110 (100, PhSH), and 109 (21, PhS).

3-Ethyl-1-(phenylthio)pent-2-ene (7d). A solution of the olefin (6d) (50 mg, 0.24 mmol) in carbon tetrachloride (1 ml) was irradiated by sunlight for several days (the reaction being followed by n.m.r.). On completion of reaction, the solution was evaporated *in vacuo* and t.l.c. [acetone—light petroleum (3 : 7)] gave 3-ethyl-1-(phenylthio)pent-2-ene (6d) as an oil (45 mg, 90%), *R*_F 0.7; ν_{\max} (film) 1 635 cm⁻¹ (C=C); $\tau(\text{CCl}_4)$ 2.65—3.02 (5 H, m, Ph), 4.8 (1 H, m, C=CH), 6.54 (2 H, d, *J* 8 Hz, CH₂SPh), 7.75—8.2 (4 H, 2 overlapping q, *J* 7 Hz, CH₂Me), and 8.85—9.25 (6 H, 2 overlapping t, *J* 7 Hz, CH₂Me) (Found: M⁺, 206.113 0. C₁₃H₁₈S requires M, 206.112 9); *m/e* 206 (8%), 110 (100, PhSH), 97 (40, M — PhS), and 96 (40, M — PhSH).

Synthesis of Geraniol/Nerol and Linalool.—**2-(Phenylthio)-1-(trimethylsilyl)butan-3-one** (9). A solution of α -phenylthioacetone¹⁵ (10.10 g, 60.7 mmol) in dry THF (50 ml) was added dropwise over 1 h to a stirred suspension of sodium

hydride [3.00 g of a 50% dispersion in oil, 62.5 mmol; which had been washed with light petroleum (3×10 ml) in dry THF (250 ml) under nitrogen]. The mixture was stirred for a further 1 h, then iodomethyltrimethylsilane ¹⁶ (14.70 g, 68.7 mmol) was added. The mixture was heated under reflux in a nitrogen atmosphere in the dark for 21 h. A standard aqueous work-up and chromatography on silica gel (500 g) gave, on elution with dichloromethane, 2-(phenylthio)-1-(trimethylsilyl)butan-3-one (11.65 g, 75%), b.p. 95–98 °C at 0.01 mmHg, R_F (dichloromethane) 0.6; ν_{\max} (film) 1 705 (C=O), 1 585 (aromatic), and 1 250 cm^{-1} (SiMe₃); $\tau(\text{CCl}_4)$ 2.79 (5 H, s, Ph), 6.36 (1 H, dd, J 9 and 6 Hz, CH), 7.87 (3 H, s, COMe), 8.88 (1 H, dd, J 9 and 15 Hz, CH_AH_B), 9.1 (1 H, dd, J 6 and 15 Hz, CH_AH_B), and 9.98 (9 H, s, SiMe₃); m/e 252 (5%, M^+), 237 (6, $M - \text{Me}$), 209 (100, $M - \text{COMe}$), 167 (31, PhSSiMe₂), 143 (83, $M - \text{PhS}$) and 110 (39, PhSH). The semicarbazone had m.p. 129–131 °C from MeOH–H₂O (Found: C, 54.5; H, 7.70; N, 13.9. $\text{C}_{14}\text{H}_{23}\text{N}_3\text{OSi}$ requires C, 54.3; H, 7.50; N, 13.6%).

5-Bromo-2-methylpent-2-ene. This was prepared by the method of Julia *et al.*¹⁷ It was purified by chromatography on silica gel using ether for elution, followed by distillation. It had b.p. and spectroscopic data as previously recorded.

3,7-Dimethyl-2-(phenylthio)-1-(trimethylsilyloct-6-en-3-ol (10). A solution of 2-(phenylthio)-1-(trimethylsilyl)butan-3-one (3.15 g, 12.5 mmol) in dry ether (10 ml) was added dropwise to the solution of the Grignard reagent, formed from magnesium (0.92 g, 37.5 mmol) and 5-bromo-2-methylpent-2-ene (6.12 g, 37.5 mmol), in dry ether (40 ml) at –78 °C under nitrogen. The mixture was maintained at –78 °C for a further 2 h, then warmed to room temperature during 1 h. A standard aqueous work-up and chromatography on silica gel (200 g), eluting with dichloromethane, gave the alcohol (10) (2.85 g, 68%) (Found: C, 68.2; H, 9.55. $\text{C}_{19}\text{H}_{32}\text{OSSi}$ requires C, 67.8; H, 9.55%), R_F (dichloromethane) 0.55, ν_{\max} (film) 3 500 (OH), 1 750 (aromatic) and 1 250 cm^{-1} (SiMe₃); $\tau(\text{CCl}_4)$ 2.58–2.93 (5 H, m, Ph), 4.97 (1 H, m, C=CH), 6.75 (1 H, dd, J 6 and 9 Hz, CHSPh), 7.85–8.13 (5 H, m, CH_2CH_2 and OH), 8.35 (3 H, s, C=CMe), 8.41 (3 H, s, C=CMe), 8.81 (3 H, s, MeCOH), 9.00–9.15 (2 H, m, CH_2 SiMe₃), and 10.03 (9 H, s, SiMe₃); m/e 336 (0.5%, M^+), 210 (81, PhSCH₂CH₂SiMe₃), 182 (69, PhSSiMe₃), 167 (30, PhSSiMe₂), 110 (74, PhSH), and 73 (100, SiMe₃).

3,7-Dimethyl-3-phenylthio-octa-1,6-diene (11). A solution of the alcohol (10) (358 mg, 1.06 mmol) was made up in dry HMPA (5 ml) saturated with lithium chloride. Triethylamine (480 mg, 4.75 mmol) was added to the mixture, followed by the dropwise addition of thionyl chloride (560 mg, 4.75 mmol) during 20 min. An aqueous work-up and t.l.c. (carbon tetrachloride) gave the diene (11) (153 mg, 58%) as an oil, R_F 0.5; ν_{\max} (film) 1 630 (C=C) and 1 570 cm^{-1} (aromatic); $\tau(\text{CCl}_4)$ 2.5–2.95 (5 H, m, Ph), 4.14 (1 H, dd, J 10 and 18 Hz, $\text{CH}=\text{CH}_2$), 4.82–5.1 (1 H, m, C=CH), 5.1 (1 H, dd, J 2 and 10 Hz, *cis*- $\text{CH}=\text{CH}_A\text{H}_B$), 5.36 (1 H, dd, J 2 and 18 Hz, *trans*- $\text{CH}=\text{CH}_A\text{H}_B$), 7.76–8.1 (4 H, m, CH_2CH_2), 8.34 (3 H, s, C=CMe), 8.4 (3 H, s, C=CMe), and 8.73 (3 H, s, MeCSPh) (Found: M^+ , 246.144 7. $\text{C}_{16}\text{H}_{22}\text{S}$ requires M , 246.144 2); m/e 246 (11%), 137 (12, $M - \text{PhS}$), 136 (90, $M - \text{PhSH}$), and 110 (100, PhSH).

Another preparation and column chromatography gave the cyclopentane (16) (20%). It could not be further purified and had R_F (carbon tetrachloride) 0.6; ν_{\max} (film) 1 640 (C=C), 1 580 (aromatic), and 1 250 cm^{-1} (SiMe₃), $\tau(\text{CCl}_4)$ 2.42–2.88 (5 H, m, Ph), 5.32 (2 H, m, C=CH₂),

7.88–8.17 (2 H, m, CH), 8.28–8.42 (7 H, m, CH_2CH_2 and C=CMe), 8.87 (3 H, s, MeCSPh), 9.08 (1 H, dd, J 5 and 15 Hz, $\text{CH}_A\text{H}_B\text{SiMe}_3$), 9.56 (1 H, dd, J 8 and 15 Hz, $\text{CH}_A\text{H}_B\text{SiMe}_3$), and 9.92 (9 H, s, SiMe₃) (Found: M^+ , 318.182 8. $\text{C}_{19}\text{H}_{30}\text{SSi}$ requires M , 318.183 7); m/e 318 (9%), 110 (78, PhSH), and 73 (100, SiMe₃).

Geraniol/nerol (12). A solution of sodium metaperiodate (230 mg, 1.07 mmol) in water (1.5 ml) was added dropwise to a stirred solution of 3,7-dimethyl-3-(phenylthio)octa-1,6-diene (11) (250 mg, 1 mmol) in methanol (26 ml) in the dark at 0 °C. After completion of the addition, the mixture was stirred in the dark at room temperature for 3 days. The mixture was poured into water (25 ml) and extracted with dichloromethane (3×25 ml). The combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo*. Chromatography on silica gel (20 g) with chloroform gave the sulphoxide (232 mg, 87%), as a mixture of diastereoisomers, R_F (chloroform) 0.2; ν_{\max} (film) 1 665 (C=C), 1 635 (C=C), and 1 050 cm^{-1} (S=O); $\tau(\text{CCl}_4)$ 2.38–2.82 (5 H, m, Ph), 4.55–6.05 (4 H, m, olefinic H), 7.8–8.04 (4 H, m, CH_2CH_2), and 8.24–8.5 (9H, m, Me); m/e 126 (29, PhSOH), 125 (47, PhSO), 110 (100, PhSH) and 109 (56, PhS). A solution of sodium thiophenolate [prepared from thiophenol (0.98 g, 8.9 mmol) and sodium hydroxide (0.36 g, 8.9 mmol)] in methanol (15 ml) was added dropwise with stirring to this sulphoxide (232 mg, 0.89 mmol) at room temperature in the dark. The mixture was stirred for 3 days and then poured into sodium carbonate solution (20 ml, saturated) and extracted with dichloromethane (3×50 ml). The combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo*. Chromatography on silica gel (14 g) gave, on elution with dichloromethane, a mixture (85 mg, 64%) of *E*-3,7-dimethylocta-2,6-dien-1-ol (geraniol); ν_{\max} (film) 3 330 (OH), and 1 670 cm^{-1} (C=C), $\tau(\text{CDCl}_3)$ 4.46–4.72 (1 H, m, C=CHR), 4.8–5.02 (1 H, m, C=CHCH₂OH), 5.85 (2 H, d, J 6 Hz, CH_2OH), 7.82–8.04 (4 H, m, CH_2CH_2), 8.12 (1 H, s, OH, removed by D₂O), and 8.32 (6 H, s, Me), and 8.39 (3 H, s, Me), and *Z*-3,7-dimethylocta-2,6-dien-1-ol (nerol); $\tau(\text{CDCl}_3)$ 4.46–4.72 (1 H, m, C=CHR), 5.26–5.37 (1 H, m, C=CHCH₂OH), 5.92 (2 H, d, J 6 Hz, CH_2OH), 7.88–8.04 (4 H, m, CH_2CH_2), 8.12 (1 H, s, OH, removed by D₂O), 8.32 (6 H, s, Me), and 8.39 (3 H, s, Me). The ratio of geraniol:nerol was 2:1 from integration of the n.m.r. spectrum. The spectroscopic data for geraniol were identical with those of an authentic sample.

3,7-Dimethyl-1-(phenylthio)octa-2,6-diene (13). A solution of 3,7-dimethyl-3-(phenylthio)octa-1,6-diene (11) (202 mg, 0.82 mmol) in carbon tetrachloride (1 ml) was irradiated by sunlight for several days (the reaction being followed by n.m.r.). On completion of reaction, the solution was evaporated *in vacuo*, and p.l.c. (carbon tetrachloride) gave 3,7-dimethyl-1-(phenylthio)octa-2,6-diene (194 mg, 97%), as a mixture of stereoisomers. This oil had R_F 0.5; ν_{\max} (film) 1 660 cm^{-1} (C=C); $\tau(\text{CCl}_4)$ 2.60–2.96 (5 H, m, Ph), 4.60–4.86 (1 H, m, C=CH), 4.86–5.10 (1 H, m, C=CH), 6.53 (2 H, d, J 7 Hz, CH_2SPh), 7.82–8.10 (4 H, m, CH_2CH_2), 8.32 (3 H, s, Me), and 8.41 (6 H, s, Me) (Found: M^+ , 246.143 8. $\text{C}_{16}\text{H}_{22}\text{S}$ requires M , 246.144 2); m/e 246 (13%), 137 (76, $M - \text{PhS}$), 136 (69, $M - \text{PhSH}$), and 110 (100, PhSH).

Linalool (14). A solution of sodium metaperiodate (230 mg, 1.07 mmol) in water (2 ml) was added dropwise to a stirred solution of 3,7-dimethyl-1-(phenylthio)octa-2,6-diene (13) (236 mg, 0.96 mmol) in methanol (30 ml) at 0 °C. The mixture was stirred at room temperature for 24 h,

poured into water (25 ml), extracted with dichloromethane (3×25 ml), and the combined extracts dried (Na_2SO_4) and evaporated *in vacuo*. Chromatography on silica gel (20 g) gave, on elution with chloroform, the sulphoxides as an oil (244 mg, 97%). The diastereoisomeric mixture had R_F (chloroform) 0.2; ν_{max} (film) 1 660 (C=C), and 1 050 cm^{-1} (S=O); $\tau(\text{CCl}_4)$ 2.4—2.8 (5 H, m, Ph), 4.8—5.45 (2 H, m, C=CH), 6.5—6.68 (2 H, m, CH_2SPh), 7.5—8.16 (4 H, m, CH_2CH_2), and 8.2—8.6 (9 H, m, Me) (Found: $M - \text{PhSO}$ 137.133 6; $\text{C}_{10}\text{H}_{17}$ requires 137.133 0. Found: PhSOH 126.013 5; $\text{C}_6\text{H}_6\text{OS}$ requires 126.013 9; m/e 137 (24%, $M - \text{PhSO}$), 126 (56, PhSOH), and 110 (100, PhSH). A solution of sodium thiophenolate [prepared from thiophenol (0.98 g, 8.9 mmol) and sodium hydroxide (360 mg, 8.9 mmol)] in methanol (15 ml), was added with stirring to this sulphoxide (179 mg, 0.68 mmol) in the dark. The mixture was heated under reflux for 7 h, then stirred for 18 h at room temperature. The mixture was poured onto sodium carbonate solution (25 ml, saturated) and extracted with dichloromethane (3×50 ml). The combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo*. Chromatography on silica gel (14 g) gave, on elution with dichloromethane, 3,7-dimethylocta-1,6-dien-3-ol (linalool) as an oil (69 mg, 66%), R_F (chloroform) 0.25; ν_{max} (film) 3 410 (OH) and 1 640 cm^{-1} (C=C); $\tau(\text{CDCl}_3)$ 4.13 (1 H, dd, J 10 and 17 Hz, $\text{CH}=\text{CH}_2$), 4.88 (1 H, dd, J 2 and 17 Hz, *trans*- $\text{HC}=\text{CH}_A\text{H}_B$), 4.82—5.04 (1 H, m, C=CH), 5.01 (1 H, dd, J 2 and 10 Hz, *cis*- $\text{HC}=\text{CH}_A\text{H}_B$), 7.86—8.17 (4 H, m, CH_2CH_2), 8.36 (3 H, s, Me), 8.44 (4 H, s, Me and OH), and 8.78 (3 H, s, MeCOH).

Rearrangements from a Tertiary Migration Origin.—2-(3-Methylbutyl)prop-2-en-1-ol (18a). Boron trifluoride-ether (*ca.* 40 mg, 0.04 ml, freshly distilled) was added to a stirred solution of the alcohol (17a) (88 mg) in light petroleum (10 ml, b.p. 60—80 °C, distilled from phosphorus pentaoxide) at room temperature under nitrogen and the mixture kept at room temperature for 5 min. An aqueous work-up and t.l.c. (silica gel, benzene) gave 2-(3-methylbutyl)prop-2-en-1-ol (26 mg, 72%) as an oil, R_F 0.2; ν_{max} (CCl_4) 3 400 (OH) and 1 650 cm^{-1} (C=C); $\tau(\text{CCl}_4)$ 5.06 (1 H, broad s, $\text{C}=\text{CH}_A\text{H}_B$), 5.21 (1 H, br s, $\text{C}=\text{CH}_A\text{H}_B$), 6.02 (2 H, s, CH_2OH), 7.8—8.8 (6 H, m, CH, CH_2 and OH), and 9.06 (6 H, d, J 6 Hz, Me).

3-(3-Methylbutyl)but-3-en-2-ol (18b). The alcohol (17b) (89 mg) in acetonitrile (2 ml, distilled from phosphorus pentaoxide) was added to a stirred solution of toluene-*p*-sulphonic acid monohydrate (26 mg) in acetonitrile (10 ml) and the mixture kept at room temperature for 15 min. An aqueous work-up and t.l.c. (silica gel, benzene) gave 3-(3-methylbutyl)but-3-en-2-ol (27 mg, 70%) as an oil, R_F 0.2; ν_{max} 3 450 (OH) and 1 650 cm^{-1} (C=C); $\tau(\text{CCl}_4)$ 5.0 (1 H, br s, $\text{C}=\text{CH}_A\text{H}_B$), 5.28 (1 H, br s, $\text{C}=\text{CH}_A\text{H}_B$), 5.84 (1 H, q, J 7 Hz, CHOH), 7.8—8.8 (9 H, m, overlain by d, J 7 Hz at 8.76, CHMe_2 ; CH_2 , MeCHOCH and OH), and 9.07 (6 H, d, J 6 Hz, CHMe_2).

Rearrangements Promoted by Thionyl Chloride.—The alcohol (17a) (30 mg), on treatment with thionyl chloride (3 drops) and triethylamine (0.1 ml) at room temperature for 1 min in dry benzene (15 ml) and work-up gave an oil (33 mg). This was a mixture of 5-methyl-1-(phenylthio)-2-(trimethylsilylmethyl)hex-2-ene (20), R_F 0.2; τ 2.7—3.0 (5 H, m, PhS), 4.8—5.0 (1 H, m, vinyl), 6.62 (2 H, s, $\text{CH}_2\text{-SPh}$), 8.2—9.3 (11 H, m, overlain by d, J 6 Hz at 9.24, CH, CH_2 , Me), and 9.98 (9 H, s, SiMe_3) (Found: M^+ , 292.167 8. $\text{C}_{17}\text{H}_{28}\text{SSi}$ requires M , 292.168 0; m/e 292 (100%), 277,

(22, $M - \text{Me}$), 248 (18), 182 (70, PhSSiMe_3), and 167 (83); and 2-chloro-5-methyl-2-trimethylsilylmethylhexyl phenyl sulphide (19); $\tau(\text{CCl}_4)$ 2.7—3.0 (5 H, PhS), 6.69 (2 H, s, CH_2SPh), 8.1—9.3 (13 H, m, CH, CH_2 and CH_3), and 9.90 (9 H, s, SiMe_3); the i.r. spectrum showed the absence of a hydroxy group. The ratio of the olefin (20) to the rearranged chloride (19) was approximately 1 : 2 in the crude mixture. When the crude mixture was heated under reflux with a solution of sodium fluoride (100 mg) in methanol (20 ml) it gave a 2 : 5 mixture of (20) and (3a).

5-Methyl-2-(phenylthiomethyl)hex-1-ene (3a). The alcohol (17a) (59 mg) was stirred at room temperature with thionyl chloride (3 drops) and triethylamine (1 ml) in hexamethylphosphoramide (15 ml) saturated with lithium chloride for 1 min at room temperature. An aqueous work-up and t.l.c. (silica gel, hexane) gave a sample of 5-methyl-2-(phenylthiomethyl)hex-1-ene (3a) as an oil (Found: C, 76.5; H, 9.40; S, 14.3. $\text{C}_{14}\text{H}_{20}\text{S}$ requires C, 76.3; H, 9.15; S, 14.5%), R_F 0.8; ν_{max} (CCl_4) 1 645 (C=C) and 1 580 cm^{-1} (aromatic), $\tau(\text{CCl}_4)$ 2.6—2.9 (5 H, m, PhS), 5.12—5.24 (2 H, m, C=CH₂), 6.50 (2 H, s, CH_2SPh), 7.7—7.9 (2 H, m, allylic CH_2), and 8.1—9.2 (overlain by doublet J 6 Hz at 9.04; total 9 H, m, CH, CH_2 and Me); m/e 220 (66%, M^+) 164 (25), 110 (100, PhSH), and 109 (43, PhS). The yield was estimated on the crude product as $60\% \pm 7\%$ from n.m.r. using dimethylformamide as an internal standard.

Reaction of 6-Methyl-3-(phenylthio)-3-(trimethylsilylmethyl)heptan-2-ol with Boron Trifluoride-Diethyl Ether.—Boron trifluoride-diethyl ether (58 mg, freshly distilled) was added dropwise with stirring to a solution of the alcohol (17b) (145 mg) in dry benzene (10 ml) and the mixture kept at room temperature for 5 min. An aqueous work-up and t.l.c. (silica gel, benzene) gave an oil (51 mg, 53%), R_F 0.7, which was a mixture of 5-methyl-2-(1-phenylthioethyl)-hexene (3b) and the two geometric isomers of 6-methyl-3-(phenylthiomethyl)hept-2-ene (21), the 1,3-shifted isomer of (3b), in a ratio of 1 : 10. The oil had $\tau(\text{CCl}_4)$ 2.6—3.0 (m, total PhS), 4.58—4.90 (m, C=CHR), 5.25 (s, C=CH₂), 6.34 (q, CHSPh), 6.52 and 6.56 (two singlets, CH_2SPh), and the usual complex multiplet in the range 7.8—9.2, overlain by doublets (J 6 Hz) at 9.06 and 9.08 (Me_2CH) (Found: M^+ , 234.143 7. $\text{C}_{15}\text{H}_{22}\text{S}$ requires M , 234.144 1; m/e 234 (64%), 110 (100, PhS), and 109 (51, PhS). The same t.l.c. gave 3-(3-methylbutyl)but-3-en-2-ol (18b) (26 mg, 44%) as an oil, R_F 0.2, with spectroscopic properties as already described.

Preparation of Starting Materials.—2-(Phenylthio)-3-trimethylsilylpropanenitrile (23). *n*-Butyl-lithium (14 ml of a 1.5M solution in hexane) was added with stirring to a solution of di-isopropylamine (10 ml, distilled from calcium hydride) in dry tetrahydrofuran (100 ml) under nitrogen. A solution of (phenylthio)acetonitrile¹⁸ (2.98 g) in dry tetrahydrofuran (25 ml) was added to the mixture with stirring at -78°C . The mixture was maintained at -78°C for 0.5 h and then a solution of chloromethyltrimethylsilane¹⁹ (2.45 g) in dry tetrahydrofuran (10 ml) was added. The mixture was maintained at -78°C for a further 3 h, allowed to warm to room temperature over 18 h, poured into water (200 ml) and extracted with ether (3×50 ml). The combined organic extracts were washed successively with sodium hydroxide solution (50 ml, 10%), dilute hydrochloric acid (50 ml), and sodium hydrogencarbonate solution (50 ml, saturated), dried (Na_2SO_4) and evaporated *in vacuo*. The black oil was chromatographed on silica gel (100 g); elution with carbon tetrachloride gave (phenylthio)(trimethylsilyl)-

methane ²⁰ (0.2 g) identical (i.r. and n.m.r.) with an authentic sample. Further elution with 10% dichloromethane-carbon tetrachloride gave 2-(phenylthio)-3-(trimethylsilyl)propanenitrile (23) (3.31 g, 71%), b.p. 79–81 °C at 0.05 Torr (Found: C, 61.2; H, 7.40; N, 5.80. C₁₂H₁₇NSSi requires C, 61.2; H, 7.30; N, 5.90%), *R*_F (30% acetone-petrol) 0.6, *v*_{max.} (liquid film) 2 240 (C≡N), 1 580 (aromatic) and 1 260 cm⁻¹ (SiMe₃); *τ*(CCl₄) 2.2–2.9 (5 H, m, PhS), 6.35 (1 H, dd, *J* 7 and 9 Hz, CHCN), 8.78 (1 H, dd, *J* 9 and 14 Hz, CH_AH_B), 8.85 (1 H, dd, *J* 7 and 14 Hz, CH_AH_B), and 9.92 (9 H, s, SiMe₃); *m/e* 235 (11%, *M*⁺), 182 (13, PhSSiMe₃), 167 (33), 110 (36, PhSH), and 73 (100, Me₃Si).

2-(Phenylthio)-3-(trimethylsilyl)propionaldehyde (24). We used the method of Marshall *et al.*²¹ In this way 1-(phenylthio)-2-(trimethylsilyl)propionitrile (4.70 g) gave, after distillation, 2-(phenylthio)-3-(trimethylsilyl)propionaldehyde (24) (1.42 g, 30%) b.p. 96–98 °C at 0.1 Torr, *R*_F (30% acetone-petrol) 0.7; *v*_{max.} (liquid film) 1 715 (C=O), 1 580 (aromatic) and 1 250 cm⁻¹ (SiMe₃); *τ*(CCl₄) 1.00 (1 H, d, *J* 5 Hz, CHO), 2.6–3.1 (5 H, m, PhS), 6.55 (1 H, ddd, *J* 5, 7, and 8 Hz), 9.00 (1 H, dd, *J* 8 and 14 Hz, CH_AH_B), 9.13 (1 H, dd, *J* 7 and 14 Hz, CH_AH_B), and 9.92 (9 H, s, SiMe₃); *m/e* 238 (31%, *M*⁺), 209 (100, *M* – CHO), 182 (27, PhSSiMe₃), 167 (31), 128 (*M* – PhSH), and 73 (SiMe₃); the semicarbazone had m.p. 147–148 °C (from MeOH–H₂O) (Found: C, 52.8; H, 7.35; N, 14.5. C₁₃H₂₁N₃OSSi requires C, 52.8; H, 7.20; N, 14.3%).

3-(Phenylthio)-4-(trimethylsilyl)butan-2-ol (4a). 2-(Phenylthio)-3-(trimethylsilyl)propionaldehyde (24) (74 mg) in dry ether (5 ml) was added dropwise to a stirred solution of methylmagnesium iodide [prepared from magnesium (24 mg) and methyl iodide (0.1 ml)] in dry ether (10 ml) under nitrogen, and the mixture refluxed for 1 h. A standard aqueous work-up and t.l.c. (silica gel, dichloromethane) gave the mixture of alcohols (4a) (74 mg, 94%) as an oil (Found: C, 61.4; H, 8.60. C₁₃H₂₂OSSi requires C, 61.4; H, 8.70%), *R*_F 0.3, identical to that of the mixture derived below from the ketone (25); *v*_{max.} 3 540 (OH), 1 580 (aromatic) and 1 250 cm⁻¹ (SiMe₃); *τ*(CCl₄) 2.5–2.9 (5 H, m, PhS), 6.30 (1 H, dq, *J* 3 and 6 Hz, CHOH), 6.81 (1 H, ddd, *J* 3, 6, and 10 Hz, CHSPh), 7.88 (1 H, s, OH), 8.90–9.20 (5 H, m, overlain by doublet, *J* 6 Hz at 8.91, CH_AH_B and *Me*), and 9.92 (9 H, s, SiMe₃); *m/e* 254 (5%, *M*⁺), 209 (47, *M* – MeOH), 182 (30, PhSSiMe₃), 167 (40), and 110 (100, PhSH).

1-Phenyl-2-(phenylthio)-3-(trimethylsilyl)propan-1-ol (4b). In a closely similar preparation to that immediately above, 2-(phenylthio)-3-(trimethylsilyl)propionaldehyde (238 mg) and phenylmagnesium bromide [prepared from magnesium (72 mg) and bromobenzene (471 mg)] in dry ether (20 ml) gave 1-phenyl-2-(phenylthio)-3-(trimethylsilyl)propan-1-ol (308 mg, 97%), as a mixture of diastereoisomers, as an oil (Found: C, 68.5; H, 7.5. C₁₈H₂₄OSSi requires C, 68.3; H, 7.6%), *R*_F 0.4; *v*_{max.} (CCl₄) 3 540 (OH), 1 600 and 1 580 (aromatic), and 1 250 cm⁻¹ (SiMe₃); *τ*(CCl₄) 2.4–2.9 (10 H, m, Ph, PhS), 5.32–5.42 (1 H, m, CHOH), 6.60 (1 H, dt, *J* 2 and 7 Hz), 7.40 (1 H, br s, OH), 9.33 (2 H, d, *J* 7 Hz), and 10.02 (9 H, s, SiMe₃); *m/e* 316 (45%, *M*⁺), 301 (7, *M* – Me), 209 (100, *M* – PhCHOH), 191 (51), 182 (56, PhSSiMe₃), and 167 (49).

3-(Phenylthio)-4-(trimethylsilyl)butan-2-one (25). 2-(Phenylthio)-3-(trimethylsilyl)propanenitrile (470 mg) in dry ether (5 ml) was added to a solution of methylmagnesium iodide [prepared from magnesium (96 mg) and methyl iodide (0.3 ml)] in dry ether (10 ml) under nitrogen and the mixture refluxed for 20 h. An aqueous work-up

and t.l.c. (silica gel, benzene) gave the ketone (25 ≡ 9) (171 mg 34%) as an oil, identical with the sample described above.

3-(Phenylthio)-4-(trimethylsilyl)butan-2-ol (4a). Sodium borohydride (20 mg) was added at room temperature to a stirred solution of the ketone (25) (141 mg) in ethanol–water (9:1) (10 ml) and the mixture kept at room temperature for 2 h. Aqueous work-up and t.l.c. (silica gel, dichloromethane) gave the mixture of alcohols (4a) (140 mg, 99%) as an oil, *R*_F 0.3; *v*_{max.} (CCl₄) 3 540 (OH), 1 580 (aromatic) and 1 250 cm⁻¹ (SiMe₃); *τ*(CCl₄) 2.5–2.9 (5 H, m, PhS), 6.34 (1 H, quintet, *J* 6 Hz, CHOH), 6.86 (1 H, quintet, *J* 6 Hz, CHSPh), 7.88 (1 H, s, OH), 8.46–9.21 (5 H, m, overlain by doublet, *J* 6 Hz at 8.81, CH_AH_B and *Me*), and 9.92 (9 H, s, SiMe₃) (Found: *M*⁺ 254.116 5. C₁₃H₂₂OSSi requires *M*, 254.115 9); *m/e* 254 (29%), 223 (6, *M* – 15), 209 (100, *M* – MeCHOH), 182 (98, PhSSiMe₃), 167 (65), and 110 (95, PhSH).

Methyl 2-(phenylthio)-3-(trimethylsilyl)propionate (26). 2-(Phenylthio)-3-(trimethylsilyl)propanenitrile (23) (117 mg) in methanol (5 ml) containing sulphuric acid (0.5 ml) was heated under reflux for 5 days. An aqueous work-up and t.l.c. (silica gel, benzene) gave methyl 2-(phenylthio)-3-(trimethylsilyl)propionate (26) (111 mg, 83%) as an oil, *R*_F 0.6; *v*_{max.} (CCl₄) 1 740 (C=O), 1 590 (aromatic), and 1 250 cm⁻¹ (SiMe₃); *τ*(CCl₄) 2.5–2.9 (5 H, m, PhS), 6.32 (1 H, dd, *J* 6 and 11 Hz, CH), 6.44 (3 H, s, OMe), 8.72 (1 H, dd, *J* 11 and 14 Hz, CH_AH_B), 8.96 (1 H, dd, *J* 6 and 14 Hz, CH_AH_B) and 10.05 (9 H, s, SiMe₃) (Found: *M*⁺, 268.095 6. C₁₃H₂₀OSSi requires *M*, 268.095 3); *m/e* 268 (12%), 253 (34, *M* – Me), 209 (24, *M* – CO₂Me), 159 (100, *M* – PhSH), 110 (42, PhSH), and 73 (74, Me₃Si).

2-Methyl-3-(phenylthio)-4-(trimethylsilyl)butan-2-ol (4c). Methyl 2-(phenylthio)-3-(trimethylsilyl)propionate (86 mg) in dry ether (3 ml) was added dropwise to a stirred solution of methylmagnesium iodide [prepared from magnesium (24 mg) and methyl iodide (0.1 ml)] in dry ether (10 ml) under nitrogen and the mixture refluxed for 1 h. An aqueous work-up and t.l.c. (silica gel, dichloromethane) gave the alcohol (76 mg, 89%) as an oil (Found: C, 63.0; H, 9.15. C₁₄H₂₄OSSi requires C, 62.8; H, 9.00%), *R*_F 0.3; *v*_{max.} 3 540 (OH), 1 590 (aromatic), and 1 250 cm⁻¹ (SiMe₃); *τ*(CCl₄) 2.5–3.0 (5 H, m, PhS), 6.78 (1 H, dd, *J* 5 and 10 Hz, CH), 7.86 (1 H, s, OH), 8.5–9.2 (8 H, m, overlain by singlet at 8.74 and by singlet at 8.80, CH_AH_B, Me_A and Me_B) and 9.88 (9 H, s, SiMe₃); *m/e* 268 (9%, *M*⁺), 236 (40), 234 (31), 209 (53, *M* – Me₂CHOH), 182 (100, PhSSiMe₃), and 110 (77, PhSH).

5-Methyl-2-(phenylthio)-2-(trimethylsilylmethyl)hexanenitrile (27). *n*-Butyl-lithium (0.25 ml of a 2.3M solution in hexane) was added with stirring to a solution of di-isopropylamine (1 ml, distilled from calcium hydride) in dry tetrahydrofuran (5 ml) under nitrogen at room temperature. A solution of 2-(phenylthio)-3-(trimethylsilyl)propanenitrile (127 mg) in dry tetrahydrofuran (2 ml) was added to the stirred mixture, which was kept at room temperature for 0.5 h. Isopentyl iodide (0.1 ml) was added dropwise to the stirred solution, which was kept at room temperature for a further 1 h. An aqueous work-up and t.l.c. (silica gel, dichloromethane) gave the nitrile (150 mg, 91%), as an oil (Found: C, 67.0; H, 8.70; N, 4.35. C₁₇H₂₇NSSi requires C, 66.8; H, 8.90; N, 4.55%), *R*_F 0.7; *v*_{max.} (CCl₄) 2 240 (C≡N), and 1 260 cm⁻¹ (SiMe₃); *τ*(CCl₄) 2.2–2.7 (5 H, m, PhS), 8.1–9.2 (13 H, m, overlain by doublet, *J* 6 Hz at 9.03, CH, CH₂ and CH₃) and 9.74 (9 H, s, SiMe₃); *m/e* 305

(0.4%, M^+), 290 (6, $M - \text{Me}$), 279 (3, $M - \text{CN}$), 184 (100), 170 (49), and 73 (54, Me_3Si).

5-Methyl-2-(phenylthio)-2-(trimethylsilylmethyl)hexanal (28). Using the method of Marshall *et al.*,²¹ 5-methyl-2-(phenylthio)-2-(trimethylsilylmethyl)hexanenitrile (1.35 g) gave, after preparative t.l.c. (silica gel, benzene), the corresponding aldehyde (1.04 g, 76%) as an oil, R_F 0.6; ν_{max} (CCl_4) 1 720 ($\text{C}=\text{O}$) and 1 255 cm^{-1} (SiMe_3); $\tau(\text{CCl}_4)$ 0.70 (1 H, s, CHO), 2.61 (5 H, s, PhS), 8.3—9.1 (13 H, m, overlain by doublet, J 6 Hz at 9.02, CH , CH_2 and CH_3), and 9.84 (9 H, s, SiMe_3); m/e 308 (2%, M^+), 293 (7, $M - \text{Me}$), 278 (18), 263 (13), 199 (18, $M - \text{PhS}$), 182 (41, PhSSiMe_3), 143 (100), 110 (44, PhSH), and 109 (55, PhS), the semicarbazone had m.p. 168—171 °C (from $\text{MeOH}-\text{H}_2\text{O}$) (Found: C, 59.0; H, 8.36; N, 11.4. $\text{C}_{18}\text{H}_{31}\text{N}_3\text{OSSi}$ requires C, 59.1; H, 8.54; N, 11.5%).

5-Methyl-2-(phenylthio)-2-(trimethylsilylmethyl)hexan-1-ol (17a). Sodium borohydride (74 mg) was added at room temperature to a stirred solution of the aldehyde (28) (308 mg) in ethanol-water (9 : 1) (20 ml) and the mixture kept at room temperature for 3 h. An aqueous work-up and t.l.c. (silica gel, benzene) gave the alcohol (306 mg, 99%) as an oil (Found: C, 66.0; H, 9.70. $\text{C}_{17}\text{H}_{30}\text{OSSi}$ requires C, 65.8; H, 9.70%), R_F 0.3; ν_{max} (CCl_4) 3 500 (OH) and 1 260 cm^{-1} (SiMe_3); $\tau(\text{CCl}_4)$ 2.5—2.8 (5 H, m, PhS), 6.79 (2 H, s, CH_2OH), 7.67 (1 H, s, OH), 8.2—9.2 (13 H, m, CH , CH_2 and CH_3), and 9.80 (9 H, s, SiMe_3); m/e (M^+ absent) 295 (1%, $M - \text{Me}$), 279 (5, $M - \text{CH}_2\text{OH}$), 110 (100, PhSH), and 109 (94, PhS).

6-Methyl-3-(phenylthio)-3-(trimethylsilylmethyl)heptan-2-ol (17b). 5-Methyl-2-(phenylthio)-2-(trimethylsilylmethyl)hexanal (308 mg) in dry ether (3 ml) was added to a stirred solution of methylmagnesium iodide [from magnesium (36 mg) and methyl iodide (0.1 ml)] in dry ether (10 ml) under nitrogen and the mixture refluxed for 0.5 h. An aqueous work-up and t.l.c. (silica gel, benzene) gave the alcohol (321 mg, 99%) as an oil, as a 1 : 1 mixture of diastereoisomers (Found: C, 66.7; H, 9.85. $\text{C}_{18}\text{H}_{32}\text{OSSi}$ requires C, 66.6; H, 9.90%), R_F 0.6; ν_{max} (CCl_4) 3 480 (OH) and 1 255 cm^{-1} (SiMe_3); $\tau(\text{CCl}_4)$ 2.4—2.8 (5 H, m, PhS), 6.2—6.6 (1 H, m, CHOH), 7.2—7.3 and 7.8—8.0 (total 1 H, two

multiplets, OH), 8.2—9.2 (16 H, m, CH , CH_2 and CH_3) and 9.81, 9.87 (9 H, two singlets, SiMe_3); m/e 281 (100%, $M - \text{MeCHOH}$), 182 (58, PhSSiMe_3), 110 (90, PhSH) and 109 (55, PhS).

[9/1980 Received, 13th December, 1979]

REFERENCES

- 1 P. Brownbridge and S. Warren, *J.C.S. Perkin I*, 1977, 2272.
- 2 P. Brownbridge, I. Fleming, A. Pearce, and S. Warren, *J.C.S. Chem. Comm.*, 1976, 751.
- 3 B. M. Trost, K. Hiroi, and S. Kurozumi, *J. Amer. Chem. Soc.*, 1975, 97, 438.
- 4 P. Brownbridge and S. Warren, *J.C.S. Perkin I*, 1976, 2125.
- 5 K. Mislow, *Rec. Chem. Progr.*, 1967, 28, 217; D. A. Evans and G. C. Andrews, *Accounts Chem. Res.*, 1974, 7, 147.
- 6 D. A. Evans, G. C. Andrews, T. T. Fujimoto, and D. Wells, *Tetrahedron Letters*, 1973, 1389.
- 7 A. H. Davidson, I. Fleming, J. I. Grayson, A. Pearce, R. L. Snowden, and S. Warren, *J.C.S. Perkin I*, 1977, 550.
- 8 P. Brownbridge and S. Warren, *J.C.S. Perkin I*, 1977, 1131.
- 9 P. Brownbridge, Ph.D. Thesis, Cambridge, 1977, p. 32.
- 10 For some recent examples in which a proton rather than a silyl group was lost in olefin-forming reactions, see: F. A. Carey and J. R. Toler, *J. Org. Chem.*, 1976, 41, 1966; R. Hillard and K. P. C. Vollhardt, *J. Amer. Chem. Soc.*, 1977, 99, 4058; R. K. Boeckman, D. M. Blum, B. Ganem, and N. Halvey, *Org. Synth.*, 1978, 58, 152; R. B. Miller and G. McGarvey, *J. Org. Chem.*, 1978, 43, 4424; E. Ehlinger and P. Magnus, *J.C.S. Chem. Comm.*, 1979, 421.
- 11 C. Juslen and T. Enkvost, *Acta Chem. Scand.*, 1958, 12, 287.
- 12 A. A. Oswald, K. Griesbaum, W. A. Thaler, and B. E. Hudson, *J. Amer. Chem. Soc.*, 1962, 84, 3897.
- 13 P. B. D. De La Mare and C. A. Vernon, *J. Chem. Soc.*, 1953, 3555.
- 14 J. Tanaka, T. Katagiri, K. Takabe, and S. Takeshita, *Yuki Gosei Kagaku Kyokai Shi*, 1971, 29, 788 (*Chem. Abs.*, 1971, 75, 140636).
- 15 A. Delisle, *Annalen*, 1890, 260, 250.
- 16 F. C. Whitmore and L. H. Sommer, *J. Amer. Chem. Soc.*, 1946, 68, 481.
- 17 M. Julia, S. Julia, and R. Guegan, *Bull. Soc. chim. France*, 1960, 1072; E. J. Corey, R. Hartmann, and R. A. Vatakencherry, *J. Amer. Chem. Soc.*, 1962, 84, 2613.
- 18 R. Dijkstra and H. J. Bacher, *Rec. Trav. chim.*, 1954, 73, 569.
- 19 J. D. Roberts and S. Dev, *J. Amer. Chem. Soc.*, 1951, 73, 1879.
- 20 G. D. Cooper, *J. Amer. Chem. Soc.*, 1954, 76, 3713.
- 21 J. A. Marshall, N. H. Anderson, and J. W. Schlicher, *J. Org. Chem.*, 1970, 35, 858.