

Some New Chalcones¹

In connection with the study of the antifungal activity of organic compounds, we have prepared fourteen new chalcones. Except as noted, these were made by the condensation of substituted benzaldehydes and acetophenones at 0–10° in ethanol–water mixture using sodium hydroxide as a catalyst,² the duration of the reaction being about eighteen hours.

ser and drying tube, dropping funnel and thermometer, was placed 4.6 g. (0.08 mole) of potassium hydroxide and 100 ml. of anhydrous ethanol. Stirring and refluxing was initiated and continued until solution was complete, after which time the heater was replaced by an ice-bath until the temperature dropped to 5°, when 8.7 g. (0.07 mole) of benzyl mercaptan in 20 ml. of absolute ethanol was added dropwise (fifteen minutes) at that temperature. At the

CHALCONES

Substituents	M. p., ^a °C.	Formula	Analyses, % ^b			
			Calculated C	H	Found C	H
2,3-Dimethoxy ^c ^d	C ₁₇ H ₁₆ O ₃	76.10	6.01	75.95	5.59
2,4'-Dichloro ^e	82–83	C ₁₆ H ₁₀ OCl ₂	65.00	3.64	64.83	3.74
3,4-Diethoxy	128	C ₁₉ H ₂₀ O ₃	77.00	6.80	76.64	6.55
3,4-Dichloro	112–113 ^f	C ₁₆ H ₁₀ OCl ₂	65.00	3.64	64.64	3.58
4-Methoxy-2'-chloro	80–81	C ₁₆ H ₁₂ O ₂ Cl	70.46	4.80	70.37	4.72
4-Isopropyl-4'-methoxy	69–70 ^g	C ₁₉ H ₂₀ O ₂	81.40	7.19	81.08	7.35
2,4,2'-Trichloro ^e	109–110	C ₁₅ H ₉ OCl ₃	75.82	2.91	57.90	3.06
2,4-Dichloro-4'-methoxy	134	C ₁₆ H ₁₂ O ₂ Cl ₂	62.56	3.94	62.61	4.18
3,4'-Dimethoxy-4-hydroxy ^h	158–159 ⁱ	C ₁₇ H ₁₆ O ₄	71.81	5.67	71.81	5.79
3-Methoxy-4-hydroxy-4'-chloro	101 ^j	C ₁₆ H ₁₃ O ₃ Cl	66.56	4.54	66.56	4.46
3,4-Diethoxy-4'-methoxy- ^j	72–74 ^k	C ₂₀ H ₂₂ O ₄	73.60	6.80	73.35	6.83
3,4-Diethoxy-4'-methyl	96	C ₂₀ H ₂₂ O ₃	77.39	7.15	77.65	7.29
3,4-Methylenedioxy-2'-chloro	97–98 ^k	C ₁₆ H ₁₁ O ₃ Cl	67.02	3.87	67.20	4.09
3,4-Methylenedioxy-2'-hydroxy-5'-chloro	145–146 ^l	C ₁₆ H ₁₁ O ₄ Cl	63.48	3.66	63.35 ^m	3.70

^a Except as noted, all compounds were crystallized from ethanol. ^b Except as noted, all analyses are by the Clark Microanalytical Laboratory. ^c Chemical Abstracts numbering. ^d Yellow liquid, b. p. 198–199° (2 mm.). ^e Reaction time four hours. ^f Recrystallized from acetone–ethanol. ^g Recrystallized from ethanol–benzene. ^h Reaction time three hours. ⁱ Reaction temperature 20–30°. ^j Reaction time two weeks. ^k Recrystallized from methanol. ^l Reaction temperature 30–40°. ^m Analysis by Micro-Tech Laboratories.

The results of the biological tests will be reported elsewhere.

(1) This paper consists of a report of work done under contract with the Medical Division, Chemical Corps, U. S. Army.

(2) Cf. "Organic Syntheses," John Wiley & Sons, Inc., New York, N. Y., Coll. Vol. I, p. 78.

DEPARTMENT OF CHEMISTRY
DUKE UNIVERSITY
DURHAM, NORTH CAROLINA

CHARLES K. BRADSHAW
FRANCES C. BROWN
WILLIS B. BLUE

RECEIVED JUNE 2, 1949

Mercaptomethylthiazole Derivatives

2-Methyl-4-mercaptomethylthiazole.—In a 250-ml. three-neck flask fitted with a sealed stirrer, reflux condenser and drying tube, and dropping funnel was placed 16 g. (0.11 mole) of 2-methyl-4-chloromethylthiazole,¹ 10.5 g. (0.14 mole) of thiourea and 32 ml. of anhydrous ethanol. The solution was refluxed with stirring for two hours, cooled, treated with a solution of 8 g. (0.2 mole) of sodium hydroxide in 75 ml. of water and refluxed for an additional hour. To the cooled mixture was added enough 10% hydrochloric acid to show slight acidity to litmus and then 10% sodium carbonate solution to slight basicity. The mixture was extracted four times with 100 ml. of ether, and the solvent phase was dried with anhydrous calcium sulfate. The desiccant was removed by filtration, the volatiles removed from the filtrate by distillation, finally at reduced pressure and the residue distilled at 89–91° (3 mm.) to give 10.3 g. (64% yield) of a colorless oil.

*Anal.*² Calcd. for C₆H₇NS₂: N, 9.64. Found: N, 9.34.

2-Methyl-4-benzylthiomethylthiazole.—In a 250-ml. three-neck flask fitted with a sealed stirrer, reflux conden-

ser and drying tube, dropping funnel and thermometer, was placed 4.6 g. (0.08 mole) of potassium hydroxide and 100 ml. of anhydrous ethanol. Stirring and refluxing was initiated and continued until solution was complete, after which time the heater was replaced by an ice-bath until the temperature dropped to 5°, when 8.7 g. (0.07 mole) of benzyl mercaptan in 20 ml. of absolute ethanol was added dropwise (fifteen minutes) at that temperature. At the end of that time, 10 g. (0.07 mole) of freshly prepared 2-methyl-4-chloromethylthiazole in 20 ml. of ethanol was added during ten minutes, causing an immediate precipitation of potassium chloride. The mixture was stirred and refluxed for one hour, cooled, diluted with 700 ml. of water and extracted four times with 200 ml. of ether. The ethereal solution was washed with a total of 200 ml. of water and then dried with calcium sulfate. After separation of the desiccant, the solvent was removed by distillation and the residue distilled at 131–135° (0.75 mm.) to give 12.1 g. (75% yield) of a colorless oil.

Anal. Calcd. for C₁₂H₁₃NS₂: N, 5.86. Found: N, 5.35.

RESEARCH LABORATORIES
AMERICAN HOME FOODS, INC.
MORRIS PLAINS, N. J.

FRANK KIPNIS³
ISIDORE LEVY
JOHN ORNFELT

RECEIVED MAY 26, 1949

(3) Present address: Oxford Products, Inc., Cleveland, Ohio.

Thiol Esters

During work on compounds containing the thiophene-2-methyl group, it became desirable to prepare a series of thiol esters derived from thiophene-2-methyl mercaptan.¹ It was found that the mercaptan reacted in the usual manner with acyl halides and with acid anhydrides to give the desired esters in acceptable yields. The new products and their properties are tabulated in Table I.

Experimental

All of the compounds were prepared by the interaction of molar quantities of the appropriate acyl halide or anhydride with the mercaptan in excess pyridine, and worked up in the usual manner.

(1) Prepared *via* the isothiuronium salt from thienyl chloride.

(1) Hooper and Johnson, *THIS JOURNAL*, **56**, 470 (1934).

(2) Analyses by Mr. H. Soloway of this Laboratory.