

# Trans- and Cis-Selective Lewis Acid Catalyzed Hydrogermylation of Alkynes


Todd Schwier and Vladimir Gevorgyan\*

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607-7061

vlad@uic.edu

Received September 1, 2005


## ABSTRACT



The first examples of Lewis acid catalyzed hydrogermylation of alkynes have been demonstrated. It was found that this method has much higher functional group compatibility compared to the known Lewis acid catalyzed hydrosilylation and hydrostannation reactions. Remarkably, the stereochemical outcome of this hydrogermylation reaction depends on the nature of the alkyne used: proceeding via a *trans*-addition pathway with simple alkynes and *cis*-addition with propiolates. Mechanistic studies strongly support the proposed rationale on the origins of *cis*-selectivity in the hydrogermylation of activated alkynes.

Hydrometalation of alkynes is a simple and efficient approach to vinylmetals, useful synthetic intermediates in organic chemistry.<sup>1</sup> Normally, transition-metal-catalyzed hydrometalation serves as a mild protocol toward *cis*-addition products.<sup>2,3</sup> Alternatively, *trans*-addition products can be obtained via Lewis acid catalyzed hydrometalations of C–C multiple bonds (eq 1).<sup>4,5</sup> Unfortunately, these methods exhibit low functional group tolerance,<sup>6,7</sup> thus justifying development

of milder *trans*-selective general hydrometalation.<sup>8</sup> Herein, we report the first Lewis acid catalyzed hydrogermylation reaction,<sup>9</sup> which not only exhibits higher functional group tolerance in comparison to the previously reported Lewis acid catalyzed hydrometalation reactions but also exhibits perfect *trans*-addition for simple alkynes and *cis*-addition for propiolates (eq 2).



During our continuing studies on B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>-catalyzed reductions with hydrosilanes,<sup>10</sup> we found this system to be

(6) The harsh conditions employed limited the functional group compatibility to bulky TBS- or TIPS-protected alcohols; see refs 4 and 5.

(7) (Me<sub>3</sub>Si)<sub>3</sub>SiH was used in the radical and Lewis acid catalyzed hydrosilylation of terminal propiolates, though the stereochemistry of addition is not established. See: Liu, Y.; Yamazaki, S.; Yamabe, S. *J. Org. Chem.* **2005**, *70*, 556.

(1) Langkopf, E.; Schinzer, D. *Chem. Rev.* **1995**, *95*, 1375.


(2) (a) Ojima, I. In *The Chemistry of Organic Silicon Compounds*; Patai, S., Rappoport, Z., Eds.; John Wiley: Chichester, 1989; p 1479. (b) Marciniak, B. *Comprehensive Handbook on Hydrosilylation*; Pergamon Press: Oxford, 1992. (c) Hiyama, T.; Kusumoto, T. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds. Pergamon Press: Oxford, 1991; Vol. 8, p 763.

(3) An isolated case of transition-metal-catalyzed *trans*-addition has recently been disclosed. See: (a) Trost, B. M.; Ball, Z. T. *J. Am. Chem. Soc.* **2001**, *123*, 12726. (b) Trost, B. M.; Ball, Z. T.; Jöge, T. *J. Am. Chem. Soc.* **2002**, *124*, 7922.

(4) For Si, see: (a) Oertle, K.; Wetter, H. *Tetrahedron Lett.* **1985**, *26*, 5511. (b) Asao, N.; Sudo, T.; Yamamoto, Y. *J. Org. Chem.* **1996**, *61*, 7654. (c) Sudo, T.; Asao, N.; Gevorgyan, V.; Yamamoto, Y. *J. Org. Chem.* **1999**, *64*, 2494. (d) Song, Y.-S.; Yoo, B. R.; Lee, G.-H.; Jung, I. N. *Organometallics* **1999**, *18*, 3109. (e) Rubin, M.; Schwier, T.; Gevorgyan, V. *J. Org. Chem.* **2002**, *67*, 1936.

(5) For Sn, see: Asao, N.; Liu, J.-X.; Sudo, T.; Yamamoto, Y. *J. Org. Chem.* **1996**, *61*, 4568.

highly efficient for the cleavage of alkyl–aryl ethers<sup>10b</sup> and for the hydrosilylation of olefins.<sup>4c</sup> While investigating the functional group compatibility of these reactions, we discovered that treatment of ambident substrate **1a** with 1 equiv of triethylsilane resulted in quantitative formation of the cleavage product **1b** with no traces of hydrosilylation products detected (eq 3). It was hypothesized that the chemoselectivity of the reaction can be affected by switching silicon to a metal of lower oxophilicity. To this end, we examined the reactivity of **1a** with triethylgermane. *To our great delight, this reaction resulted in clean hydrogermylation without disturbing the methoxy moiety, even with excess germane (eq 3)!*



Encouraged by this result, we tested hydrogermylation of different alkynes under these reaction conditions (Table 1). It was found that, regardless of the hydrogermane used, the reaction with **1c** proceeded with perfect regio- and stereoselectivity, providing the corresponding (*Z*)-vinylgermanes in virtually quantitative yield (entries 1–4). Hydrogermylation of symmetrically substituted alkynes **1d** and **1e** proceeded uneventfully to give the corresponding products in excellent yields (entries 5 and 6). Reaction with unsymmetrical tolanes **1f** and **1g** provided the stilbene derivatives with high regioselectivity and high yields (entries 7 and 8). Likewise, terminal alkyne **1h** gave (*Z*)-vinylgermane **3hb** in 96% yield (entry 9). Aryl alkynes possessing halogen at the aromatic ring reacted uneventfully to produce the corresponding vinylgermanes in quantitative yield (entries 10 and 11). Furyl-containing alkyne **1k** reacted smoothly to give the (*Z*)-vinylgermane **3kb** in 89% yield (entry 12). Remarkably, the pinacolborane moiety did not interfere with the hydrogermylation reaction: the corresponding vinylgermane was obtained quantitatively (entry 13).

It deserves mention that this hydrometalation methodology exhibits wider functional group tolerance compared to that displayed in previously known Lewis acid catalyzed methodologies. Indeed, anisole derivative **1f** and furyl derivative **1k** would not be tolerated in any previous Lewis acid catalyzed hydrometalation reactions,<sup>4</sup> whereas employment of aryl-containing metal hydride **2d** (entry 4) would lead to a decrease in yields.<sup>11</sup> Furthermore, it should be emphasized that this method offers wider functional group tolerance vs

(8) For *trans*-selective radical hydrogermylation of alkyl-substituted alkynes with triphenylgermane, see: Ichinose, Y.; Nozaki, K.; Wakamatsu, K.; Oshima, K. *Tetrahedron Lett.* **1987**, *28*, 3709.

(9) EtAlCl<sub>2</sub> was used in the functionalization of Ge(100) surfaces, though the stereoselectivity of this transformation was not investigated. See: Choi, K.; Buriak, J. M. *Langmuir* **2000**, *16*, 7737.

(10) (a) Gevorgyan, V.; Liu, J.-X.; Rubin, M.; Benson, S.; Yamamoto, Y. *Tetrahedron Lett.* **1999**, *40*, 8919. (b) Gevorgyan, V.; Rubin, M.; Benson, S.; Liu, J.-X.; Yamamoto, Y. *J. Org. Chem.* **2000**, *65*, 6179. (c) Gevorgyan, V.; Rubin, M.; Liu, J.-X.; Yamamoto, Y. *J. Org. Chem.* **2001**, *66*, 1672.

**Table 1.** *Trans*-Hydrogermylation of Simple Alkynes

#	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Product 3	% yield <sup>a</sup>
1	Ph	Me	(c) Me (a)		>99%
2	"	"	Et (b)		99%
3	"	"	Bu (c)		>99%
4	"	"	Ph (d)		>99%
5	Bu	Bu	(d) Et (b)		95%
6	Ph	Ph	(e) Et (b)		99%
7	Ph	p-OMe-Ph	(f) Et (b)		93%
8	p-CF <sub>3</sub> -Ph	Ph	(g) Et (b)		>99%
9	Ph	H	(h) Et (b)		96%
10	m-Br-Ph	Bu	(i) Et (b)		>99%
11	o-I-Ph	Bu	(j) Et (b)		>99%
12		Bu	(k) Et (b)		89%
13		Bu	(l) Et (b)		>99%

<sup>a</sup> Isolated yields.

radical hydrogermylation,<sup>8</sup> which would not tolerate aryl halides **1i** and **1j**.<sup>12</sup>


We propose the following mechanism to account for the *trans*-selective hydrogermylation of simple alkynes (Scheme 1). Equilibration between germane **2** and B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> produces ate-complex **a**, analogous to that formed from hydrosilane

(11) It was observed that employment of arylsilanes led to decreased reaction yields in the hydrosilylation of alkynes (see ref 4b) and Friedel–Crafts alkylations of the aryl groups at silicon in the hydrosilylation of olefins (see ref 4d).

(12) Nakamura, T.; Yorimitsu, H.; Shinokubo, H.; Oshima, K. *Bull. Chem. Soc. Jpn.* **2001**, *74*, 747.

and  $B(C_6F_5)_3$ , proposed by Piers.<sup>13</sup> Direct reversible addition of coordinated gerymium<sup>14</sup> to alkyne produces vinyl cation **b**. The regiochemistry of this step is governed by the stability

**Scheme 1.** Mechanism for *Trans*-Hydrogermylation



of the forming cation, which is in perfect agreement with the observed regiochemistry of the products. Vinyl cation **b** is quenched by bulky hydride,  $^-HB(C_6F_5)_3$ , from the least hindered site to give *trans*-addition product **3** and regenerate the catalyst.<sup>15</sup>

Investigating the scope of hydrogermylation, we found that the *ester* group can be perfectly tolerated in this reaction! Thus, hydrogermylation of **2** with propiolates proceeded with very good yields (Table 2). Remarkably, in striking contrast

**Table 2.** *Cis*-Hydrogermylation of Differently Substituted Propiolates

		$R^1-C\equiv C-CO_2R^2$ (1)		$R^3-GeH$ (2)		$\xrightarrow[CH_2Cl_2, rt]{B(C_6F_5)_3 - 5 mol \%}$		$R^1-CH=CH(R^2)-GeR^3$ (4)			
#	$R^1$	$R^2$	$R^3$	Product 4	% yield <sup>a</sup>						
1	H	Et	(m)	Et (b)		4mb	89%				
2	Me	Et	(n)	Me (a)		4na	>99%				
3	n-C <sub>5</sub> H <sub>11</sub>	Me	(o)	Et (b)		4ob	>99%				
4	Ph	Et	(p)	Me (a)		4pa	85%				
5	"	"		Et (b)		4pb	>99%				

<sup>a</sup> Isolated yields.

to all known Lewis acid catalyzed hydrometalations, this reaction proceeded with *cis*-selectivity! A variety of terminal (entry 1), alkyl- (entries 2–3), and aryl-substituted propiolates (entries 4–5) reacted to give vinyl germanes **4** in high yields.


(13) Parks, D. J.; Blackwell, J. M.; Piers, W. E. *J. Org. Chem.* **2000**, *65*, 3090.

(14) For a stoichiometric reaction between solvated gerymium cation and alkyne, see: Lambert, J. B.; Zhao, Y.; Wu, H. *J. Org. Chem.* **1999**, *64*, 2729.

(15) This mechanism is analogous to that proposed by Yamamoto for Zr-catalyzed hydrostannation of alkynes (see ref 5) and by us for the hydrosilylation of alkenes (see ref 4e).


It is believed that hydrogermylation of propiolates follows a different mechanistic path than that for nonactivated alkynes (Scheme 2). Reversible coordination of propiolate

**Scheme 2.** Mechanism for *Cis*-Hydrogermylation





to  $B(C_6F_5)_3$  produces zwitterionic complex **c**, which abstracts hydride from germane **2**<sup>16</sup> to produce allenolate **d**. Trapping of the latter with gerymium-type species occurs from the less hindered face, *cis* to H, thus leading to *cis* hydrogermylation products **4**.<sup>17</sup>

To gain additional support for this mechanism, we tested the hydrogermylation of deuterium analogue d-**1i** (eq 4). If the above idea on the origins of *cis*-selectivity is correct, trapping allenolate **d** generated from d-**1i** ( $R^1 = D$ ) with gerymium species should have no facial preference, thereby generating an equimolar mixture of *cis*- and *trans*-addition products d-**3ib** and d-**4ib**. Indeed, the experiment resulted in a 1:1 mixture of d-**3ib** and d-**4ib**, thus confirming the proposed rationale.



To test the synthetic utility of hydrogermylation, we pursued transformation of the obtained vinylgermanes into synthetically useful vinylhalides. As expected, halodegermylation of **3b** proceeded efficiently, producing vinyl halides **5** in excellent yields (eq 5).<sup>18</sup> Iododegermylation of **4** under slightly modified conditions proceeded smoothly, affording vinyl iodides **6** in good yields (eq 6). Importantly, all halodegermylation reactions proceeded with perfect retention of the double-bond geometry.<sup>19</sup>



(16) It was shown that hydrogermanes are excellent hydride donors for carbocations. See: Mayr, H.; Basso, N. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 1046.

(17) An analogous mechanism was proposed for the  $AlCl_3$ -catalyzed hydrosilylation of terminal propiolates. See ref 6.

In summary, we have demonstrated the first examples of Lewis acid catalyzed hydrogermylation of alkynes. The hydrogermylation reaction exhibits excellent chemoselectivity compared to the analogous hydrosilylation reaction owing to the lower oxophilicity of germanium relative to silicon. Depending on the nature of alkyne, the reaction proceeds with high *trans*- or *cis*-selectivity. It was also shown that halodegermylation of obtained vinylgermanes proceeded smoothly to afford the corresponding (*Z*)- and (*E*)-vinyl halides in high chemical yield, with complete preservation of the double-bond geometry. This methodology allowed for efficient synthesis of bifunctional vinylgermanes **3ib**, **3jb**,

---

(18) For halodegermylation reactions, see: Oda, H.; Morizawa, Y.; Oshima, K.; Nozaki, H. *Tetrahedron Lett.* **1984**, 25, 3221.

(19) GC/MS analysis of the reaction mixtures showed formation of a single product **6**. Chromatographic isolation of which results in formation of a trace amount (<5%) of *Z*-vinyl iodide.

and **3ib** which can serve as either electrophilic components in transition-metal-catalyzed cross-coupling reactions at the aryl halide moiety (**3ib** and **3jb**) or as the nucleophilic component in the Suzuki cross-coupling reaction (**3ib**) followed by transformation to the respective vinyl halides via halodegermylation.

**Acknowledgment.** The support of the Petroleum Research Fund (AC1-40691), administered by the American Chemical Society, and of the National Science Foundation (CHE 0354613) is gratefully acknowledged.

**Supporting Information Available:** Experimental details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0521026