

Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Synthesis of New Fused 1,5-Benzodiazepines, Part 3

A. Khodairy^a

^a Chemistry Department, Faculty of Science, Sohag, Egypt

Version of record first published: 18 Aug 2006.

To cite this article: A. Khodairy (2005): Synthesis of New Fused 1,5-Benzodiazepines, Part 3, Phosphorus, Sulfur, and Silicon and the Related Elements, 180:8, 1893-1907

To link to this article: <http://dx.doi.org/10.1080/104265090889611>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Synthesis of New Fused 1,5-Benzodiazepines, Part 3*

A. Khodairy

Chemistry Department, Faculty of Science, Sohag, Egypt

1,3-Dihydro-4-phenyl-1,5-benzodiazepin-2-one 1a was treated with some *ylidenecyanothioacetamides* to give the corresponding *pyrido(2,3-b)benzodiazepines 3–6*. Treatment of compound **1a** with a mixture of *thiophen-2-aldehyde* and *thiourea* or *guanidine* gave the corresponding *1,3-thiazino- and pyrimido(4,5-b)benzodiazepines 7 and 8*. *3-Arylidene derivatives 9a–e* and **10** were synthesized. Compound **10** was subject to react with *2-(1-methylthio-1'-anilinomethylidene)malononitrile* to give *oxazino-benzodiazepine 11*. *Thieno(3,2-b)benzodiazepines 12a,b* and **13** were synthesized via the reaction of compound **1b** with *sulfur* and some active nitriles. *[1,3-Dihydro-4-phenyl(1,5)-benzodiazepin-2-ylidene]malononitrile 15* was used as *synthon* to obtain novel *pyrido-, pyrano-, benzo-, and thienobenzodiazepines 16–20*, respectively. The reaction of compound **1b** with CS_2 or PhNCS along with *1,1,3-tricyano-2-aminoprop-1-ene*, *2-(1-methylthio-1'-anilinomethylidene)malononitrile*, or *1,3-dibromopropane* gave the corresponding *polyfused benzodiazepines 21–23*, respectively.

Keywords 1,3-Dihydro-4-phenyl-1,5-benzodiazepin-2-one; pyridobenzodiazepines; pyrimidobenzodiazepines; thiazinobenzodiazepines; oxazinobenzodiazepines; thienobenzodiazepines

INTRODUCTION

Benzodiazepines have been widely employed in clinical practices as anxiolytics,¹ sedative-hypnotics,² anticonvulsant vasopressin antagonists,³ and anticipated CNS-depressants.⁴ Interestingly, *benzothienobenzodiazepine (Y-931)*, *dibenzo[b,f]diazepine (clozapine)*, and *thienobenzodiazepine (olanzapine)* are known as typical effective antipsychotics.^{5,6}

This prompted us to continue our previous work^{7,8} to synthesize some new fused and spiro benzodiazepines. So, we report herein the synthesis of novel *pyridine, pyran, thiophene, pyrimidine, oxazine, thiazine, and naphthyridine rings* fused with *benzodiazepine moiety*.

Received June 6, 2004; accepted September 7, 2004.

*For Part 1 and Part 2, see references 7 and 8.

Address correspondence to A. Khodairy, Chemistry Department, Faculty of Science, Sohag, 82524 Egypt. E-mail: khodairy@yahoo.com




Downloaded by [Laurentian University] at 06:08 16 April 2013

Downloaded by [Laurentian University] at 06:08 16 April 2013

Downloaded by [Laurentian University] at 06:08 16 April 2013

Downloaded by [Laurentian University] at 06:08 16 April 2013



SCHEME 2

3-Arylidene-1,5-benzodiazepine derivatives **9a-e** and **10** were synthesized through the reaction of compound **1a** with some aromatic carbonyl reagents including *o*-, *m*-, and *p*-chlorobenzaldehyde; *p*-nitrobenzaldehyde; *p*-N,N-dimethylaminobenzaldehyde; or isatin in glacial acetic acid and fused sodium acetate as a catalyst. Compound **10** was subjected to react with 2-(1-methylthio-1'-anilinomethylidene)malononitrile¹² in *tert.* butanol to afford the corresponding 1,3-oxazine[6,5-b](1,5)benzodiazepine derivative **11**. IR spectrum of

TABLE I Analytical and Spectral Data of the New Compounds

Product no	M.P (°C) ^a cryst. solvent	Yield (%)	Mol. form. (mol. wt.)	Analytical data (cal./found) ^b				IR (Cm ⁻¹) ^c	H-NMR (δ ppm) ^d
				C	H	N	S		
2_a	166 (EtOH)	80	C ₂₆ H ₂₀ N ₄ O ₃ S (468.53)	66.50 66.77	4.30 4.65	11.96 11.70	6.84 6.65	3433, 3366, 3211(NH, NH ₂), 2214(CN), 1898 (CO), 1432(CS).	10.2(s, 1H, NH), 8.0–7.3(m, 12H, arom.), 6.3(m, 1H, CH–Ar), 7.0[br, 1H, CH(CN)], 5.5–5.2(br, 2H, NH ₂), 4.0(s, 2H, CH ₂), 3.5(d, 1H, CH _{benzodiazepine}).
2_b	187 (Dioxane)	86	C ₂₅ H ₁₉ FN ₄ OS (442.51)	67.86 67.69	4.33 4.56	12.66 12.43	7.25 7.10	3321, 3300, 3201(NH, NH ₂), 2224(CN), 1890 (CO), 1430(CS).	10.0(s, 1H, NH), 8.1–7.5(m, 13H, arom.), 6.2(t, 1H, CH–Ar), 7.2[br, 1H, CH(CN)], 5.5–5.1(br, 2H, NH ₂), 3.8(d, 1H, CH _{benzodiazepine}).
2_c	177 (Acetic acid)	90	C ₂₃ H ₁₈ N ₄ OS ₂ (430.54)	64.16 64.32	4.21 4.04	13.01 13.32	14.89 14.76	3400, 3326, 3241(NH, NH ₂), 2240(CN), 1893 (CO), 1442(CS).	10.0(s, 1H, NH), 8.2–7.7(m, 12H, arom.), 6.0(m, 1H, CH–Ar), 5.9[br, 1H, CH(CN)], 5.3–5.0(br, 2H, NH ₂), 3.7(d, 1H, CH _{benzodiazepine}).
2_d	219 (DMF)	66	C ₂₃ H ₁₉ N ₅ OS (413.50)	66.81 66.93	4.30 4.54	16.94 16.75	7.75 7.93	3344, 3312, 3265(2NH, NH ₂), 2211(CN), 1700 (CO), 1422(CS).	10.2(s, 1H, NH), 9.2(s, 1H, NH _{pyrrole}), 8.0–7.5(m, 12H, arom.), 6.1(m, 1H, CH–Ar), 5.7[br, 1H, CH(CN)], 5.3–4.9(br, 2H, NH ₂), 3.5(d, 1H, CH _{benz}).
3_a	290 (Dioxane)	85	C ₂₆ H ₁₆ N ₄ O ₂ S (448.5)	69.63 69.43	3.60 3.43	12.49 12.51	7.15 7.34	3333, 3200(NH), 2242 (CN), 1430(CS).	13.4(s, 1H, NH), 10.0(s, 1H, NH), 8.2–7.1 (m, 13H, arom.).
3_b	266 (EtOH)	80	C ₂₅ H ₁₅ FN ₄ O (422.48)	71.07 71.23	3.58 3.30	13.26 13.42	7.59 7.32	3321, 3211(NH), 2211 (CN), 1436(CS).	12.4(s, 1H, NH), 9.4(s, 1H, NH), 8.3–7.6 (m, 12H, arom.).

3_c	270 (DMF)	71	C ₂₃ H ₁₄ N ₄ S ₂ (410.00)	67.30	3.44	13.65	15.62	3300, 3241(NH), 2251 (CN), 1440(CS).	10.4(s, 1H, NH), 9.4(s, 1H, NH), 8.3–7.7 (m, 12H, arom.).
3_d	240-3 (DMF)	67	C ₂₃ H ₁₅ N ₅ S (393.46)	70.21	3.84	17.80	8.15	3341, 3231(NH), 2231 (CN), 1432(CS).	11.4(s, 1H, NH), 9.9(s, 1H, NH), 8.0–7.1 (m, 12H, arom.).
4	222 (EtOH)	68	C ₂₄ H ₂₂ N ₄ S (398.52)	72.33 72.44	5.56 5.73	14.06 14.31	8.08 8.23	3266, 3200(2NH), 2133 (CN), 1444(CS).	12.4(s, 1H, NH), 8.4(s, 1H, NH), 8.3–7.8 (m, 9H, arom.), 3.8(s, 1H, CH, CN), 3.0–1.0(m, 10H, 5CH ₂).
5_a	220 (EtOH)	69	C ₃₀ H ₂₂ N ₄ O ₄ S (534.59)	67.40	4.15	10.48	6.00	3280(NH), 2233(CN).	11.0(s, 1H, NH), 8.3–7.7(m, 12H, arom.), 4.2(s, 2H, SCH ₂), 4.0(s, 2H, CH ₂).
5_b	210 (Benzene)	64	C ₂₈ H ₁₇ N ₅ O ₂ S (487.53)	68.98 68.64	3.51 3.70	14.35 14.49	6.58 6.43	3220(NH), 2143(CN), 1734(CO).	13.0(s, 1H, NH), 8.0–7.4(m, 4H, 12H, arom.), 4.4–4.3(m, 4H, CH ₂ ester + SCH ₂), 3.9(s, 2H, CH ₂) 1.3–1.1(t, 3H, CH ₃).
5_c	222 (EtOH)	44	C ₃₄ H ₂₂ N ₄ O ₃ S (566.63)	72.07 72.33	3.91 3.72	9.89 9.71	5.66 5.87	3224(NH), 2220 (CN), 1690 (CO).	9.0(s, 1H, NH), 8.3–7.7(m, 17H, arom.), 4.0(s, 2H, SCH ₂), 3.7(s, 2H, SCH ₂).
5_d	238 (DMF)	76	C ₃₄ H ₂₃ N ₅ O ₃ S (581.65)	70.21 70.43	3.99 3.85	12.04 12.24	5.51 5.78	3210, 3122(2NH), 2218 (CN), 1690 (CO).	9.0(s, 1H, NH), 8.3–7.7(m, 17H, arom. + NH), 4.0(s, 2H, CH ₂), 3.7(s, 2H, SCH ₂).
6	>330 (DMF)	50	C ₃₀ H ₂₂ N ₄ O ₄ S (534.59)	67.40 67.66	4.15 4.33	10.48 10.57	6.00 6.12	3421, 3350, 3210(NH, NH ₂), 1734(CO).	10.0(s, 1H, NH), 8.4–7.4(m, 12H, arom.), 6.0–5.5(br, 2H, NH ₂), 4.4–4.3(q, 2H, CH ₂ ester), 3.9(s, 2H, CH ₂) 1.3–1.1(t, 3H, CH ₃).
7	305 (Dioxane)	80	C ₂₁ H ₁₆ N ₄ S ₂ (388.50)	64.92 64.62	4.15 4.36	14.42 14.48	16.50 16.81	3344, 3200, 3182(NH + NH ₂).	13.0(s, 1H, NH), 8.2–7.5(m, 12H, arom.), 5.4–5.0(br, 2H, NH ₂).

(Continued on next page)

TABLE I Analytical and Spectral Data of the New Compounds (Continued)

Product no	M.P (°C) ^a cryst. solvent	Yield (%)	Mol. form. (mol. wt.)	Analytical data (cal./found) ^b				IR (Cm ⁻¹) ^c	H-NMR (δ ppm) ^d
				C	H	N	S		
8	310 (Dioxane)	50	C ₂₁ H ₁₅ N ₅ S (39.44)	68.27 68.54	4.09 4.23	8.96 8.88	8.68 8.92	3350, 3344, 3212, 3112 (2NH + NH ₂).	11.0(s, 1H, NH), 8.3–7.7(m, 13H, arom. + NH), 6.4–6.0(br, 2H, NH ₂).
9 _a	288 (Benzene)	90	C ₂₂ H ₁₅ ClN ₂ O (358.86)	73.64	4.21	7.81		3221(NH), 1700(CO).	13.5(s, 1H, NH), 8.1–7.6(m, 13H, arom.), 6.9(s, 1H, =CH).
9 _b	300 (EtOH)	98	C ₂₂ H ₁₅ ClN ₂ O (358.86)	73.64	4.21	7.81		3201(NH), 1702(CO).	14.5(s, 1H, NH), 8.2–7.3(m, 13H, arom.), 6.8(s, 1H, =CH).
9 _c	321 (EtOH)	87	C ₂₂ H ₁₅ ClN ₂ O (358.86)	73.40	4.00	7.78		3211(NH), 1700(CO).	11.5(s, 1H, NH), 8.1–7.6(m, 13H, arom.), 6.9(s, 1H, =CH).
9 _d	278 (Acetic acid)	83	C ₂₂ H ₁₅ N ₃ O ₃ (369.38)	71.54	4.09	11.38		3231(NH), 1702(CO).	12.5(s, 1H, NH), 7.9–7.2(m, 13H, arom.), 6.9(s, 1H, =CH).
9 _e	244 (EtOH)	95	C ₂₄ H ₂₁ N ₃ O (367.44)	78.45 78.22	5.70 5.65	11.44 11.23		3221(NH), 1700(CO).	13.5(s, 1H, NH), 8.5–7.8(m, 13H, arom.), 7.4(s, 1H, =CH), 2.3(s, 6H, N(CH ₃) ₂).
10	>330 (DMF)	97	C ₂₃ H ₁₅ N ₃ O ₂ (365.39)	75.61 75.43	4.14 4.32	11.50 11.65		3321, 3231(2NH), 1720 _{isatine} , 1704(2CO).	13.5(s, 1H, NH), 10.0(s, 1H, NH _{isatine}), 8.2–7.5 (m, 13H, arom.).
11	165 (EtOH)	58	C ₃₃ H ₂₀ N ₆ O ₂ (532.5)	74.43 74.12	3.79 3.55	15.78 15.54		3221(NH), 2180(CN).	13.0(s, 1H, NH), 10.2(s, 1H, NH _{isatine}), 8.0–7.2 (m, 18H, arom.).
12 _a	181 (EtOH)	75	C ₂₀ H ₁₆ N ₄ S (344.43)	69.74 69.56	4.68 4.60	16.27 16.46	9.31 9.50	3221, 3168(NH ₂), 2100 (CN).	8.6–7.0 (m, 9H, arom.), 6.4–6.0(br, 2H, NH ₂), 2.2–2.0(q, 2H, CH ₂), 1.1–0.9(t, 3H, CH ₃).

12_b	215 (Dioxan)	65	C ₂₂ H ₂₁ N ₃ O ₂ S (391.49)	67.50 67.70	5.41 5.26	10.73 10.80	8.19 8.00	3331, 3240(NH ₂), 1730 (CO).	7.6–7.0 (m, 9H, arom.), 5.4–5.0(br, 2H, NH ₂), 4.3–4.0(q, 2H, CH ₂ ester), 2.3–2.0 (q, 2H, CH ₂), 1.1–0.9(t, 3H, CH ₃).
13	285 (EtOH)	57	C ₂₃ H ₁₈ N ₆ S (410.50)	67.30 67.43	4.42 4.66	20.47 20.65	7.81 7.99	3331, 3300, 3210, 3145 (2NH ₂), 2230(2CN).	8.6–7.0 (m, 9H, arom.), 5.1–4.6(br, 2H, NH ₂ thiophene), 4.1–3.8(br, 2H, NH ₂), 2.2–2.0(q, 2H, CH ₂), 1.1–0.9(t, 3H, CH ₃).
14	>330 (DMSO)	43	C ₃₀ H ₂₃ N ₇ S ₂ (545.8)	66.03 66.32	4.25 4.40	17.97 17.79	11.75 11.60	3350, 3310, 3230, 3185 (2NH ₂), 2170(CN), 1433 (CS).	8.6–7.0 (m, 14H, arom.), 6.7(s, 2H, NH ₂), 5.1–4.6(br, 2HNH ₂ thiophene), 2.2–2.0(q, 2H, CH ₂), 1.1–0.9(t, 3H, CH ₃).
16_a	295 (Dioxan)	77	C ₂₅ H ₁₇ N ₅ S (419.50)	71.58 71.63	4.08 4.30	1.69	7.45 7.17	3320, 3300, 3265(NH + NH ₂), 2169(CN), 1433 (CS).	12.0(s, 1H, NH), 8.0–7.5(m, 14H, arom.), 5.5–5.1(br, 2H, NH ₂).
16_b	341 (Dioxan)	68	C ₂₆ H ₁₇ N ₅ OS (447.51)	69.78 69.52	3.83 3.68	15.65 15.54	7.16 7.01	3330, 3300, 3235(NH + NH ₂), 2149(CN), 1433 (CS).	12.4(s, 1H, NH), 8.2–7.3(m, 14H, arom.), 5.4–5.0(br, 2H, NH ₂).
17	300 (EtOH)	56	C ₂₅ H ₁₆ N ₄ O (388.43)	77.31 77.56	4.15 4.34	14.42 14.23		3215, 3199(2NH), 2200 (CN), 1689(CO).	9.9(s, 1H, NH), 7.8–7.2(m, 15H, arom. + NH).
18	281 (EtOH)	56	C ₂₅ H ₁₈ N ₄ O (390.44)	76.91 76.79	4.65 4.56	14.35 14.75		3387, 3215, 3199(NH, NH ₂), 2220 (CN), 1689(CO).	10.4(s, 1H, NH), 8.0–7.5(m, 14H, arom.), 5.2–4.9(br, 2H, NH ₂).
19	311 (EtOH)	60	C ₂₇ H ₁₆ ClN ₅ (445.91)	72.73 72.55	3.62 3.76	15.71 15.57		3320, 3300, 3265(NH + NH ₂), 2169(CN).	13.4(s, 1H, NH), 8.0 – 7.5(m, 13H, arom.), 5.0–4.6(br, 2H, NH ₂).

(Continued on next page)

TABLE I Analytical and Spectral Data of the New Compounds (Continued)

Product no	M.P (°C) ^a cryst. solvent	Yield (%)	mol. form. (mol. wt.)	Analytical data (cal./found) ^b				IR (Cm ⁻¹) ^c	H-NMR (δ ppm) ^d
				C	H	N	S		
20	175 (EtOH)	85	C ₁₈ H ₁₂ N ₄ S (316.38)	68.33 68.55	3.82 3.66	17.71 17.62	10.13 10.29	3320, 3300, 3235(NH + NH ₂), 2221(CN).	10.0(s, 1H, NH), 7.9-7.2 (m, 9H, arom.), 5.1(s, 2H, NH ₂).
21 _a	301 (Benzene)	86	C ₂₄ H ₁₈ N ₆ S ₂ (454.57)	63.42 63.64	3.99 3.82	18.49 18.60	14.11 14.37	3331, 3300, 3210, 3145 (2NH ₂), 2130(CN), 1443(CS).	8.6-7.0 (m, 9H, arom.), 6.4-6.0(br, 2H, NH ₂), 5.1-4.6(br, 2H, NH ₂), 2.2-2.0(q, 2H, CH ₂), 1.1-0.9(t, 3H, CH ₃).
21 _b	295 (EtOH)	85	C ₃₀ H ₂₃ N ₇ S (513.62)	70.16 70.36	4.15 4.36	19.09 19.26	6.24 6.46	3300, 3250, 3200, 3135 (2NH ₂), 2130(CN), 1433(CS).	8.5-7.3 (m, 14H, arom.), 6.0-5.6(br, 2H, NH ₂), 5.0-4.5(br, 2H, NH ₂), 2.3-2.1(q, 2H, CH ₂), 1.1-0.8(t, 3H, CH ₃).
22 _a	285 (EtOH)	87	C ₂₈ H ₁₉ N ₅ S ₂ (489.61)	68.69 68.44	3.91 3.80	14.30 14.12	13.10 13.31	2219, 2210(2CN), 1422(CS).	8.3-7.3 (m, 9H, arom.), 2.2-2.0(q, 2H, CH ₂), 1.1-0.9(t, 3H, CH ₃).
22 _b	287 (EtOH)	77	C ₃₄ H ₂₄ N ₆ S (548.66)	74.43 74.21	4.41 4.19	15.32 15.02	5.84 5.71	2233, 2200 (2CN), 1422(CS).	8.0-7.0 (m, 14H, arom.), 2.2-2.0(q, 2H, CH ₂), 1.1-0.9(t, 3H, CH ₃).
23	311 (Dioxane)	65	C ₂₁ H ₂₀ N ₂ OS ₂ (390.52)	66.29 66.01	5.30 5.12	7.36 7.14	16.85 16.90	2966, 2877(CH), 1704 (CO).	8.3-7.3 (m, 9H, arom.), 3.5-2.4(m, 6H, 3CH ₂), 2.2-2.0(q, 2H, CH ₂), 1.1-0.9(t, 3H, CH ₃).

^aUncorrected.


^bSatisfactory microanalyses obtained; C, ± 0.3%, H, ± 0.3%, N, ± 0.45%, S, ± 0.21.

^cMeasured on Nicolet 710 FT-IR spectrophotometer.

^dMeasured with a Varian EM 360 L using TMS as internal standard.

compound **11** showed an absorption band corresponding to the CN group at 2219 cm^{-1} (c.f. Scheme 2, Table I).


1,3-Dihydro-1-ethyl-4-phenyl-1,5-benzodiazepin-2-one **1b**¹³ was allowed to react with elemental sulfur along with malononitrile, ethyl cyanoacetate, or 1,1,3-tricyano-2-aminoprop-1-ene to yield the corresponding thieno[3,2-b]benzodiazepines **12_{a,b}** and **13**, respectively. Treatment of compound **13** with phenyl isothiocyanate¹⁴ in the presence of triethylamine as a base gave the corresponding 2-amino-3-pyrimidinyl thieno(3,2-b)-1,5-benzodiazepine derivative **14**. The reaction pathway for the formation of compound **14** was assumed to proceed via a nucleophilic attack of the ethylenic NH_2 group to the $\text{C}=\text{S}$ group followed by a nucleophilic addition of the formed NH group to the CN group (c.f. Scheme 3, Table I).



SCHEME 3

[1,3-Dihydro-4-phenyl(1,5)benzodiazepin-2-ylidene]malononitrile **15**⁸ was treated with phenyl isothiocyanate,¹⁵ benzoyl isothiocyanate, benzoyl chloride,¹⁵ or benzaldehyde¹⁵ in pyridine to give the corresponding pyrido(4,3-b)benzodiazepinethiones **16_{a,b}**, pyrido(4,3-b)benzodiazepinone **17**, or pyrano(4,3-b)benzodiazepine **18**, respectively. Moreover, treatment of compound **15** with *p*-chlorobenzylidenemalononitrile¹² in the presence of triethylamine afforded the corresponding 3-amino-2,4-dicyanobenz[b]-(1,5)benzodiazepine


derivative **19**. Thieno[3,2-b]benzodiazepine **20** was synthesized through the reaction of compound **15** with elemental sulfur in the presence of triethylamine (c.f. Scheme 4, Table I).



SCHEME 4

Compound **1b** was allowed to react with a mixture of CS₂ or PhNCS and 1,1,3-tricyano-2-aminoprop-1-ene under PTC conditions (dioxane/K₂CO₃/tetrabutylammonium bromide [TBAB]) to afford pyridothiopyrano[4,3-b]-1,5-benzodiazepine **21a** and [1,5]-benzodiazepino[6,5-c][1,8]naphthyridine **21b**, respectively. The reaction pathway¹⁶ was indicated in Scheme 5. Also, treatment of compound **1b** with a mixture of CS₂ or PhNCS and 2(1-methylthio-1'-anilinomethyl-idenemalononitrile)¹² under PTC conditions afforded the corresponding 1,3-thiazino(4,5-b)-1,5- benzodiazepine **22a** and

pyrimido-1,5-benzodiazepine **22b**, derivatives, respectively. Moreover, the reaction of compound **1b** with CS₂ and 1,3-dibromopropane in a 1:1:1 molar ratio under the same PTC conditions afforded the corresponding 3[1,3-dithi-2-ylidene]benzodiazepin-2-one derivative **23** (c.f. Scheme 5, Table I).



SCHEME 5

EXPERIMENTAL

All melting points were determined on a Koffler melting points apparatus and are uncorrected. IR spectra were obtained on a Nicolet 710

FT-IR spectrometer. ^1H -NMR spectra were recorded on a Varian EM 360 at 60 MHz using TMS as an internal reference and DMSO-d_6 as a solvent. Elemental analyses were performed on a perkin-Elmer CHN-2400°C analyzer model.

Synthesis of Compounds **2_{a-d}**

A mixture of compound **1_a** (0.01 mol, 2.36 gm), the appropriate arylidenecyanothioacetamide (0.01 mol), including benz[*b*]-3,4-dioxol-methylidene- (2.32 g), *p*-fluorobenzylidene- (2.06 g), thiene-2-ylidene- (1.94 g), or pyrrol-2-ylidenecyanothioacetamide (1.77 g), and catalytic amount of piperidine (0.2 mL) was refluxed in dioxane (20 mL) for 2 h. The formed precipitate was filtered off and crystallized from an appropriate solvent to give compounds **2_{a-d}** (cf. Scheme 2, Table I).

Synthesis of Compounds **3_{a-d}** (Method A)

The appropriate compound **2_{a-d}** (0.01 mol) was suspended in sodium ethoxide solution (0.23 g Na in 30 mL absolute ethanol) and refluxed for 3 h. The solid that formed after cooling was collected by filtration, washed with water, and crystallized from the suitable solvent to give the compounds **3_{a-d}** (cf. Scheme 2, Table I).

Synthesis of Compounds **3_{a-d}** and **4** (Method B)

A mixture of compound **1_a** (0.01 mol, 2.36 gm), the appropriate ylidene cyanothioacetamide (0.01 mol), and sodium ethoxide (0.23 g of Na in 40 mL absolute ethanol) was refluxed for 2 h. The formed precipitate was filtered off, dried, washed with water, and crystallized from an appropriate solvent to give compounds **3_{a-d}** and **4**, respectively (cf. Scheme 2, Table I).

Synthesis of Compounds **5_{a-d}**

A mixture of compound **3_a** (0.005 mol, 4.48 gm), (0.005 mol) from the appropriate halo compound (chloroacetonitrile [0.31 mL], ethyl chloroacetate [0.54 mL], phenacyl bromide [0.93 g] or chloroacetanilide [0.85 g]) and sodium acetate (0.005 mol, 0.41 gm) in ethanol (20 mL) was refluxed for 2 h. The precipitate that was obtained was filtered off, washed with water, and crystallized from the appropriate solvent to give the compounds **5_{a-d}**, respectively (cf. Scheme 2, Table I).

Synthesis of Compound 6

Compound **5_a** (0.01 mol, 5.34 g) was suspended in sodium ethoxide solution (0.35 g Na in 30 mL absolute ethanol) and refluxed for 3 h. The solid that formed after cooling was filtered off, washed with water, and crystallized from the ethanol to give the compound **6** (cf. Scheme 2, Table I).

Synthesis of Compounds 7 and 8: General Procedure

To a stirred solution of compound **1_a** (0.003 mol, 0.7 g) in methanol (25 mL), thiophene-2-aldehyde (0.003 mol, 0.28 mL), thiourea (0.003 mol, 0.22 g) or guanidine hydrochloride (0.003 mol, 0.28 g) and sodium methoxide (0.23 g Na in 5 mL methanol) were added. The reaction mixture was refluxed for 2 h, evaporated *in vacuo* and the residue was triturated with pet. ether (60–80°C). The solid product was collected by filtration and crystallized (cf. Scheme 2, Table I).

Synthesis of Compounds 9 and 10: General Procedure

A mixture of compound **1_a** (4.72 g, 0.005 mol); the appropriate carbonyl reagent (0.005 mol), including *o*-chloro- (0.7 g), *m*-chloro- (0.7 g), *p*-chloro- (0.7 g), *p*-nitro- (0.75 g), *p* N,N-dimethylaminobenzaldehyde (0.73 g), or isatin (0.73 g); and fused sodium acetate (2.3 g) in glacial acetic acid (20 mL) was refluxed for 4 h. On cooling, the precipitated solid was filtered, dried, washed with water, and crystallized from appropriate solvent (cf. Scheme 2, Table I).

Synthesis of Compound 11

Equimolar amount (0.003 mol) of compound **10** (0.28 g) and 2(1-methylthio-1'-anilinomethylidene)malononitrile (0.64 g) were dissolved in *tert.* butanol (30 mL) containing few drops of triethyl amine. The reaction mixture was refluxed until the evolution of MeSH ceased (12 h), evaporated *in vacuo*, and the remaining residue was triturated with pet. ether (40–60°C). The solid formed was collected by filtration and crystallized (cf. Scheme 2, Table I).

Synthesis of Compounds 12 and 13: General Procedure

A mixture of compound **1_b** (0.002 mol, 0.53 g); elemental sulfur (0.002 mol, 0.096 g); the appropriate active methylene (0.003 mol) including malononitrile (0.1 g), ethyl cyanoacetate (0.23 mL), or 1,1,3-tricyano-2-aminoprop-1-ene (0.26 g), and triethylamine (0.003 mol,

0.32 mL) in dioxane (20 mL) was refluxed for 3 h. The solid that formed on cooling was filtered off and crystallized from an appropriate solvent (cf. Scheme 3, Table I).

Synthesis of Compound 14

A mixture of compound **13** (0.002 mol, 0.82 g), phenyl isothiocyanate (0.002 mol, 0.23 mL), and a catalytic amount of triethylamine was refluxed in dimethylformamide (20 mL) for 2 h. The formed precipitate was filtered off and crystallized to give compound **14** (cf. Scheme 3, Table I).

Synthesis of Compounds 16_{a,b}–18: General Procedure

Equimolar amounts (0.003 mol) of compound **15** (0.89 g) and phenyl isothiocyanate (0.35 mL), benzoyl isothiocyanate (0.4 mL), benzoyl chloride (0.34 mL) or benzaldehyde (0.32 mL) in pyridine (20 mL) was refluxed for 5 h. After cooling, the reaction mixture was poured into a mixture of ice water (100 mL) and HCl (5 mL). The separated solid was then collected by filtration, washed with water, and crystallized from appropriate solvent (Scheme 4, Table I).

Synthesis of Compounds 19

(0.003 Mol) of compound **15** (0.89 g), *p*-chlorobenzylidene-malononitrile (0.7 g), and triethylamine (0.2 mL) in dioxane (20 mL) was refluxed for 4 h. On cooling, the precipitated solid was filtered off, dried, and crystallized (cf. Scheme 4, Table I).

Synthesis of Compound 20

A mixture of compound **15** (0.003 mol, 0.89 g), elemental sulfur (0.003 mol, 0.1 g), and a catalytic amount of triethylamine was refluxed in dioxane (20 mL) for 2 h. The formed precipitate was filtered off and crystallized to give compound **20** (cf. Scheme 3, Table I).

Synthesis of Compounds 21–23: General Procedure

A mixture of anhydrous potassium carbonate (3 g), dry dioxane (30 mL), compound **1_b** (0.003 mol, 0.79 g), carbon disulfide (0.003 mol, 0.22 mL) or phenyl isothiocyanate (0.003 mol, 0.36 mL) and TBAB (0.003 g) was stirred for 2 h. at room temperature and stirred for 2 h at 60°C. To the

reaction mixture, the appropriate reagent (0.003 mol), namely, 1,1,3-tricyano-2-aminoprop-1-ene (0.39 g), 2(1-methylthio-1'-anilinomethylidene)malononitrile (0.64 g), or 1,3-dibromopropane (0.6 mL), was added. The reaction mixture was stirred for 5 h at 60°C until the completion of the reaction (TLC). The reaction mixture was filtered off and the filtrate evaporated in vacuo. The residual solid was washed with water and then crystallized from the appropriate solvent (cf. Scheme 5, Table I).

REFERENCES

- [1] E. Teboul and G. Chauinard, *Can. J. Psychiatr.*, **36**, 62 (1991).
- [2] S. D. Gatzonis, E. K. Angelopoulos, E. G. Daskalopoulou, V. Mantouvalos, A. Chioni, C. Zorrunas, et al., *Drug and Alcohol Dependence*, **59**, 95 (2000).
- [3] K. Matsumoto, S. I. Kohno, K. Ojima, and H. Watanabe, *Brain Res.*, **754**, 325 (1997).
- [4] Q. Huang, R. Liu, X. He, R. McKernan, T. Gan, D. W. Bennett, et al., *J. Med. Chem.*, **41**, 4130 (1998).
- [5] K. Toshiyuki, T. Hiroshi, K. Koreichi, H. Hideki, S. Kohji, A. Masafumi, et al., *J. Heterocycle Chem.*, **39**, 163 (2002).
- [6] C. M. Beasley, G. Tollefson, P. Tran, W. Satterlee, T. Sanger, and S. Hamilton, *Neuropsychopharmacology*, **14**, 111 (1996).
- [7] H. Abdel-Ghany, A. M. El-Sayed, A. Khodairy, and H. Salah, *Synthetic Comm.*, **31**, 2523 (2001).
- [8] A. Khodairy, A. M. El-Sayed, H. Abdel-Ghany, and H. Salah, *J. Chinese Chem. Soc.*, **50**, 1195 (2003).
- [9] W. Ried and P. Stahlhofen, *Chem. Ber.*, **90**, 825, (1957).
- [10] A.-B. A. G. Ghattas, A. Khodairy, M. A. Abd-Rahman, and S. Younes, *Phosphorous, Sulfur and Silicon*, **178**, 1781 (2003).
- [11] A. Perjesi, G. Batta, and A. Foldesi, *Monatsh. Chem.*, **125**, 433 (1994).
- [12] A. Khodairy, *Phosphorous, Sulfur, and Silicon*, **160**, 159 (2002).
- [13] H. Abdel-Ghany, A. M. El-Sayed, A. A. Sultan, and A. K. El-Shafei, *Synth. Comm.*, **20**, 893 (1990).
- [14] S. M. Sherif, W. W. Wardakhan, and R. M. Mohareb, *J. Chem. Res. (S)* 356 (1996), *J. Chem. Res. (M)*, 1970 (1996).
- [15] A. M. El-Sayed and H. Abdel-Ghany, *J. Hetero. Chem.*, **37**, 1233 (2000).
- [16] A. M. M. El-Saghier, *Phosphorous, Sulfur, and Silicon*, **177**, 1213 (2002).