

Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

N,N-DIBROMOBENZENESULFONAMIDE: A USEFUL REGENRABLE REAGENT FOR BROMINATION OF VARIOUS CARBANIONIC SUBSTRATES

Mahmood Tajbakhsh ^a, Ardeshir Khazaei ^b, Majid Shabani Mahalli ^a & Ramin Ghorbani Vaghi ^b

^a University of Mazadaran, Babolsar, Iran

^b University of Bu-Ali Sina, Hamadan, Iran

Published online: 16 Aug 2010.

To cite this article: Mahmood Tajbakhsh, Ardeshir Khazaei, Majid Shabani Mahalli & Ramin Ghorbani Vaghi (2004) N,N-DIBROMOBENZENESULFONAMIDE: A USEFUL REGENRABLE REAGENT FOR BROMINATION OF VARIOUS CARBANIONIC SUBSTRATES, Phosphorus, Sulfur, and Silicon and the Related Elements, 179:6, 1159-1163, DOI: [10.1080/10426500490459768](https://doi.org/10.1080/10426500490459768)

To link to this article: <http://dx.doi.org/10.1080/10426500490459768>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

N,N-DIBROMOBENZENESULFONAMIDE: A USEFUL REGENRABLE REAGENT FOR BROMINATION OF VARIOUS CARBANIONIC SUBSTRATES

Mahmood Tajbakhsh,^a Ardeshir Khazaei,^b
Majid Shabani Mahalli,^a and Ramin Ghorbani Vaghi^b
University of Mazadaran, Babolsar, Iran,^a and University
of Bu-Ali Sina, Hamadan, Iran^b

(Received September 11, 2003; accepted October 16, 2003)

N,N-Dibromobenzenesulfonamide(dibromoamine-B), which is prepared easily in high yield, has been employed as effective brominating agent for carbanionic substrates under mild conditions. β -Diketones and β -ketoesters were brominated by this reagent without using any bases. The reagent can be recovered, rebrominated, and reused several times.

Keywords: β -Dicarbonyl compounds; bromination; dibromoamine-B

Since α -bromo carbonyl compounds and 1-bromo-1-alkynes are useful synthetic intermediates in a number of widely different organic transformations, numerous methods are available for bromination of the α -position of carbonyl compounds, alkynes, and other organic compounds using various brominating reagents. These include: $\text{KMnO}_4/\text{HBr}/\text{CH}_3\text{CN}$ ¹ and $\text{KBrO}_3/\text{KBr}/\text{Dowex}$ ² for bromination of 1,3-dicarbonyl substrates, N-bromosuccinimide for bromination of carboxylic acids,³ carbonyl-conjugated alkynes,⁴ nitro compounds,⁵ N-bromosaccharin for brominating of the benzylic and α -carbonylic positions,⁶ triphenyl phosphine and CBr_4 for bromination of primary alkynes,⁷ benzyl trimethylammonium tribromide,⁸ tetra butylammonium tribromide,⁹ trimethylamine-boron bromide,¹⁰ carbon tetrabromide,¹¹ 1,2-dibromoethane,¹² quinolinium bromochromate


We are thankful to the Mazandaran and Bu-Ali Sina Universities Research Council for the partial support of this work.

Address correspondence to Mahmood Tajbakhsh, Department of Chemistry, School of Sciences, University of Mazandaran, PO Box 453, Babolsar, Iran. E-mail: tajbakhsh@umz.ac.ir

(QBC),¹³ sodium hypobromite-sodium bisulfite ($\text{NaBrO}_3/\text{NaHSO}_3$),¹⁴ and others¹⁵ also were applied for brominating organic compounds. All these reagents have some disadvantages such as instability, low yield, toxic reagents, nonselectivity, and tedious work-up.

RESULTS AND DISCUSSION

The N,N-dibromobenzenesulfonamide is a stable crystalline compound easily prepared¹⁶ by brominating directly a sodium carbonate or bicarbonate solution of the benzenesulfonamide, which gives an almost quantitative yield of the pure product. The reaction seems to proceed in two steps, $\text{C}_6\text{H}_5\text{SO}_2\text{NBrNa}$ first dissolving to give a yellow solution; then the second step occurs, and the yellow dibromide precipitates (Scheme 1).



SCHEME 1

Treatment of carbanions with N,N-dibromobenzenesulfonamide results in the transfer of bromine from nitrogen to carbon. A broad variety of anions, including β -diketones, β -ketoesters, malonates, nitroalkanes, and alkynes can be brominated in high yield (Table I). It should be mentioned that β -diketones and β -ketoesters are brominated at room temperatures and 60°C , respectively, without using any base.

The sulfonamide formed in this reaction is easily recovered, rebrominated, and reused several times.

EXPERIMENTAL

The chemicals were obtained from Aldrich Chemical Co. and were used as received without further purification. Fluka silica gel plates (F_{254}) were used for TLC. Elemental analyses (CHN) were performed on a Perkin-Elmer 2400 CHN analyzer. The IR spectra were recorded on a Perkin-Elmer 130 spectrophotometer and ^1H NMR spectra were recorded in CCl_4 on a Varian (60 and 90 MHz) spectrometer using TMS as an internal standard.

TABLE I Bromination of Carbanions with N,N-Dibromobenzenesulfonamide

Entry	Substrate	Product	Base	Temp. (°C)	Time (min)	Yield ^a (%)
1	CH ₃ COCH ₂ COCH ₃	CH ₃ COCBr ₂ COCH ₃	—	r.t.	30	85
2	CH ₃ COCH ₂ CO ₂ C ₂ H ₅	CH ₃ COCBr ₂ CO ₂ C ₂ H ₅	—	60	240	92
3	PhCOCH ₂ COPh	PhCOCBr ₂ COPh	—	60	120	85
4	CH ₂ (CO ₂ C ₂ H ₅) ₂	CBr ₂ (CO ₂ C ₂ H ₅) ₂	NaH	0	60	87
5	CH ₃ CH(CO ₂ C ₂ H ₅) ₂	CH ₃ CBr(CO ₂ C ₂ H ₅) ₂	NaH	0	60	88
6	PhCH(CO ₂ C ₂ H ₅) ₂	PhCBr(CO ₂ C ₂ H ₅) ₂	NaH	0	60	88
7	NCCCH ₂ CO ₂ C ₂ H ₅	NCCBr ₂ CO ₂ C ₂ H ₅	NaH	0	60	90
8	CH ₃ CHNO ₂ CH ₃	CH ₃ CBrNO ₂ CH ₃	NaH	0	60	91
9	PhC≡CH	PhC≡CBr	NaH	0	60	95
10	C ₄ H ₉ C≡CH	C ₄ H ₉ C≡CBr	NaH	0	60	78

^aYield refers to the isolated products that exhibited physical and spectral properties in accordance with the assigned structure.

Preparation of N,N-Dibromobenzenesulfonamide

In a 250 ml three-necked flask were placed 5 g of benzenesulfonamide, 3.6 g of KOH and 25 ml of water; then 10 g of bromine were added with vigorous stirring. The resulting precipitate of N,N-dibromobenzenesulfonamide was filtered, washed in water and dried. The yield was 9.8 (98%).

a) Bromination of Acidic Organic Compound in the Presence of a Base

A solution of carbanion in anhydrous tetrahydrofuran [prepared in conventional fashion by adding an 80% dispersion of NaH (20 mmol) in oil to the substrate (10 mmol) in THF (10 ml)] was added under dry nitrogen to a stirred slurry of the reagent (10 mmol) in CH₂Cl₂ (15 ml) at 0°C. The reaction mixture was allowed to warm to room temperature and treated with diethyl ether (50 ml). The mixture was then washed with 0.5 M oxalic acid (30 ml), 10% aqueous potassium hydrogen carbonate (30 ml), and saturated brine (30 ml), dried (MgSO₄) and evaporated under reduced pressure. The residue was washed with n-hexane and filtered. The filtrate was evaporated to give the pure product in high yield (Table I). The solid phase was removed by filtration and recrystallized with ethanol and water to give the recovered sulfonamide in 80% yield.

b) Bromination of β-Diketones and β-Ketoesters without Base

To a solution of β-diketone or β-ketoester (9 mmol) in carbon tetrachloride or cyclohexane (20 ml) was added the reagent (9 mmol), and the

mixture was stirred at room temperature and 60°C, 5 h for β -ketoesters and 1/2 h for β -diketones. The progress of the reaction was monitored by TLC (eluent: hexane/EtoAc 5:1). The reaction mixture was filtered. The filtrate was evaporated to yield the α,α -dibromo β -diketones as pale yellow oils. All products were characterized by physical and spectral methods (M.P. IR, NMR, and CHN analysis).

3,3-Dibromo-2,4-pentandione: ^1H NMR(CCl_4) δ = 2.57 (s, 6H, 2CH₃). Anal. Calc. for C₅H₆Br₂O₂: C, 23.32; H, 2.21. Found: C, 23.21; H, 2.29.

Ethyl 2,2-dibromo acetoacetate: ^1H NMR(CCl_4) δ = 1.15(t, 3H, CH₃—CH₂); 2.59 (s, 3H, CH₃); 4.37(q, 2H, CH₃—CH₂). Anal. Calc. for C₆H₈Br₂O₃: C, 25.00; H, 2.78. Found: C, 25, 10; H, 2.80.

2-Bromo-2-nitropropane: ^1H NMR(CCl_4) δ = 2.28 (s, 6H, 2CH₃). Anal. Calc. for C₃H₆BrNO₂: C, 21.43; H, 3.57; N, 8.33. Found: C, 21.68; H, 3.90; N, 8.50.

Diethyl 2,2-dibromo malonate: ^1H NMR(CCl_4) δ = 1.25 (t, 6H, 2CH₂—CH₃); 4 (q, 4H, 2CH₃—CH₂). Anal. Calc. for C₇H₁₀Br₂O₄: C, 26.41; H, 3.14. Found: C, 26.10; H, 3.00.

Diethyl 2-bromo-2-methylmalonate: ^1H NMR(CCl_4) δ = 1.3(t, 6H, 2CH₂—CH₃); 2(s, 3H, CH₃); 4.2(q, 4H, 2CH₃—CH₂). Anal. Calc. for C₈H₁₃BrO₄: C, 37.94; H, 5.14. Found: C, 37.61; H, 4.98.

Diethyl 2-bromo-2-phenylmalonate: ^1H NMR(CCl_4) δ = 1.2(t, 6H, 2CH₂—CH₃); 4(q, 4H, 2CH₃—CH₂), 7(s, 5H, aromatic H). Anal. Calc. for C₁₃H₁₅BrO₄: C, 49.52; H, 4.76. Found: C, 49.27; H, 4.71.

1-Bromo-2-phenylacetylene: ^1H NMR(CCl_4) δ = 6.8(s, 5H, aromatic H). Anal. Calc. for C₈H₅Br: C, 53.04; H, 2.76. Found: C, 52.97; H, 2.71.

Ethyl 2,2-dibromo cyanoacetate: ^1H NMR(CCl_4) δ = 1.25(t, 3H, CH₂—CH₃); 4(q, 2H, CH₃—CH₂). Anal. Calc. for C₅H₅Br₂NO₂: C, 22.14; H, 1.84. N, 5.17. Found: C, 22.52; H, 1.99; N, 5.26.

3,3-Dibromo-1,3-diphenyl-1,3-propanedione: ^1H NMR(CCl_4) δ = 7.2(d, 10H, Aromatic H). Anal. Calc. for C₁₅H₁₅Br₂O₂: C, 47.12; H, 2.62. Found: C, 47.34; H, 2.51.

REFERENCES

- [1] S. F. Cheng, C. S. Lin, and L. K. Liu, *J. Chin. Chem. Soc.*, **44**, 309 (1997).
- [2] J. Kosmrlj, M. Kocevar, and S. Polanc, *Synth. Commun.*, **26**, 3583 (1996).
- [3] L. H. Zhang, J. Duan, Yuelian Xu, and W. R. Dolbier, *Tetrahedron Lett.*, **39**, 9621 (1998).
- [4] V. L. Heasley, D. F. Shellhamer, A. E. Chappell, et al., *J. Org. Chem.*, **63**, 4433 (1998).

- [5] A. A. Majdabadadi, R. Beugelmanes, and A. Lechevallier, *Synthesis*, 826 (1986).
- [6] E. I. Sonches and M. J. Fumarola, *J. Org. Chem.*, **47**, 1588 (1982).
- [7] A. Wagner, M. P. Heitz, and C. Mioskowski, *Tetrahedron Lett.*, **31**, 3141 (1990).
- [8] S. Kajigaeshi, T. Kakinami, H. Tokiyama, T. Hirakawa, and T. Okamoto, *Bull. Chem. Soc. Jpn.*, **60**, 2667 (1987).
- [9] S. Kajigaeshi, T. Kakinami, T. Okamoto, and S. Fujisaki, *Bull. Chem. Soc. Jpn.*, **60**, 1159 (1987).
- [10] M. Le Corre, E. Gheerbrant, and H. Le Deit, *J. Chem. Soc. Chem., Commun.*, 313 (1989).
- [11] R. T. Arnold and T. Kulenovic, *J. Org. Chem.*, **43**, 3687 (1978).
- [12] J. Ibarzo and R. M. Ortuno, *Tetrahedron Lett.*, **50**, 9825 (1994).
- [13] B. Ozgun and N. Degirmenbasi, *Synth. Commun.*, **26**, 3601 (1996).
- [14] D. Kikuchi, S. Sakaguchi, and Y. Ishii, *J. Org. Chem.*, **63**, 6023 (1998).
- [15] R. Bolton, *Bromine Compounds: Chemistry and Applications*, edited by D. Price, B. Iddon, and B. J. Wakefield (Elsevier, Amsterdam, 1998), p. 145.
- [16] S. Akiyoshi and K. Okuno, *J. Am. Chem. Soc.*, **76**, 693 (1954).