# Chemical Science

# Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. A. Straube, P. Coburger, L. Duetsch and E. Hey-Hawkins, *Chem. Sci.*, 2020, DOI: 10.1039/D0SC03604H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the Information for Authors.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/chemical-science

View Article Online

View Journal

# ARTICLE

Received 00th January 20xx.

Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Triple the fun: Tris(ferrocenyl)arene-based gold(I) complexes for redox-switchable catalysis

Axel Straube,<sup>a</sup> Peter Coburger,<sup>a</sup> Luis Dütsch,<sup>b</sup> and Evamarie Hey-Hawkins<sup>\*a</sup>

The modular syntheses of  $C_3$ -symmetric tris(ferrocenyl)arene-based tris-phosphanes and their homotrinuclear gold(I) complexes are reported. Choosing the arene core allows to fine-tune the exact oxidation potentials and thus to tailor the electrochemical response. The tris[chloridogold(I)] complexes were investigated in the catalytic ring-closing isomerisation of *N*-(2-propyn-1-yl)benzamide, showing cooperative behaviour *vs.* a mononuclear chloridogold(I) complex. Adding one, two, or three equivalents of 1,1'-diacetylferrocenium[tetrakis(perfluoro-*tert*-butoxy)aluminate] as an oxidant during the catalytic reaction (*in situ*) resulted in a distinct, stepwise influence on the resulting catalytic rates. Isolation of the oxidised species is possible, and using them as (pre-)catalysts (*ex situ* oxidation) confirmed the activity trend. Proving the intactness of the P–Au–Cl motif during oxidation, the tri-oxidised benzene-based complex has been structurally characterised.

# Introduction

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 03 August 2020. Downloaded on 8/3/2020 3:40:27 PM.

Attaining control over catalytic reactions by external stimuli, preferably with as tight a grip as demonstrated in Nature's complex regulation pathways, has long been at the centre of interest for the chemical community.<sup>1-3</sup> It has been 25 years since the seminal work by the groups of Wrighton and Rebek, the former demonstrating a cobaltocene-based rhodium complex A (Chart 1) to function as either a good hydrogenation (Co<sup>II</sup>) or hydrosilylation (Co<sup>III</sup>) catalyst, the activity for the corresponding reaction being significantly lower in the respective other oxidation state.<sup>4</sup> Rebek and co-workers used light to reversibly isomerise an azobenzene-derived organo-catalyst B and thus tuning it towards an amide-forming reaction.<sup>5</sup> A vast body of work concerning artificially switchable catalysis has since been assembled,<sup>6-9</sup> with a major focus on redox-switchable catalysis (RSC).<sup>10–13</sup> Even though switching between oxidation states is usually achieved through the addition of chemical redox agents, RSC holds great potential for applications using electrodes and thus greatly reducing chemical waste.<sup>14–18</sup> Conceptually, both ligand and metal can form the centres of (reversible) electron transfer for changing the activity state of the catalyst.<sup>11</sup> For lig-

- <sup>a.</sup> M. Sc. A. Straube, Dr. P. Coburger, <sup>[#]</sup> Prof. Dr. E. Hey-Hawkins Institute of Inorganic Chemistry, Universität Leipzig Johannisallee 29, D-04103 Leipzig (Germany)
- E-mail: <u>hev@uni-leipziq.de;</u> Website; <u>https://research.uni-leipziq.de/hh</u> <sup>b</sup> M. Sc. L. Dütsch
- Institute of Inorganic Chemistry, Universität Regensburg Universitätsstr. 31, D-93053 Regensburg (Germany)
- # Present address: Laboratory of Inorganic Chemistry, ETH Zürich
- CH-8093 Zürich (Switzerland)

<sup>+</sup> Electronic Supplementary Information (ESI) available: Experimental procedures, detailed spectral and crystallographic characterisation, DFT calculations, electrochemical data, catalysis plots. See DOI: 10.1039/x0xx00000x



**Chart 1** Examples of redox- (**A** [activity indicated for hydrogenation of alkenes],<sup>4</sup> **C**<sup>21</sup> **D**<sup>29</sup> and light-/thermo-switchable (**B**,<sup>5</sup> F<sup>42</sup>) catalyst systems. The switching moiety is highlighted in blue and the catalytic sites are highlighted in pink. The most active (ON) state is labelled in green and underlined (bold), the intermediate state in yellow and underlined (dashed), and the least active state (OFF) is labelled in red.

Science Accepted Manuscri

#### **Journal Name**

and-based switching, ferrocene has proven and remains a cornerstone for ligand design owing to its amenability to synthetic modification and favourable, while modifiable, redox properties.19,20

ARTICLE

Even though one of the first redox-switchable catalysts by Long and co-workers used for the ring-opening polymerisation of raclactide, C (Chart 1),<sup>21</sup> already contained two pendant ferrocenyl groups, and a plethora of compounds featuring multiple ferrocenyl groups is available,<sup>22-28</sup> only one report by Zhao and Chen deals with exploiting the possibility of addressing more than just two catalytic states. Employing an  $\alpha$ -diimine palladium catalyst D with two pendant ferrocenyl groups, the three activity states resulting from sequential two-step oxidation were found to differ with respect to polymerisation activity, resulting in tuneable polymer molecular weight, topology, and polydispersity.<sup>29</sup> The scarcity of this concept in RSC is surprising given how multi-state switchable molecules feature prominently in molecular machines<sup>30–33</sup> and molecular electronics and logic.<sup>34–41</sup> Combining molecular machines and catalysis, Wang and Feringa have impressively demonstrated a unidirectionally light- and thermoswitchable rotor F to display three different activity states in an organo-catalysed asymmetric Michael addition, in turn also leading to different enantioselectivities (P,P-trans: racemic; M,M-cis: S enantiomer; M,M-trans: R enantiomer).42

Expanding on this idea, we sought to prepare a system with four accessible oxidation states by making use of the C<sub>3</sub>-symmetric s-tris(ferrocenyl)arene motif recently first exploited for ligand design.<sup>43</sup> So far, we have focused on the monotopic use of these tris-phosphanes 1 (Scheme 1); however, put to use as tritopic ligands renders them miniaturised and thus easier-to-study analogues of the ferrocenyl-based dendrimers we could recently show to function as redox-switchable ruthenium(II) catalysts for the isomerisation of an allylic alcohol<sup>44,45</sup> and for transfer hydrogenation of a prochiral ketone with two distinct catalytic activity states (neutral and fully oxidised).46

 $C_3$  symmetry in general has been recognised as a promising ligand design principle,<sup>47,48</sup> and adorning one ligand fragment with multiple metal centres can furthermore allow for cooperative effects to occur.<sup>49–54</sup> The underlying key feature of the catalytically active sites in close proximity has been also found crucial in the design of single-molecule magnets<sup>55,56</sup> and for molecular recognition and supramolecular assemblies.57-60

#### **Results and discussion**

#### Preparation and electrochemical characterisation of tris-phosphanes and trinuclear gold(I) complexes

The preparation of tris-phosphanes 1 (Scheme 1, top), as described for 1c,43 starts with a triple Negishi coupling of the in situ-generated 1-bromo-1'-ferrocenylenezinc halide 2, prepared from 1,1'-dibromoferrocene,<sup>61</sup> *n*-butyllithium and anhydrous zinc chloride in THF,<sup>25,62</sup> with  $C_3$ -symmetric arenes **3a–e**. Introducing electron-withdrawing (b, c) and electron-donating (e) arenes as well as a tris-benzylic arene core (d), preventing conjugation between the three ferrocenylene groups, allows for



Scheme 1 Top: Preparation of tris-phosphanes 1a–d and complexes [[1a–c]E]. ()): Pd precatalyst, THF, r. t.; II): nBuLi, Ph<sub>2</sub>PCI, THF, -80 °C to r. t.; III): [E] X ([E] X = BH<sub>3</sub>·SMe<sub>2</sub> or [AuCl(tht)]), CH<sub>2</sub>Cl<sub>2</sub>, rt; Bottom: Crystallographically char-acterised side products obtained during the syntheses of 4d (SP1), 4e (SP2), and 4b (SP3). For their solid-state structures, see Fig. S3.

2 | J. Name., 2012, 00, 1-3

This journal is C The Royal Society of Chemistry 20xx

#### Journal Name

modularly fine-tuning the system's electrochemical response (*vide infra*). Tris(1-bromo-1'-ferrocenylene)arenes **4a–e** are obtained in moderate to good yields as crystalline solids and the potential to incorporate more and highly functionalised cores is only limited by the functional group tolerance of the Negishi protocol.<sup>63</sup> **4a,b,d,e** were analysed by single crystal X-ray diffraction (XRD) analysis (Fig. S1<sup>+</sup> and Table S3), their structural parameters falling within the expected standard ranges for *C*<sub>3</sub>-symmetric tris(ferrocenyl)arenes.<sup>25,64–68</sup> In solution, **4a–e** are characterised by unhindered rotation about the C<sub>arene</sub>–C<sub>Cp</sub> bonds on the NMR timescale, thus displaying *C*<sub>3v</sub> symmetry in their <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} NMR spectra.

Depending on the choice of Pd<sup>II</sup> precatalyst, formation of bi- and triferrocenes has also been observed. Among them, triferrocene **SP1** (Scheme 1, bottom) had not been crystallographically characterised until now<sup>69</sup> and is most likely formed from hindered reductive elimination followed by a second transmetallation step of **2** onto the Pd catalyst.<sup>70</sup> The formation of pentanuclear **SP2** during the preparation of **4e** can be similarly rationalised. Together with the isolation of only di-ferrocenylated bromotrifluorobenzene **SP3**, these findings point towards the use of our synthetic protocol to access ever more complex and functionalisable redox-active structures.

Tris-phosphanes **1a–d** are obtained following the established protocol and purified by column chromatography.<sup>43</sup> Attempting the synthesis of a mesitylene-based tris-phosphane from **4e** has only resulted in the isolation of impure trace amounts, potentially due to side reactions involving the methyl protons of **4e**. Crystals suitable for XRD analysis have been obtained for **1a** and **1d** (Fig. 1 and S4). Their structural parameters are in agreement



Fig. 1 Molecular structures of tris-phosphane 1a and borane adduct  $[1a(BH_3)_3]$ . Thermal ellipsoids are set at the 50% probability level. For clarity, the phenyl rings are depicted in wireframe style, disorder in the phenyl rings of 1a has been omitted, and hydrogen atoms except for those of the BH<sub>3</sub> groups in  $[1a(BH_3)_3]$  are not depicted. with those previously reported for  $1c^{43}$  and  $other_{tic}diphe$ nylphosphanylferrocenes.<sup>71–74</sup> Similar to the hope of the second states are observed in solution, while small changes in the <sup>31</sup>P{<sup>1</sup>H} NMR chemical shifts of 1a-d reflect the electronic nature of the arene core (Table 1).

	а	b	С	d
1	-17.5	-17.6	-18.4 <sup>43</sup>	-17.1
[1(BH₃)₃]	15.7	15.5	15.6 <sup>43</sup>	-
[1(Au)₃]	28.3	28.1	27.9ª	-

ª In CDCl₃.

Borane adducts  $[1a-c(BH_3)_3]$  have been prepared  $([1c(BH_3)_3]$  has been reported before)<sup>43</sup> to study the electrochemistry of the phosphanes (*vide infra*), since direct cyclopentadienyl–phosphorus bonds usually render the oxidations irreversible due to the lone pair of electrons on phosphorus.<sup>75–77</sup> Adduct  $[1a(BH_3)_3]$  was found to crystallise with crystallographic  $C_3$  symmetry (space group  $R\overline{3}$ , Fig. 1) and compares well (Table S5) to structural parameters of similar ferrocenyl-phosphane boranes described by us<sup>78</sup> as well as by Štěpnička and co-workers.<sup>79</sup>

Recently we reported the capability of **1c** to bind coinage metal ions in a tridentate trigonal planar coordination mode;<sup>43</sup> now we have turned our focus to potential trinuclear gold(I) complexes of **1a–c**. Indeed, reacting the tris-phosphanes with the common gold(I) precursor [AuCl(tht)] (tht = tetrahydrothiophene) in slight stoichiometric excess afforded, after simple precipitation, homotrinuclear metal complexes **[1a–c(Au)<sub>3</sub>]** in good yields. Their trinuclear composition is confirmed by CHN analyses, multinuclear NMR spectroscopy, and they remain homotrinuclear in the gas phase as assessed by high-resolution electrospray-ionisation mass spectrometry (HR-ESI MS).

Representative for all three gold(I) chloride complexes, the molecular structure of  $[1c(Au)_3]$  was determined from suitable crystals (Fig. 2) and confirms the trinuclearity in the solid state as well. In line with the experimentally determined non-centrosymmetric space group P1,  $[1c(Au)_3]$  crystallises as an enantiopure compound ( $x_{Flack} = -0.008(7)$ ); however, in solution, there is no indication of a corresponding hindered rotation as assessed from NMR spectroscopy.

Open Access Article. Published on 03 August 2020. Downloaded on 8/3/2020 3:40:27 PM.



Fig. 2 Molecular structure of homotrinuclear [ $1c(Au)_3$ ] with partial atom numbering scheme. Thermal ellipsoids are set at the 50% probability level. For clarity, the phenyl rings are depicted in wireframe style, and co-crystallised solvent, disorder in the P(2)–Au(2)–Cl(2) fragment, and hydrogen atoms have been omitted.

The P-Au bond lengths (2.225(5)-2.231(6) Å, Tables 2 and S6) are similar to mononuclear analogues featuring an N-heterocyclic substituent at the 1'-position of the 1-diphenylphosphanylferrocenylene moiety described by the groups of Siemeling (d(P-Au) = 2.242 Å [2-pyridyl], 2.234 Å [3-pyridyl])<sup>80</sup> and Lang (d(P-Au) = 2.215 Å [2,2':6',2"-terpyridin-4'-yl]),<sup>81</sup> yet [1c(Au)<sub>3</sub>] is the first such complex containing ferrocene in the tris-phosphane backbone and the second to incorporate three ferrocene moieties into a trinuclear gold complex.82

The spectral data of both [1a(Au)<sub>3</sub>] and [1b(Au)<sub>3</sub>] are in line with that of [1c(Au)<sub>3</sub>] and all complexes are thus presumed to have similar structures. Notably, [1c(Au)<sub>3</sub>] does not show aurophilic interactions<sup>83–85</sup> or close ferrocene---metal contacts<sup>86</sup> in the solid state (for M···M distances see Table 2), setting it apart from related structures regularly displaying aurophilic interactions (Table S7). In the context of cooperative effects operating in multimetallic catalysis,  $^{49,52,87-90}$  [1c(Au)<sub>3</sub>] thus falls short of the proximity criterion formulated by Feringa as the gold ... gold separation exceeds 6 Å.49 Yet, owing to the flexibility of the ligand backbone, gold ... gold distances in solution might well become much closer.

Table 2 Selected bond lengths, metal...metal distances [Å], and angles [°] of [1c(Au)<sub>3</sub>], numbered according to Fig. 2 (m = 1-3). Full parameters in Table S6.

Au(m)–P(m)	2.231(6) / 2.286(7) / 2.225(5)
Au(m)–Cl(m)	2.288(6) / 2.229(4) / 2.281(5)
P(m)– M(m) – Cl(m)	179.0(2) / 177.0(2) / 176.9(2)
Au(1,2,3)…Au(2,3,1)ª	14.187(1) / 8.012(1) / 9.100(1) / 6.4204(9)
Au(m)…Fe(m)	4.302(3) / 4.080(3) / 4.544(2)

<sup>a</sup> Intramolecular distances: the shortest intermolecular distance is shown italicised.

Regarding our vision of using 1-derived complexes as "rotary" or "dimmable" switches for multi-redox-state applications, electrochemical characterisation by cyclic voltammetry (CV) was of utmost relevance. 1,3,5-Tris(ferrocenyl)arenes have previously been shown<sup>91–93</sup> to display reversible redox activity and, in supporting electrolytes (SE) containing weakly coordinating



E [mV] vs. FcH/[FcH]

Fig. 3 Cyclic voltammograms of 4a, 1a (the grey trace was recorded with a vertex potential of 245 mV vs. FcH/[FcH]<sup>+</sup>), [1a(BH<sub>3</sub>)<sub>3</sub>], and [1a–c(Au)<sub>3</sub>] at 1 mmol·L<sup>-1</sup> in 0.1 mol·L<sup>-1</sup> (*n*Bu<sub>4</sub>N)BAr<sup>4</sup><sub>4</sub>/CH<sub>2</sub>Cl<sub>2</sub> (scan rate: 100 mV·S<sup>-1</sup>, working electrode: glassy carbon, counter electrode: platinum wire). The 2<sup>nd</sup> of 3 cycles are shown for all compounds (for full voltammograms, s. Fig. S8–12).

anions such as [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup> or [B{3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>}<sub>4</sub>]<sup>-</sup> (BAr<sup>F</sup><sub>4</sub><sup>-</sup>),<sup>94,95</sup> to furthermore be oxidisable in three separate, resolved steps.<sup>25</sup> Triazine-based 4c and [1c(BH<sub>3</sub>)<sub>3</sub>] displayed a similar behaviour,43 while 1c, as expected, showed an irreversible electrochemical oxidation due to the direct cyclopentadienyl-phosphorus linkage.75-77,96 Their analogues reported herein share these redox features (s. Fig. 3, left, for 4a, 1a, and [1a(BH<sub>3</sub>)] and Fig. S8–S11). Gratifyingly, the arene core determines the exact oxidation potentials (Tables 3 and S8-S10), in line with substitution by electron-donating or -withdrawing groups. The choice of arene thus allows for electrochemically fine-tuning the whole system. The arene trend also holds true for [1a-c(Au)<sub>3</sub>] (Fig. 3, right). In a BF<sub>4</sub>-based SE, the three gold(I) complexes show partly irreversible behaviour (Fig. S12), but in all cases, their tris(ferrocenyl)arene cores are oxidised in one (quasi)reversible event. In a BAr<sup>F</sup><sub>4</sub>-based SE, the tris(ferrocen-1-yl)arene core is again oxidised in three resolved steps. In our setup (0.1 mol·L<sup>-1</sup>  $(nBu_4N)BF_4$  or  $(nBu_4N)BAr^F_4$  in  $CH_2Cl_2$ , -1.75 to 1.2 V vs. FcH/[FcH]<sup>+</sup>),<sup>97</sup> no oxidation of gold(I) in [1a-c(Au)<sub>3</sub>] was observed.

Table 3 Redox potentials for the first iron-centred oxidation E<sup>0</sup><sub>1</sub> (vs. FcH/[FcH]<sup>+</sup>) of 4a-e, 1a-d, [1a-c(BH<sub>3</sub>)<sub>3</sub>], and [1a-c(Au)<sub>3</sub>], arranged according to ascending E<sup>0</sup><sub>1</sub>.

Ε <sup>0</sup> 1 (ΔΕ <sub>p</sub> ) <sup>a</sup> [mV]							
4d	195 (85)	1d	113 <sup>b</sup>	[1a(BH₃)₃]	270 (91)		
4e	212 (87)	1a	138 (98)	[1b(BH <sub>3</sub> ) <sub>3</sub> ]	353 (87)		
4a	247 (100)	1b	206 (116)	[1c(BH₃)₃] <sup>43</sup>	421 (100)		
4b	293 (96)	<b>1c</b> <sup>43</sup>	275 (160)	[1a(Au)₃]	376 (101)		
<b>4c</b> <sup>43</sup>	386 (127)			[1b(Au)₃]	419 (93)		
				[1c(Au)₃]	535 (108)		

<sup>a</sup> Determined on 1 mmol·L<sup>-1</sup> samples in anhydrous 0.1 mol·L<sup>-1</sup> (*n*Bu<sub>4</sub>N)BAr<sup>F</sup><sub>4</sub>/CH<sub>2</sub>Cl<sub>2</sub> (working electrode: glassy carbon) at 100 mV·s<sup>-1</sup>. The difference between oxidation and reduction potential,  $\Delta E_p$ , is given in brackets. <sup>b</sup> Determined from squarewave voltammetry due to close peak-to-peak separation, leaving  $\Delta E_p$  inaccessible.

This is in line with results from DFT calculations locating the HOMO at the iron centres; in the mono-oxidised model complex [1a(Au)<sub>3</sub>]<sup>+</sup>, the spin density solely resides at the three iron centres, too (Fig. S14–S19). Among the three tris-phosphanes, 1c

Journal Name

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 03 August 2020. Downloaded on 8/3/2020 3:40:27 PM

#### Journal Name

yields the most anodically shifted redox potentials in its complexes (Fig. 3, right, and Table 3). In line with our previous findings, complexes of 1c were also found to display the least straightforward electrochemistry such as cathodically shifted reductions connected to electron transfer-induced chemical transformations (EC mechanism; cf. Fig. S13).43

#### Redox-switchable gold(I) catalysis



**Scheme 2** Gold(I)-catalysed ring-closing isomerisation of *N*-(2-propyn-1-yl)ben-zamide **5** to oxazoline **6** (top, labelled protons used for following the reaction), including alternative mononuclear (pre-)catalyst **7**, oxidant **8**, and reductant **9**.

Seeking to demonstrate the applicability of the stepwise oxidation of the s-tris(ferrocenyl)arene core, our choice fell on the gold-catalysed 5-exo-dig ring-closing isomerisation of N-(2propyn-1-yl)benzamide 5 to 5-methylene-2-phenyl-4,5-dihydrooxazole 6 (Scheme 2) as a read-out. Uncovered by Hashmi and co-workers in 2004,98 the catalytic synthesis of oxazolines has quickly developed into a standard reaction for gold(I) complexes.<sup>99–104</sup> The groups of Sarkar and Heinze, among others, have established the transformation of 5 to 6 as a platform to perform and study RSC using gold(I) catalysts.<sup>105–109</sup> Rendering the gold(I) centres reversibly more Lewis-acidic and hence more catalytically active<sup>110</sup> through oxidation of a connected ferrocenyl moiety is one way of obviating the sometimes problematic use of silver salts for activation of gold(I) pre-catalysts by halide abstraction and enables temporal control over the activity of the catalyst.<sup>111–115</sup> Since we aimed for a detailed understanding of the switching process, the reaction was performed on the NMR scale in CD<sub>2</sub>Cl<sub>2</sub>, allowing for a time-resolved study of the reaction through protons H<sub>o</sub> (reaction involving oxidised species) and H<sub>m</sub> (reactions not involving oxidised species) of oxazoline 6 (Fig. S21, S22) vs. 1,3,5-trimethoxybenzene as an internal standard.116-118

Native [1a(Au)<sub>3</sub>] (crossed light blue circles; Fig. 4), employed in a 3 mol% Au loading (referring to substrate 5), performed with low activity (TOF =  $2.0\pm0.2$  h<sup>-1</sup>; TOF fits, s. Fig. S33–S48). A control experiment without any added gold(I) complex yielded no product. While [1b(Au)<sub>3</sub>] (crossed navy squares, TOF = 2.5±0.1 h<sup>-1</sup>) performed slightly better than [1a(Au)<sub>3</sub>], poten-

#### ARTICLE

tially due to the slightly more electron-withdrawing 1,3,5, trifluorobenzene core, [1c(Au)₃] (crossed tଧାର୍ବାଧାରଣ ୩ଜନାନ୍ତି କୋଳାରା 🕯 ମାନ୍ତି କାର୍ଯ୍ୟ 🖓 ମାନ୍ତି କାର୍ଯ୍ୟ 🖓 ମାନ୍ତି କାର୍ଯ୍ୟ 🖓 ମାନ୍ତି କାର୍ଯ୍ୟ 🖓 ମାନ୍ତି କାର୍ଯ୍ୟ କାର reproducibly, not show any catalytic activity under the same conditions. Given that the only difference is in the s-triazine core, these nitrogen atoms might interact with the amide protons of substrate 5, in turn preventing the completion of the catalytic cycle by proto-deauration.98 This hypothesis is supported comparing the <sup>1</sup>H NMR spectra of [1a(Au)<sub>3</sub>] and [1c(Au)<sub>3</sub>] with 3 equivalents of 5, respectively, at -60 °C in CD<sub>2</sub>Cl<sub>2</sub>, resulting in a signal splitting for the amide proton signal of 5 in presence of [1c(Au)<sub>3</sub>] (Fig. S23).

In order to test for potential cooperative effects due to its three gold centres, [AuCl(FcPPh<sub>2</sub>)] 7 was prepared as a mononuclear gold complex analogue of [1a(Au)3].<sup>119</sup> Quite surprisingly, a catalytic evaluation of **7** has not yet been reported. As judged from <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy ([1a(Au)<sub>3</sub>]:  $\delta$ (CD<sub>2</sub>Cl<sub>2</sub>) = 28.3 ppm; 7:  $\delta$ (CDCl<sub>3</sub>) = 28.4 ppm) and CV ([1a(Au)<sub>3</sub>]: E<sup>0</sup>((*n*Bu<sub>4</sub>N)BF<sub>4</sub>) = 340 mV; **7**: E<sup>0</sup>((*n*Bu<sub>4</sub>N)PF<sub>6</sub>) = 320 mV),<sup>119</sup> **7** is a well-suited model compound with respect to the electronic properties of both gold(I) and the ferrocene unit.

At the same concentration of gold(I), [1a(Au)3] clearly outperformed the catalytically silent 7 (pink plusses, Fig. 4). This mirrors results from Mendoza-Espinosa and co-workers who have observed a similar effect in comparing tetranuclear mesoionic carbene gold(I) halide complexes and their mononuclear analogues in the hydroamination and hydrohydrazination of terminal alkynes.<sup>90</sup> Peris and co-workers found a less prominent cooperative effect for the gold(I)-catalysed hydroamination of phenylacetylene using a trinuclear gold(I) chloride complex with a triphenylene-based tris(N-heterocyclic carbene) ligand.53

Contrastingly, an anti-cooperative effect was found when [1a(Au)<sub>3</sub>] and 7 were activated by halide abstraction (Fig. S24) using NaBAr<sup>F</sup><sub>4</sub>.<sup>117,120</sup> At a 1 mol% gold(I) loading, the catalytic activity of  $7/NaBAr^{F_4}$  (TOF = 20.1±0.6 h<sup>-1</sup>) surpassed that of  $[1a(Au)_3]/NaBAr^{F_4}$  (TOF = 3.2±0.1 h<sup>-1</sup>) greatly. Following the



cience Acceb

**Bottom**: Comparison of turnover frequencies (TOF), determined from linear fits of fixed time frames (s. Fig. S33–39) for the switched reaction phases shown above (left) and for the native complexes (right).

#### ARTICLE

chloride abstraction of **[1a(Au)<sub>3</sub>]** by <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy proved inconclusive but hinted at the slow formation of a *P*,*P'*-dicoordinated gold(I) complex ( $\delta_P$  = 42.9 ppm, Fig. S25).<sup>121-123</sup> An HR-ESI mass spectrum showed signals corresponding to [M–Au–2CI]<sup>+</sup>, [M–Au–3CI]<sup>2+</sup>, and [M–2CI]<sup>2+</sup> species. The inferior performance of **[1a(Au)<sub>3</sub>]/NaBAr<sup>F</sup>**<sub>4</sub> vs. **7/NaBAr<sup>F</sup>**<sub>4</sub> might thus relate to the formation of chelated and therefore less substrate-accessible gold(I) species.

Given its favourable redox properties – the lowest oxidation potentials and three fully reversible and well-separated oxidation events in the BAr<sup>F</sup><sub>4</sub>-based SE (cf. Fig. 3) – **[1a(Au)<sub>3</sub>]** was chosen as a model (pre-)catalyst for initial tests concerning RSC. For the oxidation, 1,1'-diacetylferrocenium tetrakis(perfluoro-*tert*butoxy)aluminate (= teflonate) **8** was chosen. Next to the ready synthetic availability from silver(I) teflonate<sup>124</sup> and 1,1'-diacetylferrocene and the sufficiently high oxidation potential (E<sup>0</sup> = 490 mV vs. FcH/[FcH]<sup>+</sup> in CH<sub>2</sub>Cl<sub>2</sub>),<sup>125</sup> the highly inert non-coordinating anion informed this choice. Smaller anions such as BF<sub>4</sub><sup>-</sup> and SbCl<sub>6</sub><sup>-</sup> have been shown to form tight ion pairs with ferrocenium,<sup>105,126</sup> and anion effects in general have been found crucial for the understanding and tailoring of gold catalysts.<sup>127</sup>

After the reaction shown in Fig. 4 had reached a 25% yield (phase I), 1.0 (orange squares), 2.0 (red triangles), or 3.0 (maroon circles) equivalents of **8** were added. The catalytic activity increased considerably (phase II; cf. Fig. 4, bottom left), in line with findings for similar systems.<sup>105–109</sup> Gratifyingly, the activity differed according to the amount of oxidant added: one equivalent of oxidant resulted in a 5.5-fold, two in a 10-fold, and three in a 13-fold increase of the experimentally determined TOFs (Fig. 4, bottom). Upon addition of decamethylferrocene (**9**) as reductant (phase III), less-than-initial activity (TOF $\varphi$  = 0.67±0.06 h<sup>-1</sup>) was restored after some delay. In control experiments, employing only **8** or **8+9** did not trigger product formation (Fig. S20, S21).

The amplified activity is usually explained by an increase in Lewis acidity at the gold(I) site,<sup>105–109</sup> and DFT calculations for mono-oxidised **[1a(Au)<sub>3</sub>]**<sup>+</sup> showed a decrease of energy of unoccupied gold-centred orbitals by over 1.9 eV (Fig. S14–S19). Such a decrease in LUMO energy can cautiously be understood in terms of increased electrophilicity.<sup>128</sup>

The different activities, which might correspond to distinct activity states in line with findings of Zhao and Chen,<sup>29</sup> were found to be easier to distinguish at 1 mol% Au catalyst loading (Fig. 5, top), decelerating the reaction in so much as to allow for an **OFF–ON–OFF–ON** switching sequence. Moreover, adding two equivalents of **8** to a catalytic run previously oxidised by one equivalent of **8** (VII to VIII, rose squares), a distinct increase of activity was found. Due to the negligible initial activity (TOF <  $0.05 h^{-1}$ ), the great increase in activity is difficult to quantify by numbers (Fig. 5, centre left). It is, however, evident that the TOFs effected by the re-oxidised species (VI to VII) are slightly larger than those originating from the first oxidation for one (rose squares) and two (pink triangles), yet not for the second oxidation with three equivalents of **8** (purple circles). Oxidising



Fig. 5 Top: Yield-over-time graphs for redox-switchable gold(1)-catalysed cyclisation of 5 to 6 using  $[1a(Au)_3]$  (1 mol% Au,  $[5]_0 = 60 \text{ mmol}\cdot\text{L}^{-1}$ , CD<sub>2</sub>Cl<sub>2</sub>, 25 °C). Arrows indicate the addition of additives. IV: initial OFF phase; V: 1<sup>st</sup> ON phase after addition of 1.0–3.0 eq. oxidant 8; VI: 2<sup>nd</sup> OFF phase after addition of 1.1–3.3 eq. oxidant 9; VII: 2<sup>nd</sup> ON phase after addition of 1.1–3.3 eq. oxidant 8; VII: 2<sup>nd</sup> OFF phase after addition of 2.2 eq. 8 to 1 eq.-switched reaction (rose squares).

**Centre**: Comparison of turn-over frequencies TOF determined from linear fits (cf. Fig. S40–48 for regression plots) of reaction phases **IV** to **VIII** (top), **IX** (bottom left), and **X** (bottom right).

**Bottom**: Left: Yield-over-time graphs for gold(1)-catalysed cyclisation of **5** to **6** using **[1b(Au)<sub>3</sub>]** (red) and **[1c(Au)<sub>3</sub>]** (yellow; conditions as above), oxidised *in situ* with 2 eq. of **8**. Right: Yield-over-time graphs for gold(1)-catalysed cyclisation of **5** to **6** using isolated oxidised complexes **[1a(Au)<sub>3</sub>](TE)**<sub>n</sub> (n = 1–3, conditions as above). Dashed lines represent the chosen timeframe for linear fitting.

what we assume to be the mono-oxidised species  $[1a(Au)_3]^+$  to presumably tri-oxidised  $[1a(Au)_3]^{3+}$  (VII to VIII) leads to an even larger reaction rate (TOF =  $4.3\pm0.2$  h<sup>-1</sup>). The ON-switching can also be used on a macroscopic level, to which end two 5 mLscale reactions (57 mg 5, 2.2 mg  $[1a(Au)_3]$ , *i.e.*, 1 mol% Au,  $[5]_0$ = 60 mmol·L<sup>-1</sup>) were compared. One run was oxidised using 3.0 eq. 8 at 12.5 h after the start and led to a quantitative yield *vs.* only 20% conversion for the non-switched case after 42.0 h (29.5 h after the ON-switch).<sup>129</sup>

These findings are invariant to the order of addition, that is, the presence of substrate during oxidation;<sup>130</sup> when **[1a(Au)<sub>3</sub>]** is oxidised with 1–3 equivalents of **8** before the addition of **5**, similar reaction profiles and TOFs result (Fig. S26, Table S13). In all cases, the first **ON**-switch is accompanied by an induction period which is absent for the re-oxidation (*e.g.*, **IV** to **V** vs. **VI** to **VII** in Fig. 5). When **[1b(Au)<sub>3</sub>]** (red triangles) is used in the cyclisation (Fig. 5, bottom right) and oxidised using two equivalents of **8**, the same behaviour, including a similar TOF (2.3±0.1 h<sup>-1</sup>) as for **[1a(Au)<sub>3</sub>]**, is found. Oxidising **[1c(Au)<sub>3</sub>]** (yellow triangles) with two equivalents of **8** converted the otherwise silent complex (*vide supra*) into an active catalyst with

**Journal Name** 

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 03 August 2020. Downloaded on 8/3/2020 3:40:27 PM.

#### Journal Name

higher TOF (3.3±0.1 h<sup>-1</sup>). Even though other redox-switchable gold(I) catalysts perform with sometimes significantly higher activity, complexes **[1a–c(Au)**<sub>3</sub>] were intended as models demonstrating the feasibility of tailorable multi-oxidation-state applications. Lower observed activities for **[1a–c(Au)**<sub>3</sub>] are hence an acceptable trade-off prior to further optimisation.<sup>105,109</sup> In the same way, the significantly higher reaction rate of using oxidised **7** (1 mol% Au, TOF = 10.6±0.4 h<sup>-1</sup>, Fig. S24) comes at the expense of not being principally able to address separate states with differing catalytic activity.

In order to gain more insight into the switching process, the oxidation of **[1a(Au)<sub>3</sub>]** in the absence of substrate was first followed by <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy (Fig. S27, S28). The <sup>31</sup>P{<sup>1</sup>H} NMR resonance shifted upfield and broadened with each added equivalent of **8**, while the subsequent addition of **9** reversed this trend. In line with the findings from the catalytic runs, the reduction took quite long to take its full effect.

Given these encouraging results, we sought to isolate the individually switched complexes [1a(Au)<sub>3</sub>](TEF)<sub>n</sub> (n = 1-3), attempting to dismiss the possibility that oxidising [1a(Au)3] during the catalytic reaction (in situ) might lead to mixtures of different oxidation states (e.g. by disproportionation). Furthermore, we wanted to ascertain the integrity of the oxidised species, since Nataro and co-workers observed loss of a chlorido ligand (G, Chart 2) from dinuclear gold(I) complexes based on 1,1'bis(phosphanyl)ferrocenes after chemical oxidation using tris(pbromophenyl)ammoniumyl tetrakis(pentafluorophenyl)-borate ("Magic Blue").131 While the group of Peris found oxidation-induced protonation (H, Chart 2) of the ferrocenyl-imidazolylidene backbone in their gold(I) chloride complexes upon addition of acetylferrocenium tetrafluoroborate,126 Heinze and coworkers described valence isomerisation from, initially, Fe<sup>III</sup>/Au<sup>I</sup> to Fe<sup>II</sup>/Au<sup>II</sup> (J, Chart 2) assisted by both the SbCl<sub>6</sub><sup>-</sup> anion of their

Nataro et al. 2016 tBu *t*Bu tBu [N(p-BrC<sub>6</sub>H<sub>4</sub>)<sub>3</sub>]tBu  $[B(C_6F_5)_4]$ CIAL AuC CH<sub>2</sub>Cl<sub>2</sub>  $[B(C_6F_5)_4]$ ťBu tBu tBu *t*Bu G Peris et al. 2016 *n*Bu *n*Bu [AcFc]BF AuCl AuCl CH<sub>2</sub>Cl<sub>2</sub> Ĥ  $BF_4$ 'nΒι 'nBu н Heinze et al. 2019 [N(p-BrC<sub>6</sub>H<sub>4</sub>)<sub>3</sub>]-**OEt** OEt [SbCl<sub>6</sub>] <sup>∼</sup>Au<sup>II</sup>CI Au<sup>I</sup>CI THE ĊI−SbCl₅ Fc

Chart 2 Behaviour of ferrocene-derived chloridogold(I) complexes upon chemical oxidation (Compounds G and H have been characterised by XRD in the solid state, while information about the structure of J has been gathered from DFT calculations and various analytical techniques).

This journal is © The Royal Society of Chemistry 20xx

oxidant and propargylic amide 5.105 In the last two instances, the respective Au–Cl fragment was found የው የይሰባቆጠባበት አርጅ 604H Gratifyingly, [1a(Au)3](TEF)n can indeed by obtained as analytically pure compounds from the reaction of [1a(Au)<sub>3</sub>] with 1, 2, or 3 equivalents of  ${\bf 8}$  as evidenced by CHN analyses. While [1a(Au)<sub>3</sub>](TEF) is a pale green powder, both [1a(Au)<sub>3</sub>](TEF)<sub>2</sub> and [1a(Au)<sub>3</sub>](TEF)<sub>3</sub> are dark-green microcrystalline solids. In contrast to [1a(Au)<sub>3</sub>], they are readily soluble in diethyl ether which allows for their purification by precipitation with pentanes, as 1,1'-diacetylferrocene is soluble under these conditions and can hence be extracted. The effective magnetic moments  $\mu_{eff}$  of [1a(Au)<sub>3</sub>](TEF)<sub>n</sub> in solution were determined using Evans' method (Fig. S99, S103, S107) and match the expected spin-only values for one (2.03  $\mu_B$ , expected: 1.73  $\mu_B$ ), two (2.98  $\mu_B$ , expected: 2.83  $\mu_B$ ), and three (3.52  $\mu_B$ , expected: 3.82  $\mu_B$ ) unpaired electrons reasonably well.<sup>132,133</sup> The <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} NMRspectroscopic features (Fig. S98-109) match those of the previously mentioned stepwise oxidation (Fig. S27, S28). HR-ESI mass spectra show peaks for the three different cations respectively, the tri-cation [1a(Au)<sub>3</sub>]<sup>3+</sup> apparently stabilised by a tight contact to the teflonate anion as {[1a(Au)3](TEF)}<sup>2+</sup> under these conditions; similarly, the <sup>19</sup>F resonance of [1a(Au)<sub>3</sub>](TEF)<sub>3</sub> ( $\omega_{1/2}$  = 17 Hz) is slightly broadened with respect to that of the monoand dioxidised species ( $\omega_{1/2}$  = 4 Hz). IR spectra (Fig. S29) of [1a(Au)<sub>3</sub>](TEF)<sub>n</sub> display both the signature of the teflonate anion and a band at 860 cm<sup>-1</sup>, characteristic for the  $\delta$ (C–H) vibration of ferrocenium.<sup>134,135</sup> In line with previous reports,<sup>136</sup> the ferrocenylene v(C-H) stretches shift to higher wavenumbers upon increasing degree of oxidation and do not indicate the presence of native or lower oxidation states as expected for disproportionation. Neither can this be inferred from UV/Vis spectra of [1a(Au)<sub>3</sub>](TEF)<sub>n</sub> (Fig. 6 top right, Fig. S30), as they differ in the position of their bands at long wavelength, most likely related to LMCT, inner ferrocenyl, and potentially even to Aul-Felli MMCT transitions,<sup>106,137,138</sup> between 500 and 1000 nm.

As final proof for the site of oxidation, we were able to isolate single crystals of the tri-oxidised complex as {[1a(Au)<sub>3</sub>]}<sub>2</sub>(TEF)<sub>6</sub> (Fig. 6, left and bottom right), thus unambiguously confirming the oxidation state of three times Fe<sup>III</sup> through the presence of six, albeit only partly crystallographically describable, teflonate anions per asymmetric unit. Notably, all three P-Au-Cl fragments are still intact and form dimers of an overall sixfold positive charge through a set of two aurophilic interactions (d(Au(1)–Au(6)) = 2.989(1) Å, d(Au(2)–Au(4)) = 3.188(1) Å) and a less-close Au…Au contact (d(Au(3)…Au(5)) = 3.397(1) Å).<sup>139</sup> While the poor crystal quality precludes detailed metric analyses, generally larger C₅H₄(centroid)–Fe distances of over 1.68 Å and, accordingly, Fe–C bond lengths of up to 2.15(2) Å, typical for ferrocenium species, 136, 140 are discernible (Table S8). Other structural parameters, particularly with respect to the P-Au-Cl are, within error, comparable to the molecular structure of [1c(Au)<sub>3</sub>]. While ferrocenium cations are frequently encountered as counter cations for metallate anions and several homoand heteromultinuclear bridged metallocenes featuring ferrocenium units have been listed in the CSD, {[1a(Au)3]}2(TEF)6 is a

View Article Online DOI: 10.1039/D0SC03604H

# ARTICLE



Fig. 6 Left: Molecular structure of {[1a(Au)<sub>3</sub>]}<sub>2</sub>(TEF)<sub>6</sub> with partial atom numbering scheme. Thermal ellipsoids are set at the 50% probability level. For clarity, *P*-bound phenyl rings are depicted in wireframe style, and co-crystallised solvent, hydrogen atoms, and anions have been omitted. Bottom right: Asymmetric unit of {[1a(Au)<sub>3</sub>]}(TEF)<sub>6</sub> with the four localisable teflonate anions in ball-and-stick style. The presence of severely disordered teflonate anions TEF(5) and TEF(6) has been confirmed through their electron density and the unoccupied volume. Top right: Stacked UV/Vis spectra of [1a(Au)<sub>3</sub>](TEF)<sub>n</sub> (n = 0–3) in CH<sub>2</sub>Cl<sub>2</sub>. The insert shows a magnification for the absorptions at long wavelength.

very rare example of a crystallographically characterised metal complex containing ferrocenium in its ligand backbone. To the best of our knowledge, a rhenium(0) carbonyl complex by Nataro and co-workers is the only other example to contain both ferrocenium and a phosphane-bound metal fragment,<sup>141</sup> while the +III oxidation state of iron in reported bis[1,1'-bis(diphenylphosphanyl)ferrocenium] hexadecachlorotetraantimony(III) by the group of Kasim<sup>142</sup> has been questioned by both Nataro and Connick.<sup>143</sup> Not backed up by a solid-state molecular structure, Grandberg and co-workers also reported a phosphanylferrocenium-based gold(III) bromide complex in 1977.<sup>144</sup> Last but not least, a very recent report by the Lapinte group details a related tetranuclear iron(II/III) half-sandwich array with potential use in molecular electronics.<sup>145</sup>

 $[1a(Au)_3](TEF)_n$  appear to be air- and moisture-insensitive in the solid state for short periods. They can be weighed out in ambient conditions but quickly turn from green to yellow solutions in wet solvents. Their stability in dry CD<sub>2</sub>Cl<sub>2</sub> was monitored by <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy (Fig. S31–33). Surprisingly, a solution of  $[1a(Au)_3](TEF)_3$  did show little change over three days while  $[1a(Au)_3](TEF)$  and  $[1a(Au)_3](TEF)_2$  slowly decom-

posed under the same conditions (room temperature, protected from light). <sup>31</sup>P{<sup>1</sup>H} resonances attributable to *P*,*P*'-diccordinate species at around 40 ppm appeared, and [**1a(Au)**<sub>3</sub>](**TEF)**<sub>2</sub> produced a metallic mirror of presumably elemental gold on the wall of the NMR tube. Furthermore, the <sup>1</sup>H NMR spectral evolution of [**1a(Au)**<sub>3</sub>](**TEF)**<sub>2</sub> (Fig. S32) indicates at least some degree of disproportionation to the mono- and trioxidised species, making it the least solution-stable of the three isolated oxidation products. Although a mass spectrum of the crystals of [**1a(Au)**<sub>3</sub>](**TEF)**<sub>3</sub> does not contain signals for dimeric species, aurophilic interactions in solution might play a role in stabilising the trication. When fresh solutions of [**1a(Au)**<sub>3</sub>](**TEF)**<sub>n</sub> (n = 1–3) are reacted with one, two, or three equivalents of reductant **9**, native [**1a(Au)**<sub>3</sub>] is re-obtained respectively (Fig. S34– S36).

The isolated oxidised species  $[1a(Au)_3](TEF)_n$  have finally also been tested for their catalytic performance (*ex situ*; Fig. 5, bottom right). Their TOFs show the previously noted distinct dependence on the degree of oxidation, with tri-oxidised  $[1a(Au)_3](TEF)_3$  (light blue circles,  $5.8\pm0.1$  h<sup>-1</sup>) performing with about threefold activity than mono-oxidised  $[1a(Au)_3](TEF)$ (navy squares,  $1.8\pm0.1$  h<sup>-1</sup>). All of them outperform the *in situ*-

#### Journal Name

emical Science Accepted Manus

oxidised species, most notably  $[1a(Au)_3](TEF)_3$ . This points towards redox equilibria between 8 and the complexes in higher oxidation states which might be overcome using stronger oxidants, thus ensuring full conversion upon addition. Even though we cannot fully exclude potential disproportionation or the presence of mixtures under these catalytic conditions, the distinctly different activities in the read-out catalytic conversion strongly suggest that  $[1a(Au)_3]$  and its analogues can function as a molecular switches with four addressable states.

Similar to the *in situ*-generated species, induction periods are observed for  $[1a(Au)_3](TEF)_n$ . Notably, the addition of a fresh batch of 5 to an almost-completed reaction after 24 h (Fig. S37) using  $[1a(Au)_3](TEF)_3$  did not result in another induction period but led to a slight loss of activity  $(3.2\pm0.1 vs. 5.0\pm0.1 h^{-1})$ . Mixing isolated  $[1a(Au)_3](TEF)_3$  with 5 (Fig. S38) in CH<sub>2</sub>Cl<sub>2</sub> led to an appreciably slow colour change which was followed by time-resolved UV/Vis spectroscopy. We thus speculate that the catalytically active species are, in general, formed from a chemical process involving 5. Bearing in mind the aforementioned results from Heinze and co-workers and in accordance with loss of the Fe<sup>III</sup>-associated absorptions of  $[1a(Au)_3](TEF)_3$  at long wavelength,<sup>105</sup> a coordination-assisted valence isomerisation from Fe<sup>III</sup>/Au<sup>I</sup> to Fe<sup>II</sup>/Au<sup>II</sup> might be at the heart of this behaviour.

### Conclusions

In summary, we have demonstrated the modular syntheses of a new class of tris(ferrocenyl)arene-based tris-phosphanes 1 which can be used to form well-defined,  $C_3$ -symmetric homotrinuclear gold(I) complexes. Four oxidation states relating to the tris(ferrocenyl)arene backbone - non-, mono-, di-, and tri-oxidised - have been identified by cyclic voltammetry. Stoichiometric oxidation of [1a(Au)<sub>3</sub>] produces isolable products [1a(Au)<sub>3</sub>](TEF)<sub>n</sub>. This redox behaviour can be advantageously used in redox-switchable catalysis ex and in situ, as we were able to show for the proof-of-principle ring-closing isomerisation of N-(2-propyn-1-yl)benzamide (5) forming oxazoline 6. The arene cores determine the exact redox potential and were also found to influence the catalytic performance of the native and oxidised species, a feature which we are currently investigating in more detail. Metal complexes of 1 and its analogues thus hold great promise for applications in molecular electronics and logic, possibly extending binary to quaternary signal processing, as four different oxidation states can be addressed and isolated.

## **Conflicts of interest**

There are no conflicts to declare.

## Acknowledgements

Financial support by the Studienstiftung des deutschen Volkes (doctoral fellowships to P. C. and A. S.), the DFG (He 1376/51-1) and the Graduate School BuildMoNa is gratefully acknowledged. L. D. acknowledges funding from DFG Sche 384/36-1.

The authors thank Dr. Andy Schmied for the XRD measurement of 4a, Dr. Peter Lönnecke for help with the structure solution of {[1a(Au)<sub>3</sub>]}<sub>2</sub>(TEF)<sub>6</sub>, and Ramona Oehme (MS Core Facility of Leipzig University) for HR-ESI MS measurements.

## Notes and references

- 1 M. D. Wodrich and X. Hu, Nat. Rev. Chem., 2018, 2, 12236.
- 2 H. H. Kung and M. C. Kung, Catal. Lett., 2014, 144, 1643.
- 3 M. Desage-El Murr, *ChemCatChem*, 2019, **12**, 53.
- 4 I. M. Lorkovic, R. R. Duff and M. S. Wrighton, J. Am. Chem. Soc., 1995, 117, 3617.
- 5 F. Würthner and J. Rebek, Angew. Chem. Int. Ed., 1995, **34**, 446.
- 6 J. Choudhury, *Tetrahedron Lett.*, 2018, **59**, 487.
- 7 A. J. Teator, D. N. Lastovickova and C. W. Bielawski, Chem. Rev., 2016, 116, 1969.
- 8 V. Blanco, D. A. Leigh and V. Marcos, Chem. Soc. Rev., 2015, 44, 5341.
- 9 E. Peris, Chem. Rev., 2018, **118**, 9988.
- 10 J. Wei and P. L. Diaconescu, Acc. Chem. Res., 2019, **52**, 415.
- 11 A. M. Allgeier and C. A. Mirkin, *Angew. Chem. Int. Ed.*, 1998, **37**, 894.
- 12 Y. Ryu, G. Ahumada and C. W. Bielawski, *Chem. Commun.*, 2019, **55**, 4451.
- O. R. Luca and R. H. Crabtree, *Chem. Soc. Rev.*, 2013, 42, 1440.
   A. J. D. Magenau, N. C. Strandwitz, A. Gennaro and K. Matyjaszewski, *Sci*-
- ence, 2011, **332**, 81.
- 15 E. J. Horn, B. R. Rosen and P. S. Baran, *ACS Cent. Sci.*, 2016, **2**, 302.
- 16 C. Schotten, T. P. Nicholls, R. A. Bourne, N. Kapur, B. N. Nguyen and C. E. Willans, *Green Chem.*, 2020, **149**, S21.
- 17 Y. Yuan and A. Lei, Nat. Commun., 2020, 11, 802.
- 18 B. A. Frontana-Uribe, R. D. Little, J. G. Ibanez, A. Palma and R. Vasquez-Medrano, Green Chem., 2010, 12, 2099.
- 19 D. Astruc, Eur. J. Inorg. Chem., 2017, 2017, 6.
- 20 P. Štěpnička, ed., Ferrocenes. Ligands, Materials and Biomolecules, J. Wiley, Chichester, England, Hoboken, NJ, 2008.
- C. K. A. Gregson, V. C. Gibson, N. J. Long, E. L. Marshall, P. J. Oxford and A. J. P. White, J. Am. Chem. Soc., 2006, **128**, 7410.
- 22 M. S. Inkpen, S. Scheerer, M. Linseis, A. J. P. White, R. F. Winter, T. Albrecht and N. J. Long, *Nature Chem.*, 2016, **8**, 825.
- 23 D. Astruc, C. Ornelas and J. Ruiz, Acc. Chem. Res., 2008, **41**, 841.
- 24 A. K. Diallo, C. Absalon, J. Ruiz and D. Astruc, J. Am. Chem. Soc., 2011, 133,
- 629.
   U. Pfaff, A. Hildebrandt, D. Schaarschmidt, T. Hahn, S. Liebing, J. Kortus and H. Lang. Organometallics. 2012. 31. 6761.
- 26 U. Pfaff, G. Filipczyk, A. Hildebrandt, M. Korb and H. Lang, *Dalton Trans.*, 2014. **43**, 16310.
- 27 B. Topolinski, B. M. Schmidt, S. Schwagerus, M. Kathan and D. Lentz, *Eur. J. Inorg. Chem.*, 2014, 2014, 5391.
- 28 Y. Yu, A. D. Bond, P. W. Leonard, U. J. Lorenz, T. V. Timofeeva, K. P. C. Vollhardt, G. D. Whitener and A. A. Yakovenko, *Chem. Commun.*, 2006, 2572.
- 29 M. Zhao and C. Chen, ACS Catal., 2017, 7, 7490.
- 30 G. Haberhauer, Angew. Chem. Int. Ed., 2011, 50, 6415.
- 31 L. van Dijk, M. J. Tilby, R. Szpera, O. A. Smith, H. A. P. Bunce and S. P. Fletcher, *Nat. Rev. Chem.*, 2018, **2**, 22.
- 32 M. Baroncini, L. Casimiro, C. de Vet, J. Groppi, S. Silvi and A. Credi, *ChemistryOpen*, 2018, **7**, 169.
- 33 M. Baroncini, S. Silvi and A. Credi, *Chem. Rev.*, 2020, **120**, 200.
- A. P. de Silva, T. P. Vance, B. Wannalerse and M. E. S. West, in *Molecular Switches*, ed. B. L. Feringa and W. R. Browne, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2<sup>nd</sup> edn., 2011, pp. 669–696.
- 35 H. Keisar, M. Lahav and M. E. van der Boom, *ChemPhysChem*, 2019, **20**, 2403.
- 36 K. Szaciłowski, Chem. Eur. J., 2004, 10, 2520.
- 37 C. Sporer, I. Ratera, D. Ruiz-Molina, Y. Zhao, J. Vidal-Gancedo, K. Wurst, P. Jaitner, K. Clays, A. Persoons, C. Rovira and J. Veciana, *Angew. Chem. Int. Ed.*, 2004, **43**, 5266.
- 38 F. Pina, A. Roque, M. J. Melo, M. Maestri, L. Belladelli and V. Balzani, Chem. Eur. J., 1998, 4, 1184.

Nanu

Accepted

emical science

#### Journal Name

- H. J. Chandler, M. Stefanou, E. E. B. Campbell and R. Schaub, Nat. Com-39 mun., 2019, 10, 1.
- W. Auwärter, K. Seufert, F. Bischoff, D. Ecija, S. Vijayaraghavan, S. Joshi, F. 40 Klappenberger, N. Samudrala and J. V. Barth, Nature Nanotech., 2012, 7,
- 41 S. Silvi, E. C. Constable, C. E. Housecroft, J. E. Beves, E. L. Dunphy, M. Tomasulo, F. M. Raymo and A. Credi, Chem. Eur. J., 2009, 15, 178.
- 42 J. Wang and B. L. Feringa, Science, 2011, 331, 1429.
- A. Straube, P. Coburger, M. R. Ringenberg and E. Hey-Hawkins, Chem. Eur. 43 J., 26, 2020, 5758.
- P. Neumann, H. Dib, A.-M. Caminade and E. Hev-Hawkins, Angew. Chem. 44 Int. Ed., 2015, 54, 311.
- P. Neumann, H. Dib, A. Sournia-Saguet, T. Grell, M. Handke, A.-M. 45 Caminade and E. Hey-Hawkins, Chem. Eur. J., 2015, 21, 6590.
- 46 J. Popp, A.-M. Caminade and E. Hey-Hawkins, Eur. J. Inorg. Chem., 2020, 17,
- 47 S. E. Gibson and M. P. Castaldi, Angew. Chem. Int. Ed., 2006, 45, 4718.
- C. Moberg, Angew. Chem. Int. Ed., 1998, 37, 248. 48
- E. K. van den Beuken and B. L. Feringa, Tetrahedron, 1998, 54, 12985. 49
- 50 R. Peters, Cooperative Catalysis, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2015.
- X. Liu, P. Du and R. Cao, Nat. Commun., 2013, 4, 2375. 51
- 52 L.-I. Rodríguez, T. Roth, J. Lloret Fillol, H. Wadepohl and L. H. Gade, Chem. Eur. J., 2012, 18, 3721.
- S. Gonell, M. Poyatos and E. Peris, Angew. Chem. Int. Ed., 2013, 52, 7009. 53
- R. Maity, A. Mekic, M. van der Meer, A. Verma and B. Sarkar, Chem. Commun., 2015, 51, 15106.
- T. Glaser, M. Heidemeier, T. Weyhermüller, R.-D. Hoffmann, H. Rupp and P. Müller, Anaew. Chem. Int. Ed., 2006, 45, 6033.
- C.-G. Freiherr von Richthofen, A. Stammler, H. Bögge, M. W. DeGroot, J. R. Long and T. Glaser, Inorg. Chem., 2009, 48, 10165.
- 57 S. Naik, M. Kumaravel, J. T. Mague and M. S. Balakrishna, Inorg. Chem., 2014. 53. 1370.
- 58 G. S. Ananthnag, J. T. Mague and M. S. Balakrishna, Inorg. Chem., 2015, 54, 10985.
- 59 J. Zhang, P. W. Miller, M. Nieuwenhuyzen and S. L. James, Chem. Eur. J., 2006. 12. 2448.
- C. S. A. Fraser, M. C. Jennings and R. J. Puddephatt, Chem. Commun., 2001, 1310.
- M. S. Inkpen, S. Du, M. Driver, T. Albrecht and N. J. Long, Dalton Trans., 2013. 42. 2813.
- S. Steffens, M. H. Prosenc, J. Heck, I. Asselberghs and K. Clays, Eur. J. Inorg. Chem., 2008, 1999.
- D. Haas, J. M. Hammann, R. Greiner and P. Knochel, ACS Catal., 2016, 6, 1540.
- 64 K. Schlögl and H. Soukup, Tetrahedron Lett., 1967, 8, 1181.
- 65 V. Mamane, I. Ledoux-Rak, S. Deveau, J. Zyss and O. Riant, Synthesis, 2003, **3**. 455
- Y.-K. Lim, S. Wallace, J. C. Bollinger, X. Chen and D. Lee, Inorg. Chem., 2007, 66 **46**, 1694.
- 67 A. Donoli, A. Bisello, R. Cardena, C. Prinzivalli and S. Santi, Organometallics, 2013, 32, 1029.
- 68 S. Santi, A. Bisello, R. Cardena and A. Donoli, Dalton Trans., 2015, 44, 5234.
- 69 T.-Y. Dong and D. N. Hendrickson, Bull. Inst. Chem., Acad. Sin., 1987, 34, 67. 70 V. Leigh, W. Ghattas, H. Mueller-Bunz and M. Albrecht, J. Organomet.
- Chem, 2014, 771, 33. 71
- J. Kühnert, M. Dusek, J. Demel, H. Lang and P. Štěpnička, Dalton Trans., 2007. 2802.
- J. Schulz, I. Císařová and P. Štěpnička, Organometallics, 2012, 31, 729. 72
- K. Škoch, I. Císařová, J. Schulz, U. Siemeling and P. Štěpnička, Dalton Trans., 73 2017, 46, 10339.
- 74 U. Siemeling, T. Klemann, C. Bruhn, J. Schulz and P. Štěpnička, Dalton Trans., 2011, 40, 4722.
- 75 M. J. Verschoor-Kirss, O. Hendricks, C. M. Verschoor, R. Conry and R. U. Kirss, Inorg. Chim. Acta, 2016, 450, 30.
- 76 F. Barrière, R. U. Kirss and W. E. Geiger, Organometallics, 2005, 24, 48.
- 77 D. A. Durfey, R. U. Kirss, C. Frommen and W. Feighery, Inorg. Chem., 2000,
- J. Popp, S. Hanf and E. Hey-Hawkins, ACS omega, 2019, 4, 22540. 78
- P. Štěpnička and I. Císařová, Dalton Trans., 2013, 42, 3373. 79

- A. Hildebrandt, N. Wetzold, P. Ecorchard, B. Walfort, T. Rüffer and H. Lang, 80 Eur. J. Inorg. Chem., 2010, 2010, 3615. 
   Eur. J. Inorg. Chem., 2010, 2010, 3615.
   DOI: 10.1039/D0SC03604H

   U. Siemeling, T. Klemann, C. Bruhn, J. Schulz and P. Štěpnička, Z. anorg.
- 81 allg. Chem., 2011, 637, 1824.
- T. M. Dau, J. R. Shakirova, A. Doménech, J. Jänis, M. Haukka, E. V. Grachova, 82 T. A. Pakkanen, S. P. Tunik and I. O. Koshevoy, Eur. J. Inorg. Chem., 2013, 28, 4976-4983.
- B. Assadollahzadeh and P. Schwerdtfeger, Chem. Phys. Lett., 2008, 462, 83 222.
- Q. Zheng, S. Borsley, G. S. Nichol, F. Duarte and S. L. Cockroft, Angew. 84 Chem. Int. Ed., 2019, 58, 12617.
- 85 M. J. Katz, K. Sakai and D. B. Leznoff, Chem. Soc. Rev., 2008, 37, 1884.
- M. R. Ringenberg, Chem. Eur. J., 2019, 25, 2396. 86
- 87 P. Kalck, Polyhedron, 1988, 7, 2441.
- P. Kalck, ed., Homo- and Heterobimetallic Complexes in Catalysis. Coopera-88 tive Catalysis, Springer International Publishing, Cham, 1st edn., 2016.
- J. A. Mata, F. E. Hahn and E. Peris, Chem. Sci., 2014, 5, 1723. 89
- 90 M. Flores-Jarillo, D. Mendoza-Espinosa, V. Salazar-Pereda and S. González-Montiel, Organometallics, 2017, 36, 4305.
- 91 Y. Li, E. M.W. Tsang, A. Y.C. Chan and H.-Z. Yu, Electrochem. Commun., 2006. 8. 951.
- 92 J. Heck and M. Dede, in Ferrocenes. Ligands, Materials and Biomolecules, ed. P. Štěpnička, J. Wiley, Chichester, England, Hoboken, NJ, 2008, pp. 319-392.
- 93 M. Iyoda, T. Kondo, T. Okabe, H. Matsuyama, S. Sasaki and Y. Kuwatani, Chem. Lett., 1997, 26, 35.
- 94 F. Barrière and W. E. Geiger, J. Am. Chem. Soc., 2006, 128, 3980.
- 95 R. J. LeSuer, C. Buttolph and W. E. Geiger, Anal. Chem., 2004, 76, 6395.
- P. Zanello, A. Cinquantini, M. Fontani, M. Giardiello, G. Giorgi, C. R. Landis 96 and B. F.M. Kimmich, J. Organomet. Chem., 2001, 637-639, 800.
- 97 G. Gritzner and J. Kuta. Pure Appl. Chem., 1984, 56, 461.
- A. S. K. Hashmi, J. P. Weyrauch, W. Frey and J. W. Bats, Org. Lett., 2004, 6, 98 4391.
- 99 A. S. K. Hashmi and G. J. Hutchings, Angew. Chem. Int. Ed., 2006, 45, 7896.
- 100 D. Canseco-Gonzalez, A. Petronilho, H. Mueller-Bunz, K. Ohmatsu, T. Ooi and M. Albrecht, J. Am. Chem. Soc., 2013, 135, 13193.
- 101 R. Pretorius, M. R. Fructos, H. Müller-Bunz, R. A. Gossage, P. J. Pérez and M. Albrecht, Dalton Trans., 2016, 45, 14591.
- 102 L. Hettmanczyk, D. Schulze, L. Suntrup and B. Sarkar, Organometallics, 2016, 35, 3828.
- 103 M. Rigo, L. Hettmanczyk, F. J. L. Heutz, S. Hohloch, M. Lutz, B. Sarkar and C. Müller, Dalton Trans., 2016, 46, 86.
- 104 A. S. K. Hashmi, M. C. Blanco Jaimes, A. M. Schuster and F. Rominger, J. Org. Chem., 2012, 77, 6394.
- 105 P. Veit, C. Volkert, C. Förster, V. Ksenofontov, S. Schlicher, M. Bauer and K. Heinze, Chem. Commun., 2019, 55, 4615.
- 106 L. Hettmanczyk, S. Manck, C. Hoyer, S. Hohloch and B. Sarkar, Chem. Commun., 2015, **51**, 10949,
- 107 L. Hettmanczyk, L. Suntrup, S. Klenk, C. Hoyer and B. Sarkar, Chem. Eur. J., 2017, 23, 576.
- 108 S. Klenk, S. Rupf, L. Suntrup, M. van der Meer and B. Sarkar, Organometallics, 2017, 36, 2026.
- 109 S. Vanicek, M. Podewitz, J. Stubbe, D. Schulze, H. Kopacka, K. Wurst, T. Müller, P. Lippmann, S. Haslinger, H. Schottenberger, K. R. Liedl, I. Ott, B. Sarkar and B. Bildstein, Chem. Eur. J., 2018, 24, 3742.
- 110 R. Dorel and A. M. Echavarren, Chem. Rev., 2015, 115, 9028.
- 111 D. Weber and M. R. Gagné, Org. Lett., 2009, 11, 4962.
- 112 A. Homs, J. Escofet and A. M. Echavarren, Org. Lett., 2013, 15, 5782.
- 113 A. Zhdanko and M. E. Maier, ACS Catal., 2015, 5, 5994.
- 114 D. Wang, R. Cai, S. Sharma, J. Jirak, S. K. Thummanapelli, N. G. Akhmedov, H. Zhang, X. Liu, J. L. Petersen and X. Shi, J. Am. Chem. Soc., 2012, 134, 9012
- 115 Y. Sota, M. Yamamoto, M. Murai, J. Uenishi and M. Uemura, Chem. Eur. J., 2015, 21, 4398.
- 116 M. Chen, B. Yang and C. Chen, Angew. Chem. Int. Ed., 2015, 54, 15520.
- 117 E. Deck, H. E. Wagner, J. Paradies and F. Breher, Chem. Commun., 2019, 55, 5323
- 118 K. Arumugam, C. D. Varnado, S. Sproules, V. M. Lynch and C. W. Bielawski, Chem. Eur. J., 2013, 19, 10866.

63



ARTICLE

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 03 August 2020. Downloaded on 8/3/2020 3:40:27 PM

- 119 K. Rößler, T. Rüffer, B. Walfort, R. Packheiser, R. Holze, M. Zharnikov and H. Lang, J. Organomet. Chem., 2007, **692**, 1530.
- 120 C. García-Morales, B. Ranieri, I. Escofet, L. López-Suarez, C. Obradors, A. I. Konovalov and A. M. Echavarren, *J. Am. Chem. Soc.*, 2017, **139**, 13628.
- 121 J. E. Aguado, S. Canales, M. C. Gimeno, P. G. Jones, A. Laguna and M. D. Villacampa, *Dalton Trans.*, 2005, 3005.
- 122 M. C. Gimeno, A. Laguna, C. Sarroca and P. G. Jones, *Inorg. Chem.*, 1993, 32, 5926.
- 123 J. Wolf, F. Huber, N. Erochok, F. Heinen, V. Guerin, C. Legault, S. Kirsch and S. M. Huber, *Angew. Chem. Int. Ed.*, **2020**, DOI: 10.1002/anie.202005214.
- 124 I. Krossing, *Chem. Eur. J.*, 2001, **7**, 490.
- 125 N. G. Connelly and W. E. Geiger, *Chem. Rev.*, 1996, 96, 877.
  126 S. Ibáñez, M. Poyatos, L. N. Dawe, D. Gusev and E. Peris, *Organometallics*,
- 2016, **35**, 2747.
  127 J. Schießl, J. Schulmeister, A. Doppiu, E. Wörner, M. Rudolph, R. Karch and A. S. K. Hashmi, *Adv. Synth. Catal.*, 2018, **360**, 2493.
- 128 L.-G. Zhuo, W. Liao and Z.-X. Yu, *Asian J. Org. Chem.*, 2012. **1**. 336.
- 129 The increased rate for the non-switched reaction is likely linked to the stirring (500 rpm) which, in contrast to the NMR experiment, was applied.
- L. A. Brown, J. L. Rhinehart and B. K. Long, ACS Catal., 2015, 5, 6057.
   S. F. Hartlaub, N. K. Lauricella, C. N. Ryczek, A. G. Furneaux, J. D. Melton, N.
- A. Piro, W. S. Kassel and C. Nataro, *Eur. J. Inorg. Chem.*, 2017, **2017**, 424. 132 D. F. Evans, *J. Chem. Soc.*, 1959, 2003.
- 133 G. A. Bain and J. F. Berry, J. Chem. Educ., 2008, 85, 532.
- 134 R. J. Webb, S. J. Geib, D. L. Staley, A. L. Rheingold and D. N. Hendrickson, J. Am. Chem. Soc., 1990, **112**, 5031.
- 135 J. A. Kramer and D. N. Hendrickson, Inorg. Chem., 1980, 19, 3330.
- 136 K. Kowalski, M. Linseis, R. F. Winter, M. Zabel, S. Záliš, H. Kelm, H.-J. Krüger,
   B. Sarkar and W. Kaim, *Organometallics*, 2009, 28, 4196.
- 137 R. Prins, J. Chem. Soc. D, 1970, 280b.
- H. B. Gray, Y. S. Sohn and N. Hendrickson, J. Am. Chem. Soc., 1971, 93, 3603.
- 139 H. Schmidbaur and A. Schier, Chem. Soc. Rev., 2012, 41, 370.
- B. Artetxe, A. Iturrospe, P. Vitoria, E. Ruiz-Bilbao, J. S Garitaonandia and J. M. Gutiérrez-Zorrilla, *Molecules*, 2018, 23.
- 141 A. G. Furneaux, N. A. Piro, R. Hernández Sánchez, K. M. Gramigna, N. Fey, M. J. Robinson, W. S. Kassel and C. Nataro, *Dalton Trans.*, 2016, **45**, 4819.
- 142 I. Abdul-Razak, A. Usman, H.-K. Fun, B. M. Yamin and N. A. Kasim, Acta Crystallogr. Sect. C, 2002, **58**, m162-164.
- 143 S. Chatterjee, J. A. Krause, W. B. Connick, C. Genre, A. Rodrigue-Witchel and C. Reber, *Inorg. Chem.*, 2010, **49**, 2808.
- 144 A. N. Nesmeyanov, E. G. Perevalova, O. B. Afanasova and K. I. Grandberg, Izv. Akad. Nauk, Ser. Khim., 1977, 2166.
- 145 R. Makhoul, P. Hamon, T. Roisnel, J.-R. Hamon and C. Lapinte, *Chem. Eur. J.*, 2020, **26**, 8368.

ARTICLE

Trinuclear gold(I) complexes of  $C_3$ -symmetric tris(ferrocenyl)arene-based tris-phosphanes with four Article Online accessible oxidation states catalyse the ring-closing isomerisation of *N*-(2-propyn-1-yl)benzamide with different rates depending on their redox state.



71x39mm (300 x 300 DPI)